

5 References

- Airaksinen MS, Koltzenburg M, Lewin GR, Masu Y, Helbig C, Wolf E, Brem G, Toyka KV, Thoenen K, Meyer M (1996). Specific subtypes of cutaneous mechanoreceptors require neurotrophin-3 following peripheral target innervation. *Neuron* **16**, 287-295.
- Akopian AN, Chen CC, Ding Y, Cesare P, Wood JN (2000). A new member of the acid-sensing ion channel family. *Neuroreport* **11**: 2217-2222.
- Barbacid M (1995). Neurotrophic factors and their receptors. *Curr Opin Cell Biol* **7**: 148-155.
- Barde YA, Edgar D, Thoenen H (1982). Purification of a new neurotrophic factor from mammalian brain *EMBO J.* **1**(5):549-53.
- Bassilana F, Champigny G, Waldmann R, deWeille JR, Heurteaux C, Lazdunski M (1997). The acid-sensitive ionic channel subunit ASIC and the mammalian degenerin subunit MDEG form a heteromultimeric H⁺-gated Na⁺ channel with novel properties. *J Biol Chem* **272**: 28819-28822.
- Bergmann I, Priestley JV, McMahon SB, Bröcker EB, Toyka KV, Koltzenburg M. (1997). Analysis of cutaneous sensory neurons in transgenic mice lacking the low affinity neurotrophin receptor p75. *Eur. J. Neurosci.* **9**:18–28
- Berridge MJ, Lipp P, Bootman MD (2000). The versatility and universality of calcium signalling. *Nat Rev Mol Cell Biol.* **1**(1):11-21. Review.
- Bezprozvanny I, Tsien RW (1995). Voltage-dependent blockade of diverse types of voltage-gated Ca²⁺ channels expressed in Xenopus oocytes by the Ca²⁺ channel antagonist mibepradil (Ro 40-5967). *Mol Pharmacol.* **48**(3):540-9
- Bordier C (1981). Phase separation of integral membrane proteins in Triton X-114 solution *J Biol Chem.* **256**(4):1604-7
- Brennan C, Rivas-Plata K, Landis SC (1999). The p75 neurotrophin receptor influences NT-3 responsiveness of sympathetic neurons in vivo. *Nat. Neurosci.* **2**:699–705
- Brown A.G. *Organization of the Spinal Cord.* (1981) New York: Springer-Verlag
- Brown AG, Iggo A (1967). A quantitative study of cutaneous receptors and afferent fibres in the cat and rabbit. *J.Physiol.* **193**; 707-733
- Brown AG, Iggo A, Miller S (1967) Myelinated afferent nerve fibers from the skin of the rabbit ear. *Exp Neurol* **18**(3): 338-49.

- Burgess PR, Petit D, Warren RM (1968). Receptor types in cat hairy skin supplied by myelinated fibers. *J Neurophysiol.* **31**(6):833-48.
- Canessa CM, Horisberger JD, Rossier BC (1993). Epithelial sodium channel related to proteins involved in neurodegeneration. *Nature* **361**: 467-470.
- Canessa CM, Merillat AM, Rossier BC (1994). Membrane topology of the epithelial sodium channel in intact cells. *Am J Physical* **267**: C1682-C1690.
- Canessa CM, Schild L, Buell G, Thorens B, Gautschi I, Horisberger JD, Rossier BC (1994). Amiloride-sensitive epithelial Na⁺ channel is made of three homologous subunits. *Nature* **367**: 412-413.
- Carroll P, Lewin GR, Koltzenburg M, Toyka KV, Thoenen K (1998). A role for BDNF in mechanotransduction. *Nature Neurosci* **1**, 42-46.
- Carter BD, Lewin GR (1997) Neurotrophins live or let die: does p75NTR decide? *Neuron*. **18**(2):187-90. Review
- Catterall WA (2000). Structure and regulation of voltage gated Ca channels. *Annu. Rev. Cell Dev. Biol.* **16**, 521-555
- Catton WT (1970). Mechanoreceptor function. *Physiol Rev.* **50**(3):297-318.
- Chalfie M, Driscoll M, Huang M (1993). Degenerin similarities. *Nature* **361**: 504-509.
- Chalfie M, Jorgensen EM (1998). *Caenorhabditis elegans* neuroscience: genetics to genome. *TIGs* **14**: 506-512.
- Chalfie M, Sulston JE (1981). Developmental genetics of the mechanosensory neurons of *Caenorhabditis elegans*. *Dev Biol* **82**; 2558-270.
- Chalfie M, Thomson NJ (1979). Organization of neuronal microtubules in the nematode *Caenorhabditis elegans*. *J Cell Biol* **82**: 278-289.
- Chalfie M, Wolinsky E (1990). The identification and suppression of inherited neurodegeneration in *Caenorhabditis elegans*. *Nature* **345**: 410-416
- Chang SS, Gründer S, Hanukoglu A, Rosler A, Mathew PM, Hanukodlu I, Schild L, Lu Y, Shimkets RA, Nelson-Williams C, Rossier BC, Lifton RP (1996). Mutations in the subunits of the epithelial sodium channel cause salt wasting with hyperkalaemic acidosis, pseudohypoaldosteronism type 1. *Nat Genet* **12**: 248-253.
- Chao MV (1994). The p75 neurotrophin receptor. *J. Neurobiol.* **25**:1373–85
- Chen CC, England S, Akopian A, Wood J (1998). A sensory neuron-specific, proton-gated ion channel. *Proc Natl Acad Sci USA* **95**, 10240–10245.
- Clapham DE (1995). Calcium signaling. *Cell* **80**, 259-268

- Conover JC, Erickson JT, Katz DM, Bianchi LM, Poueymirou WT, McClain J, Pan L, Helgren M, Ip NY, Boland P, Friedman B, Wiegand S, Vejsada R, Kato AC, DeChiara TM, Yancopoulos GD (1995). Neuronal deficits, not involving motor neurons, in mice lacking BDNF and/or NT-4. *Nature* **375**: 235-238.
- Conover JC, Yancopoulos GD (1997). Neurotrophin regulation of the developing nervous system: analyses of knockout mice. *Rev Neurosci*. **8**(1):13-27. Review
- Corey DP Hudspeth AJ (1979) Ionic basis of the receptor potential in a vertebrate hair cell *Nature* **281**(5733):675-7
- Cribbs LL, Lee JH, Yang J, Satin J, Zhang Y, Daud A, Barclay J, Williamson MP, Fox M, Rees M, Perez-Reyes E (1998). Cloning and characterization of alpha1H from human heart, a member of the T-type Ca²⁺ channel gene family. *Circ Res*. **83**(1):103-9.
- Davies AM, Minichiello L, Klein R (1995) Developmental changes in NT-3 signaling via trkB and trkA in embryonic neurons. *EMBO J*. **14**: 4482-4489.
- Davies, A. (1996) The neurotrophic hypothesis: Where does it stand? *Phil. Trans. R. Soc. Lond. B* **351**: 389-394
- Driscoll M, Chalfie M (1991). The *mec-4* gene is a member of a family of *Caenorhabditis elegans* genes that can mutate to induce neuronal degeneration. *Nature* **349**: 588-593.
- Drummond HA, Abboud FM, Welsh MJ (2000). Localization of β and γ subunits of ENaC in sensory nerve endings in the rat footpad. *Brain Research* **884**: 1-12.
- Du H, Gu G, William C, Chalfie M (1996). Extracellular proteins needed for *C. elegans* mechanosensation. *Neuron* **10**: 183-194.
- Duc C, Farman N, Canessa CM, Bonvalet JP, Rossier BC (1994). Cell specific expression of epithelial sodium channel α , β and γ subunits in aldosterone responsive epithelia from the rat: localization by in situ hybridization and immunocytochemistry. *J. Cell Biol.* **127**:1907– 21
- Erlanger J, Gasser HS (1924). The compound nature of the action current of nerve as disclosed by cathode ray oscillograph. *Am J Physiol* **70**: 624-666.
- Erlanger J, Gasser HS (1930). The action potential in fibers of slow conduction in spinal roots and somatic nerves. *Am J Physiol* **92**: 43-82.
- Ernfors P, Lee KF, Jaenisch R (1994). Mice lacking brain-derived neurotrophic factor develop sensory neuron deficits. *Nature* **368**: 147-150
- Ernfors P, Lee KF, Jaenisch R (1994). Target derived and putative local actions of neurotrophins in the peripheral nervous system. *Prog Brain Res* **103**:43-54

- Ernfors P, Lee KF, Kucera J, Jaenisch R (1994). Lack of neurotrophin-3 leads to deficiencies in the peripheral nervous system and loss of limb proprioceptive afferents. *Cell* **77**: 503-512.
- Farinas I, Jones KR, Backus C, Wang XY, Reichardt LF (1994). Severe sensory and sympathetic deficits in mice lacking neurotrophin-3. *Nature* **369**: 658-661.
- Farinas I, Yoshida CK, Backus C, Reichardt LF (1996). Lack of neurotrophin-3 results in death of spinal sensory neurons and premature differentiation of their precursors. *Neuron* **17**: 1065-1078.
- Fleischer E, Handwerker HO, Joukhadar S. (1983) Unmyelinated nociceptive units in two skin areas of the rat. *Brain Res.* **267**(1):81-92.
- Fox, A. P., Nowycky, M. C., and Tsien, R. W. (1987). Kinetic and pharmacological properties distinguishing three types of calcium current in chick sensory neurons. *J. Physiol. (Lond.)* **394**: 149-172
- Frade JM, Barde YA. (1998). Nerve growth factor: two receptors, multiple functions. *BioEssays* **20**:137–45
- French AS (1992). Mechanotransduction. *Annu Rev Physiol* **54**: 135-152.
- Frey M von (1897). Untersuchungen über die Sinnesfunktionen der menschlichen Haut. I. Druckempfindung und Schmerz. *Königl Sächs Ges Wiss Math Phys Kl* **23**: 169-266.
- Fricke B, Lints R, Stewart G, Drummond H, Dodt G, Driscoll M, von Düring M (2000). Epithelial sodium channels and stomatin are expressed in rat trigeminal mechanosensory neurons. *Cell Tissue Res* **299**: 327-334.
- Friedman W. (2000). Neurotrophins induce death of hippocampal neurons via the p75 receptor. *J. Neurosci.* **20**:6340–46
- Friedman WJ, Greene LA (1999). Neurotrophin signaling via trks and p75. *Exp Cell Res* **253**: 131-142.
- Fuller CM, Awayda MS, Arrate MP, Bradford AL, Morris RG, et al. 1995. Cloning of a bovine renal epithelial Na^+ channel subunit. *Am. J. Physiol.* **269**:C641–54
- García-Anoveros J, Corey DP (1997). The molecules of mechanosensation. *Annu Rev Neurosci* **20**: 567-594.
- García-Anoveros J, Derfler B, Neville-Golden J, Hyman BT, Corey DP (1997). BNaC1 and BNaC2 constitute a new family of human neuronal sodium channels related to degenerins and epithelial sodium channels. *Proc Natl Acad Sci USA* **94**, 1459–1464.

- Garty H, Edelman IS Amiloride-sensitive trypsinization of apical sodium channels. Analysis of hormonal regulation of sodium transport in toad bladder (1983) *J Gen Physiol.* **81**(6):785-803.
- Garty H, Palmer L (1997). Epithelial sodium channels: function, structure and regulation. *Physiol Rev* **77**: 359-396.
- Gasser HS, Erlanger J (1927). The role played by the size of the constituent fibers of a nerve trunk in determining the form of its action potential wave. *Am J Physiol* **80**: 522-547.
- Gillespie PG, Walker RG (2001) Molecular basis of mechanosensory transduction. *Nature.* **413**(6852):194-202.
- Goldstein BJ, Kulaga HM, Reed RR (2002). Cloning and Characterization of SLP3: a Novel Member of the Stomatin Family Expressed by Olfactory Receptor Neurons (2002). *J Assoc Res Otolaryngol.* Sep 23 (epub ahead of print)
- Goldstein O, Asher C, Garty H (1997). Cloning and induction by low NaCl intake of avian intestine Na^+ channel subunits. *Am. J. Physiol.* **272**:C270–77
- Goodman MB, Ernstrom GG, Chelur DS, O'Hagan R, Yao CA, Chalfie M (2002) *Mec-2/C. elegans* DEG/ENaC channels needed for mechanosensation. *Nature.* **415**(6875):1039-42.
- Gotz R, Koster R, Winkler C, Raulf F, Lottspeich F, Schartl M, Thoenen H. (1994) Neurotrophin-6 is a new member of the nerve growth factor family. *Nature.* **372**(6503):266-9.
- Gutnick MJ, Yarom Y (1989) Low threshold calcium spikes, intrinsic neuronal oscillation and rhythm generation in the CNS. *J Neurosci Methods.* **28**(1-2):93-9. Review.
- Hamill OP, Martinac B (2001) Molecular basis of mechanotransduction in living cells. *Physiol Rev.* **81**(2):685-740.
- Harteneck, C, Plant, TD and Schultz, G (2000) From worm to man: three subfamilies of TRP channels. *Trends Neurosci.* **23**, 159-166
- Hiebl-Dirschmied CM, Adolf GR, Prohaska R (1991) Isolation and partial characterization of the human erythrocyte band 7 integral membrane protein. *Biochim Biophys Acta.* **1065**(2):195-202
- Hille B (1984) “Ionic channels of excitable membrane” Sinauer Associates, Sunderland, MA

- Hodgkin AL, Huxley AF (1952) Currents carried by sodium and potassium ion through the membrane of the giant axon of *Loligo*. *J. Physiol.* (London) **116**, 449-472
- Hodgkin J (1991). Nematodes on a nervous decline. *Nature* **349**: 564-565.
- Hong K, Driscoll M (1994). A transmembrane domain of the putative channel subunit MEC-4 influences mechanotransduction and neurodegeneration in *C. elegans*. *Nature* **367**, 470-473.
- Hong K, Mano I, Driscoll M (2000). In vivo structure/function analyses of *C. elegans* MEC-4, a candidate mechanosensory ion channel subunit. *J Neurosci* **20**: 2575-2588.
- Housley GD *et al.*(1999) Expression of the P2X₂ receptor subunit of the ATP-gated ion channel in the cochlea: implications for sound transduction and auditory neurotransmission. *J. Neurosci.* **19**, 8377-8388
- Huang EJ, Reichardt LF (2001) Neurotrophins: roles in neuronal development and function. *Annu Rev Neurosci.* **24**:677-736. Review.
- Huang M, Chalfie M (1994). Genetic interactions affecting mechanosensory transduction in *Caenorhabditis elegans*. *Nature* **376**: 467-470.
- Huguenard JR (1998) Low-voltage-activated (T-type) calcium-channel genes identified. *Trends Neurosci.* **21**(11):451-2. Review.
- Ichikawa H, Matsuo S, Silos-Santiago I, Jacquin MF, Sugimoto T (2001) Developmental dependency of Merkel endings on trks in the palate. *Brain Res Mol Brain Res.* **88**: 171-5.
- Iggo A, Muir AR (1969). The structure and function of slowly adapting touch corpuscle in hairy skin. *J Physiol (Lond)* **200**: 763-796.
- Jentsch TJ (1994). Trinity of cation channels. *Nature* 367(6462):412-3
- Jones KR, Farinas I, Backus C, Reichardt LF (1994). Targeted disruption of the BDNF gene perturbs brain and sensory neuron development but not motor neuron development. *Cell* **76**: 989-999.
- Jönsson U, Fägerstam L, Ivarsson B, Johnsson B, Karlsson R, Lundh K, Löfas S, Persson B, Roos H, Rönnberg I, Sjölander S, Sternberg E, Stahlberg R, Urbaniczky C, Östlin H, Malmqvist M (1991). Real-time biospecific interaction analysis using surface plasmon resonance and a sensor chip technology. *BioTechniques* **11**, 520-527
- Joseph R, Dou D, Tsang W (1994) Molecular cloning of a novel mRNA (neuronatin) that is highly expressed in neonatal mammalian brain. *Biochem Biophys Res Commun.* **201**(3):1227-34
- Julius D, Basbaum AI (2001) Molecular mechanisms of nociception. *Nature* **413**(6852):203-10.

- Kaplan JM (1996). Sensory signaling in *Caenorhabditis elegans*. *Curr Opin Neurobiol* **6**: 494-499.
- Karchewski LA, Kim FA, Johnston J, McKnight RM, Verge VM (1999) Anatomical evidence supporting the potential for modulation by multiple neurotrophins in the majority of adult lumbar sensory neurons. *J Comp Neurol*. **413**:327-341.
- Kawai F and Miyachi E (2001) Enhancement by T-type Ca²⁺ currents of odor sensitivity in olfactory receptor cells. *J. Neurosci.* **21**, RC144.
- Kernan M, Cowan D, Zuker C (1994). Genetic dissection of mechanosensory transduction: mechanoreception-defective mutations of *Drosophila*. *Neuron* **12**, 1195-1206
- Kernan M, Zuker C (1995). Genetic approaches to mechanosensory transduction. *Curr Opin Neurobiol* **5**: 443-448.
- Kinkelin I, Stucky CL, Koltzenburg M (1999). Postnatal loss of Merkel cells, but not of slowly adapting mechanoreceptors in mice lacking the neurotrophin receptor p75. *Eur. J. Neurosci.* **11**:3963–69
- Koltzenburg M, Stucky CL, Lewin GR (1997). Receptive properties of mouse sensory neurons innervating hairy skin. *J Neurophysiol.* **78**(4):1841-50.
- Krasagakis K, Almond-Roesler B, Geilen C, Fimmel S, Krengel S, Chatzaki E, Gravanis A, Orfanos CE (2001). Growth and characterization of a cell line from a human primary neuroendocrine carcinoma of the skin (Merkel cell carcinoma) in culture and as xenograft. *J Cell Physiol.* **187**(3):386-91
- Kros CJ, Rüsch A, Richardson GP (1992). Mechano-electrical transducer currents in hair cells of the cultured neonatal mouse cochlea. *Proc. R. Soc. Lond. B* **249**, 185-193
- Kucera J, Fan G, Jaenisch R, Linnarson S, Ernfors P (1995). Dependence of developing group Ia afferents on neurotrophin-3. *J Comp Neurol* **363**: 307-320.
- Kunzelmann K, Schreiber R, Boucherot A. (2001) Mechanisms of the inhibition of epithelial Na(+) channels by CFTR and purinergic stimulation. *Kidney Int.* **60**(2):455-61. Review
- Lai CC, Hong K, Kinnell M, Chalfie M, Driscoll M (1996). Sequence and membrane topology of MEC-4, an ion channel subunit required for mechanotransduction in *Caenorhabditis elegans*. *J Cell Biol* **133**: 1071-1081.
- Lämmli UK (1970). Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* **227**, 680-685.

- Lee JH, Daud AN, Cribbs LL, Lacerda AE, Pereverzev A, Klockner U, Schneider T, Perez-Reyes E (1999). Cloning and expression of a novel member of the low voltage-activated T-type calcium channel family. *J Neurosci.* **19**(6):1912-21.
- Lee JH, Gomora JC, Cribbs LL, Perez-Reyes E (1999). Nickel block of three cloned T-type calcium channels: low concentrations selectively block alpha1H. *Biophys J.* **77**(6):3034-42
- Leibrock J, Lottspeich F, Hohn A, Hofer M, Hengerer B, Masiakowski P, Thoenen H, Barde YA. Molecular cloning and expression of brain-derived neurotrophic factor (1989). *Nature* **341**(6238):149-52.
- LeMaster AM, Krimm RF, Davis BM, Noel T, Forbes ME, Johnson JE, Albers KM (1999). Overexpression of brain-derived neurotrophic factor enhances sensory innervation and selectively increases neuron number. *J Neurosci.* **19**:5919-5931.
- Leonard JH, Bell JR (1997) Insights into the Merkel cell phenotype from Merkel cell carcinoma cell lines. *Australas J Dermatol.* **38** Suppl 1:S91-8. Review.
- Leung MS, Wong CC (2000). Expressions of putative neurotransmitters and neuronal growth related genes in Merkel cell-neurite complexes of the rats. *Life Sci.* **66**(16):1481-90
- Levi-Montalcini R and Angeletti P (1968). Nerve growth factor. *Physiol. Rev.* **48**: 534-569
- Lewin GR (1996). Neurotrophins and the specification of neuronal phenotype. *Phil Trans R Soc Lond B Biol Sci* **351**: 405-411.
- Lewin GR (1999). Society of Neuroscience meeting, vol. 32 abstract number 162.6
- Lewin GR, Barde YA (1996). Physiology of the neurotrophins. *Ann Rev Neurosci* **19**, 289-317.
- Lewin GR, Mendell LM (1993) Nerve growth factor and nociception *Trends Neurosci.* Sep; **16**(9):353-9
- Lewin GR, Mendell LM (1994). Regulation of cutaneous C-fiber heat nociceptors by nerve growth factor in the developing rat. *J Neurophysiol* **71**:941-949.
- Lewin GR, Ritter AM, Mendell LM (1992). On the role of nerve growth factor in the development of myelinated nociceptors. *J Neurosci* **12**:1896-1905.
- Lewin GR, Stucky CL (2000). Sensory neurons mechanotransduction: Regulation and underlying molecular mechanisms. In: Molecular basis of pain induction. (Wood, J.N. Ed.) pp 129-. Wiley-Liss, Inc.

- Lewin, GR (1996) Neurotrophins and the specification of neuronal phenotype. *Philos Trans R Soc Lond B Biol Sci.*; **351**(1338):405-11.
- Lindemann B (1996). Taste reception. *Physiol. Rev.* **76**:719–66
- Lingueglia E, deWeille JR, Bassilana F, Heurteaux C, Sakai H, Waldmann R, Lazdunski M. (1997). A modulatory subunit of acid sensing ion channels in brain and dorsal root ganglion cells. *J Biol Chem* **272**, 29778–29783.
- Lingueglia E, Renard S, Waldmann R, Voilley N, Champigny G, Plass H, Lazdunski M, Barbry P. (1994). Different homologous subunits of the amiloride-sensitive Na^+ channel are differently regulated by aldosterone. *J Biol Chem* **269**: 13736–13739.
- Lingueglia E, Voilley N, Waldmann R, Lazdunski M, Barbry P (1993) Expression cloning of an epithelial amiloride-sensitive Na^+ -channel. *FEBS Lett* **318**: 95-99.
- Liu X, Ernfors P, Wu H, Jaenisch R (1995) Sensory but not motor neuron deficits in mice lacking NT4 and BDNF. *Nature*. **375**(6528): 238-41
- Llinas R, Yarom Y. Electrophysiology of mammalian inferior olivary neurones in vitro. Different types of voltage-dependent ionic conductances. *J Physiol.* **315**:549-67
- Loewenstein WR (1959). The generation of electrical activity in a nerve ending. *Ann NY Acad Sci* **81**: 367-387.
- Loewenstein WR (1971). Mechanoelectrical transduction in the Pacinian corpuscle. Initiation of sensory impulses in mechanoreceptors. In: Handbook of sensory physiology. (Loewenstein, W.R. Ed.) pp. 269-290. Springer-Verlag, Berlin.
- Low MG (1989). Glycosyl-phosphatidylinositol: a versatile anchor for cell surface proteins. *FASEB J.* **3**(5):1600-8. Review
- Maissonpierre PC, Belluscio L, Squinto P, Ip NY; Furth ME, Lindsay RM, Yancopoulos GD (1990). Neurotrophin-3 – A neurotrophic factor related to NGF and BDNF. *Science* **247**: 1446-1451.
- Mannsfeldt AG (1999). The role of stomatin and nstom1 in sensory neuron mechanotransduction in mammals. Dissertation, Fakultät für Biologie, Universität Konstanz.
- Mannsfeldt AG, Carroll P, Stucky CL, Lewin GR (1999). Stomatin, a MEC-2 like protein, is expressed by mammalian sensory neurons. *Mol Cell Neurosci* **13**: 391-404.
- Mano I, Driscoll M (1999). DEG/ENaC channels: a touchy superfamily that watches its salt. *BioEssays* **21**: 568-578.

- Martin RL, Lee JH, Cribbs LL, Perez-Reyes E, Hanck DA (2000). Mibepradil block of cloned T-type calcium channels. *J. Pharmacol. Exp. Ther.* **295**, 302-8
- McDonald FJ, Price MP, Snyder PM, Welsh MJ. (1995) Cloning and expression of the beta- and gamma-subunits of the human epithelial sodium channel *Am. J. Physiol.* **268**:C1157-63
- Mendell LM (1996). Neurotrophins and sensory neurons: role in development, maintenance and injury. A thematic summary. *Phil. Trans. R. Soc. London. B* **351**, 463-467.
- Millard CL, Woolf CJ (1988) Sensory innervation of the hairs of the rat hindlimb: a light microscopic analysis. *J Comp Neurol.* **277**(2):183-94.
- Minichiello L, Casagranda F, Tatché RS, Stucky CL, Postigo A, Lewin GR, Davies AM, Klein R (1998). Point mutation in trkB causes loss of NT4-dependent neurons without major effects on diverse BDNF responses. *Neuron.* 1998 **21**(2):335-45.
- Mintz IM, Adams ME, Bean BP (1992). P-type calcium channels in rat central and peripheral neurons. *Neuron* 9: 85-95
- Mu X, Silos-Santiago I, Carroll SL, Snider WD (1993). Neurotrophin receptor genes are expressed in distinct patterns in developing dorsal root ganglia. *J Neurosci.* **13**:4029-4041.
- Mullins et al (1998) Life threatening interactions of mibepradil and beta-blockers with dihydropyridine calcium channel blockers. *JAMA.*, **280**:157-158
- Ockel M, Lewin GR, Barde YA (1996) In vivo effects of neurotrophin-3 during sensory neurogenesis. *Development.* **122**(1):301-7
- Oppenheim R (1996). Neurotrophic survival molecules for motoneurons: An embarrassment of riches. *Neuron* **17**: 195-197
- Payne J, Middleton J, Fitzgerald M (1991). The pattern and timing of cutaneous hair follicle innervation in the rat pup and human fetus. *Brain Res Dev Brain Res.* **61**(2):173-82.
- Perez-Reyes E (1999). Three for T: molecular analysis of the low voltage-activated calcium channel family. *Cell Mol Life Sci.* **56**(7-8):660-9. Review
- Perez-Reyes E, Cribbs LL, Daud A, Lacerda AE, Barclay J, Williamson MP, Fox M, Rees M, Lee JH (1998). Molecular characterization of a neuronal low-voltage-activated T-type calcium channel. *Nature.* **391**(6670):896-900.
- Perez-Reyes E, Schneider T (1995). Molecular biology of calcium channels. *Kidney Int.* **48**(4):1111-24. Review

- Perl ER (1992). Function of dorsal root ganglion neurons: An overview. In: Sensory Neurons Diversity, Development, and Plasticity. (Scott, S.A. Ed.) pp 3-23. Oxford Univ Press, NY.
- Price MP, Lewin GR, McIlwrath SL, Cheng C, Xie J, Heppenstall PA, Stucky CL, Mannsfeldt AG, Brennan TJ, Drummond HA, Qiao J, Benson CJ, Tarr DE, Hrstka RF, Yang B, Williamson RA, Welsh MJ (2000). The mammalian sodium channel BNC1 is required for normal touch sensation. *Nature* **407**: 1007-1011.
- Price MP, McIlwrath SL, Xie J, Cheng C, Qiao J, Tarr DE, Sluka KA, Brennan TJ, Lewin GR, Welsh MJ (2001). The DRASIC cation channel contributes to the detection of cutaneous touch and acid stimuli in mice. *Neuron*. **32**(6):1071-83.
- Price MP, Snyder PM, Welsh MJ (1996). Cloning and expression of a novel brain Na^+ channel. *J Biol Chem* **271**, 7879–7882.
- Puoti A, May A, Canessa CM, Horisberger JD, Schild L (1995). The highly selective low-conductance epithelial Na channel of Xenopus laevis A6 kidney cells *Am. J. Physiol.* **269**:C188–97
- Reeh PW (1986). Sensory receptors in mammalian skin in an *in vitro* preparation. *Neurosci Lett* **66**:141-146.
- Reichardt LF, Fariñas I (1997). Neurotrophic factors and their receptors. Roles in neuronal development and function. In *Molecular and Cellular Approaches to Neural Development*, ed. WM Cowan, TM Jessell, SL Zipursky, pp. 220–63. New York: Oxford Univ. Press
- Renard S, Voilley N, Bassilana F, Lazdunski M, Barbry P (1995). Localization and regulation by steroids of the alpha, beta and gamma subunits of the amiloride-sensitive Na^+ channel in colon, lung and kidney. *Pflügers Arch.* **430**:299–307
- Reuter H (1967) The dependence of slow inward current in Purkinje fibres on the extracellular calcium-concentration. *J. Physiol.* **192**: 479-92
- Reuter H (1979) Properties of two inward membrane currents in the heart. *Annu. Rev. Physiol.* **41**: 413-24
- Riordan JR, Rommens JM, Kerem B, Alon N, Rozmahel R, Grzelczak Z, Zielenski J, Lok S, Plavsic N, Chou JL, et al. (1989) Identification of the cystic fibrosis gene: cloning and characterization of complementary DNA. *Science* **245**(4922):1066-73.
- Rusch A Hummler E (1999) Mechano-electrical transduction in mice lacking the alpha-subunit of the epithelial sodium channel. *Hear Res.* **131**, 170-176

- Savage C, Hamelin M, Culotti JG, Coulson A, Albertson D, Chalfie M (1989). Mec-7 is a beta tubulin gene required for the production of 15-protofilament microtubules in *Caenorhabditis elegans*. *Genes Dev* **3**: 870-881.
- Schaeren-Wiemers, Gerfin-Moser (1993). A single protocol to detect transcripts of various types and expression levels in neural tissue and cultured cells: in situ hybridization using digoxigenin-labelled cRNA probes. *Histochemistry*. **100**(6):431-40.
- Schmidt R, Schmelz M, Forster C, Ringkamp M, Torebjork E, Handwerker H (1995). Novel classes of responsive and unresponsive C nociceptors in human skin. *J Neurosci*. **15**:333-41
- Schroeder JE, Fischbach PS, McCleskey EW (1990). T-type calcium channels: heterogeneous expression in rat sensory neurons and selective modulation by phorbol esters. *J. Neurosci*. **10**: 947-51
- Scopes RK (1982). Protein purification – Principles and practice. In: Springer advanced texts in chemistry series. (Cantor C.R. Series Ed.). Springer-Verlag, New York.
- Scroggs RS, Fox AP (1992). Calcium current variation between acutely isolated adult rat dorsal root ganglion neurons of different size. *J. Physiol*. **445**:639-58
- Seidel G, Prohaska R (1998). Molecular cloning of hSLP-1, a novel human brain-specific member of the band 7/MEC-2 family similar to *Caenorhabditis elegans* UNC-24 *Gene*. **225**(1-2):23-9.
- Sherrington CS (1906). The integrative action of the nervous system. Scribner, New York.
- Shimkets RA, Warnock DG, Bositis CM, Nelson-Williams C, Hansson JH, Schambelan M, Gill JRJ, Ulick S, Milora RV, Findling JW, Canessa CM, Rossier BC, Lifton RP (1994). Liddle's syndrome: heritable human hypertension caused by mutations in the α subunit of the epithelial sodium channel. *Cell* **79**: 407-414.
- Snyers L, Thinés-Semproux D, Prohaska R (1997). Colocalization of stomatin (band 7.2b) and actin filaments in UAC epithelial cells. *Eur J Cell Biol* **73**: 281-285.
- Sossey-Alaoui K, Su G, Malaj E, Roe B, Cowell JK (2002). WAVE3, an actin-polymerization gene, is truncated and inactivated as a result of a constitutional t(1;13)(q21;q12) chromosome translocation in a patient with ganglioneuroblastoma. *Oncogene*. **21**(38):5967-74
- Staub O, Gautschi I, Ishikawa T, Breitschopf K, Ciechanover A, Schild L, Rotin D. (1997) Regulation of stability and function of the epithelial Na⁺ channel (ENaC) by ubiquitination. *EMBO J*. **16**(21):6325-36.

- Steen KH, Reeh PW (1993). Sustained graded pain and hyperalgesia from harmless experimental tissue acidosis in human skin. *Neurosci Lett* **154**: 113-116.
- Steen KH, Reeh PW, Anton F, Handwerker HO (1992). Protons selectively induce lasting excitation and sensitization to mechanical stimulation of nociceptors in rat skin, *in vitro*. *J Neurosci* **12**:86-95
- Stewart GW, Hepworth-Jones BE, Keen JN, Dash BCJ, Argent AC, Casimir CM (1992). Isolation of cDNA coding for an ubiquitous membrane protein deficient in high Na⁺, low K⁺ stomatocytic erythrocytes. *Blood* **79**: 1593-1601.
- Stucky CL, DeChiara T, Lindsay RM, Yancopoulos GD, Koltzenburg M (1998). Neurotrophin-4 is required for the survival of a subclass of hair follicle receptors. *J Neurosci* **18**, 7040-7046.
- Stucky CL, Koltzenburg M (1997). The low-affinity neurotrophin receptor p75 regulates the function but not the selective survival of specific subpopulations of sensory neurons. *J. Neurosci.* **17**:4398–4405
- Stucky CL, Lewin GR (1999). Isolectin B4-positive and -negative nociceptors are functionally distinct. *J Neurosci* **19**, 6497-6505.
- Stucky CL, Shin JB, Lewin GR (2002). Neurotrophin-4: A survival factor for adult sensory neurons. *Curr. Biol.* **12** (16) 1401-1405
- Swett JE, Woolf CJ (1985) The somatotopic organization of primary afferent terminals in the superficial laminae of the dorsal horn of the rat spinal cord. *J Comp Neurol.* **231**(1):66-77.
- Talley EM, Cribbs LL, Lee JH, Daud A, Perez-Reyes E, Bayliss D (1999). Differential distribution of three members of a gene family encoding low voltage-activated (T-type) calcium channels. *J. Neurosci.* **19**, 1895-911
- Tavernarakis N, Driscoll M (1997) Molecular modeling of mechanotransduction in the nematode *Caenorhabditis elegans*. *Annu Rev Physiol.* **59**:659-89. Review.
- Tsien RW, Lipscombe D, Madison DV, Bley KR, Fox AP (1988). Multiple types of neuronal calcium channels and their selective modulation. *Trends Neurosci* **11**(10):431-8. Review.
- Vallet V, Chraibi A, Gaeggeler HP, Horisberger JD, Rossier BC (1997). An epithelial serine protease activates the amiloride-sensitive sodium channel. *Nature* **389**: 607-610.
- Verdi JM, Birren SJ, Ibanez CF, Persson H, Kaplan DR, et al.(1994). p75LNGFR regulates Trk signal transduction and NGF-induced neuronal differentiation in MAH cells. *Neuron* **12**:733–45

- Von Krosigk M, Bal T, McCormick DA (1993). Cellular mechanisms of a synchronized oscillation in the thalamus. *Science*. **261**(5119):361-4.
- Vuagniaux G, Vallet V, Jaeger NF, Hummler E, Rossier BC (2002). Synergistic Activation of ENaC by Three Membrane-bound Channel-activating Serine Proteases (mCAP1, mCAP2, and mCAP3) and Serum- and Glucocorticoid-regulated Kinase (Sgk1) in Xenopus Oocytes. *J Gen Physiol*. **120**(2):191-201
- Vuagniaux G, Vallet V, Jaeger NF, Pfister C, Bens M, Farman N, Courtois-Coutry N, Vandewalle A, Rossier BC, Hummler E (2000). Activation of the amiloride-sensitive epithelial sodium channel by the serine protease mCAP1 expressed in a mouse cortical collecting duct cell line. *J Am Soc Nephrol*. **11**(5):828-34.
- Waldmann R, Bassilana F, deWeille J, Champigny G, Heurteaux C, Lazdunski M (1997a). Molecular cloning of a non-inactivating proton-gated Na^+ channel specific for sensory neurons. *J Biol Chem* **272**, 20975– 20978.
- Waldmann R, Champigny G, Bassilana F, Heurteaux C, Lazdunski M. (1997b). A proton-gated cation channel involved in acid-sensing. *Nature* **386**, 173–177.
- Waldmann R, Champigny G, Bassilana F, Voilley N, Lazdunski M. (1995). Molecular cloning and functional expression of a novel amiloride-sensitive Na^+ channel. *J Biol Chem* **270**, 27411–27414.
- Waldmann R, Champigny G, Voilley N, Lauritzen I, Lazdunski M. (1996). The mammalian degenerin MDEG, an amiloride-sensitive cation channel activated by mutations causing neurodegeneration in *Caenorhabditis elegans*. *J Biol Chem* **271**, 10433–10436.
- Walker RG, Willingham AT, Zuker CS (2000). A Drosophila mechanosensory transduction channel. *Science* **287**, 2229-2234
- Wang D, Mentzer WC, Cameron T, Johnson RM (1991). Purification of band 7.2b, a 31-kDa integral phosphoprotein absent in hereditary stomatocytosis. *J Biol Chem*. **25**;266(27):17826-31.
- White G, Lovinger DM and Weight FF (1989). Transient low-threshold calcium current triggers burst firing through an afterdepolarizing potential in an adult mammalian neuron. *Proc. Natl. Acad. Sci. USA* **86**: 6802–6806
- Willis WD, Coggeshall RE (1991). *Sensory Mechanisms of the Spinal Cord*. New York: Plenum
- Wolfe JT, Wang H, Perez-Reyes E, Barrett PQ (2002). Stimulation of recombinant Ca(v)3.2, T-type, $\text{Ca}(2+)$ channel currents by CaMKIIGamma(C). *J. Phys.* **538**: 343-355

-
- Woodbury CJ, Ritter AM, Koerber HR (2001). Central anatomy of individual rapidly adapting low-threshold mechanoreceptors innervating the "hairy" skin of newborn mice: early maturation of hair follicle afferents. *J. Comp. Neurol.* **436**, 304-323