

## VI LITERATURVERZEICHNIS

1. Richtlinien des Bundesausschusses der Ärzte und Krankenkassen über die ärztliche Betreuung während der Schwangerschaft und nach der Entbindung („Mutterschafts-Richtlinien“) in der Fassung vom 10.12.1985 (veröffentlicht im Bundesanzeiger Nr. 60 a vom 27. März 1986), zuletzt geändert am 28.10.2002 (veröffentlicht im Bundesanzeiger Nr. 242 vom 31.12.2002), in Kraft getreten am 1.1.2003.
2. *Aantaa K, Forss M.* Growth of the fetal biparietal diameter in different types of pregnancies. *Radiology* 1980; 137: 167-170
3. *Aardema MW, Lander M, Oosterhof H, De Wolf BT, Aarnoudse JG.* Doppler ultrasound screening predicts recurrence of poor pregnancy outcome in subsequent pregnancies, but not the recurrence of PIH or preeclampsia. *Hypertens Pregnancy* 2000; 19: 281-288
4. *Abramovich DR.* The volume of amniotic fluid in early pregnancy. *J Obstet Gynaecol Br Commonw* 1968; 75: 728-731
5. *Adkison LR, Andrews RW, Vowell NL, Koontz WL.* Improved detection of fetal cells from maternal blood with polymerase chain reaction. *Am J Obstet Gynecol* 1994; 170: 952-955
6. *Adomßent S, Sadenwasser W.* Influence of parental physical characteristics and parity on weight of the newborn infant. *Zentralbl Gynakol* 1986; 108: 26-35
7. *Albertsson-Wikland K, Boguszewski M, Karlberg J.* Children born small-for-gestational age: postnatal growth and hormonal status. *Horm Res* 1998; 49(suppl 2): 7-13
8. *Alexander GR, Himes JH, Kaufman RB, Mor J, Kogan M.* A United States national reference for fetal growth. *Obstet Gynecol* 1996; 87: 163-168
9. *Alkalay AL, Graham JM, Pomerance JJ.* Evaluation of neonates born with intrauterine growth retardation: review and practice guidelines. *Journal of Perinatology* 1998; 18: 142-151

10. *Altman D, Hytten F.* Intrauterine growth retardation: let's be clear about it. *Br J Obstet Gynecol* 1989; 96: 1127-1128
11. *Aoki M.* Fetal weight calculation; Osaka University method. In: Yoshihide C, ed. *Ultrasound in obstetrics and gynaecology*. 2nd ed. Kyoto: Kinpodo, 1990: 95-107
12. *Apuzzio J, Iffy L, Weiss G.* Mode of term singleton breech delivery. *Obstet Gynecol* 2002; 99(6): 1131-3; discussion 1131-1133
13. *Ariel I, Lerer I, Yagel S, Cohen R, Ben-Neriah Z, Abeliovich D.* Trisomy 2: Confined placental mosaicism in a fetus with intrauterine growth retardation. *Prenatal Diagnosis* 1997; 17(2): 180-183
14. *Bader B, Graham D, Stinson S.* Significance of ultrasound measurements of the head of the breech fetus. *J Ultrasound Med* 1987; 6: 437-439
15. *Bancher-Todesca D, Lee A, Dadak C, Weninger M, Kainz C.* Morbidität und Mortalität bei retardierten und normalgewichtigen Frühgeborenen. *Z Geburtshilfe Neonatol* 1995; 199: 54-57
16. *Banks EH, Miller DA.* Perinatal risks associated with borderline amniotic fluid index. *Am J Obstet Gynecol* 1999; 180: 1461-1463
17. *Barnhard Y, Bar-Hava I, Divon MY.* Is polyhydramnios in an ultrasonographically normal fetus an indication for genetic evaluation? *Am J Obstet Gynecol* 1995; 173: 1523-1527
18. *Baron C, Morgan MA, Garite TJ.* The impact of amniotic fluid volume assessed intrapartum on perinatal outcome. *Am J Obstet Gynecol* 1995; 173: 167-174
19. *Baschat AA, Harman CR.* Antenatal assessment of the growth restricted fetus. *Curr Opin Obstet Gynecol* 2001; 13: 161-168
20. *Baschat AA, Weiner CP.* Umbilical artery Doppler screening for the small for gestational age fetus in need of antenatal surveillance. *Am J Obstet Gynecol* 2000; 182: 154-158

21. *Bates JA, Evans JA, Mason G.* Differentiation of growth retarded from normally grown fetuses and prediction of intrauterine growth retardation using Doppler ultrasound. *Br J Obstet Gynaecol* 1996; 103: 670-675
22. *Becker RH, Brinnel C, Arabin B.* Der Kopfindex: ein vernachlässigter Parameter der fetalen Standardbiometrie. *Ultraschall in Med* 1992; 13: 2-6
23. *Becker R, Vonk R, Vollert W, Entezami M.* Doppler sonography of uterine arteries at 20-23 weeks: risk assessment of adverse pregnancy outcome by quantification of impedance and notch. *J Perinat Med* 2002; 30: 388-394
24. *Beinder E, Schlußner E.* Fallen die Würfel für die spätere Schwangerschaft bereits in utero? Welche Auswirkungen hat die "Fetal-Origin"-Hypothese von David Barker auf die Tätigkeit des Frauenarztes? *Frauenarzt* 44; 2003: 276-284
25. *Benoit B.* Early fetal gender determination. *Ultrasound Obstet Gynecol* 1999; 13: 299-300
26. *Benson CB, Belville JS, Lentini JF, Saltzman DH, Doubilet PM.* Intrauterine growth retardation: diagnosis based on multiple parameters – a prospective study. *Radiology* 1990; 177: 499-502
27. *Benson CB, Doubilet PM, Saltzman DH.* Intrauterine growth retardation: predictive value of ultrasound criteria for antenatal diagnosis. *Radiology* 1986; 160: 415-417
28. *Berg D, Albrecht H, Dudenhausen JW, Hochuli E, Neuhäuser G, Versmold HT, Brand M, Eskes TKAB, Kubli F, Staudach A, Wulf H.* Bericht der Standardkommission "Beckenendlage". *Z Geburtsh Perinat* 1984; 188: 100-103
29. *Berg TG, Devoe LD.* Amniotic fluid assessment. Methods and role in fetal assessment. *Clin Perinatol* 1994; 21: 809-822
30. *Berkus MD, Conway D, Langer O.* The large Fetus. *Clin Obstet Gynecol* 1999; 42: 766-784
31. *Biggio JR Jr, Wenstrom KD, Dubard MB, Cliver SP.* Hydramnios prediction of adverse perinatal outcome. *Obstet Gynecol* 1999; 94: 773-777

32. *Biswas A, Leung WC, Pun TC, Premru-Srsen T, Uchide K, Murakami K, Ponzzone R, Sismondi P, Cunha-Filho JS.* Term breech trial (Comments & Discussion). *Lancet*, 2001; 357: 225-228
33. *Boos R, Hendrik HJ, Schmidt W.* Das fetale Lageverhalten in der zweiten Schwangerschaftshälfte bei Geburten aus Beckenendlage und Schädellage. *Geburtsh. u. Frauenheilk.* 1987; 47: 341-345
34. *Boos R, Rabe D, Hendrik HJ, Schmidt W.* Geburten aus Beckenendlage – geburts-hilfliche und antepartale ultrasonographische Befunde. *Z Geburtsh und Perinat* 1985; 189: 130-135
35. *Brace RA.* Physiology of amniotic fluid volume regulation. *Clin Obstet Gynecol* 1997; 40: 280-289
36. *Brace RA, Wolf EJ.* Normal amniotic fluid volume changes throughout pregnancy. *Am J Obstet Gynecol* 1989; 161: 382-388
37. *Brenner WE, Bruce RD, Heinrichs CH.* The characteristics and perils of breech presentation. *Am J Obstet Gynecol* 1974; 118: 700-712
38. *Brenner WE, Edelmann DA, Hendricks CH.* A standard of fetal growth for the United States of America. *Am J Obstet Gynecol* 1976; 126: 555-564
39. *Brodzki J, Marsal K.* Management of pregnancies with suspected intrauterine growth retardation in Sweden. *Acta Obstet Gynecol Scand* 2000; 79: 723-728
40. *Bronshtein M, Rottem S, Yoffe N, Blumenfeld Z, Brandes JM.* Early determination of fetal sex using transvaginal sonography: technique and pitfalls. *J Clin Ultrasound* 1990; 18: 302-306
41. *Bulic M, Vratar M.* Ultrasonic planimetry of the gestation sac as a biometric method in early pregnancy. *J Clin Ultrasound* 1978; 6: 228-232
42. *Burke G, Stuart B, Crowley P, Scanail SN, Drumm J.* Is intrauterine growth retardation with normal umbilical artery blood flow a benign condition? *Br Med J* 1990; 300: 1044-1045

43. *Burke CJ, Tannenbergs AE, Payton DJ.* Ischaemic cerebral injury, intrauterine growth retardation, and placental infarction. *Dev Med Child Neurol* 1997; 39: 726-730
44. *Campbell D, Hall M, Lemon J, Carr-Hill R, Pritchard C, Samphier M.* Clinical birth-weight standards for a total population in the 1980s. *Br J Obstet Gynaecol* 1993; 100: 436-445
45. *Campbell S, Thoms A.* Ultrasound measurement of the fetal head to abdomen circumference ratio in the assessment of growth retardation. *Br J Obstet Gynaecol* 1977; 84: 165-174
46. *Campbell S, Wilkin D.* Ultrasonic measurement of fetal abdomen circumference in the estimation of fetal weight. *Br J Obstet Gynaecol* 1975; 82: 689-697
47. *Carlson DE, Platt LD, Medearis AL, Horenstein J.* Quantifiable Polyhydramnios: Diagnosis and management. *Obstet Gynecol* 1990; 75: 989-992
48. *Chamberlain PF, Manning FA, Morrison I, Harman CR, Lange IR.* Ultrasound evaluation of amniotic fluid volume. I. The relationship of marginal and decreased amniotic fluid volumes to perinatal outcome. *Am. J. Obstet. Gynecol.* 1984; 150: 245-249
49. *Chamberlain PF, Manning FA, Morrison I, Harman CR, Lange IR.* Ultrasound evaluation of amniotic fluid volume. II. The relationship of increased amniotic fluid volume to perinatal outcome. *Am J Obstet Gynecol* 1984; 150: 250-253
50. *Chandra RK.* Fetal malnutrition and postnatal immunocompetence. *Am J Dis Child* 1975; 129: 450-454
51. *Chang TC, Robson SC, Spencer JAD, Gallivan S.* Identification of fetal growth retardation; comparison of Doppler waveform indices and serial ultrasound measurements of abdominal circumference and fetal weight. *Obstet Gynecol* 1993; 82: 230-236
52. *Chard T, Yoong A, Macintosh M.* The myth of growth retardation at term. *Br J Obstet Gynaecol* 1993; 100: 1076-1081
53. *Charles D, Jacobi HE, Burgess F.* Amniotic fluid volumes in the second half of pregnancy. *Am J Obstet Gynecol* 1965; 93: 1042-1047

54. *Charles D, Jacobi HE.* Preliminary data on use of sodium aminohippurate to determine amniotic fluid volumes. *Am J Obstet Gynecol* 1966; 95: 266-269
55. *Chauhan SP.* Amniotic fluid index before and after amnioinfusion of a fixed volume of normal saline. *J Reprod Med* 1991; 36: 801-802
56. *Chauhan SP, Magann EF, Morrison JC, Whitworth NS, Hendrix NW, Devoe LD.* Ultrasonographic assessment of amniotic fluid does not reflect actual amniotic fluid volume. *Am J Obstet Gynecol* 1997; 177: 291-297
57. *Chauhan SP, Scardo JA, Hendrix NW, Magann EF, Morrison JC.* Accuracy of sonographically estimated fetal weight with and without oligohydramnios. A case-control study. *J Reprod Med* 1999; 44: 969-973
58. *Chauhan SP, West DJ, Scardo JA, Boyd JM, Joiner J, Hendrix NW.* Antepartum detection of makrosomic fetus: clinical versus sonographic, including soft-tissue measurements. *Obstet Gynecol* 2000; 95: 639-642
59. *Chien PFW, Owen P, Khan KS.* Validity of ultrasound estimation of fetal weight. *Obstet Gynecol* 2000; 95: 856-860
60. *Chinn DH, Filly R, Callen PW.* Prediction of intrauterine growth retardation by sonographic estimation of total intrauterine volume. *JCU* 1981; 9: 175-179
61. *Chowen JA, Evain-Brion D, Pozo J, Alsat E, García-Segura LM, Argente J.* Decreased expression of placental growth hormone in intrauterine growth retardation. *Pediatr Res* 1996; 39: 736-739
62. *Colley NV, Tremble JM, Henson GL, Cole TJ.* Head circumference/ abdominal circumference ratio, ponderal index and fetal malnutrition. Should head circumference/ abdominal circumference ratio be abandoned? *Br J Obstet Gynaecol* 1991; 98: 524-527
63. *Combs CA, Jaekle RK, Rosenn B, Pope M, Miodovnik M, Siddiqi TA.* Sonographic estimation of fetal weight based on a model of fetal volume. *Obstet Gynecol* 1993; 82: 365-370
64. *Croom CS, Baines BB, Ramos-Santos E.* Do semiquantitative amniotic fluid indexes reflect actual volume? *Am J Obstet Gynecol* 1992; 167: 995-999

65. *Croughan-Minihane MS, Petitti DB, Gordis L, Golditch I.* Morbidity among breech infants according to method of delivery. *Obstet Gynecol* 1990; 75: 821-825
66. *Dahri S, Snoek A, Reusens-Billen B, Remacle C, Hoet JJ.* Islet function in offspring of mothers on low-protein diet during gestation. *Diabetes* 1991; 40(Suppl 2): 115-120
67. *Dar P, Gross SJ.* Macrosomia: A genetic perspective. *Clin Obstet Gynecol* 2000; 43: 298-308
68. *Dashe JS, McIntire DD, Lucas MJ, Leveno KJ.* Effects of symmetric and asymmetric fetal growth on pregnancy outcomes. *Obstet Gynecol* 2000; 96: 321-327
69. *Dashe JS, McIntire DD, Ramus RM, Santos-Ramos R, Twickler DM.* Hydramnios: Anomaly prevalence and sonographic detection. *Obstet Gynecol* 2002; 100: 134-139
70. *David C, Tagliavini G, Pilu G, Rudenholz A, Bovicelli L.* Receiver-operator characteristic curves for the ultrasonographic prediction of small-for-gestational-age fetuses in low-risk pregnancies. *Am J Obstet Gynecol* 1996; 174: 1037-1042
71. *Denbow ML, Sepulveda W, Ridout D, Fisk NM.* Relationship between change in amniotic fluid index and volume of fluid removed at amnioreduction. *Obstet Gynecol* 1997; 90: 529-532
72. *Dieckmann WJ, Davies ME.* The volumetric determination of amniotic fluid with congo red. *Am J Obstet Gynecol* 1933; 25: 623-627
73. *Dildy GA, Clark SL.* Shoulder dystocia: Risk identification. *Clin Obstet Gynecol* 2000; 43: 265-282
74. *Dildy GA, Lira N, Moise KJ Jr, Riddle GD, Deter RL.* Amniotic fluid volume assessment: comparison of ultrasonographic estimates versus direct measurements with a dye-dilution technique in human pregnancy. *Am J Obstet Gynecol* 1992; 167: 986-994
75. *Divon MY, Guidetti DA, Braverman JJ, Oberlander E, Langer O, Merkatz IR.* Intrauterine growth retardation – a prospective study of the diagnostic value of real-time sonography combined with umbilical artery flow velocimetry. *Obstet Gynecol* 1988; 72: 611-614

76. *Divon MY, Marks AD, Henderson CE.* Longitudinal measurement of amniotic fluid index in postterm pregnancies and its association with fetal outcome. *Am J Obstet Gynecol* 1995; 172: 142-146
77. *Doubilet PM, Benson CB.* Sonographic evaluation of intrauterine growth retardation. *Am J Roentgenol* 1995; 164(3): 709-717
78. *Dubinsky T, Lau M, Powell F, Garcia J, Mastobattista J, Parvey HR, Sickler K, Maklad M.* Predicting poor neonatal outcome: A comparative study of noninvasive antenatal testing methods. *Am J Roentgenol* 1997; 168: 827-831
79. *Dulitzki M, Soriano D, Schiff E, Chetrit A, Mashiach S, Seidman DS.* Effect of very advanced maternal age on pregnancy outcome and rate of cesarean delivery. *Obstet Gynecol* 1998; 92: 935-939
80. *Efrat Z, Akinfenwa OO, Nicolaidis KH.* First-trimester determination of fetal gender by ultrasound. *Ultrasound Obstet Gynecol* 1999; 13: 305-307
81. *Emerson DS, Felker RE, Brown DL.* The sagittal sign. An early second trimester sonographic indicator of fetal gender. *J Ultrasound Med* 1989; 8: 293-297
82. *Feige A, Krause M, Lenz A.* Entscheidungskriterien zur vaginalen Beckenendlagegeburt – Einfluß auf die kindliche Früh- und Spätmorbidität. *Z Geburtshilfe Neonatol* 1997; 201 Suppl 1: 35-42
83. *Feige A, Douros A.* Mortalität und Morbidität kleiner Frühgeborener (<1500g) in Abhängigkeit von Poleinstellung und Entbindungsmodus. *Z Geburtshilfe Neonatol* 1996; 200: 50-55
84. *Feldman KW, Smith DW.* Fetal phyllic growth and penile standards for newborn male infants. *J Pediatr* 1975; 86: 395-398
85. *Ferber A.* Maternal complications of fetal macrosomia. *Clin Obstet Gynecol* 2000; 43: 335-339
86. *Fianu S, Vaclavinkova V.* External cephalic version in the management of breech presentation with special references to the placental location. *Acta Obstet Gynecol Scand* 1979; 58: 209-210



87. *Fischer Th, Krause M, Feige A.* Beckenendlage – Informations-, Angst- und Erwartungsmuster von niedergelassenen Frauenärzten und Schwangeren. Teil 2: Ergebnisse einer Umfrage unter Frauen mit einer BEL-Schwangerschaft. *Z Geburtsh Neonatol* 1996; 200: 61-65
88. *Fitzhardinge PM, Steven EM.* The small-for-date infant. II. Neurological and intellectual sequelae. *Pediatrics (United States)* 1972; 50: 50-57
89. *Flock F, Stoz F, Paulus W, Scheurle B, Kreienberg R.* Äußere Wendung aus Beckenendlage in Schädellage: Einflußfaktoren, Nutzen und Risiken. *Zentralbl Gynakol* 1998; 120: 60-65
90. *Frenzel J, Krause W, Sander I, Michels W.* Zur Früh- und Spätmorbidität mindergewichtiger Neugeborener (LBWI) nach Beckenendlagen in Abhängigkeit vom Entbindungsmodus. *Z Geburtshilfe Perinatol* 1984; 188: 261-268
91. *Fuchs F.* Volume of amniotic fluid at various stages of pregnancy. *Clin Obstet Gynecol* 1966; 9: 449-460
92. *Gadd RL.* The volume of the liquor amnii in normal and abnormal pregnancies. *J Obstet Gynaec Br Commonw* 1966; 73: 11-22
93. *Gardosi J.* Customized growth curves. *Clin Obstet Gynecol* 1997; 40: 715-722
94. *Gardosi J, Mul T, Mongelli M.* Application of a fetal weight standard to study associations between growth retardation and stillbirth. *Ultrasound Obstet Gynecol* 1995; 6 (2 Suppl): 66
95. *Geirsson RT, Christie AD, Patel N.* Ultrasound volume measurements comparing a prolate ellipsoid method with a parallel planimetric area method against a known volume. *JCU* 1982; 10: 329-332
96. *Geirsson RT, Ogston SA, Patel NB, Christie AD.* Growth of total intrauterine, intra-amniotic and placental volume in normal singleton pregnancy measured by ultrasound. *Br J Obstet Gynaecol* 1985; 92: 46-53
97. *Geirsson RT, Patel NB, Christie AD.* Efficacy of intrauterine volume, fetal abdominal area and biparietal diameter measurements with ultrasound in the prediction of small-for-dates babies. *Br J Obstet Gynaecol* 1985; 92: 929-935

98. *Geirsson RT, Patel NB, Christie AD.* Intrauterine volume, fetal abdominal area and biparietal diameter measurements with ultrasound in the prediction of small-for-dates babies in a high-risk obstetric population. *Br J Obstet Gynaecol* 1985; 92: 936-940
99. *Geirsson RT, Patel NB, Christie AD.* In-vivo accuracy of ultrasound measurements of intrauterine volume in pregnancy. *Br J Obstet Gynaecol* 1984; 91: 37-40
100. *Gifford DS, Morton SC, Fiske M, Kahn K.* A meta-analysis of infant outcomes after breech delivery. *Obstet Gynecol* 1995; 85: 1047-1054
101. *Giles WB, Trudinger BJ, Baird PJ.* Fetal umbilical artery flow velocity waveforms and placental resistance: pathological correlation. *Br J Obstet Gynaecol* 1985; 92: 31-38
102. *Gillibrand PN.* Changes in amniotic fluid volume with advancing pregnancy. *J Obstet Gynaecol Br Commonw* 1969; 76: 527-529
103. *Gluckman PD, Harding JE.* The physiology and pathophysiology of intrauterine growth retardation. *Horm Res* 1997; 48(suppl 1): 11-16
104. *Göttlicher S, Madjaric J.* Morgens KL – mittags BEL: Ein Ammen-märchen? *Geburtsh. u. Frauenheilk.* 1989; 49: 363-366
105. *Göttlicher S, Madjaric J.* Die Lage der menschlichen Frucht im Verlauf der Schwangerschaft und die Wahrscheinlichkeit einer spontanen Drehung in die Kopflage bei Erst- und Mehrgebärenden. *Geburtsh. u. Frauenheilk.* 1985; 45: 534-538
106. *Göttlicher S, Madjaric J, Krone HA.* Über die Lage des menschlichen Feten und die Wahrscheinlichkeit. *Z. Geburtsh. u. Perinat.* 1981; 185: 288-292
107. *Gohari P, Berkowitz RL, Hobbins JC.* Prediction of intrauterine growth retardation by determination of total intrauterine volume. *Am J Obstet Gynecol* 1977; 127: 255-260
108. *Goldenberg RL, Cutter GR, Hoffman HJ, Foster JM, Nelson KG, Hauth JC.* Intrauterine growth retardation: Standards for diagnosis. *Am J Obstet Gynecol* 1989; 161: 271-277
109. *Goldstein RB, Filly RA.* Sonographic estimation of amniotic fluid volume. Subjective assessment versus pocket measurements. *J Ultrasound Med* 1988; 7: 363-369

110. *Golfier F, Vaudoyer F, Ecochard R, Champion F, Audra P, Raudrant D.* Planned vaginal delivery versus elective caesarean section in singleton term breech presentation: a study of 1116 cases. *Eur J Obstet Gynecol Reprod Biol* 2001; 98(2): 186-192
111. *Gray DL, Songster GS, Parvin CA, Crane JP.* Cephalic index: a gestational age-dependent biometric parameter. *Obstet Gynecol* 1989; 74: 600-603
112. *Greene MF.* Vaginal breech delivery is no longer justified. *Obstet Gynecol* 2002; 99(6): 1113-1115
113. *Grossman M, Flynn JJ, Aufrichtig D, Handler CR.* Pitfalls in ultrasonic determination of total intrauterine volume. *J Clin Ultrasound* 1982; 10: 17-20
114. *Grover J, Mentakis EA, Ross MG.* Three-dimensional method for determination of amniotic fluid volume in intrauterine pockets. *Obstet Gynecol* 1997; 90: 1006-1010
115. *Haber Kern CM, Smith DW, Jones KL.* The "breech head" and its relevance. *Am J Dis Child* 1979; 133: 154-156
116. *Hadlock FP, Deter RL, Carpenter RJ, Park SK.* Estimating fetal age: effect of head shape on BPD. *AJR* 1980; 137: 83-85
117. *Hadlock FP, Harrist RB, Martinez-Poyer J.* In-utero analysis of fetal growth: a sonographic weight standard. *Radiology* 1991; 181: 129-133
118. *Hadlock FP, Harrist RB, Sharman RS, Deter RL, Park SK.* Estimation of fetal weight with the use of head, body, and femur measurements: a prospective study. *Am J Obstet Gynecol* 1985; 151: 333-337
119. *Haller U, Hepp H, Winter R.* Sectio nach Wunsch oder elektive Sectio: Aufforderung zum Umdenken. *Gynäkol Geburtshilfliche Rundsch* 2002; 42: 1-3
120. *Hannah ME, Hannah WJ, Hewson SA, Hodnett ED, Saigal S, Willian AR, for the Term breech Trial Collaborative Group.* Planned caesarean section versus planned vaginal birth for breech presentation at term: A randomised multicentre trial. *Lancet* 2000; 356: 1375-1383
121. *Haram K, Pirhonen J, P Bergsjø.* Suspected big baby: a difficult clinical problem in obstetrics. *Acta Obstet Gynecol Scand* 2002; 81: 185-194

122. *Harding JE, Bauer MK, Kimble RM.* Antenatal therapy for intrauterine growth retardation. *Acta Paediatr Suppl* 1997; 423: 196-201
123. *Harrington K, Armstrong V, Freeman J, Aquilina J, Campbell S.* Fetal sexing by ultrasound in the second trimester: maternal preferences and professional ability. *Ultrasound Obstet Gynecol* 1996; 8: 318-321
124. *Haswell GL, Morris JA.* Amniotic fluid volume studies. *Obstet Gynecol* 1973; 42: 725-732
125. *Hauth JC, Cunningham FG.* Vaginal breech delivery is still justified. *Obstet Gynecol* 2002; 99(6): 1115-1116
126. *Hellmeyer L, Meyer-Wittkopf M, Ziegler A, Ballast A, Sierra F, Schmidt S.* Comparison of sonographic weight estimation to actual birth weight: a retrospective analysis. *Ultraschall Med* 2001; 22: 167-171
127. *Henner H, Wolf-Zimpel O, Rüttgers H, Haller U, Kubli F.* Häufigkeit und Verteilung von Beckenendlagen in der Schwangerschaft und bei Geburt. *Z Geburtsh Perinat* 1975; 179: 17-23
128. *HersHKovitz R, Kingdom JC, Geary M, Rodeck CH.* Fetal cerebral blood flow redistribution in late gestation: identification of compromise in small fetuses with normal umbilical artery Doppler. *Ultrasound Obstet Gynecol* 2000; 15: 209-212
129. *Hickl EJ.* Wandlungen in der Kaiserschnittindikation. *Gynäkol Geburtshilfliche Rundsch* 2002; 42: 15-18
130. *Hill LM.* Prevalence of breech presentation by gestational age. *Am J Perinat* 1990; 7(1): 92-93
131. *Hill LM, Breckle R, Thomas ML, Fries JK.* Polyhydramnios: Ultrasonically detected prevalence and neonatal outcome. *Obstet Gynecol* 1987; 69: 21-25
132. *Hill LM, Breckle R, Wolfgram KR, O'Brien PC.* Oligohydramnios: ultrasonically detected incidence and subsequent fetal outcome. *Am J Obstet Gynecol* 1983; 147: 407-410

133. *Hobbins JC, Grannum PAT, Berkowitz RL, Silverman R, Mahoney MJ.* Ultrasound in the diagnosis of congenital anomalies. *Am J Obstet Gynecol* 1979; 134: 331-345
134. *Hodnett E, Hannah M.* Term breech trial (Letters). *Birth* 2002; 29(3): 217-219
135. *Hofman PL, Cutfield WS, Robinson EM, Bergman RN, Menon RK, Sperling MA, Gluckman PD.* Insulin resistance in short children with intrauterine growth retardation. *Journal of Clinical Endocrinology and Metabolism* 1997; 82: 402-406
136. *Hofmann D, Carle-Grewe G.* Ergebnisse bei überwiegend vaginal durchgeführten Beckenendlagenentbindungen. *Zentralbl Gynakol* 1990; 112: 1525-1540
137. *Hofmeyr GJ, Sadan O, Myer IG, Galal KC, Simko G.* External cephalic version and spontaneous version rates: ethnic and other determinants. *Br J Obstet Gynaecol* 1986; 93: 13-16
138. *Hohenauer L.* Intrauterine Wachstumskurven für den deutschen Sprachraum. *Z Geburtshilfe Perinatol* 1980; 184: 167-179
139. *Hohenauer L.* Intrauterines Längen- und Gewichtswachstum. Standardwerte aus Österreich. *Pädiatrie und Pädologie* 1973; 195-205
140. *Hohlfeld P.* Cesarean Section on Request: A Case for Common Sense. *Gynäkol Geburtshilfliche Rundsch* 2002; 42: 19-21
141. *Horsager R, Nathan L, Leveno KJ.* Correlation of measured amniotic fluid volume and sonographic predictions of oligohydramnios. *Obstet Gynecol* 1994; 83: 955-958
142. *Hoskins IA, McGovern PG, Ordorica SA, Frieden FJ, Young BK.* Amniotic fluid index: correlation with amniotic fluid volume. *Am J Perinatol* 1992; 9: 315-318
143. *Hsieh TT, Hung TH, Chen KC, Hsieh CC, Lo LM, Chiu TH.* Perinatal outcome of oligohydramnios without associated premature rupture of membranes and fetal anomalies. *Gynecol Obstet Invest* 1998; 45: 232-236
144. *Husslein P.* Elektive Sectio. *Gynäkol Geburtshilfliche Rundsch* 2002; 42: 22-24
145. *Irion O, Hirsbrunner Almagbaly P, Morabia A.* Planned vaginal delivery versus elective caesarean section: a study of 705 singleton term breech presentations. *Br J Obstet Gynaecol* 1998; 105: 710-717

146. *Irvine LM*. Patient education is way to influence maternal requests for caesarean section. *BMJ* 1999; 319: 190
147. *Ismail MA, Nagib N, Ismail T, Cibilis LA*. Comparison of vaginal and cesarean section delivery for fetuses in breech presentation. *J Perinat Med* 1999; 27: 339-351
148. *Jeng CJ, Lee JF, Wang KG, Yang YC, Lan CC*. Decreased amniotic fluid index in term pregnancy. Clinical significance. *J Reprod Med* 1992; 37: 789-792
149. *Joern H; Rath W*. Comparison of Doppler sonographic examinations of the umbilical and uterine arteries in high-risk pregnancies. *Fetal Diagn Ther* 1998; 13:150-153
150. *Jones RAK, Robertson NRC*. Small for dates babies: are they really a problem? *Arch Dis Child* 1986; 61: 877-880
151. *Jones TB, Price RR, Gibbs SJ*. Volumetric determination of placental and uterine growth relationships from B-mode ultrasound by serial area-volume determinations. *Invest Radiol* 1981; 16: 101-106
152. *de Jong CLD, Gardosi J, Dekker GA, Colenbrander GJ, van Geijn HP*. Application of a customised birthweight standard in the assessment of perinatal outcome in a high risk population. *Br J Obstet Gynaecol* 1998; 105: 531-535
153. *Jovanovic L*. What is so bad about a big baby? *Diabetes Care* 2001; 24: 1317-1318
154. *Kasby CB, Poll V*. The breech head and its ultrasound significance. *Br J Obstet Gynaecol* 1982; 89: 106-110
155. *Kawasaki N, Nishimura H, Yoshimura T, Okamura H*. A diminished intrapartum amniotic fluid index is a predictive marker of possible adverse neonatal outcome when associated with prolonged labor. *Gynecol Obstet Invest* 2002; 53: 1-5
156. *Keirse MJNC*. Evidence-based childbirth only for breech babies? *Birth* 2002; 29: 55-59
157. *Kirkinen P, Ylöstalo P*. Ultrasonic examination before external version of breech presentation. *Gynecol Obstet Invest* 1982; 13: 90-97

158. *Kirschbaum M, Bödeker RH, Münstedt K, Künzel W.* Der Stellenwert der präpartualen sonographischen Gewichtsschätzung bei Beckenendlagen. *Geburtshilfe Frauenheilkd* 1992; 52: 264-269
159. *Klein M, Waldhor T, Vutuc C, Beck A.* Die Häufigkeit von Kaiserschnitten in Österreich (1996-1998). *Gynaekol Geburtshilfliche Rundsch* 2000; 40: 125-129
160. *Kofinas AD, Penry M, Hatjis CG.* Compliance-weight deficit index. Combining umbilical artery resistance and growth deficit for predicting intrauterine growth retardation and poor neonatal outcome. *J Reprod Med* 1994; 39: 595-600
161. *Kouam L, Miller EC.* Zum Problem der perinatalen Mortalität bei Beckenendlagenkindern mit Geburtsgewichten über 2500g. *Zentralbl Gynakol* 1980; 102: 1375-1382
162. *Kramer MS, Morin I, Yang H, Platt RW, Usher R, McNamara H, Joseph KS, Wen SW.* Why are babies getting bigger? Temporal trends in fetal growth and its determinants. *J Pediatr* 2002; 141: 538-542
163. *Kramer MS, Olivier M, McLean FH, Willis DM, Usher RH.* Impact of intrauterine growth retardation and body proportionality on fetal and neonatal outcome. *Pediatrics* 1990; 86: 707-713
164. *Krause M, Fischer Th, Feige A.* Beckenendlage – Informations-, Angst- und Erwartungsmuster von niedergelassenen Frauenärzten und Schwangeren. Teil 1: Ergebnisse einer Umfrage unter niedergelassenen Frauenärzten. *Z Geburtsh Neonatol* 1994; 198: 113-116
165. *Krause M, Fischer Th, Feige A.* Vaginale Geburt aus Beckenendlage erhöht nicht die kindliche Frühmorbidity – Ergebnisse von 423 aus Beckenendlage geborenen Kindern aus den Jahren 1988-1992. *Z Geburtsh u Perinat* 1994; 198: 88-95
166. *Kropp S, Schmidt S, Saling E.* Einflüsse auf Entscheidungen von Schwangeren mit BEL zur äußeren Wendung oder zur Schnittentbindung – Aufklärungsstatus und Angstmuster. *Z Geburtshilfe Perinatol* 1989; 193: 84-91
167. *Kubli F.* Geburtseinleitung bei Beckenendlagen. *Gynäkologe* 1975; 8: 48-97
168. *Kumari AS.* Pregnancy outcome in women with morbid obesity. *Int J Gynecol Obstet* 2001; 73: 101-107

169. *Kurmanavicius J, Wright EM, Royston P, Wisser J, Huch R, Huch A, Zimmermann R.* Fetal ultrasound biometry: 1. Head reference values. *Br J Obstet Gynaecol* 1999; 106: 126-135
170. *Landy HJ, Isada NB, Larsen JW.* Genetic implications of idiopathic hydramnios. *Am J Obstet Gynecol* 1987; 157: 114-117
171. *Langer O.* Fetal macrosomia: Etiologic factors. *Clin Obstet Gynecol* 2000; 43: 283-297
172. *Lapunzina P, Camelo JSL, Rittler M, Castilla EE.* Risks of congenital anomalies in large for gestational age infants. *J Pediatr* 2002; 140: 200-204
173. *Larsen T, Larsen JF, Petersen S, Greisen G.* Detection of small-for-gestational-age fetuses by ultrasound screening in a high risk population: a randomized control study. *Br J Obstet Gynaecol* 1992; 99: 469-474
174. *Lazebnik N, Many A.* The severity of polyhydramnios, estimated fetal weight and pre-term delivery are independent risk factors for the presence of congenital malformations. *Gynecol Obstet Invest* 1999; 48: 28-32
175. *Lepercq J, Mahieu-Caputo D.* Diagnosis and management of intrauterine growth retardation. *Horm Res* 1998; 49(suppl 2): 14-19
176. *Leucht W, Rabe D, Hendrik HJ, Grischke EM, Schmidt W.* Sono-graphische Beurteilung der Fruchtwassermenge I. Polyhydramnion – Bedeutung für den Schwangerschafts- und Geburtsverlauf. *Geburtsh. u. Frauenheilk.* 1986; 46: 157-161
177. *Levine S, Filly RA, Creary RK.* Identification of fetal growth retardation by ultrasonographic estimation of total intrauterine volume. *J Clin Ultrasound* 1979; 7: 21-26
178. *Lituania M, Passamonti U, Esposito V.* Genetic factors and fetal anomalies in intrauterine growth retardation. *J Perinat Med* 1994; 22: 79-83
179. *Lu GC, Rouse DJ, DuBard M, Cliver S, Kimberlin D, Hauth JC.* The effect of the increasing prevalence of maternal obesity on perinatal morbidity. *Am J Obstet Gynecol* 2001; 185: 845-849



180. *Lubchenco LO, Hausmann C, Dressler M, Boyd E.* Intrauterine growth as estimated from liveborn birth weight data at 24 to 42 weeks of gestation. *Pediatrics* 1963; 32: 793-800
181. *Macfarlane A.* At last: maternity statistics for England. *BMJ* 1998; 316: 566-567
182. *Magann EF, Bass D, Chauhan SP, Young RA, Whithworth NS, Morrison JC.* Amniotic fluid volume in normal singleton pregnancies. *Obstet Gynecol* 1997; 90: 524-528
183. *Magann EF, Chauhan SP, Barrilleaux PS, Whitworth NS, McCurley S, Martin JN.* Ultrasound estimate of amniotic fluid volume: color Doppler overdiagnosis of oligohydramnios. *Obstet Gynecol* 2001; 98: 71-74
184. *Magann EF, Chauhan SP, Whitworth NS, Isler C, Wiggs C, Morrison JC.* Subjective versus objective evaluation of amniotic fluid volume of pregnancies of less than 24 weeks' gestation: how can we be accurate? *J Ultrasound Med* 2001; 20: 191-195
185. *Magann EF, Chauhan SP, Barrilleaux PS, Withworth NS, Martin JN.* Amniotic fluid index and single deepest pocket: Weak indicators of abnormal amniotic volumes. *Obstet Gynecol* 2000; 96: 737-740
186. *Magann EF, Chauhan SP, Kinsella MJ, McNamara MF, Whitworth NS, Morrison JC.* Antenatal testing among 1001 patients at high risk: The role of ultrasonographic estimate of amniotic fluid volume. *Am J Obstet Gynecol* 1999; 180: 1330-1336
187. *Magann EF, Kinsella MJ, Chauhan SP, McNamara MF, Gehring BW, Morrison JC.* Does an amniotic fluid index of  $\leq 5$ cm necessitate delivery in high-risk pregnancies? A case-control study. *Am J Obstet Gynecol* 1999; 180: 1354-1359
188. *Magann EF, Morton ML, Nolan TE, Martin JN, Whithworth NS, Morrison JC.* Comparative efficacy of two sonographic measurements for the detection of aberrations in the amniotic fluid volume and the effect of amniotic fluid volume on pregnancy outcome. *Obstet Gynecol* 1994; 83: 959-962
189. *Magann EF, Nolan TE, Hess LW, Martin RW, Whitworth NS, Morrison JC.* Measurement of amniotic fluid volume: Accuracy of ultrasonography techniques. *Am J Obstet Gynecol* 1992; 167: 1533-1537

190. *Magann EF, Sanderson M, Martin JN, Chauhan S.* The amniotic fluid index, single deepest pocket and two-diameter pocket in normal human pregnancy. *Am J Obstet Gynecol* 2000; 182: 1581-1588
191. *Maher JE, Wenstrom KD, Hauth JC, Meis PJ.* Amniotic fluid embolism after saline amnioinfusion: two cases and review of the literature. *Obstet Gynecol* 1994; 83: 851-854
192. *Manning FA, Hill LM, Platt LD.* Qualitative amniotic fluid volume determination by ultrasound: Antepartum detection of intrauterine growth retardation. *Am J Obstet Gynecol.* 1981; 139: 254-258
193. *Manning FA, Platt LD, Sipos L.* Antepartum fetal evaluation: development of a fetal biophysical profile score. *Am J Obstet Gynecol* 1980; 136: 787-795
194. *Many A, Hill LM, Lazebnik N, Martin JG.* The association between polyhydramnios and preterm delivery. *Obstet Gynecol* 1995; 86: 389-391
195. *Marsden D, Huntingford PJ.* An appraisal of the coomassie blue dilution technique for measuring the volume of liquor amnii in late pregnancy. *J Obstet Gynaecol Br Commonw* 1965; 72: 65-68
196. *Martin WL, Durrant LG, Liu DTY.* Non-invasive fetal cell isolation from maternal blood. *Br J Obstet Gynaecol* 1998; 105: 576-583
197. *Mau G.* Somatogramme für westdeutsche Neugeborene. *Klin Pädiatr* 1976; 188: 42-50
198. *Mazor M, Ghezzi F, Maymon E, Shoham-Vardi I, Vardi H, Hershkowitz R, Leiberman JR.* Polyhydramnios is an independent risk factor for perinatal mortality and intrapartum morbidity in preterm delivery. *Eur J Obstet Gynecol Reprod Biol* 1996; 70: 41-47
199. *Mazza V, Contu G, Falcinelli C, Battafarano S, Cagnacci A, Vito G, Forabosco A, Volpe A.* Biometrical threshold of biparietal diameter for certain fetal sex assignment by ultrasound. *Ultrasound Obstet Gynecol* 1999; 13: 308-311
200. *Meagher S, Davison G.* Early second-trimester determination of fetal gender by ultrasound. *Ultrasound Obstet Gynecol* 1996; 8: 322-324

201. *Mercer LJ, Brown LG, Petres RE, Messer RH.* A survey of pregnancies complicated by decreased amniotic fluid. *Am J Obstet Gynecol* 1984; 149: 355-361
202. *Mellor DJ, Matheson IC.* Daily changes in the curved crown-rump length of individual sheep fetuses during the last 60 days of pregnancy and effects of different levels of maternal nutrition. *Q J Exp Physiol* 1979; 64: 119-131
203. *F Mikulandra, M Perisa, E Stojni.* When is fetal macrosomia (> or = 4500 g) an indication for caesarean section? *Zentralblatt für Gynäkologie* 1996; 118(8): 441-447
204. *Miller JM, Kissling GA, Brown HL, Gabert HA.* Estimated fetal weight: applicability to small- and large-for-gestational-age fetus. *J Clin Ultrasound* 1988; 16: 95-97
205. *Miller EC, Kouam L.* Zur Häufigkeit von Beckenendlagen im Verlauf der Schwangerschaft und zum Zeitpunkt der Geburt. *Zbl. Gynäkol.* 1981; 103: 105-109
206. *Moise KJ.* Polyhydramnios. *Clin Obstet Gynecol* 1997; 40: 266-279
207. *Moore TR.* Clinical assessment of amniotic fluid. *Clin Obstet Gynecol* 1997; 40: 303-313
208. *Moore TR.* Superiority of the four-quadrant sum over the single-deepest-pocket technique in ultrasonographic identification of abnormal amniotic fluid volumes. *Am J Obstet Gynecol* 1990; 163: 762-767
209. *Morales-Rosello J.* Physiologic restriction versus genetic weight potential: study in normal fetuses and in fetuses with intrauterine growth retardation. *J Ultrasound Med* 1999; 18: 343-347
210. *Murray SR.* Hydramnios. A study of 846 cases. *Am J Obstet Gynecol* 1964; 88: 65-67
211. *Natsuyama E.* Sonographic determination of fetal sex from twelve weeks of gestation. *Am J Obstet Gynecol* 1984; 149: 748-757
212. *Nelson MM.* Amniotic fluid volumes in early pregnancy. *J Obstet Gynaecol Br Commonw* 1972; 79: 50-53
213. *Nohe G, Hartmann W, Klapproth CE.* Äußere Wendung des Feten als ambulanter Eingriff. *Geburtshilfe Frauenheilkd* 1996; 56: 328-330

214. *Normington EA*. The use of sodium amino hippurate as an alternative to Coomassie Blue to determine amniotic fluid volume. *J Obstet Gynaecol Br Commonw* 1970; 77: 156-157
215. *O'Brien JM, Mercer BM, Friedman SA, Sibai BM*. Amniotic fluid index in hospitalized hypertensive patients managed expectantly. *Obstet Gynecol* 1993; 82: 247-250
216. *Obwegeser R, Ulm M, Simon M, Ploeckinger B, Gruber W*. Breech infants: vaginal or cesarean delivery? *Acta Obstet Gynecol Scand* 1996; 75: 912-916
217. *Oral E, Cagas A, Gezer A, Kaleli S, Aydinli K, Öcer F*. Perinatal and maternal outcomes of fetal macrosomia. *Eur J Obstet Gynecol Reprod Biol* 2001; 99: 167-171
218. *O'Reilly-Green C, Divon M*. Sonographic and clinical methods in the diagnosis of macrosomia. *Clin Obstet Gynecol* 2000; 43: 309-320
219. *Ott WJ*. Sonographic diagnosis of intrauterine growth restriction. *Clin Obstet Gynecol* 1997; 40: 787-795
220. *Ott WJ, Doyle S, Flamm S*. Accurate ultrasonic estimation of fetal weight: III Prospective analysis of new ultrasonic formulae. *Am J Perinatol* 1986; 3: 307-311
221. *Ott WJ, Doyle S, Flamm S*. Accurate ultrasonic estimation of fetal weight. *Am J Perinatol* 1985; 2: 178-182
222. *Ott WJ, Doyle S*. Ultrasonic diagnosis of altered fetal growth by use of a normal ultrasonic fetal weight curve. *Obstet Gynecol* 1984; 63: 201-204
223. *Ott WJ, Doyle S*. Normal ultrasonic fetal weight curve. *Obstet Gynecol* 1982; 59: 603-606
224. *Owen P, Khan KS, Howie P*. Single and serial estimates of amniotic fluid volume and umbilical artery resistance in the prediction of intrauterine growth restriction. *Ultrasound Obstet Gynecol* 1999; 13: 415-419
225. *Pajntar M, Verdenik I, Pestevsek M*. Caesarean section in breech by birth weight. *Eur J Obstet Gynecol Reprod Biol* 1994; 54: 181-184

226. *Palo P, Erkkola R, Piironen O, Ruotsalainen P.* Accuracy of ultrasonic fetal weight estimation and detection of small for gestational age fetuses. *Am J Perinatol* 1989; 6: 400-404
227. *Panting-Kemp A, Nguyen T, Chang E, Quillen E, Castro L.* Idiopathic polyhydramnios and perinatal outcome. *Am J Obstet Gynecol* 1999; 181: 1079-1082
228. *Patterson RM, Prihoda TJ, Pouliot MR.* Sonographic amniotic fluid measurement and fetal growth retardation: a reappraisal. *Am J Obstet Gynecol* 1987; 157: 1406-1410
229. *Phelan JP, Ahn MO, Smith CV, Rutherford SE, Anderson E.* Amniotic fluid volume measurements during pregnancy. *J Reprod Med* 1987; 32: 601-604
230. *Phelan JP, Smith CV, Broussard P, Small M.* Amniotic fluid volume assessment with the four-quadrant technique at 36-42 weeks' gestation. *J Reprod Med* 1987; 32: 540-542
231. *Philipson EH, Sokol RJ, Williams T.* Oligohydramnios: clinical associations and predictive value for intrauterine growth retardation. *Am J Obstet Gynecol* 1983; 146: 271-278
232. *Piper JM, Xenakis EMJ, McFarland M, Elliot BD, Berkus MD, Langer O.* Do growth-retarded premature infants have different rates of perinatal mortality than appropriately grown premature infants? *Obstet Gynecol* 1996; 87: 169-174
233. *Queenan JT, Thompson W, Whitfield CR, Shah SI.* Amniotic fluid volumes in normal pregnancies. *Am. J. Obstet. Gynecol.* 1972; 114: 34-38
234. *Rabe D, Leucht W, Hendrik HJ, Boos R, Schmidt W.* Sonographische Beurteilung der Fruchtwassermenge II. Oligohydramnion – Bedeutung für den Schwangerschafts- und Geburtsverlauf. *Geburtsh. u. Frauenheilk.* 1986; 46: 422-426
235. *Ragosch V, Entezami M, Hundertmark S, Mutz A.* Accuracy of ultrasound weight assessment – comparison of vertex vs. breech presentation. *Ultraschall in der Medizin* 1997; 18: 19-25
236. *Rayburn WF, Motley ME, Stempel LE, Gendreau M.* Antepartum prediction of the postmature infant. *Obstet Gynecol* 1982; 60-2: 148-153

237. *Remy N, Jaluvka V, Weitzel HK.* Mortalität und Letalität nach Schnittentbindung in West-Berlin 1975 bis 1989. *Zentralblatt für Gynäkologie* 1993; 115(1): 7-12
238. *Reynolds RM, Phillips DIW.* Long-term consequences of intrauterine growth retardation. *Horm Res* 1998; 49(suppl 2): 28-31
239. *Rhodes P.* The volume of liquor amnii in early pregnancy. *J Obstet Gynaec Br Commonw* 1966; 73: 23-26
240. *Richmond JR, Morin L, Benjamin A.* Extremely preterm vaginal breech delivery en Caul. *Obstet Gynecol* 2002; 99(6): 1025-1030
241. *Roberts CL, Algert CS, Peat B, Henderson-Smart D.* Small fetal size: a risk factor for breech birth at term. *Int J Gynaecol Obstet* 1999; 67:1-8
242. *Robinson HP.* „Gestation sac“ volumes as determination by sonar in the first trimester of pregnancy. *Brit J Obstet Gynaecol* 1975; 82: 100-107
243. *Robson SC, Gallivan S, Walkinshaw SA, Vaughan J, Rodeck CH.* Ultrasonic estimation of fetal weight: use of targeted formulas in small for gestational age fetuses. *Obstet Gynecol* 1993; 82: 359-364
244. *Roemer VM, Bühler K, Kieback DG.* Gestationszeit und Geburtsgewicht. 1.Mitteilung: Intrauterine Wachstumskurven. *Z Geburtsh Perinatol* 1990; 194: 241-253
245. *Rossi L, Branca F, Cianfarani S.* Collagen cross-links and early postnatal growth in newborns with intrauterine growth retardation. *Metabolism* 2000; 49(11): 1467-1472
246. *Rouse DJ, Owen J.* Sonography, suspected macrosomia, and prophylactic cesarean: A limited partnership. *Clin Obstet Gynecol* 2000; 43: 326-334
247. *Rutherford SE, Smith CV, Phelan JP, Kawakami K, Ahn MO.* Four-quadrant assessment of amniotic fluid volume. *J Reprod Med* 1987; 32: 587-589
248. *Rutherford SE, Phelan JP, Smith CV, Jacobs N.* The four-quadrant assessment of amniotic fluid volume: An adjunct to antepartum fetal heart rate testing. *Obstet Gynecol* 1987; 70: 353-356
249. *Sachs BP, Kobelin C, Castro MA, Frigoletto F.* The risks of lowering the cesarean-delivery rate. *N Engl J Med* 1999; 340: 54-57

250. *Sadenwasser W, Adomßent S.* Influence of the mother's age on the weight of the newborn and the duration of pregnancy. *Zentralbl Gynakol* 1986; 108: 36-43
251. *Sanderson DA, Wilcox MA, Johnson IR.* Relative macrosomia identified by the individualised birthweight ratio (IBR). A better method of identifying the at risk fetus. *Acta Obstet Gynecol Scand* 1994;73: 246-249
252. *Sanderson DA, Wilcox MA, Johnson IR.* The individualised birthweight ratio: A new method of identifying intrauterine growth retardation. *Br J Obstet Gynaecol* 1994; 101: 310-314
253. *Scheer K, Nubar J.* Variation of fetal presentation with gestational age. *Am J Obstet Gynecol* 1976; 125(2): 269-270
254. *Schiff E, Ben-Baruch G, Kushnir O, Mashiach S.* Standardized measurement of amniotic fluid volume by correlation of sonography with dye dilution technique. *Obstet Gynecol* 1990; 76: 44-46
255. *Schmidt W, Leucht W, Boos R, Tariverdian S, Rabe D, Walter Ch, Heberling D.* Sonographische Diagnostik schwerer fetaler Fehlbildungen. *Geburtsh u. Frauenheilk* 1985; 45: 511-524
256. *Schneider H.* Die Wunschsectio – Eine gleichwertige Alternative zur Spontangeburt? *Gynäkol Geburtshilfliche Rundsch* 2002; 42: 4-11
257. *Schneider KTM.* Sectio nach Wunsch – Muss man hinsichtlich der Morbiditätsrisiken umdenken? *Gynäkol Geburtshilfliche Rundsch* 2002; 42: 12-14
258. *Schucker JL, Mercer BM, Audibert F, Lewis RL, Friedman SA, Sibai BM.* Serial amniotic fluid index in severe preeclampsia: A poor predictor of adverse outcome. *Am J Obstet Gynecol* 1996; 175: 1018-1023
259. *Seeds JW.* Impaired fetal growth: definition and clinical diagnosis. *Obstet Gynecol* 1984; 64: 303-310
260. *Seeds JW, Peng T.* Impaired growth and risk of fetal death: is the tenth percentile the appropriate standard? *Am J Obstet Gynecol* 1998; 178: 658-667

261. *Sepulveda W, Flack NJ, Fisk NM.* Direct volume measurement at midtrimester amnioinfusion in relation to ultrasonographic indexes of amniotic fluid volume. *Am J Obstet Gynecol* 1994; 170: 1160-1163
262. *Shepard MJ, Richards VA, Berkowitz RL, Warsof SL, Hobbins JC.* An evaluation of two equations for predicting fetal weight by ultrasound. *Am. J. Obstet. Gynecol.* 1982; 142: 47-54
263. *Simchen MJ, Beiner ME, Strauss-Liviathan N, Dulitzky M, Kuint J, Mashiach S, Schiff E.* Neonatal outcome in growth-restricted versus appropriately grown preterm infants. *Am J Perinatol* 2000; 17: 187-192
264. *Sinha R, Carlton M.* The volume and composition of amniotic fluid in early pregnancy. *J Obstet Gynaecol Br Commonw* 1970; 77: 211-214
265. *Smith CV, Plambeck RD, Rayburn WF, Albaugh KJ.* Relation of mild idiopathic polyhydramnios to perinatal outcome. *Obstet Gynecol* 1992; 79: 387-389
266. *Soothill PW, Ajayi RA, Campbell S, Nicolaidis KH.* Prediction of morbidity and normally grown fetuses by fetal heart rate variability, biophysical profile score and umbilical artery Doppler studies. *Br J Obstet Gynaecol* 1993; 100: 742-745
267. *Spinillo A, Capuzzo E, Egbe TO, Fazzi E, Colonna L, Nicola S.* Pregnancies complicated by idiopathic intrauterine growth retardation. Severity of growth failure, neonatal morbidity and two-year infant neurodevelopmental outcome. *J Reprod Med* 1995; 40: 209-215
268. *Strong TH Jr, Hetzler G, Paul RH.* Amniotic fluid volume increase after amnioinfusion of a fixed volume. *Am J Obstet Gynecol* 1990; 162: 746-748
269. *Sudik R.* Results of an ultrasonic screening programme in the detection of intrauterine growth retardation. *Z Geburtshilfe Perinatol* 1982; 186: 119-124
270. *Tadmor OP, Rabinowitz R, Alon L, Mostoslavsky V, Aboulafia Y, Diamant YZ.* Can breech presentation at birth be predicted from ultrasound examinations during the second or third trimesters? *Int J Gynecol Obstet* 1994; 46: 11-14



271. *Van HJ, Dekker G, Jeffries B.* Gestational diabetes versus obesity as risk factors for pregnancy-induced hypertensive disorders and fetal macrosomia. *Aust N Z J Obstet Gynaecol* 2002; 42: 29-34
272. *Van Otterlo LC, Wladimirnoff JW, Wallenburg HC.* Relationship between fetal urine production and amniotic fluid volume in normal pregnancy and pregnancy complicated by diabetes. *Br J Obstet Gynaecol* 1977; 84: 205-209
273. *Van Roosmalen J.* Elective caesarean section on request. Unnecessary caesarean sections should be avoided. *BMJ* 1999; 318: 121
274. *Voigt M, Eggers H.* Newborn-percentiles for the GDR – 1985. Birth weight and length percentiles. *Zentralbl Gynakol* 1988; 110: 927-943
275. *Voigt M, Eggers H, Jährig K, Grauel EL, Zwahr C, Plesse R.* Newborn-percentiles for the GDR – 1985. Correlations between age, parity, body weight and height of the mother and weight of the newborn. *Zentralbl Gynakol* 1989; 111: 337-349
276. *Voigt HJ, Grünsteidel W, Faschingbauer C.* Oligo-Anhydramnie. *Gynäkologische Praxis* 1990; 14: 11-22
277. *Voigt M, Schneider KTM, Jährig K.* Analyse des Geburtenguts des Jahrgangs 1992 der Bundesrepublik Deutschland. Teil 1: Neue Perzentilwerte für die Körpermaße von Neugeborenen. *Geburtshilfe Frauenheilkd* 1996; 56: 550-558
278. *Vorherr H.* Factors influencing fetal growth. *Am J Obstet Gynecol* 1982; 142: 577-588
279. *Wagner G, Fuchs F.* The volume of amniotic fluid in the first half of human pregnancy. *J Obstet Gynaecol Br Commonw* 1962; 69: 131-136
280. *Walkinshaw S, Cameron H, MacPhail S, Robson S.* The prediction of fetal compromise and acidosis by biophysical profile scoring in the small for gestational age fetus. *J Perinat Med* 1992; 20: 227-232
281. *Wallenburg HCS, Wladimirnoff JW.* The amniotic fluid. II. Poly-hydramnios and oligohydramnios. *J Perinat. Med.* 1977; 5: 233-243
282. *Warsof SL, Gohari B, Berkowitz RL, Hobbins JC.* The estimation of fetal weight by computer assisted analysis. *Am J Obstet Gynecol* 1977; 128: 881-892

283. *Webb AL, Sturgiss S, Warwicker P, Robson SC, Goodship JA, Wolsten-holme J.* Maternal uniparental disomy for chromosome 2 in association with confined placental mosaicism for trisomy 2 and severe intrauterine growth retardation. *Prenatal Diagnosis* 1996; 16: 958-962
284. *Weiner Z, Ben-Shlomo I, Beck-Fruchter R, Goldberg Y, Shalev E.* Clinical and ultrasonographic weight estimation in large for gestational age fetus. *Eur J Obstet Gynecol Reprod Biol* 2002; 105: 20-24
285. *Weiner Z, Peer E, Zimmer EZ.* Fetal testing in growth restriction. *Clin Obstet Gynecol* 1997; 40: 804-813
286. *Welsch H.* Mütterliche Sectio-Sterblichkeit – eine kritische Bilanz. *Arch Gynecol Obstet* 1995; 257: 206-215
287. *Wennergren M, Wennergren G, Vilbergsson G.* Obstetric characteristics and neonatal performance in a four-year small for gestational age population. *Obstet Gynecol* 1988; 72: 615-620
288. *Westgren M, Edvall H, Nordström L, Svalenius E, Ranstam J.* Spontaneous cephalic version of breech presentation in the last trimester. *Br J Obstet Gynecol* 1985; 92: 19-22
289. *Whitlow BJ, Lazanakis MS, Economides DL.* The sonographic identification of fetal gender from 11 to 14 weeks of gestation. *Ultrasound Obstet Gynecol* 1999; 13: 301-304
290. *Wieacker P, Wilhelm C, Fürste H, Schillinger H.* Polyhydramnie bei kongenitaler myotonischer Dystrophie. *Zentralblatt für Geburtshilfe und Perinatologie* 1988; 192: 36-37
291. *Wilcox MA, Newton CS, Johnson IR.* Paternal influences on birthweight. *Acta Obstet Gynecol Scand* 1995; 74: 15-18
292. *Wilcox MA, Johnson IR, Maynard PV, Smith SJ, Chilvers CE.* The individualised birth-weight ratio: A more logical outcome measure of pregnancy than birthweight alone. *Br J Obstet Gynaecol* 1993; 100: 342-347

293. *Wilhelm C, Wieacker P, Danneger K, Schillinger H.* Semiquantitative Fruchtwassermengenbestimmung bei normalen Schwangerschaften. *Geburtsh. u. Frauenheilk.* 1991b; 51: 341-344
294. *Wolf H, Schaap AH, Bruinse HW, Smolders-de Haas H, van Ertbruggen I, Treffers PE.* Vaginal delivery compared with caesarean section in early preterm breech delivery: a comparison of long term outcome. *Br J Obstet Gynaecol* 1999; 106: 486-491
295. *Wolf G, Wenzel U, Stahl RA, Huneke B.* Hypertensive Erkrankungen in der Schwangerschaft. *Med Klin* 2001; 96: 78-86
296. *Wollmann HA.* Intrauterine growth restriction: Definition and etiology. *Horm Res* 1998; 49(suppl 2): 1-6
297. *Woo JS, Wan MC.* An evaluation of fetal weight prediction using a simple equation containing the fetal femur length. *J Ultraouns Med* 1986; 5: 453-457
298. *Woodall SM, Johnston BM, Breier BH, Gluckman PD.* Chronic maternal undernutrition in the rat leads to delayed postnatal growth and elevated blood pressure of offspring. *Pediatr Res* 1996; 40(13): 438-443
299. *Wright RC.* Reduction of perinatal mortality and morbidity in breech delivery through routine use of cesarean section. *Obstet Gynecol* 1959; 14: 758-763
300. *Xiong X, Demianczuk NN, Buekens P, Saunders LD.* Association of preeclampsia with high birth weight for gestational age. *Am J Obstet Gynecol* 2000; 183: 148-155
301. *Yudkin PL, Aboualfa M, Eyre JA, Redman CWG, Wilkinson AR.* New birth weight and head circumference centiles for gestational ages 24 to 42 weeks. *Early Hum Dev* 1987; 15: 45-52
302. *Zamah NM, Gillieson MS, Walters JH, Hall PF.* Sonographic detection of polyhydramnios: A five-year-experience. *Am J Obstet Gynecol* 1982; 143: 523-527
303. *Zander J, Holzmann K, Selbmann HK.* Data from the Bavarian West Germany inquiry on the problem of the cesarean section frequency rate. *Geburtshilfe und Frauenheilkunde* 1989; 49: 328-336

304. *Zeitlin J, Ancel PY, Saurel-Cubizolles MJ, Papiernik E.* The relationship between intrauterine growth restriction and preterm delivery: an empirical approach using data from a European case-control study. *BJOG* 2000; 107: 750-758
305. *Zhang L.* The effects of essential fatty acids preparation in the treatment of intrauterine growth retardation. *Am J Perinat* 1997; 14: 535-537
306. *Zidar N, Cavic MA, Kenda RB, Koselj M, Ferluga D.* Effect of intrauterine growth retardation on the clinical course and prognosis of IgA Glomerulonephritis in children. *Nephron* 1998; 79: 28-32