

8. Literaturverzeichnis

- Acharya, R., Fry, E., Stuart, D., Fox, G., Rowlands, D., and Brown, F. (1989). The three-dimensional structure of foot-and-mouth disease virus at 2.9 Å resolution. *Nature* **337**, 709-716.
- Ali, M., Lemoine, N. R., and Ring, C. J. (1994). The use of DNA viruses as vectors for gene therapy. *Gene Ther.* **1**, 367-384.
- Andersson, B., Tomko, R. P., Edwards, K., Mirza, M., Darban, H., Öncü, D., Sonnhammer, E., Sollerbrant, K. and Philipson, L. (2000). Putative regulatory domains in the human and mouse CVADR genes. *Gene Function & Disease* **1**, 4-8
- Arnberg, N., Edlund, K., Kidd, A. H., and Wadell, G. (2000). Adenovirus type 37 uses sialic acid as a cellular receptor. *J.Viro.* **74**, 42-48.
- Arnold, B. L., Itakura, K., and Rossi, J. J. (1992). PCR-based quantitation of low levels of HIV-1 DNA by using an external standard. *Genet.Anal.Tech.Appl.* **9**, 113-116.
- Asaoka, K., Tada, M., Sawamura, Y., Ikeda, J., and Abe, H. (2000). Dependence of efficient adenoviral gene delivery in malignant glioma cells on the expression levels of the Coxsackievirus and adenovirus receptor. *J.Neurosurg.* **92**, 1002-1008.
- Bai, M., Campisi, L., and Freimuth, P. (1994). Vitronectin receptor antibodies inhibit infection of HeLa and A549 cells by adenovirus type 12 but not by adenovirus type 2. *J.Viro.* **68**, 5925-5932.
- Bailey, A. and Mautner, V. (1994). Phylogenetic relationships among adenovirus serotypes. *Virology* **205**, 438-452.
- Balague, C., Zhou, J., Dai, Y., Alemany, R., Josephs, S. F., Andreason, G., Hariharan, M., Sethi, E., Prokopenko, E., Jan, H. Y., Lou, Y. C., Hubert-Leslie, D., Ruiz, L., and Zhang, W. W. (2000). Sustained high-level expression of full-length human factor VIII and restoration of clotting activity in hemophilic mice using a minimal adenovirus vector. *Blood* **95**, 820-828.
- Barillari, G., Gendelman, R., Gallo, R. C., and Ensoli, B. (1993). The Tat protein of human immunodeficiency virus type 1, a growth factor for AIDS Kaposi sarcoma and cytokine-activated vascular cells, induces adhesion of the same cell types by using integrin receptors recognizing the RGD amino acid sequence. *Proc.Natl.Acad.Sci.U.S.A.* **90**, 7941-7945.
- Bartholomew, R. M., Esser, A. F., and Muller-Eberhard, H. J. (1978). Lysis of oncornaviruses by human serum. Isolation of the viral complement (C1) receptor and identification as p15E. *J.Exp.Med.* **147**, 844-853.
- Basak, A. K., Gouet, P., Grimes, J., Roy, P., and Stuart, D. (1996). Crystal structure of the top domain of African horse sickness virus VP7: comparisons with bluetongue virus VP7. *J.Viro.* **70**, 3797-3806.
- Becker-Andre, M. and Hahlbrock, K. (1989). Absolute mRNA quantification using the polymerase chain reaction (PCR). A novel approach by a PCR aided transcript titration assay (PATTY). *Nucleic.Acids.Res.* **17**, 9437-9446.

- Benhamou, P. Y., Moriscot, C., Richard, M. J., Beatrix, O., Badet, L., Pattou, F., Kerr-Conte, J., Chroboczek, J., Lemarchand, P., and Halimi, S. (1998). Adenovirus-mediated catalase gene transfer reduces oxidant stress in human, porcine and rat pancreatic islets. *Diabetologia* **41**, 1093-1100.
- Bergelson, J. M. (1999). Receptors mediating adenovirus attachment and internalization. *Biochem.Pharmacol.* **57**, 975-979.
- Bergelson, J. M., Cunningham, J. A., Drogue, G., Kurt-Jones, E. A., Krithivas, A., Hong, J. S., Horwitz, M. S., Crowell, R. L., and Finberg, R. W. (1997). Isolation of a common receptor for Coxsackie B viruses and adenoviruses 2 and 5. *Science* **275**, 1320-1323.
- Bergelson, J. M., Krithivas, A., Celi, L., Drogue, G., Horwitz, M. S., Wickham, T., Crowell, R. L., and Finberg, R. W. (1998). The murine CAR homolog is a receptor for coxsackie B viruses and adenoviruses. *J.Viro.* **72**, 415-419.
- Bergelson, J. M., Mohanty, J. G., Crowell, R. L., St John, N. F., Lublin, D. M., and Finberg, R. W. (1995). Coxsackievirus B3 adapted to growth in RD cells binds to decay-accelerating factor (CD55). *J.Viro.* **69**, 1903-1906.
- Bett, A. J., Prevec, L., and Graham, F. L. (1993). Packaging capacity and stability of human adenovirus type 5 vectors. *J.Viro.* **67**, 5911-5921.
- Bewley, M. C., Springer, K., Zhang, Y. B., Freimuth, P., and Flanagan, J. M. (1999). Structural Analysis of the Mechanism of Adenovirus Binding to Its Human Cellular Receptor, CAR. *Science* **286**, 1579-1583.
- Bouaboula, M., Legoux, P., Pessegue, B., Delpech, B., Dumont, X., Piechaczyk, M., Casellas, P., and Shire, D. (1992). Standardization of mRNA titration using a polymerase chain reaction method involving co-amplification with a multispecific internal control. *J.Biol.Chem.* **267**, 21830-21838.
- Bowles, K. R., Gibson, J., Wu, J., Shaffer, L. G., Towbin, J. A., and Bowles, N. E. (1999). Genomic organization and chromosomal localization of the human Coxsackievirus B-adenovirus receptor gene. *Hum.Genet.* **105**, 354-359.
- Brenner, M. (1999). Gene transfer by adenovectors. *Blood* **94**, 3965-3967.
- Brillanti, S., Garson, J. A., Tuke, P. W., Ring, C., Briggs, M., Masci, C., Miglioli, M., Barbara, L., and Tedder, R. S. (1991). Effect of alpha-interferon therapy on hepatitis C viraemia in community-acquired chronic non-A, non-B hepatitis: a quantitative polymerase chain reaction study. *J.Med.Viro.* **34**, 136-141.
- Brody, S. L. and Crystal, R. G. (1994). Adenovirus-mediated in vivo gene transfer. *Ann.N.Y.Acad.Sci.* **716:90-101; discussion 101-3.**, 90-101.
- Byrne, B. C., Li, J. J., Sninsky, J., and Poiesz, B. J. (1988). Detection of HIV-1 RNA sequences by in vitro DNA amplification. *Nucleic.Acids.Res.* **16**, 4165.
- Cao, B., Pruchnic, R., Ikezawa, M., Xiao, X., Li, J., Wickham, T. J., Kovacs, I., Rudert, W. A., and Huard, J. (2001). The role of receptors in the maturation-dependent adenoviral transduction of myofibers. *Gene Ther.* **8**, 627-637.

- Capecci, M. R. (1980). High efficiency transformation by direct microinjection of DNA into cultured mammalian cells. *Cell* **22**, 479-488.
- Carson, S. D., Chapman, N. N., and Tracy, S. M. (1997). Purification of the putative coxsackievirus B receptor from HeLa cells. *Biochem.Biophys.Res.Commun.* **233**, 325-328.
- Carson, S. D., Hobbs, J. T., Tracy, S. M., and Chapman, N. M. (1999). Expression of the coxsackievirus and adenovirus receptor in cultured human umbilical vein endothelial cells: regulation in response to cell density. *J.Virol.* **73**, 7077-7079.
- Carson, S. D. (2000). Limited proteolysis of the coxsackievirus and adenovirus receptor (CAR) on HeLa cells exposed to trypsin. *FEBS Lett.* **484**, 149-152.
- Carson, S. D. (2001). Receptor for the group B coxsackieviruses and adenoviruses: CAR. *Rev.Med.Viro.* **11**, 219-226.
- Chaisomchit, S., Tyrrell, D. L., and Chang, L. J. (1997). Development of replicative and nonreplicative hepatitis B virus vectors. *Gene Ther.* **4**, 1330-1340.
- Chauvet, A. E., Kesava, P. P., Goh, C. S., and Badie, B. (1998). Selective intraarterial gene delivery into a canine meningioma. *J.Neurosurg.* **88**, 870-873.
- Chiu, C. Y., Mathias, P., Nemerow, G. R., and Stewart, P. L. (1999). Structure of adenovirus complexed with its internalization receptor, alphavbeta5 integrin. *J.Viro.* **73**, 6759-6768.
- Cho, W. K., Ebihara, S., Nalbantoglu, J., Gilbert, R., Massie, B., Holland, P., Karpati, G., and Petrof, B. J. (2000). Modulation of Starling forces and muscle fiber maturity permits adenovirus-mediated gene transfer to adult dystrophic (mdx) mice by the intravascular route. *Hum.Gene Ther.* **11**, 701-714.
- Chou, P. Y. and Fasman, G. D. (1974). Conformational parameters for amino acids in helical, beta-sheet, and random coil regions calculated from proteins. *Biochemistry* **13**, 211-222.
- Chretien, I., Marcuz, A., Courtet, M., Katevuo, K., Vainio, O., Heath, J. K., White, S. J., and Du Pasquier, L. (1998). CTX, a Xenopus thymocyte receptor, defines a molecular family conserved throughout vertebrates. *Eur.J.Immunol.* **28**, 4094-4104.
- Chroboczek, J., Ruigrok, R. W., and Cusack, S. (1995). Adenovirus fiber. *Curr.Top.Microbiol.Immunol.* **199**, 163-200.
- Clementi, M., Menzo, S., Bagnarelli, P., Manzin, A., Valenza, A., and Varaldo, P. E. (1993). Quantitative PCR and RT-PCR in virology. *PCR.Methods Appl.* **2**, 191-196.
- Cohen, C. J., Gaetz, J., Ohman, T., and Bergelson, J. M. (2001). Multiple regions within the Coxsackievirus and Adenovirus receptor cytoplasmic domain are required for basolateral sorting. *J.Biol.Chem.* ..
- Cornetta, K., Moen, R. C., Culver, K., Morgan, R. A., McLachlin, J. R., Sturm, S., Selegue, J., London, W., Blaese, R. M., and Anderson, W. F. (1990). Amphotropic murine leukemia retrovirus is not an acute pathogen for primates. *Hum.Gene Ther.* **1**, 15-30.
- Cotten, M. and Wagner, E. (1993). Non-viral approaches to gene therapy. *Curr.Opin.Biotechnol.* **4**, 705-710.

- Cotten, M. and Weber, J. M. (1995). The adenovirus protease is required for virus entry into host cells. *Virology* **213**, 494-502.
- Cottrez, F., Auriault, C., Capron, A., and Groux, H. (1994). Quantitative PCR: validation of the use of a multispecific internal control. *Nucleic Acids Res.* **22**, 2712-2713.
- Cross, N. C. (1995). Quantitative PCR techniques and applications. *Br.J.Haematol.* **89**, 693-697.
- Croyle, M. A., Walter, E., Janich, S., Roessler, B. J., and Amidon, G. L. (1998). Role of integrin expression in adenovirus-mediated gene delivery to the intestinal epithelium. *Hum.Gene Ther.* **9**, 561-573.
- Crystal, R. G., McElvaney, N. G., Rosenfeld, M. A., Chu, C. S., Mastrangeli, A., Hay, J. G., Brody, S. L., Jaffe, H. A., Eissa, N. T., and Danel, C. (1994). Administration of an adenovirus containing the human CFTR cDNA to the respiratory tract of individuals with cystic fibrosis. *Nat.Genet.* **8**, 42-51.
- Cunningham, B. A., Hemperly, J. J., Murray, B. A., Prediger, E. A., Brackenbury, R., and Edelman, G. M. (1987). Neural cell adhesion molecule: structure, immunoglobulin-like domains, cell surface modulation, and alternative RNA splicing. *Science* **236**, 799-806.
- Curtis, S., Wilkinson, G. W., and Westmoreland, D. (1998). An outbreak of epidemic keratoconjunctivitis caused by adenovirus type 37. *J.Med.Microbiol.* **47**, 91-94.
- Dai, Y., Schwarz, E. M., Gu, D., Zhang, W. W., Sarvetnick, N., and Verma, I. M. (1995). Cellular and humoral immune responses to adenoviral vectors containing factor IX gene: tolerization of factor IX and vector antigens allows for long-term expression. *Proc.Natl.Acad.Sci.U.S.A.* **92**, 1401-1405.
- Damke, H., Baba, T., Warnock, D. E., and Schmid, S. L. (1994). Induction of mutant dynamin specifically blocks endocytic coated vesicle formation. *J.Cell Biol.* **127**, 915-934.
- Davison, E., Kirby, I., Elliott, T., and Santis, G. (1999). The human HLA-A*0201 allele, expressed in hamster cells, is not a high-affinity receptor for adenovirus type 5 fiber. *J.Viro.* **73**, 4513-4517.
- de Kant, E., Rochlitz, C. F., and Herrmann, R. (1994). Gene expression analysis by a competitive and differential PCR with antisense competitors. *Biotechniques* **17**, 934-942.
- Defer, C., Belin, M. T., Caillet-Boudin, M. L., and Boulanger, P. (1990). Human adenovirus-host cell interactions: comparative study with members of subgroups B and C. *J.Viro.* **64**, 3661-3673.
- DeMatteo, R. P., McClane, S. J., Fisher, K., Yeh, H., Chu, G., Burke, C., and Raper, S. E. (1997). Engineering tissue-specific expression of a recombinant adenovirus: selective transgene transcription in the pancreas using the amylase promoter. *J.Surg.Res.* **72**, 155-161.
- Di Guilmi, A. M., Barge, A., Kitts, P., Gout, E., and Chroboczek, J. (1995). Human adenovirus serotype 3 (Ad3) and the Ad3 fiber protein bind to a 130-kDa membrane protein on HeLa cells. *Virus Res.* **38**, 71-81.

- Diamond, M. S. and Springer, T. A. (1994). The dynamic regulation of integrin adhesiveness. *Curr.Biol.* **4**, 506-517.
- Diviacco, S., Norio, P., Zentilin, L., Menzo, S., Clementi, M., Biamonti, G., Riva, S., Falaschi, A., and Giacca, M. (1992). A novel procedure for quantitative polymerase chain reaction by coamplification of competitive templates. *Gene* **122**, 313-320.
- Dmitriev, I., Kashentseva, E., Rogers, B. E., Krasnykh, V., and Curiel, D. T. (2000). Ectodomain of coxsackievirus and adenovirus receptor genetically fused to epidermal growth factor mediates adenovirus targeting to epidermal growth factor receptor-positive cells. *J.Virol.* **74**, 6875-6884.
- Dmitriev, I., Krasnykh, V., Miller, C. R., Wang, M., Kashentseva, E., Mikheeva, G., Belousova, N., and Curiel, D. T. (1998). An adenovirus vector with genetically modified fibers demonstrates expanded tropism via utilization of a coxsackievirus and adenovirus receptor-independent cell entry mechanism. *J.Virol.* **72**, 9706-9713.
- Dostal, D. E., Rothblum, K. N., and Baker, K. M. (1994). An improved method for absolute quantification of mRNA using multiplex polymerase chain reaction: determination of renin and angiotensinogen mRNA levels in various tissues. *Anal.Biochem.* **223**, 239-250.
- Edgell, C. J., McDonald, C. C., and Graham, J. B. (1983). Permanent cell line expressing human factor VIII-related antigen established by hybridization. *Proc.Natl.Acad.Sci.U.S.A* **80**, 3734-3737.
- Eiz, B. and Pring-Akerblom, P. (1997). Molecular characterization of the type-specific gamma-determinant located on the adenovirus fiber. *J.Virol.* **71**, 6576-6581.
- Eizema, K., Fechner, H., Bezstarosti, K., Schneider-Rasp, S., van der Laarse, A., Wang, H., Schultheiss, H. P., Poller, W. C., and Lamers, J. M. (2000). Adenovirus-based phospholamban antisense expression as a novel approach to improve cardiac contractile dysfunction: comparison of a constitutive viral versus an endothelin-1-responsive cardiac promoter. *Circulation* **101**, 2193-2199.
- Elder, P. K., French, C. L., Subramaniam, M., Schmidt, L. J., and Getz, M. J. (1988). Evidence that the functional beta-actin gene is single copy in most mice and is associated with 5' sequences capable of conferring serum- and cycloheximide-dependent regulation. *Mol.Cell Biol.* **8**, 480-485.
- Engelhardt, J. F., Ye, X., Doranz, B., and Wilson, J. M. (1994). Ablation of E2A in recombinant adenoviruses improves transgene persistence and decreases inflammatory response in mouse liver. *Proc.Natl.Acad.Sci.U.S.A.* **91**, 6196-6200.
- Fallaux, F. J., Bout, A., van, der Velde, I., van den Wollenberg, D. J., Hehir, K. M., Keegan, J., Auger, C., Cramer, S. J., van Ormondt, H., van der Eb, A. J., Valerio, D., and Hoeben, R. C. (1998). New helper cells and matched early region 1-deleted adenovirus vectors prevent generation of replication-competent adenoviruses. *Hum.Gene Ther.* **9**, 1909-1917.
- Fechner, H., Haack, A., Wang, H., Wang, X., Eizema, K., Pauschinger, M., Schoemaker, R., Veghel, R., Houtsmuller, A., Schultheiss, H. P., Lamers, J., and Poller, W. (1999). Expression of Coxsackie adenovirus receptor and alphav-integrin does not correlate with adenovector targeting in vivo indicating anatomical vector barriers. *Gene Ther.* **6**, 1520-1535.

- Fechner, H., Wang, X., Wang, H., Jansen, A., Pauschinger, M., Scherubl, H., Bergelson, J. M., Schultheiss, H. P., and Poller, W. (2000). Trans-complementation of vector replication versus Coxsackie-adenovirus-receptor overexpression to improve transgene expression in poorly permissive cancer cells. *Gene Ther.* **7**, 1954-1968.
- Felgner, P. L., Gadek, T. R., Holm, M., Roman, R., Chan, H. W., Wenz, M., Northrop, J. P., Ringold, G. M., and Danielsen, M. (1987). Lipofection: a highly efficient, lipid-mediated DNA-transfection procedure. *Proc.Natl.Acad.Sci.U.S.A.* **84**, 7413-7417.
- Fender, P., Kidd, A. H., Brebant, R., Oberg, M., Drouet, E., and Chroboczek, J. (1995). Antigenic sites on the receptor-binding domain of human adenovirus type 2 fiber. *Virology* **214**, 110-117.
- Ferre F., Marchese A., Pezzoli P., Griffin S., Buxton E., and Boyer V. (1994). Quantitative PCR: an overview. In 'The polymerase chain reaction'. (Mullis K.B., Ferre F., and Gibbs R.A. Eds.) pp. 67-88. (Birkhäuser Boston: Basel, Berlin.)
- Ferre, F. (1992). Quantitative or semi-quantitative PCR: reality versus myth. *PCR.Methods Appl.* **2**, 1-9.
- Foley, K. P., Leonard, M. W., and Engel, J. D. (1993). Quantitation of RNA using the polymerase chain reaction. *Trends.Genet.* **9**, 380-385.
- Fox, G., Parry, N. R., Barnett, P. V., McGinn, B., Rowlands, D. J., and Brown, F. (1989). The cell attachment site on foot-and-mouth disease virus includes the amino acid sequence RGD (arginine-glycine-aspartic acid). *J.Gen.Viro.* **70**, 625-637.
- Freimuth, P., Springer, K., Berard, C., Hainfeld, J., Bewley, M., and Flanagan, J. (1999). Coxsackievirus and adenovirus receptor amino-terminal immunoglobulin V-related domain binds adenovirus type 2 and fiber knob from adenovirus type 12 [In Process Citation]. *J.Viro.* **73**, 1392-1398.
- Galdiero, M., Whiteley, A., Bruun, B., Bell, S., Minson, T., and Browne, H. (1997). Site-directed and linker insertion mutagenesis of herpes simplex virus type 1 glycoprotein H. *J.Viro.* **71**, 2163-2170.
- Geller, A. I. and Breakefield, X. O. (1988). A defective HSV-1 vector expresses Escherichia coli beta-galactosidase in cultured peripheral neurons. *Science* **241**, 1667-1669.
- Gilliland, G., Perrin, S., Blanchard, K., and Bunn, H. F. (1990). Analysis of cytokine mRNA and DNA: detection and quantitation by competitive polymerase chain reaction. *Proc.Natl.Acad.Sci.U.S.A.* **87**, 2725-2729.
- Goodman, C. S. and Shatz, C. J. (1993). Developmental mechanisms that generate precise patterns of neuronal connectivity. *Cell* **72 Suppl:77-98.**, 77-98.
- Graham, F. L., Smiley, J., Russell, W. C., and Nairn, R. (1977). Characteristics of a human cell line transformed by DNA from human adenovirus type 5. *J.Gen.Viro.* **36**, 59-74.
- Grassi, G., Zentilin, L., Tafuro, S., Diviacco, S., Ventura, A., Falaschi, A., and Giacca, M. (1994). A rapid procedure for the quantitation of low abundance RNAs by competitive reverse transcription-polymerase chain reaction. *Nucleic Acids Res.* **22**, 4547-4549.

- Greber, U. F., Willetts, M., Webster, P., and Helenius, A. (1993). Stepwise dismantling of adenovirus 2 during entry into cells. *Cell* **75**, 477-486.
- Grodums E.L. and Dempster G. (1959). The age factor in experimental coxsackie B-3 infection. *Can.J.Microbiol.* **5**, 595-604.
- Grodums E.L. and Dempster G. (1961). Encephalitis in experimental coxsackie B-3 infection. *Can.J.Microbiol.* **7**, 175-184.
- Grotewiel, M. S., Beck, C. D., Wu, K. H., Zhu, X. R., and Davis, R. L. (1998). Integrin-mediated short-term memory in *Drosophila*. *Nature* **391**, 455-460.
- Haack, A., Poller, W., Schneider-Rasp, S., Thalheimer, P., Schmitt, C., Hanfland, P., Brackmann, H. H., and Schwaab, R. (1999). Highly sensitive and species-specific assay for quantification of human transgene expression levels. *Haemophilia*. **5**, 334-339.
- Haberhausen, G., Pinsl, J., Kuhn, C. C., and Markert-Hahn, C. (1998). Comparative study of different standardization concepts in quantitative competitive reverse transcription-PCR assays. *J.Clin.Microbiol.* **36**, 628-633.
- Hackett N.R. and Crystal R.G. (2000). Adenoviral vectors for gene therapy. In 'Gene therapy: therapeutic mechanisms and strategies'. (Templeton N.S. and Lasic D.D. Eds.) pp. 17-59. (Marcel Dekker, Inc.: New York, Basel.)
- Hajjar, R. J., Schmidt, U., Matsui, T., Guerrero, J. L., Lee, K. H., Gwathmey, J. K., Dec, G. W., Semigran, M. J., and Rosenzweig, A. (1998). Modulation of ventricular function through gene transfer in vivo. *Proc.Natl.Acad.Sci.U.S.A.* **95**, 5251-5256.
- Halank, M., Wiemer, M., Tschope, C., Poller, W., Schwimmbeck, P., Horstkotte, D., and Hoffken, G. (1998). [Precapillary pulmonary hypertension of uncertain etiology]. *Dtsch.Med.Wochenschr.* **123**, 861-865.
- Hamilton, T. E., McClane, S. J., Baldwin, S., Burke, C., Patel, H., Rombeau, J. L., and Raper, S. E. (1997). Efficient adenoviral-mediated murine neonatal small intestinal gene transfer is dependent on alpha(v) integrin expression. *J.Pediatr.Surg.* **32**, 1695-1703.
- Hautala, T., Grunst, T., Fabrega, A., Freimuth, P., and Welsh, M. J. (1998). An interaction between penton base and alpha v integrins plays a minimal role in adenovirus-mediated gene transfer to hepatocytes in vitro and in vivo. *Gene Ther.* **5**, 1259-1264.
- Hay, J. G., McElvaney, N. G., Herena, J., and Crystal, R. G. (1995). Modification of nasal epithelial potential differences of individuals with cystic fibrosis consequent to local administration of a normal CFTR cDNA adenovirus gene transfer vector. *Hum.Gene Ther.* **6**, 1487-1496.
- Heath, J. K., White, S. J., Johnstone, C. N., Catimel, B., Simpson, R. J., Moritz, R. L., Tu, G. F., Ji, H., Whitehead, R. H., Groenen, L. C., Scott, A. M., Ritter, G., Cohen, L., Welt, S., Old, L. J., Nice, E. C., and Burgess, A. W. (1997). The human A33 antigen is a transmembrane glycoprotein and a novel member of the immunoglobulin superfamily. *Proc.Natl.Acad.Sci.U.S.A* **94**, 469-474.
- Hehir, K. M., Armentano, D., Cardoza, L. M., Choquette, T. L., Berthelette, P. B., White, G. A., Couture, L. A., Everton, M. B., Keegan, J., Martin, J. M., Pratt, D. A., Smith, M. P.,

- Smith, A. E., and Wadsworth, S. C. (1996). Molecular characterization of replication-competent variants of adenovirus vectors and genome modifications to prevent their occurrence. *J.Viro.* **70**, 8459-8467.
- Helenius, A., Kartenbeck, J., Simons, K., and Fries, E. (1980). On the entry of Semliki forest virus into BHK-21 cells. *J.Cell Biol.* **84**, 404-420.
- Hemmi, S., Geertsen, R., Mezzacasa, A., Peter, I., and Dummer, R. (1998). The presence of human coxsackievirus and adenovirus receptor is associated with efficient adenovirus-mediated transgene expression in human melanoma cell cultures. *Hum.Gene Ther.* **9**, 2363-2373.
- Henrard, D. R., Mehaffey, W. F., and Allain, J. P. (1992). A sensitive viral capture assay for detection of plasma viremia in HIV-infected individuals. *AIDS Res.Hum.Retroviruses* **8**, 47-52.
- Henry, L. J., Xia, D., Wilke, M. E., Deisenhofer, J., and Gerard, R. D. (1994). Characterization of the knob domain of the adenovirus type 5 fiber protein expressed in Escherichia coli. *J.Viro.* **68**, 5239-5246.
- Hewson, R. (2000). RNA viruses: emerging vectors for vaccination and gene therapy. *Mol.Med.Today* **6**, 28-35.
- Hidaka, C., Milano, E., Leopold, P. L., Bergelson, J. M., Hackett, N. R., Finberg, R. W., Wickham, T. J., Kovesdi, I., Roelvink, P., and Crystal, R. G. (1999a). CAR-dependent and CAR-independent pathways of adenovirus vector-mediated gene transfer and expression in human fibroblasts. *J.Clin.Invest.* **103**, 579-587.
- Hierholzer, J. C. (1973). Further subgrouping of the human adenoviruses by differential hemagglutination. *J.Infect.Dis.* **128**, 541-550.
- Higuchi, R., Dollinger, G., Walsh, P. S., and Griffith, R. (1992). Simultaneous amplification and detection of specific DNA sequences. *Biotechnology (N.Y.)* **10**, 413-417.
- Higuchi, R., Fockler, C., Dollinger, G., and Watson, R. (1993). Kinetic PCR analysis: real-time monitoring of DNA amplification reactions. *Biotechnology (N.Y.)* **11**, 1026-1030.
- Hirsch, E., Gullberg, D., Balzac, F., Altruda, F., Silengo, L., and Tarone, G. (1994). Alpha v integrin subunit is predominantly located in nervous tissue and skeletal muscle during mouse development. *Dev.Dyn.* **201**, 108-120.
- Holzinger, A., Trapnell, B. C., Weaver, T. E., Whitsett, J. A., and Iwamoto, H. S. (1995). Intraamniotic administration of an adenoviral vector for gene transfer to fetal sheep and mouse tissues. *Pediatr.Res.* **38**, 844-850.
- Honda, T., Saitoh, H., Masuko, M., Katagiri-Abe, T., Tominaga, K., Kozakai, I., Kobayashi, K., Kumanishi, T., Watanabe, Y. G., Odani, S., and Kuwano, R. (2000). The coxsackievirus-adenovirus receptor protein as a cell adhesion molecule in the developing mouse brain. *Brain Res.Mol.Brain Res.* **77**, 19-28.
- Hong, S. S., Karayan, L., Tournier, J., Curiel, D. T., and Boulanger, P. A. (1997). Adenovirus type 5 fiber knob binds to MHC class I alpha2 domain at the surface of human epithelial and B lymphoblastoid cells. *EMBO J.* **16**, 2294-2306.

- Horikoshi, T., Danenberg, K. D., Stadlbauer, T. H., Volkenandt, M., Shea, L. C., Aigner, K., Gustavsson, B., Leichman, L., Frosing, R., and Ray, M. (1992). Quantitation of thymidylate synthase, dihydrofolate reductase, and DT-diaphorase gene expression in human tumors using the polymerase chain reaction. *Cancer Res.* **52**, 108-116.
- Horwitz, M. S. (1996). In 'Fields Virology'. (Ed. K. D. H. P. Fields BN.) p. 2149. (Lippincott-Raven: Philadelphia, New York.)
- Hotta, Y., Masuko, M., Hanyu T., Abe, T. K., Honda T., Tominaga, K., Odani, S., and Kuwano, R. (1998). Immunohistochemical analysis of new adhesion molecule gmp46. *Neurosci.Res.* **22 (suppl.)**, 313.
- Howell, J. M., Lochmuller, H., O'Hara, A., Fletcher, S., Kakulas, B. A., Massie, B., Nalbantoglu, J., and Karpati, G. (1998). High-level dystrophin expression after adenovirus-mediated dystrophin minigene transfer to skeletal muscle of dystrophic dogs: prolongation of expression with immunosuppression. *Hum.Gene Ther.* **9**, 629-634.
- Huang, S., Kamata, T., Takada, Y., Ruggeri, Z. M., and Nemerow, G. R. (1996). Adenovirus interaction with distinct integrins mediates separate events in cell entry and gene delivery to hematopoietic cells. *J.Virol.* **70**, 4502-4508.
- Huang, S., Reddy, V., Dasgupta, N., and Nemerow, G. R. (1999). A single amino acid in the adenovirus type 37 fiber confers binding to human conjunctival cells. *J.Virol.* **73**, 2798-2802.
- Hutchin, M. E., Pickles, R. J., and Yarbrough, W. G. (2000). Efficiency of adenovirus-mediated gene transfer to oropharyngeal epithelial cells correlates with cellular differentiation and human coxsackie and adenovirus receptor expression [In Process Citation]. *Hum.Gene Ther.* **11**, 2365-2375.
- Hynes, R. O. (1992). Integrins: versatility, modulation, and signaling in cell adhesion. *Cell* **69**, 11-25.
- Hynes, R. O. and Lander, A. D. (1992). Contact and adhesive specificities in the associations, migrations, and targeting of cells and axons. *Cell* **68**, 303-322.
- Ito, M., Kodama, M., Masuko, M., Yamaura, M., Fuse, K., Uesugi, Y., Hirono, S., Okura, Y., Kato, K., Hotta, Y., Honda, T., Kuwano, R., and Aizawa, Y. (2000). Expression of coxsackievirus and adenovirus receptor in hearts of rats with experimental autoimmune myocarditis. *Circ.Res.* **86**, 275-280.
- Jackson, T., Sharma, A., Ghazaleh, R. A., Blakemore, W. E., Ellard, F. M., Simmons, D. L., Newman, J. W., Stuart, D. I., and King, A. M. (1997). Arginine-glycine-aspartic acid-specific binding by foot-and-mouth disease viruses to the purified integrin alpha(v)beta3 in vitro. *J.Virol.* **71**, 8357-8361.
- Jolly, D. (1994). Viral vector systems for gene therapy. *Cancer Gene Ther.* **1**, 51-64.
- Kaneko, S., Hallenbeck, P., Kotani, T., Nakabayashi, H., McGarrity, G., Tamaoki, T., Anderson, W. F., and Chiang, Y. L. (1995). Adenovirus-mediated gene therapy of hepatocellular carcinoma using cancer-specific gene expression. *Cancer Res.* **55**, 5283-5287.
- Kaner, R. J., Worgall, S., Leopold, P. L., Stolze, E., Milano, E., Hidaka, C., Ramalingam, R., Hackett, N. R., Singh, R., Bergelson, J., Finberg, R., Falck-Pedersen, E., and Crystal, R. G.

(1999). Modification of the Genetic Program of Human Alveolar Macrophages by Adenovirus Vectors In Vitro Is Feasible but Inefficient, Limited in Part by the Low Level of Expression of the Coxsackie/Adenovirus Receptor. *Am.J.Respir.Cell Mol.Biol.* **20**, 361-370.

Kaplan, J. M., St George, J. A., Pennington, S. E., Keyes, L. D., Johnson, R. P., Wadsworth, S. C., and Smith, A. E. (1996). Humoral and cellular immune responses of nonhuman primates to long-term repeated lung exposure to Ad2/CFTR-2. *Gene Ther.* **3**, 117-127.

Kasono, K., Blackwell, J. L., Douglas, J. T., Dmitriev, I., Strong, T. V., Reynolds, P., Kropf, D. A., Carroll, W. R., Peters, G. E., Bucy, R. P., Curiel, D. T., and Krasnykh, V. (1999). Selective gene delivery to head and neck cancer cells via an integrin targeted adenoviral vector [In Process Citation]. *Clin.Cancer Res.* **5**, 2571-2579.

Kass-Eisler, A., Falck-Pedersen, E., Elfenbein, D. H., Alvira, M., Buttrick, P. M., and Leinwand, L. A. (1994). The impact of developmental stage, route of administration and the immune system on adenovirus-mediated gene transfer. *Gene Ther.* **1**, 395-402.

Kaufman, J. F., Auffray, C., Korman, A. J., Shackelford, D. A., and Strominger, J. (1984). The class II molecules of the human and murine major histocompatibility complex. *Cell* **36**, 1-13.

Kellogg, D. E., Sninsky, J. J., and Kowk, S. (1990). Quantitation of HIV-1 proviral DNA relative to cellular DNA by the polymerase chain reaction. *Anal.Biochem.* **189**, 202-208.

Kirby, I., Davison, E., Beavil, A. J., Soh, C. P., Wickham, T. J., Roelvink, P. W., Kovacs, I., Sutton, B. J., and Santis, G. (2000). Identification of contact residues and definition of the CAR-binding site of adenovirus type 5 fiber protein. *J.Viro.* **74**, 2804-2813.

Knowles, M. R., Hohneker, K. W., Zhou, Z., Olsen, J. C., Noah, T. L., Hu, P. C., Leigh, M. W., Engelhardt, J. F., Edwards, L. J., and Jones, K. R. (1995). A controlled study of adenoviral-vector-mediated gene transfer in the nasal epithelium of patients with cystic fibrosis. *N Engl.J.Med.* **333**, 823-831.

Koi, H., Zhang, J., Makrigiannakis, A., Getsios, S., MacCalman, C. D., Kopf, G. S., Strauss, J. F., and Parry, S. (2001). Differential expression of the coxsackievirus and adenovirus receptor regulates adenovirus infection of the placenta. *Biol.Reprod.* **64**, 1001-1009.

Krasnykh, V. N., Mikheeva, G. V., Douglas, J. T., and Curiel, D. T. (1996). Generation of recombinant adenovirus vectors with modified fibers for altering viral tropism. *J.Viro.* **70**, 6839-6846.

Kunin C.M. (1962). Virus-tissue union and the pathogenesis of enterovirus infections. *J.Immunol.* **88**, 556-589.

Kunin C.M. and Halmagyi N.E. (1961). The relative abundance of viral receptors: an explanation of the differential susceptibility of suckling and adult mice to coxsackie B1 infection. *J.Clin.Invest.* **40**, -1055.

Kyte, J. and Doolittle, R. F. (1982). A simple method for displaying the hydropathic character of a protein. *J.Mol.Biol.* **157**, 105-132.

Larochelle, N., Lochmuller, H., Zhao, J., Jani, A., Hallauer, P., Hastings, K. E., Massie, B., Prescott, S., Petrof, B. J., Karpati, G., and Nalbantoglu, J. (1997). Efficient muscle-specific

transgene expression after adenovirus-mediated gene transfer in mice using a 1.35 kb muscle creatine kinase promoter/enhancer. *Gene Ther.* **4**, 465-472.

Ledley, F. D. (1994). Non-viral gene therapy. *Curr.Opin.Biotechnol.* **5**, 626-636.

Lee, B., Dennis, J. A., Healy, P. J., Mull, B., Pastore, L., Yu, H., Aguilar-Cordova, E., O'Brien, W., Reeds, P., and Beaudet, A. L. (1999). Hepatocyte gene therapy in a large animal: a neonatal bovine model of citrullinemia. *Proc.Natl.Acad.Sci.U.S.A.* **96**, 3981-3986.

Leon, R. P., Hedlund, T., Meech, S. J., Li, S., Schaack, J., Hunger, S. P., Duke, R. C., and DeGregori, J. (1998). Adenoviral-mediated gene transfer in lymphocytes. *Proc.Natl.Acad.Sci.U.S.A.* **95**, 13159-13164.

Li, D., Duan, L., Freimuth, P., and O'Malley, B. W. Jr (1999). Variability of adenovirus receptor density influences gene transfer efficiency and therapeutic response in head and neck cancer [In Process Citation]. *Clin.Cancer Res.* **5**, 4175-4181.

Li, E., Stupack, D., Bokoch, G. M., and Nemerow, G. R. (1998a). Adenovirus endocytosis requires actin cytoskeleton reorganization mediated by Rho family GTPases. *J.Viro.* **72**, 8806-8812.

Li, E., Stupack, D., Klemke, R., Cheresh, D. A., and Nemerow, G. R. (1998b). Adenovirus endocytosis via alpha(v) integrins requires phosphoinositide-3-OH kinase. *J.Viro.* **72**, 2055-2061.

Li, E., Stupack, D. G., Brown, S. L., Klemke, R., Schlaepfer, D. D., and Nemerow, G. R. (2000). Association of p130CAS with phosphatidylinositol-3-OH kinase mediates adenovirus cell entry. *J.Biol.Chem.* **275**, 14729-14735.

Li, Y., Pong, R. C., Bergelson, J. M., Hall, M. C., Sagalowsky, A. I., Tseng, C. P., Wang, Z., and Hsieh, J. T. (1999). Loss of adenoviral receptor expression in human bladder cancer cells: a potential impact on the efficacy of gene therapy. *Cancer Res.* **59**, 325-330.

Lochmuller, H., Jani, A., Huard, J., Prescott, S., Simoneau, M., Massie, B., Karpati, G., and Acsadi, G. (1994). Emergence of early region 1-containing replication-competent adenovirus in stocks of replication-defective adenovirus recombinants (delta E1 + delta E3) during multiple passages in 293 cells. *Hum.Gene Ther.* **5**, 1485-1491.

Lonberg-Holm, K., Crowell, R. L., and Philipson, L. (1976). Unrelated animal viruses share receptors. *Nature* **259**, 679-681.

Loser, P., Hillgenberg, M., Arnold, W., Both, G. W., and Hofmann, C. (2000). Ovine adenovirus vectors mediate efficient gene transfer to skeletal muscle. *Gene Ther.* **7**, 1491-1498.

Louis, N., Fender, P., Barge, A., Kitts, P., and Chroboczek, J. (1994). Cell-binding domain of adenovirus serotype 2 fiber. *J.Viro.* **68**, 4104-4106.

Luque, F., Caruz, A., Pineda, J. A., Torres, Y., Larder, B., and Leal, M. (1994). Provirus load changes in untreated and zidovudine-treated human immunodeficiency virus type 1-infected patients. *J.Infect.Dis.* **169**, 267-273.

Lusky, M., Christ, M., Rittner, K., Dieterle, A., Dreyer, D., Mourot, B., Schultz, H., Stoeckel, F., Pavirani, A., and Mehtali, M. (1998). In vitro and in vivo biology of recombinant adenovirus vectors with E1, E1/E2A, or E1/E4 deleted. *J.Virol.* **72**, 2022-2032.

Mack, C. A., Song, W. R., Carpenter, H., Wickham, T. J., Kovacs, I., Harvey, B. G., Magovern, C. J., Isom, O. W., Rosengart, T., Falck-Pedersen, E., Hackett, N. R., Crystal, R. G., and Mastrangeli, A. (1997). Circumvention of anti-adenovirus neutralizing immunity by administration of an adenoviral vector of an alternate serotype. *Hum.Gene Ther.* **8**, 99-109.

Maddon, P. J., Dalgleish, A. G., McDougal, J. S., Clapham, P. R., Weiss, R. A., and Axel, R. (1986). The T4 gene encodes the AIDS virus receptor and is expressed in the immune system and the brain. *Cell* **47**, 333-348.

Maitland, N. J. and McDougall, J. K. (1977). Biochemical transformation of mouse cells by fragments of herpes simplex virus DNA. *Cell* **11**, 233-241.

Marienfeld, U., Haack, A., Thalheimer, P., Schneider-Rasp, S., Brackmann, H. H., and Poller, W. (1999). 'Autoreplication' of the vector genome in recombinant adenoviral vectors with different E1 region deletions and transgenes. *Gene Ther.* **6**, 1101-1113.

Martin, A. B., Webber, S., Fricker, F. J., Jaffe, R., Demmler, G., Kearney, D., Zhang, Y. H., Bodurtha, J., Gelb, B., and Ni, J. (1994). Acute myocarditis. Rapid diagnosis by PCR in children. *Circulation* **90**, 330-339.

Martino, T. A., Petric, M., Brown, M., Aitken, K., Gauntt, C. J., Richardson, C. D., Chow, L. H., and Liu, P. P. (1998). Cardiovirulent coxsackieviruses and the decay-accelerating factor (CD55) receptor. *Virology* **244**, 302-314.

Martino, T. A., Petric, M., Weingartl, H., Bergelson, J. M., Opavsky, M. A., Richardson, C. D., Modlin, J. F., Finberg, R. W., Kain, K. C., Willis, N., Gauntt, C. J., and Liu, P. P. (2000). The coxsackie-adenovirus receptor (CAR) is used by reference strains and clinical isolates representing all six serotypes of coxsackievirus group B and by swine vesicular disease virus. *Virology* **271**, 99-108.

Mathias, P., Wickham, T., Moore, M., and Nemerow, G. (1994). Multiple adenovirus serotypes use alpha v integrins for infection. *J.Virol.* **68**, 6811-6814.

Mayr, G. A. and Freimuth, P. (1997). A single locus on human chromosome 21 directs the expression of a receptor for adenovirus type 2 in mouse A9 cells. *J.Virol.* **71**, 412-418.

McCabe, D., Swain, W., Martinell, P., and Christou, B. (1988). Stable transformation of soy bean (*Glycine max*) by particle acceleration. *Biotechnology* **6**, 923-926

McCray, P. B. Jr, Armstrong, K., Zabner, J., Miller, D. W., Koretzky, G. A., Couture, L., Robillard, J. E., Smith, A. E., and Welsh, M. J. (1995). Adenoviral-mediated gene transfer to fetal pulmonary epithelia in vitro and in vivo. *J.Clin.Invest.* **95**, 2620-2632.

McCulloch, R. K., Choong, C. S., and Hurley, D. M. (1995). An evaluation of competitor type and size for use in the determination of mRNA by competitive PCR. *PCR.Methods Appl.* **4**, 219-226.

McDonald, D., Stockwin, L., Matzow, T., Zajdel, M. B., and Blair, G. (1999). Coxsackie and adenovirus receptor (CAR)-dependent and major histocompatibility complex (MHC) class I-

independent uptake of recombinant adenoviruses into human tumour cells. *Gene Ther.* **6**, 1512-1519.

Mei Y.F. and Wadell, G. (1995). Molecular determinants of adenovirus tropism. In 'The molecular repertoire of adenoviruses'. (Doerfler W. and Böhm P. Eds.) pp. 213-228. (Springer Verlag KG: Berlin.)

Mei, Y. F. and Wadell, G. (1996). Epitopes and hemagglutination binding domain on subgenus B:2 adenovirus fibers. *J.Viro.* **70**, 3688-3697.

Melnick J.L. (1996). Polioviruses, Coxsackieviruses, Echoviruses, and newer Enteroviruses. In 'Fields Virology'. (Fields B.N., Knipe D.M., and Howley P.M. Eds.) pp. 655-712. (Lippincott-Raven: Philadelphia.)

Mena, I., Fischer, C., Gebhard, J. R., Perry, C. M., Harkins, S., and Whitton, J. L. (2000). Coxsackievirus infection of the pancreas: evaluation of receptor expression, pathogenesis, and immunopathology. *Virology* **271**, 276-288.

Mendelsohn, C. L., Wimmer, E., and Racaniello, V. R. (1989). Cellular receptor for poliovirus: molecular cloning, nucleotide sequence, and expression of a new member of the immunoglobulin superfamily. *Cell* **56**, 855-865.

Mentel, R., Dopping, G., Wegner, U., Seidel, W., Liebermann, H., and Dohner, L. (1997). Adenovirus-receptor interaction with human lymphocytes. *J.Med.Viro.* **51**, 252-257.

Merrick, A. F., Shewring, L. D., Sawyer, G. J., Gustafsson, K. T., and Fabre, J. W. (1996). Comparison of adenovirus gene transfer to vascular endothelial cells in cell culture, organ culture, and in vivo. *Transplantation* **62**, 1085-1089.

Michel, G. (1986). Kompendium der Embryologie der Haustiere. 4., überarbeitete Auflage, Gustav Fischer Verlag Jena

Miller, C. R., Buchsbaum, D. J., Reynolds, P. N., Douglas, J. T., Gillespie, G. Y., Mayo, M. S., Raben, D., and Curiel, D. T. (1998). Differential susceptibility of primary and established human glioma cells to adenovirus infection: targeting via the epidermal growth factor receptor achieves fiber receptor-independent gene transfer. *Cancer Res.* **58**, 5738-5748.

Mocharla, H., Mocharla, R., and Hodes, M. E. (1990). Coupled reverse transcription-polymerase chain reaction (RT-PCR) as a sensitive and rapid method for isozyme genotyping. *Gene* **93**, 271-275.

Mondain, M., Restituito, S., Vincenti, V., Gardiner, Q., Uziel, A., Delabre, A., Mathieu, M., Bousquet, J., and Demoly, P. (1998). Adenovirus-mediated in vivo gene transfer in guinea pig middle ear mucosa. *Hum.Gene Ther.* **9**, 1217-1221.

Montgomery, A. M., Becker, J. C., Siu, C. H., Lemmon, V. P., Cheresh, D. A., Pancook, J. D., Zhao, X., and Reisfeld, R. A. (1996). Human neural cell adhesion molecule L1 and rat homologue NILE are ligands for integrin alpha v beta 3. *J.Cell Biol.* **132**, 475-485.

Morral, N., O'Neal, W., Rice, K., Leland, M., Kaplan, J., Piedra, P. A., Zhou, H., Parks, R. J., Velji, R., Aguilar-Cordova, E., Wadsworth, S., Graham, F. L., Kochanek, S., Carey, K. D., and Beaudet, A. L. (1999). Administration of helper-dependent adenoviral vectors and

sequential delivery of different vector serotype for long-term liver-directed gene transfer in baboons. *Proc.Natl.Acad.Sci.U.S.A.* **96**, 12816-12821.

Morrison, C. and Gannon, F. (1994). The impact of the PCR plateau phase on quantitative PCR. *Biochim.Biophys.Acta* **1219**, 493-498.

Morsy, M. A. and Caskey, C. T. (1999). Expanded-capacity adenoviral vectors--the helper-dependent vectors. *Mol.Med.Today* **5**, 18-24.

Morsy, M. A., Gu, M., Motzel, S., Zhao, J., Lin, J., Su, Q., Allen, H., Franklin, L., Parks, R. J., Graham, F. L., Kochanek, S., Bett, A. J., and Caskey, C. T. (1998). An adenoviral vector deleted for all viral coding sequences results in enhanced safety and extended expression of a leptin transgene. *Proc.Natl.Acad.Sci.U.S.A.* **95**, 7866-7871.

Muckelbauer, J. K., Kremer, M., Minor, I., Diana, G., Dutko, F. J., Groarke, J., Pevear, D. C., and Rossmann, M. G. (1995). The structure of coxsackievirus B3 at 3.5 Å resolution. *Structure*. **3**, 653-667.

Murphy, L. D., Herzog, C. E., Rudick, J. B., Fojo, A. T., and Bates, S. E. (1990). Use of the polymerase chain reaction in the quantitation of mdr-1 gene expression. *Biochemistry* **29**, 10351-10356.

Muzyczka, N. (1992). Use of adeno-associated virus as a general transduction vector for mammalian cells. *Curr.Top.Microbiol.Immunol.* **158:97-129.**, 97-129.

Nalbantoglu, J., Larochelle, N., Wolf, E., Karpati, G., Lochmuller, H., and Holland, P. C. (2001). Muscle-specific overexpression of the adenovirus primary receptor car overcomes low efficiency of gene transfer to mature skeletal muscle. *J.Viro.* **75**, 4276-4282.

Nalbantoglu, J., Pari, G., Karpati, G., and Holland, P. C. (1999). Expression of the primary coxsackie and adenovirus receptor is downregulated during skeletal muscle maturation and limits the efficacy of adenovirus-mediated gene delivery to muscle cells. *Hum.Gene Ther.* **10**, 1009-1019.

Nardelli, L., Lodetti, E., Gualandi, G. L., Burrows, R., Goodridge, D., Brown, F., and Cartwright, B. (1968). A foot and mouth disease syndrome in pigs caused by an enterovirus. *Nature* **219**, 1275-1276.

Nemerow, G. R. (2000). Cell receptors involved in adenovirus entry. *Virology* **274**, 1-4.

Nemerow, G. R. and Stewart, P. L. (1999). Role of alpha(v) integrins in adenovirus cell entry and gene delivery. *Microbiol.Mol.Biol.Rev.* **63**, 725-734.

Neubauer, A., Neubauer, B., and Liu, E. (1990). Polymerase chain reaction based assay to detect allelic loss in human DNA: loss of beta-interferon gene in chronic myelogenous leukemia. *Nucleic.Acids.Res.* **18**, 993-998.

Neumann, E., Schaefer-Ridder, M., Wang, Y., and Hofschneider, P. H. (1982). Gene transfer into mouse lymphoma cells by electroporation in high electric fields. *EMBO J.* **1**, 841-845.

Neumann, R., Chroboczek, J., and Jacrot, B. (1988). Determination of the nucleotide sequence for the penton-base gene of human adenovirus type 5. *Gene* **69**, 153-157.

Noonan, K. E., Beck, C., Holzmayer, T. A., Chin, J. E., Wunder, J. S., Andrulis, I. L., Gazdar, A. F., Willman, C. L., Griffith, B., and Von Hoff, D. D. (1990). Quantitative analysis of MDR1 (multidrug resistance) gene expression in human tumors by polymerase chain reaction. *Proc.Natl.Acad.Sci.U.S.A.* **87**, 7160-7164.

Noutsias, M., Fechner, H., de Jonge, H., Wang, X., Dekkers, D., Houtsmuller, A. B., Pauschinger, M., Bergelson, J., Warraich, R., Yacoub, M., Hetzer, R., Lamers, J., Schultheiss, H. P., and Poller, W. (2001). Human Coxsackie-Adenovirus Receptor Is Colocalized With Integrins alpha(v)beta(3) and alpha(v)beta(5) on the Cardiomyocyte Sarcolemma and Upregulated in Dilated Cardiomyopathy: Implications for Cardiotropic Viral Infections. *Circulation* **104**, 275-280.

O'Neal, W. K., Zhou, H., Morral, N., Aguilar-Cordova, E., Pestaner, J., Langston, C., Mull, B., Wang, Y., Beaudet, A. L., and Lee, B. (1998). Toxicological comparison of E2a-deleted and first-generation adenoviral vectors expressing alpha1-antitrypsin after systemic delivery. *Hum.Gene Ther.* **9**, 1587-1598.

Okegawa, T., Li, Y., Pong, R. C., Bergelson, J. M., Zhou, J., and Hsieh, J. T. (2000). The dual impact of coxsackie and adenovirus receptor expression on human prostate cancer gene therapy. *Cancer Res.* **60**, 5031-5036.

Okegawa, T., Pong, R. C., Li, Y., Bergelson, J. M., Sagalowsky, A. I., and Hsieh, J. T. (2001). The mechanism of the growth-inhibitory effect of coxsackie and adenovirus receptor (CAR) on human bladder cancer: a functional analysis of car protein structure. *Cancer Res.* **61**, 6592-6600.

Orlando, C., Pinzani, P., and Pazzaglia, M. (1998). Developments in quantitative PCR. *Clin.Chem.Lab.Med.* **36**, 255-269.

Orr, H. T., Lancet, D., Robb, R. J., Lopez, de Castro JA, and Strominger, J. L. (1979). The heavy chain of human histocompatibility antigen HLA-B7 contains an immunoglobulin-like region. *Nature* **282**, 266-270.

Palese, P., Zheng, H., Engelhardt, O. G., Pleschka, S., and Garcia-Sastre, A. (1996). Negative-strand RNA viruses: genetic engineering and applications. *Proc.Natl.Acad.Sci.U.S.A.* **93**, 11354-11358.

Park, O. K. and Mayo, K. E. (1991). Transient expression of progesterone receptor messenger RNA in ovarian granulosa cells after the preovulatory luteinizing hormone surge. *Mol.Endocrinol.* **5**, 967-978.

Parks, R., Evelegh, C., and Graham, F. (1999). Use of helper-dependent adenoviral vectors of alternative serotypes permits repeat vector administration. *Gene Ther.* **6**, 1565-1573.

Pastoret, P. P. and Brochier, B. (1992). Development of a recombinant vaccinia-rabies vaccine for oral vaccination of foxes against rabies. *Dev.Biol.Stand.* **79:105-11**., 105-111.

Pauschinger, M., Doerner, A., Kuehl, U., Schwimmbeck, P. L., Poller, W., Kandolf, R., and Schultheiss, H. P. (1999). Enteroviral RNA Replication in the Myocardium of Patients With Left Ventricular Dysfunction and Clinically Suspected Myocarditis. *Circulation* **99**, 889-895.

- Pearson, A. S., Koch, P. E., Atkinson, N., Xiong, M., Finberg, R. W., Roth, J. A., and Fang, B. (1999). Factors limiting adenovirus-mediated gene transfer into human lung and pancreatic cancer cell lines. *Clin.Cancer Res.* **5**, 4208-4213.
- Perales, J. C., Ferkol, T., Molas, M., and Hanson, R. W. (1994). An evaluation of receptor-mediated gene transfer using synthetic DNA-ligand complexes. *Eur.J.Biochem.* **226**, 255-266.
- Philipson, L., Lonberg-Holm, K., and Pettersson, U. (1968). Virus-receptor interaction in an adenovirus system. *J.Virol.* **2**, 1064-1075.
- Piatak, M. Jr, Luk, K. C., Williams, B., and Lifson, J. D. (1993). Quantitative competitive polymerase chain reaction for accurate quantitation of HIV DNA and RNA species. *Biotechniques* **14**, 70-81.
- Pickles, R. J., Fahrner, J. A., Petrella, J. M., Boucher, R. C., and Bergelson, J. M. (2000). Retargeting the coxsackievirus and adenovirus receptor to the apical surface of polarized epithelial cells reveals the glycocalyx as a barrier to adenovirus-mediated gene transfer. *J.Virol.* **74**, 6050-6057.
- Pickles, R. J., McCarty, D., Matsui, H., Hart, P. J., Randell, S. H., and Boucher, R. C. (1998). Limited entry of adenovirus vectors into well-differentiated airway epithelium is responsible for inefficient gene transfer. *J.Virol.* **72**, 6014-6023.
- Porcher, C., Malinge, M. C., Picat, C., and Grandchamp, B. (1992). A simplified method for determination of specific DNA or RNA copy number using quantitative PCR and an automatic DNA sequencer. *Biotechniques* **13**, 106-114.
- Raeymaekers, L. (1993). Quantitative PCR: theoretical considerations with practical implications. *Anal.Biochem.* **214**, 582-585.
- Raphael, Y., Frisancho, J. C., and Roessler, B. J. (1996). Adenoviral-mediated gene transfer into guinea pig cochlear cells in vivo. *Neurosci.Lett.* **207**, 137-141.
- Rebel, V. I., Hartnett, S., Denham, J., Chan, M., Finberg, R., and Sieff, C. A. (2000). Maturation and lineage-specific expression of the coxsackie and adenovirus receptor in hematopoietic cells. *Stem.Cells* **18**, 176-182.
- Rekhter, M. D., Shah, N., Simari, R. D., Work, C., Kim, J. S., Nabel, G. J., Nabel, E. G., and Gordon, D. (1998a). Graft permeabilization facilitates gene therapy of transplant arteriosclerosis in a rabbit model. *Circulation* **98**, 1335-1341.
- Rekhter, M. D., Simari, R. D., Work, C. W., Nabel, G. J., Nabel, E. G., and Gordon, D. (1998b). Gene transfer into normal and atherosclerotic human blood vessels. *Circ.Res.* **82**, 1243-1252.
- Robbins, P. D., Tahara, H., and Ghivizzani, S. C. (1998). Viral vectors for gene therapy. *Trends.Biotechnol.* **16**, 35-40.
- Roelvink, P. W., Kovesdi, I., and Wickham, T. J. (1996). Comparative analysis of adenovirus fiber-cell interaction: adenovirus type 2 (Ad2) and Ad9 utilize the same cellular fiber receptor but use different binding strategies for attachment. *J.Virol.* **70**, 7614-7621.

- Roelvink, P. W., Lizonova, A., Lee, J. G., Li, Y., Bergelson, J. M., Finberg, R. W., Brough, D. E., Kovesdi, I., and Wickham, T. J. (1998). The coxsackievirus-adenovirus receptor protein can function as a cellular attachment protein for adenovirus serotypes from subgroups A, C, D, E, and F. *J.Virol.* **72**, 7909-7915.
- Roelvink, P. W., Mi, Lee G., Einfeld, D. A., Kovesdi, I., and Wickham, T. J. (1999). Identification of a Conserved Receptor-Binding Site on the Fiber Proteins of CAR-Recognizing Adenoviridae. *Science* **286**, 1568-1571.
- Roessler, B. J., Allen, E. D., Wilson, J. M., Hartman, J. W., and Davidson, B. L. (1993). Adenoviral-mediated gene transfer to rabbit synovium in vivo. *J.Clin.Invest.* **92**, 1085-1092.
- Rovainen, M., Hyypia, T., Piirainen, L., Kalkkinen, N., Stanway, G., and Hovi, T. (1991). RGD-dependent entry of coxsackievirus A9 into host cells and its bypass after cleavage of VP1 protein by intestinal proteases. *J.Virol.* **65**, 4735-4740.
- Rovainen, M., Piirainen, L., and Hovi, T. (1996). Efficient RGD-independent entry process of coxsackievirus A9. *Arch.Virolog.* **141**, 1909-1919.
- Rovainen, M., Piirainen, L., Hovi, T., Virtanen, I., Riikonen, T., Heino, J., and Hyypia, T. (1994). Entry of coxsackievirus A9 into host cells: specific interactions with alpha v beta 3 integrin, the vitronectin receptor. *Virology* **203**, 357-365.
- Rothmann, T., Katus, H. A., Hartong, R., Perricaudet, M., and Franz, W. M. (1996). Heart muscle-specific gene expression using replication defective recombinant adenovirus. *Gene Ther.* **3**, 919-926.
- Ruigrok, R. W., Barge, A., Albiges-Rizo, C., and Dayan, S. (1990). Structure of adenovirus fibre. II. Morphology of single fibres. *J.Mol.Biol.* **215**, 589-596.
- Russell, W. C. (2000). Update on adenovirus and its vectors. *J.Gen.Virol.* **81 Pt 11:2573-604**, 2573-2604.
- Ruster, B., Zeuzem, S., and Roth, W. K. (1995). Quantification of hepatitis C virus RNA by competitive reverse transcription and polymerase chain reaction using a modified hepatitis C virus RNA transcript. *Anal.Biochem.* **224**, 597-600.
- Rüger R (1997). Methoden der Genübertragung. In 'Handbuch der molekularen Medizin'. (Ed. Ganter D & Ruckpaul K.) pp. 197-241. (Springer: Berlin, Heidelberg.)
- Saiki, R. K., Scharf, S., Faloona, F., Mullis, K. B., Horn, G. T., Erlich, H. A., and Arnheim, N. (1985). Enzymatic amplification of beta-globin genomic sequences and restriction site analysis for diagnosis of sickle cell anemia. *Science* **230**, 1350-1354.
- Sambrook, J., Fritsch, E.F., Maniatis, T. (1989). Molecular cloning: A laboratory manual. 2nd ed. Cold Spring Harbor Laboratory Press, New York
- Santis, G., Legrand, V., Hong, S. S., Davison, E., Kirby, I., Imler, J. L., Finberg, R. W., Bergelson, J. M., Mehtali, M., and Boulanger, P. (1999). Molecular determinants of adenovirus serotype 5 fibre binding to its cellular receptor CAR. *J.Gen.Virol.* **80**, 1519-1527.

- Saphire, A. C., Guan, T., Schirmer, E. C., Nemerow, G. R., and Gerace, L. (2000). Nuclear import of adenovirus DNA in vitro involves the nuclear protein import pathway and hsc70. *J.Biol.Chem.* **275**, 4298-4304.
- Schachtner, S., Buck, C., Bergelson, J., and Baldwin, H. (1999). Temporally regulated expression patterns following in utero adenovirus-mediated gene transfer. *Gene Ther.* **6**, 1249-1257.
- Schanke, J. T., Quam, L. M., and Van Ness, B. G. (1994). Flip-PCR for DNA sequence motif inversion. *Biotechniques* **16**, 414-416.
- Schmidt, E. E. and Merrill, G. F. (1991). Changes in dihydrofolate reductase (DHFR) mRNA levels can account fully for changes in DHFR synthesis rates during terminal differentiation in a highly amplified myogenic cell line. *Mol.Cell Biol.* **11**, 3726-3734.
- Schmidt, M. R., Piekos, B., Cabatingan, M. S., and Woodland, R. T. (2000). Expression of a human coxsackieadenovirus receptor transgene permits adenovirus infection of primary lymphocytes. *J.Immunol.* **165**, 4112-4119.
- Schoehn, G., Fender, P., Chroboczek, J., and Hewat, E. A. (1996). Adenovirus 3 penton dodecahedron exhibits structural changes of the base on fibre binding. *EMBO J.* **15**, 6841-6846.
- Scimenti, C. R. and Calos, M. P. (1998). Epstein-Barr virus vectors for gene expression and transfer. *Curr.Opin.Biotechnol.* **9**, 476-479.
- Seidman, M. A., Hogan, S. M., Wendland, R. L., Worgall, S., Crystal, R. G., and Leopold, P. L. (2001). Variation in adenovirus receptor expression and adenovirus vector-mediated transgene expression at defined stages of the cell cycle. *Mol.Ther.* **4**, 13-21.
- Sekhon, H. S. and Larson, J. E. (1995). In utero gene transfer into the pulmonary epithelium. *Nat.Med.* **1**, 1201-1203.
- Sestini, R., Orlando, C., Peri, A., Tricarico, C., Pazzagli, M., Serio, M., Pagani, A., Bussolati, G., Granchi, S., and Maggi, M. (1996). Quantitation of somatostatin receptor type 2 gene expression in neuroblastoma cell lines and primary tumors using competitive reverse transcription-polymerase chain reaction. *Clin.Cancer Res.* **2**, 1757-1765.
- Sestini, R., Orlando, C., Zentilin, L., Gelmini, S., Pinzani, P., Bianchi, S., Sell, C., Giacca, M., and Pazzagli, M. (1994). Measuring c-erbB-2 oncogene amplification in fresh and paraffin-embedded tumors by competitive polymerase chain reaction. *Clin.Chem.* **40**, 630-636.
- Shayakhmetov, D. M., Papayannopoulou, T., Stamatoyannopoulos, G., and Lieber, A. (2000). Efficient gene transfer into human CD34(+) cells by a retargeted adenovirus vector. *J.Viro.* **74**, 2567-2583.
- Shears, L. L. 2nd, Kibbe, M. R., Murdock, A. D., Billiar, T. R., Lizonova, A., Kovesdi, I., Watkins, S. C., and Tzeng, E. (1998). Efficient inhibition of intimal hyperplasia by adenovirus- mediated inducible nitric oxide synthase gene transfer to rats and pigs in vivo. *J.Am.Coll.Surg.* **187**, 295-306.

Shenk T. (1996). Adenoviridae: The viruses and their replication. In 'Fields Virology'. (Fields BN, Knipe DM, and Howley PMEds.) pp. 2111-2148. (Lippincott-Raven: Philadelphia, New York.)

Siebert, P. D. and Fukuda, M. (1985). Induction of cytoskeletal vimentin and actin gene expression by a tumor-promoting phorbol ester in the human leukemic cell line K562. *J.Biol.Chem.* **260**, 3868-3874.

Siebert, P. D. and Lerrick, J. W. (1992). Competitive PCR. *Nature* **359**, 557-558.

Simmonds, P., Balfé, P., Peutherer, J. F., Ludlam, C. A., Bishop, J. O., and Brown, A. J. (1990). Human immunodeficiency virus-infected individuals contain provirus in small numbers of peripheral mononuclear cells and at low copy numbers. *J.Virol.* **64**, 864-872.

Smollich, A., Drost, J., and Michel, G. (1985). In 'Mikroskopische Anatomie der Haustiere'. (A. Smollich and G. Michel Eds.) pp. 13-177. (Gustav Fischer Verlag: Jena.)

Sprengel, J., Schmitz, B., Heuss-Neitzel, D., Zock, C., and Doerfler, W. (1994). Nucleotide sequence of human adenovirus type 12 DNA: comparative functional analysis. *J.Virol.* **68**, 379-389.

Staunton, D. E., Merluzzi, V. J., Rothlein, R., Barton, R., Marlin, S. D., and Springer, T. A. (1989). A cell adhesion molecule, ICAM-1, is the major surface receptor for rhinoviruses. *Cell* **56**, 849-853.

Stevenson, S. C., Rollence, M., Marshall-Neff, J., and McClelland, A. (1997). Selective targeting of human cells by a chimeric adenovirus vector containing a modified fiber protein. *J.Virol.* **71**, 4782-4790.

Stevenson, S. C., Rollence, M., White, B., Weaver, L., and McClelland, A. (1995). Human adenovirus serotypes 3 and 5 bind to two different cellular receptors via the fiber head domain. *J.Virol.* **69**, 2850-2857.

Stewart, P. L., Burnett, R. M., Cyrklaff, M., and Fuller, S. D. (1991). Image reconstruction reveals the complex molecular organization of adenovirus. *Cell* **67**, 145-154.

Stewart, P. L., Fuller, S. D., and Burnett, R. M. (1993). Difference imaging of adenovirus: bridging the resolution gap between X-ray crystallography and electron microscopy. *EMBO J.* **12**, 2589-2599.

Summerford, C., Bartlett, J. S., and Samulski, R. J. (1999). AlphaVbeta5 integrin: a coreceptor for adeno-associated virus type 2 infection. *Nat.Med.* **5**, 78-82.

Svensson, U. (1985). Role of vesicles during adenovirus 2 internalization into HeLa cells. *J.Virol.* **55**, 442-449.

Sykes, P. J., Neoh, S. H., Brisco, M. J., Hughes, E., Condon, J., and Morley, A. A. (1992). Quantitation of targets for PCR by use of limiting dilution. *Biotechniques* **13**, 444-449.

Takayama, K., Ueno, H., Pei, X. H., Nakanishi, Y., Yatsunami, J., and Hara, N. (1998). The levels of integrin alpha v beta 5 may predict the susceptibility to adenovirus-mediated gene transfer in human lung cancer cells. *Gene Ther.* **5**, 361-368.

Tamai, H., Kim, H. S., Arai, H., Inoue, K., and Mino, M. (1998). Developmental changes in the expression of alpha-tocopherol transfer protein during the neonatal period of rat. *Biofactors* **7**, 87-91.

Tan, P. K., Michou, A. I., Bergelson, J. M., and Cotten, M. (2001). Defining CAR as a cellular receptor for the avian adenovirus CELO using a genetic analysis of the two viral fibre proteins. *J.Gen.Viro.* **82**, 1465-1472.

Telenti, A., Imboden, P., and Germann, D. (1992). Competitive polymerase chain reaction using an internal standard: application to the quantitation of viral DNA. *J.Virol.Methods* **39**, 259-268.

Thoelen, I., Magnusson, C., Tagerud, S., Polacek, C., Lindberg, M., and Van Ranst, M. (2001a). Identification of alternative splice products encoded by the human coxsackie-adenovirus receptor gene. *Biochem.Biophys.Res.Commun.* **287**, 216-222.

Thoelen, I., Keyarts, E., Lindberg, M. and Van Ranst, M. (2001b). Charakterisation of a cDNA encoding the bovine coxsackie and adenovirus receptor. *Biochem.Biophys.Res.Commun.* **288**, 805-808

Tomko, R. P., Johansson, C. B., Totrov, M., Abagyan, R., Frisen, J., and Philipson, L. (2000). Expression of the adenovirus receptor and its interaction with the fiber knob. *Exp.Cell Res.* **255**, 47-55.

Tomko, R. P., Xu, R., and Philipson, L. (1997). HCAR and MCAR: the human and mouse cellular receptors for subgroup C adenoviruses and group B coxsackieviruses. *Proc.Natl.Acad.Sci.U.S.A.* **94**, 3352-3356.

Torres, J. M., Alonso, C., Ortega, A., Mittal, S., Graham, F., and Enjuanes, L. (1996). Tropism of human adenovirus type 5-based vectors in swine and their ability to protect against transmissible gastroenteritis coronavirus. *J.Viro.* **70**, 3770-3780.

Tsubota, K., Inoue, H., Ando, K., Ono, M., Yoshino, K., and Saito, I. (1998). Adenovirus-mediated gene transfer to the ocular surface epithelium [In Process Citation]. *Exp.Eye Res.* **67**, 531-538.

Turturro, F., Seth, P., and Link, C. J. Jr (2000). In vitro adenoviral vector p53-mediated transduction and killing correlates with expression of coxsackie-adenovirus receptor and alpha(nu)beta5 integrin in SUDHL-1 cells derived from anaplastic large-cell lymphoma. *Clin.Cancer Res.* **6**, 185-192.

van Raaij, M. J., Chouin, E., van der Zandt, H., Bergelson, J. M., and Cusack, S. (2000). Dimeric structure of the coxsackievirus and adenovirus receptor D1 domain at 1.7 Å resolution. *Structure.Fold.Des.* **8**, 1147-1155.

van Raaij, M. J., Louis, N., Chroboczek, J., and Cusack, S. (1999). Structure of the human adenovirus serotype 2 fiber head domain at 1.5 Å resolution. *Virology* **262**, 333-343.

van Rhee, F., Marks, D. I., Lin, F., Szydlo, R. M., Hochhaus, A., Treleaven, J., Delord, C., Cross, N. C., and Goldman, J. M. (1995). Quantification of residual disease in Philadelphia-positive acute lymphoblastic leukemia: comparison of blood and bone marrow. *Leukemia* **9**, 329-335.

- Van, den Veyver, I., Ni, J., Bowles, N., Carpenter, R. J. Jr, Weiner, C. P., Yankowitz, J., Moise, K. J. Jr, Henderson, J., and Towbin, J. A. (1998). Detection of intrauterine viral infection using the polymerase chain reaction. *Mol. Genet. Metab.* **63**, 85-95.
- Vanderkwaak, T. J., Wang, M., Gomez-Navarro, J., Rancourt, C., Dmitriev, I., Krasnykh, V., Barnes, M., Siegal, G. P., Alvarez, R., and Curiel, D. T. (1999). An advanced generation of adenoviral vectors selectively enhances gene transfer for ovarian cancer gene therapy approaches. *Gynecol. Oncol.* **74**, 227-234.
- Varga, M. J., Weibull, C., and Everitt, E. (1991). Infectious entry pathway of adenovirus type 2. *J. Virol.* **65**, 6061-6070.
- Vincent, M. C., Trapnell, B. C., Baughman, R. P., Wert, S. E., Whitsett, J. A., and Iwamoto, H. S. (1995). Adenovirus-mediated gene transfer to the respiratory tract of fetal sheep in utero. *Hum. Gene Ther.* **6**, 1019-1028.
- Viswalingam, N. D. (1993). Adenovirus keratoconjunctivitis: an enigma. *Eye* **7**, 5-7.
- Wadell, G. (1990). Adenoviruses. In 'Principles and practices of clinical virology'. (Zuckerman A.J., Banatvala J.E., and Pattison J.R. Eds.) pp. 267-287. (John Wiley & Sons: Chichester.)
- Wadell, G., Allard, A., Johansson, M., Svensson, L., and Uhnoo, I. (1987). Enteric adenoviruses. *Ciba Found. Symp.* **128:63-91.**, 63-91.
- Wadell, G., Hammarckjold, M. L., Winberg, G., Varsanyi, T. M., and Sundell, G. (1980). Genetic variability of adenoviruses. *Ann. N.Y. Acad. Sci.* **354:16-42.**, 16-42.
- Wahlfors, J. J., Zullo, S. A., Loimas, S., Nelson, D. M., and Morgan, R. A. (2000). Evaluation of recombinant alphaviruses as vectors in gene therapy. *Gene Ther.* **7**, 472-480.
- Walters, R. W., Grunst, T., Bergelson, J. M., Finberg, R. W., Welsh, M. J., and Zabner, J. (1999). Basolateral localization of fiber receptors limits adenovirus infection from the apical surface of airway epithelia. *J. Biol. Chem.* **274**, 10219-10226.
- Wan, Y. Y., Leon, R. P., Marks, R., Cham, C. M., Schaack, J., Gajewski, T. F., and DeGregori, J. (2000). Transgenic expression of the coxsackie/adenovirus receptor enables adenoviral-mediated gene delivery in naive T cells. *Proc. Natl. Acad. Sci. U.S.A.* **97**, 13784-13789.
- Wang, A. M., Doyle, M. V., and Mark, D. F. (1989). Quantitation of mRNA by the polymerase chain reaction. *Proc. Natl. Acad. Sci. U.S.A.* **86**, 9717-9721.
- Wang, G., Williamson, R., Mueller, G., Thomas, P., Davidson, B. L., and McCray, P. B., Jr. (1998). Ultrasound-guided gene transfer to hepatocytes in utero. *Fetal Diagn. Ther.* **13**, 197-205.
- Wang, K., Guan, T., Cheresh, D. A., and Nemerow, G. R. (2000). Regulation of adenovirus membrane penetration by the cytoplasmic tail of integrin beta5. *J. Virol.* **74**, 2731-2739.
- Wang, X. and Bergelson, J. M. (1999). Coxsackievirus and adenovirus receptor cytoplasmic and transmembrane domains are not essential for coxsackievirus and adenovirus infection [In Process Citation]. *J. Virol.* **73**, 2559-2562.

- Watzka, M., Waha, A., Koch, A., Schmutzler, R. K., Bidlingmaier, F., von Deimling, A., Klingmuller, D., and Stoffel-Wagner, B. (1997). An optimized protocol for mRNA quantification using nested competitive RT-PCR. *Biochem.Biophys.Res.Commun.* **231**, 813-817.
- Wenstrom, K. D., Andrews, W. W., Bowles, N. E., Towbin, J. A., Hauth, J. C., and Goldenberg, R. L. (1998). Intrauterine viral infection at the time of second trimester genetic amniocentesis. *Obstet.Gynecol.* **92**, 420-424.
- Wickham, T. J. (2000). Targeting adenovirus. *Gene Ther.* **7**, 110-114.
- Wickham, T. J., Filardo, E. J., Cheresh, D. A., and Nemerow, G. R. (1994). Integrin alpha v beta 5 selectively promotes adenovirus mediated cell membrane permeabilization. *J.Cell Biol.* **127**, 257-264.
- Wickham, T. J., Mathias, P., Cheresh, D. A., and Nemerow, G. R. (1993). Integrins alpha v beta 3 and alpha v beta 5 promote adenovirus internalization but not virus attachment. *Cell* **73**, 309-319.
- Wickham, T. J., Roelvink, P. W., Brough, D. E., and Kovesdi, I. (1996a). Adenovirus targeted to heparan-containing receptors increases its gene delivery efficiency to multiple cell types. *Nat.Biotechnol.* **14**, 1570-1573.
- Wickham, T. J., Segal, D. M., Roelvink, P. W., Carrion, M. E., Lizonova, A., Lee, G. M., and Kovesdi, I. (1996b). Targeted adenovirus gene transfer to endothelial and smooth muscle cells by using bispecific antibodies. *J.Viro.* **70**, 6831-6838.
- Wickham, T. J., Tzeng, E., Shears, L. L., Roelvink, P. W., Li, Y., Lee, G. M., Brough, D. E., Lizonova, A., and Kovesdi, I. (1997). Increased in vitro and in vivo gene transfer by adenovirus vectors containing chimeric fiber proteins. *J.Viro.* **71**, 8221-8229.
- Wiesner, R. J., Beinbrech, B., and Ruegg, J. C. (1993). Quantitative PCR. *Nature* **366**, 416.
- Wigler, M., Silverstein, S., Lee, L. S., Pellicer, A., Cheng, Yc, and Axel, R. (1977). Transfer of purified herpes virus thymidine kinase gene to cultured mouse cells. *Cell* **11**, 223-232.
- Williams, A. F. and Barclay, A. N. (1988). The immunoglobulin superfamily--domains for cell surface recognition. *Annu.Rev.Immunol.* **6:381-405.**, 381-405.
- Wold, W. S. and Gooding, L. R. (1991). Region E3 of adenovirus: a cassette of genes involved in host immunosurveillance and virus-cell interactions. *Virology* **184**, 1-8.
- Worgall, S., Wolff, G., Falck-Pedersen, E., and Crystal, R. G. (1997). Innate immune mechanisms dominate elimination of adenoviral vectors following in vivo administration. *Hum.Gene Ther.* **8**, 37-44.
- Xia, D., Henry, L. J., Gerard, R. D., and Deisenhofer, J. (1994). Crystal structure of the receptor-binding domain of adenovirus type 5 fiber protein at 1.7 Å resolution. *Structure*. **2**, 1259-1270.
- Xu, R. and Crowell, R. L. (1996). Expression and distribution of the receptors for coxsackievirus B3 during fetal development of the Balb/c mouse and of their brain cells in culture. *Virus Res.* **46**, 157-170.

Yarden, Y., Escobedo, J. A., Kuang, W. J., Yang-Feng, T. L., Daniel, T. O., Tremble, P. M., Chen, E. Y., Ando, M. E., Harkins, R. N., and Francke, U. (1986). Structure of the receptor for platelet-derived growth factor helps define a family of closely related growth factor receptors. *Nature* **323**, 226-232.

You, Z., Fischer, D. C., Tong, X., Hasenburg, A., Aguilar-Cordova, E., and Kieback, D. G. (2001). Coxsackievirus-adenovirus receptor expression in ovarian cancer cell lines is associated with increased adenovirus transduction efficiency and transgene expression. *Cancer Gene Ther.* **8**, 168-175.

Yukawa, H., Miyatake, S. I., Saiki, M., Takahashi, J. C., Mima, T., Ueno, H., Nagata, I., Kikuchi, H., and Hashimoto, N. (1998). In vitro growth suppression of vascular smooth muscle cells using adenovirus-mediated gene transfer of a truncated form of fibroblast growth factor receptor [In Process Citation]. *Atherosclerosis* **141**, 125-132.

Zabner, J., Couture, L. A., Gregory, R. J., Graham, S. M., Smith, A. E., and Welsh, M. J. (1993). Adenovirus-mediated gene transfer transiently corrects the chloride transport defect in nasal epithelia of patients with cystic fibrosis. *Cell* **75**, 207-216.

Zabner, J., Ramsey, B. W., Meeker, D. P., Aitken, M. L., Balfour, R. P., Gibson, R. L., Launspach, J., Moscicki, R. A., Richards, S. M., and Standaert, T. A. (1996). Repeat administration of an adenovirus vector encoding cystic fibrosis transmembrane conductance regulator to the nasal epithelium of patients with cystic fibrosis. *J.Clin.Invest.* **97**, 1504-1511.

Zhang, L. Q., Simmonds, P., Ludlam, C. A., and Brown, A. J. (1991). Detection, quantification and sequencing of HIV-1 from the plasma of seropositive individuals and from factor VIII concentrates. *AIDS* **5**, 675-681.

Zhou, L., Lim, L., Costa, R. H., and Whitsett, J. A. (1996). Thyroid transcription factor-1, hepatocyte nuclear factor-3beta, surfactant protein B, C, and Clara cell secretory protein in developing mouse lung. *J.Histochem.Cytochem.* **44**, 1183-1193.

Zimmermann, K. and Mannhalter, J. W. (1996). Technical aspects of quantitative competitive PCR. *Biotechniques* **21**, 268-269.

Zinn, K. R., Douglas, J. T., Smyth, C. A., Liu, H. G., Wu, Q., Krasnykh, V. N., Mountz, J. D., Curiel, D. T., and Mountz, J. M. (1998). Imaging and tissue biodistribution of 99mTc-labeled adenovirus knob (serotype 5). *Gene Ther.* **5**, 798-808.