

8 Literaturverzeichnis

Adams, J. C.; Tucker, R. P.; Lawler, J.: The Thrombospondin Gene Family. *R. G., Landes, Austin, Texas and Springer-Verlag*, 1995; S.195

Adams, R. H.; Wilkinson, G. A.; Weiss, C. et al.: Roles of ephrinB ligands and EphB receptors in cardiovascular development: demarcation of arterial / venous domains, vascular morphogenesis and sprouting angiogenesis. *Genes.Dev.*, 1999; 13: 295-306

Albo, D.; Arnoletti, J. P.; Castiglioni, A.; Granick, M. S.; Solomon, M. P.; Rothman, V. L.; Tuszynski, G. P.: Thrombospondin (TSP) and transforming growth factor beta 1 (TGF- β) promote human A549 lung carcinoma cell plasminogen activator inhibitor type 1 (PAI-1) production and stimulate tumor cell attachment in vitro. *Biochem.Biophys.Res.Comm.*, 1994; 203(1): 857-865

Arnoletti, J. P.; Albo, D.; Granick, M. S.; Solomon, M. P.; Castiglioni, A.; Rothman, V. L.; Tuszynski, G.P.: Thrombospondin and transforming growth factor beta 1 increase expression of urokinase-type plasminogen activator and plasminogen activator inhibitor-1 in human MDA-MB-231 breast cancer cells. *Cancer*, 1995; 76(6): 998-1005

Baenziger, N. L.; Brodie, G. N.; Majerus, P. W.: A thrombin-sensitive protein of human platelet membranes. *Proc.Natl.Acad.Sci.U.S.A*, 1971; 68: 240-249

Bagavandoss, P. et Wilks, J. W.: Specific inhibition of endothelial cell proliferation by thrombospondin. *Biochem.Biophys.Res.Comm.*, 1990; 170: 867-872

Bartling, B; Tostlebe, H.; Darmer, D; Holtz, J; Silber, R. E.; Morawietz, H.: Shear stress-dependent expression of apoptosis-regulating genes in endothelial cells. *Biochem.Biophys.Res.Comm.*, 2000; 278 (3): 740-746

Bhattacharya, S.; Michels, C. L.; Leung, M. K.; Arany, Z. P.; Kung, A. L.; Livingston, D. M.: Functional role of p35srj, a novel p300/CBP binding protein, during transactivation by HIF-1. *Genes Dev.*, 1999; 13: 64-75

Bauersachs, J., Popp, R., Hecker, M., Sauer, E., Fleming, I., und Busse, R.: Nitric oxide attenuates the release of endothelium-derived hyperpolarizing factor, *Circulation*, 1996; 94: 3341-3347

Bein, K.; Simons, M.: Thrombospondin type 1 repeats interact with matrix metalloproteinase 2. Regulation of metalloproteinase activity. *J.Biol.Chem.*, 2000; 275: 32167-32173

Belperio, J. A. ; Keane, M. P.; Arenberg, D. A. et al.: CXC chemokine in angiogenesis. *J.Leukocyte.Biol.*, 2000; 68: 1-8

- BenEzra, D.; Griffin, B. W.; Maftzir, G.; Aharonov, O.: Thrombospondin and *in vivo* angiogenesis induced by basic fibroblast growth factor or lipopolysaccharide. *Invest. Ophthalmol. Vis. Sci.*, 1993; 34: 3601-3608
- Berthiaume, F. et Frangos, J. A.: Flow-induced prostacyclin production is mediated by a pertussis toxin-sensitive G protein. *FEBS Lett.*, 1992; 308: 272- 279
- Bongrazio, M.; Zakrzewicz, A.; Knöchel, J.; Baumann, C.; Gaehtgens, P.; Pries, A.R.: Differential expression of thrombospondin repeat type1-containing proteins in flow exposed endothelial cells. *Eur. Heart J.*, 2000; 21, Abstr. Suppl.: 359.
- Bongrazio, M.; Baumann, C.; Zakrzewicz, A.; Pries, A.R.; Gaehtgens, P.: Evidence for modulation of genes involved in vascular adaptation by prolonged exposure of endothelial cells to shear stress. *Cardiovasc. Res.*, 2000; 47: 384-393.
- Bongrazio, M.; Pries, A.R.; Zakrzewicz, A.: The endothelium as physiological source of proendothelin: role of wall shear stress. *Mol. Immunol.*, 2003; 39: 669-675.
- Bornstein, P.: Thrombospondins as matricellular modulators of cell function. *J. Clin. Invest.*, 2001; 107: 929-934
- Braddock, M.; Schwachtgen, J. L.; Houston, P.; Dickson, M. C.; Lee, M. J.; Campbell, C. J.: Fluid shear stress modulation of gene expression in endothelial cells. *News Physiol. Sci.*, 1998; 13: 241-246
- Brew, K.; Dinakarandian, D.; Nagase, H.: Tissue inhibitors of metalloproteinases: evolution, structure and function. *Biochem. Biophys. Acta*, 2000; 1477: 267-283
- Brooks, P. C.; Clark, R.; Cheresh, D. A.: Requirement of vascular integrin $\alpha_v\beta_3$ for angiogenesis. *Science*, 1994a; 264: 569-571
- Brooks, P. C.; Montgomery, A. M. P., Rosenfeld, M.; Reisfeld, R. A.; Hu, T.; Klier, G.; Cheresh, D. A.: Integrin $\alpha_v\beta_3$ antagonists promote tumor regression by inducing apoptosis of angiogenic blood vessels. *Cell*, 1994b; 79: 5-17
- Buri, P. H. et Djonov, V.: Intussusceptive angiogenesis – the alternative to capillary sprouting. *Molecular Aspects of Medicine*, 2002; 23: S2-S27
- Buschmann, I. et Schaper, W.: Arteriogenesis versus Angiogenesis: Two mechanisms of vessel growth. *News Physiol. Sci.*, 1999; 14: 121-125
- Canfield, A. E.; Boot-Hanford, R. P.; Schor, A. M.: Thrombospondin gene expression by endothelial cells in culture is modulated by cell proliferation, cell shape and the substratum. *Biochem. J.*, 1990; 268: 225-230

Carmeliet, P.; Mackman, N; Moons,L.; Luther, T.; Gressens, P.; Van Vlaenderen, I.; Demunck, H.; Kasper, M.; Breier, G.; Evrard, P.; Muller; M.; Risau, W.; Edgington, T.; Collen, D.: Role of tissue factor in embryonic blood vessel development. *Nature*, 1996; 383 (6595): 73-75

Carmeliet, P.; Dor, Y.; Herbert, J. M. et al.: Role of HIF-alpha in hypoxia-mediated apoptosis, cell proliferation and tumor angiogenesis. *Nature*, 1998; 394: 485-490

Carmeliet, P.: Mechanisms of angiogenesis and arteriogenesis. *Nat.Med.*, 2000; 6: 389-395

Carmeliet, P.; Jain, R. K.: Angiogenesis in cancer and other diseases. *Nature*, 2000; 407: 249-257

Castle, V.; Varani, J.; S.; Fligel, S.; Prochownik, E. V.; Dixit, V.: Antisense-mediated reduction in thrombospondin reverses the malignant phenotype of a human squamous carcinoma. *J.Clin.Invest.*, 1991; 87: 1883-1888

Chen, D.; Asahara, T.; Krasinski, K.; Witzenbilcher, B.; Yang, J.; Magner, M.; Kearney, M.; Frazier, W. A.; Isner, J. M.; Andres, V.: Antibody blockade of thrombospondin accelerates reendothelialization and reduces neointima formation in balloon-injured rat carotid artery. *Circulation*, 1999; 100 (8): 849-854

Chun, T. H.; Itoh, H.; Ogawa, Y.; Tamura, N.; Takaya, K.; Igaki, T.; Yamashita, J.; Doi, K.; Inoue, M.; Masatsugu, K.; Korenaga, R.; Ando, J.; Nakao, K.: Shear stress augments expression of C-type natriuretic peptide and adrenomedullin. *Hypertension*, 1997; 29: 1296-1302

Cohen, R. A. et Vanhoute, P. M.: Endothelium-dependent hyperpolarization. Beyond nitric oxid and cyclic GMP, *Circulation*, 1995; 92: 3337-3349

Conway, E. M.; Collen, D.; Carmeliet, P.: Molecular mechanisms of blood vessel growth. *Cardiovasc.Res.*49, 2001; 507-521

Cooke, J., Rossitch, E., Andon, N., Loscalzo, J.,und Dzau, V.: Flow activates an endothelial potassium channel to release an endogenous nitrovasodilator , *J.Clin.Invest.*, 1991; 88: 1663-1671

Crawford, S. E.; Stellmach, V.; Murphy-Ullrich, J. E.; Ribeiro, S. M. F.; Lawler, J.; Hynes, R. O.; Boivin, G. P.; Bouck, N: Thrombospondin-1 is a major activator of TGF- β 1 in vivo. *Cell*, 1998; 93: 1159-1170

Dameron, K. M.; Volpert, O. V.; Tainsky, M. A.; Bouck, N.: Control of angiogenesis in fibroblasts by p53 regulation of thrombospondin-1. *Science*, 1994; 265: 1582-1584

Davies, S. et Yancopoulos, G. D.: The angiopoietins: yin and yang in angiogenesis. *Curr.Top.Microbiol.Immunol.*, 1999; 237: 173-185

Dawson, D. W.; Pearce, S. F. A.; Zhong, R.; Silverstein, R. L.; Frazier, W. A.; Bouck, N.: CD36 mediates the in vitro inhibitory effects of thrombospondin-1 on endothelial cells. *J.Cel.Biol.*, 1997; 183: 707-717

Dawson, D. W. et Bouck, N. P.: Thrombospondin as an inhibitor of angiogenesis. *Antiangiogenetic Agents in Cancer Therapy*, edited by Teicher, B. A.; Totowa N. J.: Human Press Inc., 1999: 185-203

Diamond, S. L.; Sharefkin, J. B.; Dieffenbach, C.; Frasier-Scott, K.; McIntire, L. V.; Eskin, S. G.: Tissue plasminogen activator messenger RNA levels increase in cultured human endothelial cells exposed to laminar shear stress. *J.Cell.Physiol.*, 143(2): 364

DiPietro, L. A. et Polverini, P. J.: Angiogenic macrophages produce the angiogenic inhibitor thrombospondin-1. *Am.J.Pathol.*, 143: 678-684

Drake, C. J.; Davis, L. A.; Little, C. D.: Antibodies to β_1 -Integrins cause alterations of aortic vasculogenesis, in vivo. *Dev.Dyn.*, 1992;193: 83-91

Ebert, B. L.; Firth, J. D.; Ratcliffe, P. J.: Hypoxia and mitochondrial inhibitors regulate expression of glucose transporter-1 *via* distinct *cis*-acting sequences. *J.Biol.Chem.*, 1995; 270: 29083-29089

Eckhart, A. D.; Yang, N.; Xin, X.; Faber, J. E.: Characterization of the α_{1B} -adrenergic receptor gene promoter region and hypoxia regulatory elements in vascular smooth muscle. *Proc.Natl.Acad.Sci. USA*, 1997; 94: 9487-9492

Edgell, C. J.; McDonald, C. C.; Graham, J. B.: Permanent cell line expressing human factor VIII-related antigen established by hybridization. *Proc.Natl.Acad.Sci.USA*, 1983; 80(12): 3734-3737

Egginton, S.; Zhou, A.-L.; Brown, M. D.; Hudlicka, O.: Unorthodox angiogenesis in skeletal muscle. *Cardiovasc.Res.*, 2001; 49: 634-646

Estes, S. D.; Stoler, D. L.; Anderson, G. R.: Anoxic induction of a sarcoma virus-related VL30 retrotransposon is mediated by a *cis*-acting element which binds hypoxia-inducible factor 1 and an anoxia-inducible factor. *J.Virol.*, 1995; 69: 6335-6341

Febbraio, M.; Hajjar, D. P.; Silverstein, R. L.: CD36: A class B scavenger receptor involved in angiogenesis, atherosclerosis, inflammation, and lipid metabolism. *J.Clin.Invest.*, 2001; 108: 785-791

Ferrara, N.; Houck, K.; Jakeman, L.; Leung, D.W.:Molecular and biological properties of the vascular endothelial growth factor family of proteins. *Endocr.Rev.*,1992; 13:18-32

Firth, J. D.; Ebert, B. L.; Pugh, C. W.; Ratcliffe, P. J.: Oxygen-regulated control elements in the phosphoglycerat kinase 1 and lactate dehydrogenase A genes: similarities with the erythropoietin 3' enhancer. *Proc.Nat.Acad.Sci. USA*, 1994; 91: 6496-6500

Firth, J. D.; Ebert, B. L.; Ratcliffe, P. J.: Hypoxic regulation of lactate dehydrogenase A. Interaction between hypoxia-inducible factor 1 and cyclic AMP response elements. *J.Biol.Chem.*, 1995; 270: 21021-21027

Folkman, J. et Shing, Y.: Control of angiogenesis by heparin and other sulfated polysaccharides. *Adv.Exp.Med.Biol.*, 1992; 313: 355-364

Fong, G. H.; Rossant, J.; Gerstenstein, M.; Breitman, M. L.: Role of Flt-1 receptor tyrosine kinase in regulating the assembly of vascular endothelium. *Nature*, 1995; 376: 66-70

Forsythe, J. A.; Jiang, B. H.; Iyer, N. V.; Agani, F.; Leung, S. W.; Koos, R. D.; Semenza, G. L.: Activation of vascular endothelial growth factor gene transcription by hypoxia-inducible factor-1. *Mol.Cell. Biol.*, 1996; 16: 4604-4613

Frater-Schröder, M.; Risau, W.; Hallmann, R.; Gautschi, P.; Böhlen, P.: Tumor necrosis factor type α , a potent inhibitor of endothelial cell growth in vitro, is angiogenic in vivo. *P.N.A.S. USA*, 1987; 84: 5277-5281

Freyberg, M. A.; Kaiser, D.; Graf, R.; Vischer, P.; Friedl, P.: Integrin-associated protein and thrombospondin-1 as endothelial mechanosensitive death mediators. *Biochem.Biophys.Res.Comm.*, 2000; 271(3): 584-588

Freyberg, M. A.; Kaiser, D.; Graf, R.; Buttenbender, J.; Friedl, P.: Proatherogenic flow conditions initiate endothelial apoptosis via thrombospondin-1 and the integrin-associated protein. *Biochem.Biophys.Res.Comm.*, 2001 a; 286(1): 141-149

Feyberg, M. A.; Kaiser, D.; Graf, R.; Friedl, P.: Vascular endothelial cells express a functional fas-receptor due to lack of hemodynamic forces. *Apoptosis*, 2001 b; 6(5): 339-343

Frazier, W. A.; Gao, A. G.; Dimitry, J.; Chung, J.; Brown, E. J.; Lindberg, F. P.; Linder, M. E.: The thrombospondin receptor integrin-associated protein (CD47) functionally couples to heterotrimeric Gi. *J.Biol.Chem.*, 1999; 274: 8554-8560

Friedlander, M.; Brooks, P. C.; Shaffer, R. W.; Kincaid, C.M.; Varner, J. A.; Cheresch, D. A.: Definition of two angiogenetic pathways by distinct α_v integrins. *Science*, 1995; 270: 1500-1502

Gahtan, V.; Wang, X. J; Willis, A. I.; Tuszynski, G. P.; Sumpio, B. E.: Thrombospondin-1 regulation of smooth muscle cell chemotaxis extracellular signal-regulated protein kinases 1/2 dependent. *Surgery*, 1999; 126: 203-207

- Gahtan, V.; Wang, X. J.; Ikeda, M.; Willis, A. I.; Tuszynski, G. P.; Sumpio, B. E.: Thrombospondin-1 induces activation of focal adhesion kinase in vascular smooth muscle cells. *J.Vasc.Surg.*, 1999; 29: 1031-1036
- Gendron, R. L.; Adams, L. C.; Paradis, H.: Tubedown-1, a novel acetyltransferase associated with blood vessel development. *Dev.Dyn.*, 2000; 218: 300-315
- Gentz, R.; Rauscher, F. J.; Abate, C.; Curran, T.: Parallel association of Fos and Jun leucine zippers juxtaposes DNA binding domains. *Science*, 1989; 243: 1695-1699
- Gerber, H. P.; Condorelli, F.; Park, J.; Ferrara, N.: Differential transcriptional regulation of two vascular endothelial growth factor receptor genes. Flt-1, but not Flk-1/KDR, is up-regulated by hypoxia. *J.Biol.Chem.*, 1997; 272: 23659-23667
- Good, D. J., Polverini, P. J., Rastinejad, F., LeBeau, M. M., Lemons, R. S., Frazier, W. A., Bouck, N. P.: A tumor suppressor-dependent inhibitor of angiogenesis is immunologically and functionally indistinguishable from a fragment of thrombospondin. *Proc.Natl.Acad.Sci.*, 1990; 87: 6624-6628
- Gong, W.; Hao, B.; Mansy, S. S.; Gonzalez, G.; Gilles-Gonzalez, M. A.; Chan, M. K.: Structure of a biological oxygen sensor: a new mechanism for heme-driven signal transduction. *Proc.Natl.Acad.Sci.USA*, 1998; 95: 15177-15182
- Graven, K. K.; Yu, Q.; Pan, D.; Roncarati, J. S.; Farber, H. W.: Identification of an oxygen responsive enhancer element in the glyceraldehyde-3-phosphate dehydrogenase gene. *Biochem.Biophys.Acta*, 1999; 1447: 208-218
- Greenwalt, D. E., Lipsky, R. H., Ockenhouse, C. F., Ikeda, H., Tandon, H. H., Jamieson, G. A.: Membrane glycoprotein CD36: A review of its roles in adherence, signal transduction, and transfusion medicine. *Blood*, 1992; 80: 1005-1015
- Grossfeld, G. D.; Ginsberg, D. A.; Stein, J. P.; Bochner, B. H.; Esrig, D.; Groshen, S.; Dunn, M.; Nichols, P. W.; Taylor, C. R.; Skinner, D. G.; Cote, R. J.: Thrombospondin-1 expression in bladder cancer: association with p53 alterations, tumor angiogenesis, and tumor progression. *J.Natl.Cancer Inst.*, 1997; 89(3): 219-227
- Guo, N.; Krutzsch, H. C.; Inman, J. K.; Roberts, D. D.: Thrombospondin 1 and type I repeat peptides of thrombospondin 1 specifically induce apoptosis of endothelial cells. *Cancer Res.*, 1997; 57(9): 1735-1742
- Hanahan, D.: Signaling vascular morphogenesis and maintenance. *Science*, 1997; 277: 48-50
- Hiraoka, N.; Allen, E.; Apel, E. J.; Gyetko, M. R. and Weiss, S. J.: Matrix metalloproteinases regulate neovascularization by acting as pericellular fibrinolysins. *Cell*, 1998; 95: 365-377

- Honda, H. M.; Wortham, C.; Navab, M.; Demer, L. L.: Disturbed flow induces heat shock protein-70 mRNA in bovine and human aortic endothelial cells. *Circulation*, 1992; 86: Suppl.1/4: 224
- Hsieh, H.-J.; Li, N.-Q.; Frangos, J. A.: Shear stress increases endothelial platelet-derived growth factor mRNA levels. *Am.J.Phys.*, 1991; 260: 642-644
- Hsieh, H. -J.; Li, N. Q.; Frangos, J. A.: Pulsatile and steady flow induces c-fos expression in human endothelial cells. *J.Cell Physiol.*, 1993; 154: 143-151
- Hu, J.; Discher, D. J.; Bishopric, N. H.; Webster, K. A.: Hypoxia regulates expression of the endothelin-1 gene through a proximal hypoxia-inducible factor-1 binding site on the antisense strand. *Biochem.Biophys.Res.Commun.*, 1998; 245: 894-899
- Hudlicka, O.; Brown, M.; Egginton, S.: Angiogenesis in skeletal and cardiac muscle. *Physiol.Rev.*, 1992; 72: 369-417
- Ilan, M.; Mahooti, S.; Rimm, D.L.; Madri, J. A.: PECAM-1 (CD31) functions as a reservoir for and a modulator of tyrosine-phosphorylated β -catenin. *J.Cell.Sci.*, 1999; 112(18): 3005-3014
- Ingber, D. E.; Dike, L.; Hansen, L.; Karp, S.; Liley, H.; Maniotis, A.; McNamee, H.; Mooney, D.; Plopper, G.; Sims, J.: Cellular tensegrity: exploring how mechanical changes in the cytoskeleton regulate cell growth, migration, and tissue pattern during morphogenesis. *Int.Rev.Cytol.*, 1994; 150: 173-224
- Iruela-Arispe, M. L.; Bornstein, P.; Sage, H.: Thrombospondin exerts an antiangiogenic effect on cord formation by endothelial cells *in vitro*. *Proc.Natl.Acad.Sci. USA*, 88: 5026-5030
- Iruela-Arispe, M. L.; Porter, P.; Bornstein, P.; Sage, E. H.: Thrombospondin-1, an Inhibitor of Angiogenesis, is regulated by Progesterone in the Human Endometrium. *J.Clin.Invest.*, 1996; 97: 403-412
- Ishida, T.; Takahashi, M.; Corson, M. A.; Berk, B. C.: Fluid shear stress-mediated signal transduction: how do endothelial cells transduce mechanical force into biological responses? *Ann.N.Y.Acad.Sci.*, 1997; 811: 12-13
- Iyer, N. V.; Kotch, L. E.; Agani, F.; Leung, S. W.; Laughner, E.; Wenger, R. H.; Gassmann, M.; Gearhart, G. D.; Lawler, A. M.; Yu, A. J.; Semenza, G. L.: Cellular and developmental control of O₂-homeostasis by hypoxia-inducible factor 1 α . *Genes Dev.*, 1998; 12: 149-162
- Jaffe, E. A.; Nachman, R. L.; Becker, C. G.; Minick, R. C.: Culture of human endothelial cells derived from umbilical veins. Identification by morphologic and immunologic criteria. *J.Clin.Invest.*, 1973; 52: 2745-2756

Jimenez, B.; Volpert, O. V.; Crawford, S. E.; Febbraio, M.; Silverstein, R. L.; Bouck, N.: Signals leading to apoptosis-dependent inhibition of neovascularization by thrombospondin-1. *Nat.Med.*, 2000; 6: 41-48

Kaiser, D.; Freyberg, M. A.; Schrimpf, G.; Friedl, P.: Apoptosis induced by lack of hemodynamic forces is a general endothelial feature even occurring in immortalized cell lines. *Endothelium*, 1999; 6(4): 325-334

Kanda, S.; Shono, T.; Tomasini-Johansson, B.; Klint, P.; Saito, Y.: Role of thrombospondin-1 derived peptide, 4NIK, in FGF-2 induced angiogenesis. *Exp.Cell.Res.*, 1999; 252: 262-272

Khachigian, L. M.; Anderson, K. R.; Halnon, N. J.; Gimbrone, M. A. Jr.; Resnick, N.; Collins, T.: Egr-1 is activated in endothelial cells exposed to fluid shear stress and interacts with a novel shear-stress-response element in the PDGF-A chain promoter. *Arterioscler.Thromb.Vasc.Biol.*, 1997; 17 : 2280-2286

Kietzmann, T.; Roth, U.; Jungermann, K.: Induction of the plasminogen activator inhibitor-1 gene expression by mild hypoxia *via* a hypoxia response element binding the hypoxia-inducible factor-1 in rat hepatocytes. *Blood*, 1999; 94: 4177-4185

Kimura, H., Weisz, A., Kurashima, Y.: Hypoxia response element of human vascular endothelial growth factor gene mediates transcriptional regulation by nitric oxide; control of hypoxia-inducible factor-1 activity by nitric oxide. *Blood*, 2000; 95: 189-197

Knedler, A. et Ham, R. G.: Optimized medium for clonal growth of human microvascular endothelial cells with minimal serum. *In Vitro Cell Dev.Biol.*, 1987; 23 (7): 481-491

Knudsen, K. A.; Frankowski, C.; Johnson, K. R.; Wheelock, M. J.: A role for cadherins in cellular signaling and differentiation. *J.Cell.Biochem.Suppl.*, 1998; 31: 168-176

Koch, A. E.; Polverini, P. J; Kunkel, S. L.; Harlow, L. A.; DiPietro, L. A.; Elnor, V. M.; Elnor, S. G.; Strieter, R. M.: Interleukin-8 as a macrophage-derived mediator of angiogenesis. *Science*, 1992; 258 (5089): 1798-1801

Kosaki, K.; Ando, J.; Korenaga, R.; Kurokawa, T.; Kamiya, A.: Fluid Shear Stress increases the production of Granulocyte-Macrophage Colony-Stimulating Factor by endothelial cells *via* mRNA stabilization. *Circ.Res.*, 1998; 82: 794-802

Kotch, L. E.; Iyer, N. V.; Laughner, E.; Semenza, G. L.: Defective vascularization of HIF-1 α -embryos is not associated with VEGF deficiency but with mesenchymal cell death. *Dev.Biol.*, 1999; 209: 254-267

Lan, Q.; Mercurius, K. O.; Davies, P. F.: Stimulation of transcription factors NF kappa B and AP 1 in endothelial cells subjected to shear stress. *Biochem.Biophys.Res.Comm.*, 1994; 201: 950-956

- Lawler, J.; Sunday, M.; Tilbert, V.; Duquette, M.; George, E. L.; Rayburn, H.; Hynes, R. O.: Thrombospondin-1 is required for normal murine pulmonary homeostasis and its absence causes pneumonia. *J.Clin.Invest.*, 1998; 101: 982-992
- Lawler, J.: The functions of thrombospondin-1 and 2. *Curr.Opin.Cell.Biol.*, 2000; 12: 634-640
- Lee, P. J.; Jiang, B. H.; Chin, B. Y.; Iyer, N. V.; Alam, J.; Semenza, G. L.; Choi, A. M.: Hypoxia-inducible factor-1 mediates transcriptional activation of the heme oxygenase-1 gene in response to hypoxia. *J.Biol.Chem.*, 1997; 272: 5375-5381
- Lesoon-Wood, L. A.; Kim, W. H.; Kleinman, H. K.; Weintraub, B. D.; Mixon, A. J.: Systemic gene therapy with p53 reduces growth and metastasis of a malignant human breast cancer in nude mice. *Hum.Gene Ther.*, 1995; 6: 395-405
- Lelièvre, E.; Lionneton, F.; Soncin, F.; Vandebunder, B.: The Ets family contains transcriptional activators and repressors involved in angiogenesis. *Int.J.Biochem.Cell.Biol.*, 2001; 33: 391-407
- Li, Y. S.; Shyy, J. Y.; Li, S.; Lee, J.; Su, B.; Karin, M.; Chien, M.: The Ras-JNK pathway is involved in shear stress-induced gene expression. *Mol.Cell.Biol.*, 1996; 16(11): 5947
- Lin, M. C.; Almus-Jacobs, F.; Chen, H. H.; Parry, G. C. N.; Mackman, N.; Shyy, J. Y. J.; Chien, S.: Shear stress induction of the tissue factor gene. *J.Clin.Invest.*, 1997; 99: 737
- Lindahl, P.; Bostrom, H.; Karlsson, L.; Hellstrom, M.; Kalen, M.; Betsholtz, C.: Role of platelet -derived growth factors in angiogenesis and alveogenesis. *Curr.Top.Pathol.*, 1999; 93: 27-33
- Liu, Y.; Cox, S. R.; Morita, T.; Kourembanas, S.: Hypoxia regulates vascular endothelial growth factor gene expression in endothelial cells. Identification of a 5' enhancer. *Circ.Res.*, 1995; 77: 638-643
- Lok, C. N. et Ponka, P.: Identification of a hypoxia response in the transferrin receptor gene. *J.Biol.Chem.*, 1999; 274: 24147-24152
- Lymn, J. S.; Rao, S. J.; Clunn, G. F.; Gallagher, K. L.; O'Neil, C. O.; Thompson, N. T.; Hughes, A. D.: Phosphatidylinositol 3-kinase and focal adhesion kinase are early signals in the growth factor-like responses to thrombospondin-1 seen in human vascular smooth muscle. *Arterioscler.Thromb.Vasc.Biol.*, 1999; 19: 2133-2140
- Maeda, K.; Nishiguchi, Y.; Yashiro, M.; Yamada, S.; Onoda, N.; Sawada, T.; Kang, S. M.; Hirakawa, K.: Expression of vascular endothelial growth factor and thrombospondin-1 in colorectal carcinoma. *Int.J.Mol.Med.*, 2000; 5(4): 373-378

- Maione, T. E.; Gray, G. S.; Petro, G.; Hunt, A. J.; Donner, A. L.; Bauer, S. I.; Carson, H. F.; Sharpe, R. J.: Inhibition of angiogenesis by recombinant human platelet factor-4 and related peptides. *Science*, 1990; 247 (4938): 77-79
- Maisonpierre, P. C.; Suri, C.; Jones, P. F. et al.: Angiopoietin-2, a natural antagonist for Tie-2 that disrupts in vivo angiogenesis. *Science*, 1997; 277: 55-60
- Majack, R. A.; Goodman, L. V.; Dixit, V. M.: Cell surface thrombospondin is functionally essential for vascular smooth muscle cell proliferation. *J.Cell.Biol.*, 106: 415-422
- Majack, R. A.; Majesky, M. W.; Goodman, L. V.: Role of PDGF-A expression in the control of vascular smooth muscle cell growth by transforming growth factor-beta. *J.Cell.Biol.*, 1990; 111: 239-247
- Malek, A. et Izumo, S.: Physiological fluid shear stress causes downregulation of endothelin-1 mRNA in bovine aortic endothelium. *Am.J.Phys.*, 1992; 263: C389-C396
- Malek, A.; Gibbons, G. H.; Dzau, V. J.; Izumo, S.: Fluid shear stress differentially modulates expression of genes encoding basic fibroblast growth factor and platelet-derived growth factor B chain in vascular endothelium. *J.Clin.Invest.*, 1993a; 92(4): 2013
- Malek, A. M.; Jackman, R.; Rosenberg, R. D.; Izumo, S.: Endothelial expression of thrombomodulin is reversibly regulated by fluid shear stress. *Circ.Res.*, 1994; 74: 852-860
- Mansfield, P. J. et Suchard, S. J.: Thrombospondin promotes chemotaxis and haptotaxis of human peripheral blood monocytes. *J.Immunol.*, 1994; 153: 4219-4229
- Mateo, V.; Lagneaux, L.; Bron, D.; Biron, G.; Armant, M.; Delespesse, G.; Sarfati, M.: CD47 ligation induces caspase-independent cell death in chronic lymphocytic leukemia. *Nat.Med.*, 1999; 5: 1277-1284
- Maxwell, P. H.; Wiesener, M. S.; Chang, G. W.; Clifford, S. C.; Vaux, E. C.; Cockman, M. E.; Wykoff, C. C.; Pugh, C. W.; Maher, E. R.; Ratcliffe, P. J.: The tumour suppressor protein VHL targets hypoxia-inducible factors for oxygen-dependent proteolysis. *Nature*, 1999; 399: 271-275
- Melillo, G.; Musso, T.; Sica, A.; Taylor, L. S.; Cox, G. W.; Varesio, L.: A hypoxia-responsive element mediates a novel pathway of activation of the inducible nitric oxide synthase promoter. *J.Exp.Med.*, 1995; 182: 1683-1693
- Metivier, F.; Marchais, S. J.; Guerin, A. P.; Pannier, B.; London, G. M.: Pathophysiology of anaemia: focus on the heart and blood vessels. *Nephrol.Dial.Transplant.*, 2000; 15 Suppl. 3: 14-18

Mettouchi, A.; Cabon, F.; Montreau, N.; Vernier, P.; Mercier, G.; Bangy, D.; Tricoire, H.; Vigier, P.; Binetruy, B.: SPARC and thrombospondin genes are repressed by the c-jun oncogene in rat embryo fibroblasts. *EMBO.J.*, 1994; 13: 5668-5578

Milkiewicz, M.; Brown, M. D.; Egginton, S.; Hudlicka, O.: Association between shear stress, angiogenesis, and VEGF in skeletal muscle in vivo. *Microcirculation*, 2001; 8(4): 229-241

Moncada, S., Gryglewski, R. Bunting, S., und Vane, J. R.: An enzyme isolated from arteries transforms prostaglandin endoperoxides to an unstable substance that inhibits platelet aggregation, *Nature*, 21-10-76; 263, 663-665

Morandi, V.; Fauvel-Lafeve, F.; Legrand, C.; Legrand, Y. J.: Role of thrombospondin in the adhesion of human endothelial cells in primary culture. *In Vitro Cell.Dev.Biol.*, 1993; 29 A: 585-591

Morawietz, H.; Talanow, R.; Szibor, M.; Rueckschloss, U.; Schubert, A.; Bartling, B.; Darmer, D.; Holtz, J.: Regulation of the endothelin system by shear stress in human endothelial cells. *J.Physiol.*, 2000; 525 Pt 3: 761-770

Murphy-Ullrich, J. E. et Hook, M.: Thrombospondin modulates focal adhesions in endothelial cells. *J.Cell.Biol.*, 1989; 109: 1309-1319

Murphy-Ullrich, J. E.; Gurusiddappa, S.; Frazier, W. A.; Höök, M.: Heparin-binding peptides from thrombospondin 1 and 2 contain focal adhesion-labilizing activity. *J.Biol.Chem.*, 1993; 268: 26784-26789

Nagel, T.; Atkinson, W. J.; Dewey, C. F.; Gimbrone, M. A.: Shear stress selectively upregulates intercellular adhesion molecule-1 expression in cultured human vascular endothelial cells. *J.Clin.Invest.*, 1994; 94: 885-891

Nesselroth, S. M.; Willis, A. I.; Fuse, S.; Olson, E. T.; Lawler, J.; Sumpio, B. E.; Gahtan, V.: The C-terminal domain of thrombospondin-1 induces vascular smooth muscle cell chemotaxis. *J.Vasc.Surg.*, 2001; 33(3): 595-600

Nguyen, S. V. et Claycomb, W. C.: Hypoxia regulates the expression of the adrenomedullin and HiF-1 genes in cultured HL-1 cardiomyocytes. *Biochem.Biophys.Res.Comm.*, 1999; 265: 382-386

Nicosia, R. F.; Tuszynski, G. P.: Matrix-bound thrombospondin promotes angiogenesis *in vitro*. *J.Cell Biol.*, 1994; 124: 183-193

Nor, J. E.; Christensen, J.; Mooney, D. J.; Polverini, P. J.: Vascular endothelial growth factor (VEGF-) mediated angiogenesis is associated with enhanced endothelial cell survival and induction of Bcl-2-expression. *Am.J.Pathol.*, 1999; 154(2): 375-384

- Nor, J. E.; Mitra, R. S.; Sutorik, M. M.; Mooney, D. J.; Castle, V. P.; Polverini, P. J.: Thrombospondin-1 induces endothelial cell apoptosis and inhibits angiogenesis by activating the caspase death way. *J.Vasc.Res.*, 2000; 37(3): 209-218
- Ohno, M.; Cooke, J. P.; Dzau, V. L.; Gibbons, G. H.: Fluid shear stress induces endothelial transforming growth factor beta-1 transcription and production. Modulation by potassium channel blockade. *J.Clin.Invest.*, 1995; 95: 1363-1369
- Ohtsuka, A.; Ando, J.; Korenaga, R.; Kamiya, A.; Toyama-Sorimachi, N.; Miyasaka, M.: The effect of flow on the expression of vascular adhesion molecule-1 by cultured mouse endothelial cells. *Biochem.Biophys.Res.Comm.*, 1993; 193: 303-310
- Okahara, K.; Sun, B.; Kawasaki, T.; Monden, M.; Kambayashi, J.: Expression of platelet-activating factor receptor transcript-2 is induced by shear stress in HUVEC. *Prostaglandins Other Lipid Mediat.*, 1998; 55: 323-329
- Olesen, S. P.; Clapham, D. E.; Davies, P. F.: Haemodynamic shear stress activates a K^+ current in vascular endothelial cells. *Nature*, 1988; 331: 168-170
- O'Reilly, M. S.; Holmgren, L.; Shing, Y. et al.: Angiostatin: a novel angiogenesis inhibitor that mediates the suppression of metastases by a Lewis lung carcinoma. *Cell*, 1994; 79:315-328
- O'Reilly, M. S.; Boehm, T.; Shing, Y. et al.: Endostatin: an endogenous inhibitor of angiogenesis and tumor growth. *Cell*, 1997; 88: 277-285
- Palmer, L. A.; Semenza, G. L.; Stoler, M. H.; Johns, R. A.: Hypoxia induces type II NOS gene expression in pulmonary artery endothelial cells *via* HIF-1. *Am.J.Physiol.*, 1998; 274: L 212-L 219
- Pepper, M. S.; Transforming growth factor-beta: vasculogenesis, angiogenesis and vessel wall integrity. *Cytokine Growth Factor Rev.*, 1997; 8: 21-43
- Phan, S. H.; Dillon, R. G.; McGarry, B. M.; Dixit, V. M.: Stimulation of fibroblast proliferation by thrombospondin. *Biochem.Biophys.Res.Comm.*, 1989; 163: 56-63
- Phelan, M. W.; Forman, L. W.; Perrine, S. P.; Faller, D. V.: Hypoxia increases thrombospondin-1 transcript and protein in cultured endothelial cells. *J.Lab.Clin.Med.*, 1998; 132(6): 519-529
- Pries, A. R.; Reglin, B.; Secomb, T. W.: Structural adaption of microvascular networks: functional roles of adaptive responses. *Am.J.Physiol.Heart Circ.Physiol.*, 2000; 281: H1015-H1025
- Qian, X.; Nicosia, R. F.; Bochenek, K. M.; Rothman, V. L.; Tuszynski, G. P.: The effect of thrombospondin on endothelial cell tube formation *in vitro*. *Mol.Biol.Cell* 5, 1994; 179 a

- Qian, X.; Wang, T. N.; Rothman, V. L.; Nicosia, R. F.; Tuszynski, G.P.: Thrombospondin-1 modulates angiogenesis in vitro by up-regulation of matrix metalloproteinase-9 in endothelial cells. *Exp.Cell.Res.*, 1997; 235: 403-412
- Raugi, G. J., Olerud, J. E., Gown, A. M. : Thrombospondin in early human wound tissue. *J.Invest.Dermatol.*, 1987 ; 89 : 551-554
- RayChaudhury, A. ; Frazier, W. A. ; D`Amore, P. A. : Comparison of normal and tumorigenic endothelial cells : Differences in thrombospondin production and responses to transforming growth factor-beta. *J.Cell Sci.*, 1994 ; 107 : 39-46
- Resnick, N.; Collins, T.; Atkinson, W.; Bonthron, D. T.; Dewey, C. F. Jr.; Gimbron, M. A. Jr.: Platelet-derived growth factor B chain promoter contains a cis-acting fluid shear-stress-responsive element. *Proc.Natl.Acad.Sci. U.S.A.*, 1993 ; 90 : 7908
- Richard, D. E.; Berra, E.; Gothie, E.; Roux, D.; Pouyssegur, J.: p42/p44 mitogen-activated protein kinases phosphorylate hypoxia-inducible factor 1 α (HIF-1 α) and enhance the transcriptional activity of HIF-1. *J.Biol.Chem.*, 1999; 274: 32631-32637
- Rieder, M. J.; Carmona, R.; Krieger, J. E.; Pritchard, K. A.; Greene, A. S.: Suppression of angiotensin-converting enzyme expression and activity by shear stress. *Circ.Res.*, 1997; 80: 312-319
- Rieger, H. et Gaethgens, P.: Pathophysiologie der Gewebsdurchblutung und ihre Regulation: Durchblutungsstörungen in der Pathophysiologie des Menschen, Chapman & Hall, 1994; 17.3.
- Risau, W.: Mechanisms of angiogenesis. *Nature*, 1997; 386(6626): 671-674
- Rolfs, A.; Kvietikova, I.; Gassmann, M.; Wenger, R. H.: Oxygen-regulated transferrin expression is mediated by hypoxia-inducible factor-1. *J.Biol.Chem.*, 1997; 272: 20055-20062
- Ryan, H. E.; Lo, J.; Johnson, R. S.: HIF-1 α is required for solid tumor formation and embryonic vascularization. *EMBO.J.*, 1998; 17: 3005-3015
- Sage, E. H. et Bornstein, P.: Extracellular proteins that modulate cell-matrix interactions: SPARC, Tenascin and Thrombospondin. *J.Biol.Chem.*, 1991; 266: 14831-14834
- Sargiannidou, I.; Zhou, J.; Tuszynski, G. P.: The role of thrombospondin-1 in tumor progression. *Exp.Biol.Med.*, 2001; 226(8): 726-733
- Savill, J.; Hogg, N.; Ren, Y.; Hasslet, C.: Thrombospondin cooperates with CD36 and the vitronectin receptor in macrophage recognition of neutrophils undergoing apoptosis. *J.Clin.Invest.*, 1992; 90: 1513-1522

Savill, J.: Recognition and phagocytosis of cells undergoing apoptosis. *Br.Med.Bull.*, 1997; 53: 491-508

Schubert, A.; Cattaruzza, M.; Hecker, M.; Darmer, D.; Holtz, J.; Morawietz, H.: Shear stress-dependent regulation of the human beta-tubulin folding cofactor D gene. *Circ.Res.*, 2000; 87 (12): 1188-1194

Schwachtgen, J.-L.; Houston, P.; Campbell, C.; Sukhatme, V.; Braddock, M.: Fluid shear stress activation of egr-1 transcription in cultured human endothelial and epithelial cells is mediated *via* the extracellular signal-related kinase 1/2 mitogen-activated protein kinase pathway. *J.Clin.Invest.*, 1998; 101: 2540-2549

Sdougos, H. P.; Bussolari, S. R.; Dewey, C. F.: Secondary flow and turbulence in a cone-and-plate device. *J.Fluid.Mech.*, 1984; 138: 397-404

Semenza, G. L.; Roth, P. H.; Fang, H. M.; Wang, G. L.: Transcriptional regulation of genes encoding glycolytic enzymes by hypoxia-inducible factor 1. *J.Biol.Chem.*, 1994; 269: 23757-23763

Semenza, G. L.; Jiang, B. H.; Leung, S. W.; Passantino, R.; Concordet, J. P.; Maire, P.; Giallongo, A.: Hypoxia response elements in the aldolase A, enolase 1 and lactate dehydrogenase A gene promoters contain essential binding sites for hypoxia-inducible factor 1. *J.Biol.Chem.*, 1996; 271: 32529-32537

Semenza, G. L.: HIF-1: mediator of physiological and pathophysiological responses to hypoxia. *J.Appl.Physiol.*, 2000; 88: 1474-1480

Senger, D. R., Ledbetter, S. R., Claffey, K. P., Papadopoulos-Sergiou, A., Peruzzi, C. A., Detmar, M.: Stimulation of endothelial cell migration by vascular permeability factor / vascular endothelial growth factor through cooperative mechanisms involving the $\alpha v \beta 3$ integrin, osteopontin and thrombin. *Am.J.Pathol.*, 1996; 149: 293-306

Shattil S. J.; Brugge, J. S.: Protein tyrosine phosphorylation and the adhesive functions of platelets. *Curr.Opin.Cell.Biol.*, 1991; 3: 869-879

Sheibani, N. et Frazier, W. A.: Thrombospondin-1 expression in transformed endothelial cells restores a normal phenotype and suppresses their tumorigenesis. *Proc.Natl.Acad.Sci. USA*, 1995; 92: 6788-6792

Shyy, Y, J.; Hsieh, J.-H.; Usami, S.; Chien, S.: Fluid shear stress induces a biphasic response of human monocyte chemotactic protein 1 gene expression in vascular endothelium. *Proc.Natl.Acad.Sci. USA*, 1994; 91: 4678-4682

Shyy, J. Y. J. et Chien, S.: Role of integrins in cellular responses to mechanical stress and adhesion. *Curr.Opinion.Cell.Biol.*, 1997; 707-713

Simon, M. I.; Strathmann, M. P.; Gautam, N.: Diversity of G proteins in signal transduction. *Science*, 1991; 252: 802-808

Skalak, T. C. et Price, R. J.: The role of mechanical stresses in microvascular remodeling. *Microcirculation*, 1996; 3: 143-165

Smith, L. E. H.; Kopchick, J. J.; Chen, W.; Knapp, J.; Kinose, F.; Daley, D.; Foley, E.; Smith, R. G.; Schaeffer, J. M.: Essential role of Growth Hormone in Ischemia-Induced Retinal Neovascularization. *Science*, 1997; 276 (5319): 1706

Stamler, J. Mendelsohn, M. E., Amarante, P., Smick, D., Andon, N., Davies, P. F., Cooke, J. P. und Loscalzo, J.: N-acetylcysteine potentiates inhibition by endothelium-derived relaxing factor, *Circ.Res.*, 1989; 65, 789-795

Streit, M., Velasco, P., Brown, L. F., Skobe, M., Richard, L., Riccardi, L., Lawler, J., Detmar, M: Overexpression of thrombospondin-1 decreases angiogenesis and inhibits the growth of human cutaneous squamous cell carcinomas. *Am.J.Pathol.*, 1999; 155: 441-452

Streit, M., Riccardi, L., Velasco, P., Brown, L. F., Hawighorst, T., Bornstein, P., Detmar, M.: Thrombospondin-2: a potent endogenous inhibitor of tumor growth and angiogenesis. *Proc.Natl.Acad.Sci.USA*, 1999; 96: 14888-14893

Suri, C.; Jones, P. F.; Patan, S.; Bartunkova, S.; Maisonpierre, P. C; Davis, S.; Sato, T. N. and Yancopoulos, G. D.: Requisite role of angiopoietin-1, a ligand for the TIE-2 receptor, during embryonic angiogenesis. *Cell*, 1997; 1171-1180

Suri, C.; McClain, J.; Thurston, G.; McDonald, D. M.; Zhou, H.; Oldmixon, T. N. S.; Yancopoulos: Increased vascularization in mice overexpressing Angiopoietin-1. *Science*, 1998; 282: 468-471

Suzuma, K.; Takagi, H.; Otani, A.; Oh, A.; Honda, Y.: Expression of thrombospondin-1 in ischemia-induced retinal neovascularization. *Am.J.Pathol.*, 1999: 154(2): 343-354

Swerlick, R. A.; Lee, K. H.; Wick, T. M.; Lawley, T. J.: Human dermal microvascular endothelial but not human umbilical vein endothelial cells express CD 36 in vivo and in vitro. *J.Immunol.*, 1992; 148: 78-83

Tacchini, L.; Bianchi, L.; Bernelli-Zazzera, A.; Cairo, G.: Transferrin receptor induction by hypoxia. HIF-1 mediated transcriptional activation and cell-specific post-transcriptional regulation. *J.Biol.Chem*, 1999; 274: 24142-24146

Takahashi, M et Berk, B. C.: Mitogen-activated protein kinase (ERK 1/2) activation by shear stress and adhesion in endothelial cells – essential role for a herbimycin-sensitive kinase. *J.Clin.Invest.*, 1996; 98: 2623-2631

Taraboletti, G.; Roberts, D.; Liotta, L. A.; Giavazzi, R.: Platelet thrombospondin modulates endothelial cell adhesion, motility, and growth: a potential angiogenesis regulatory factor. *J.Cell.Biol.*, 1990; 111: 765-772

- Tazuke, S. I.; Mazure, N. M.; Sugawara, J.; Carland, G.; Faessen, G. H.; Suen, L. F.; Irwin, J. C.; Powell, D. R.; Giaccia, A. J.; Giudice, L. C.: Hypoxia stimulates insulin-like growth factor binding protein 1 (ICFBP-1) gene expression in HepG2 cells: a possible model for ICFBP-1 expression in fetal hypoxia. *Proc.Natl.Acad.Sci. USA*, 1998; 10188-10193
- Tenan, M.; Fulci, G.; Albertoni, M.; Diserens, A.-C.; Hamou, M.-F.; El Atifi-Borel, M.; Feige, J.-J.; Pepper, M. S.; Van Meir, E. G.: Thrombospondin-1 is downregulated by anoxia and suppresses tumorigenicity of human glioblastoma cells. *J.Exp.Med.*,2000; 191(10): 1789-1797
- Tolsma, S. S., Volpert, O. V., Good, D. J., Frazier, D. A., Polverini, P. J., Bouck, N: Peptides derived from two separate domains of the matrix protein thrombospondin-1 have anti-angiogenetic activity. *J.Cell.Biol.*, 1993; 122: 497-511
- Topper, J. N.; Cai, J.; Falb, D.; Gimbrone, M. A.: Identification of vascular endothelial genes differentially responsive to fluid mechanical stimuli: cyclooxygenase-2, manganese superoxide dismutase, and endothelial cell nitric oxide synthase are selectively up-regulated by steady laminar shear stress. *Proc.Natl.Acad.Sci. USA*, 1996; 93(19): 10417-10422
- Topper, J. N.; Cai, J.; Qiu, Y.; Anderson, K. R.; Xu, Y. Y.; Deeds, J. D.; Feeley, R.; Gimeno, C. J.; Woolf, E. A.; Tayber, O.; Mays, G. G.; Sampson, B. A.; Schoen, F. J.; Gimbrone, M. A. Jr.; Falb, D.: Vascular MADs: two novel MAD-related genes selectively inducible by flow in human vascular endothelium. *Proc.Natl.Acad.Sci, USA*, 1997; 94 (17): 9314-9319
- Tuszynski, G. P.; Rothman, V.; Murphy, A.; Siegler, K.; Smith, L.; Smith, S.; Karczewski, J.; Knudsen, K. A.: Thrombospondin promotes cell-substratum adhesion. *Science*, 1987; 236: 1570-1573
- Tuszynski, G. P. et Nicosia, R. F.: Localization of Thrombospondin and its cysteine-serine-valine-threonine-cysteine-glycine-specific receptor in human breast carcinoma. *Lab.Invest.*, 1994; 70: 228-233
- Tuszynski, P. et Nicosia, R. F.: The role of thrombospondin in tumor progression and angiogenesis. *BioEssays* (ICSU Press), 1996; 18(1):71-76
- Vandenbunder, B.; Pardanaud, L.; Jaffredo, T.; Mirabel, M. A.; Stehelin, D.: Complementary patterns of expression of c-ets 1, c-myb and myc in the blood-forming system of the chick embryo. *Development*, 1989; 107(2): 265-274
- Vane, J. R. et Botting, R. M.: The role of chemical mediators released by the endothelium in the control of the cardiovascular system, *Int.J.Tissue.React.*, 1992; 14(2): 55-64

- Vazquez, F.; Hastings, G.; Ortega, M. A.; Lane, T.F.; Oikemus, S.; Lombardo, M.; Iruela-Arispe, M. L.: METH-1, a human ortholog of ADAMTS-1, and METH-2 are members of a new family of proteins with angio-inhibitory activity. *J.Biol.Chem.*, 1999; 274 (33): 23349-57
- Vernon, R. B.; Lara, S. L.; Drake, C. J.; Iruela-Arispe, M. L.; Angello, J. C.; Little, C. D.; Wight, T. N.; Sage, E. H.: Organized type I collagen influences endothelial patterns during spontaneous angiogenesis *in vitro*: planar cultures as models of vascular developmet. *In vitro Cell.Dev.Biol.*, 1995; 31:120-131
- Volpert, O. V.; Stellmach, V.; Bouck, N.: The modulation of thrombospondin and other naturally occurring inhibitors of angiogenesis during tumor progression. *Breast Cancer Res.Treat.*, 1995; 36(2): 119-126
- Wang, H. U.; Chen, Z. F.; Anderson, D. J.: Molecular distinction and angiogenetic interaction between embryonic arteries and veins revealed by ephrin-B2 and its receptor Eph-B4. *Cell*, 1998; 93: 741-753
- Wang, X. Q. et Frazier, W. A.: The thrombospondin receptor CD47(IAP) modulates and associates with alpha 2 beta 1 integrin in vascular smooth muscle cells. *Mol.Biol.Cell.*, 1998; 9: 865-874
- Wang, X.-Q.; Lindberg, F.; Frazier, W.: Integrin-associated protein stimulates $\alpha_2\beta_1$ -dependent chemotaxis via Gi-mediated inhibition of adenylate-cyclase and extracellular-regulated kinases. *J.Cell Biol.*, 1999; 147: 389-399
- Wenger, R. H. et Gassmann, M.: Oxygen and the hypoxia-inducible factor-1. *Biol.Chem.*, 1997; 378: 609-616
- Wenger, R. H.; Rolfs, A.; Spielmann, P.; Zimmermann, D. R.; Gassmann, M.: Mouse hypoxia-inducible factor-1 α is encoded by two different mRNA isoforms – expression from a tissue-specific and a housekeeping-type promoter. *Blood*, 1998c; 91: 3471-3480
- Wenger, R. H.: Mammalian oxygen sensing, signalling and gene regulation. *J.Exp.Biol.*, 2000; 203: 1253-1263
- Wernert, N.; Raes, M. B.; Lassalle, P.; Dehouck, M. P.; Gosselin, B.; Vandebunder, B.; Stehelin, D.: C-ets 1 proto-oncogene is a transcription factor expressed in endothelial cells during tumor vascularization and other forms of angiogenesis in humans. *Am.J.Pathol.*, 1992; 140(1): 119-127
- Willis, A. L.; Wang, X.; Kito, H.; Tuszynski, G. P.; Sumpio, B. E.; Gahtan, V.: Ras N17 transfection inhibits thrombospondin-1 induced vascular smooth muscle cell migration. *Surgical Forum*, 1999; 50:454-455

Witzenbichler, B.; Maisonpierre, P. C.; Jones, P.; Yancopoulos, G. D.; Isner, J. M.: Chemotactic properties of Angiopoietin-1 and-2, ligands for the endothelial-specific receptor tyrosine kinase Tie2. *J.Biol.Chem.*, 1998; 273: 18514-18524

Xiang, J.; Chao, D. T.; Korsmeyer, S. J.: Bax-induced cell death may not require interleukin-1 β -converting enzyme-like proteases. *Proc.Natl.Acad.Sci. USA*, 1996; 93: 14559-14563

Xiao, F.; Wei, Y.; Yang, L.; Zhao, X.; Tian, L.; Ding, Z.; Yuan, S.; Lou, Y.; Liu, F.; Wen, Y.; Li, J.; Deng, H.; Kang, B.; Mao, Y.; Lei, S.; He, Q.; Su, J.; Lu, Y.; Niu, T.; Hou, J.; Huang, M. J.: A gene therapy for cancer based on the angiogenesis inhibitor, vasostatin. *Gene Ther.*, 2002; 9 (18): 1207-1213

Yao, L.; Zhao, Y. L.; Itoh, S.; Wada, S.; Yue, L.; Furuta, I.: Thrombospondin-1 expression in oral squamous cell carcinomas: correlations with tumor vascularity, clinicopathological features and survival. *Oral.Oncol.*, 2000; 36(6): 539-544

Yehualaeshet, T.; O'Connor, R.; Green-Johnson, J.; Mai, S.; Silverstein, R.; Murphy-Ullrich, J. E.; Khalil, N.: Activation of rat alveolar macrophage-derived latent transforming growth factor β 1 by plasmin requires interaction with thrombospondin-1 and its cell surface receptor, CD 36. *Am.J.Pathol.*, 1999; 155: 841-851

Yu, A. Y.; Shimoda, L. A.; Iyer, N. V.; Huso, D. L.; Sun, X.; McWilliams, R.; Beaty, T.; Sham, J. S.; Wiener, C. M.; Sylvester, J. T.; Semenza, G. L.: Impaired physiological responses to chronic hypoxia in mice partially deficient for hypoxia-inducible factor 1 α . *J.Clin.Invest.*, 1999; 103: 691-696

Zabrenetzky, V.; Harris, C. C.; Steeg, P. S.; Roberts, D. D.: Expression of the extracellular matrix molecule thrombospondin inversely correlates with malignant progression in melanoma, lung and breast carcinoma cell lines. *Int.J.Cancer*; 59: 191-195

Zakrzewicz, A.; Gräfe, M.; Terbeek, D.; Bongrazio, M.; Auch-Schwelk, W.; Walzog, B.; Graf, K.; Fleck, E.; Ley, K.; Gaehtgens, P.: L-Selectin-dependent leukocyte adhesion to microvascular but not to macrovascular endothelial cells of the human coronary system. *Blood*, 1997; 89: 3228-3235

Zakrzewicz, A.; Bongrazio, M.; Baum, O.; Baumann, C.; Knöchel, J.; Walter, F.; Doster, P.; Bergmann, E. C.; Pries, A. R.: Wall shear stress regulates the expression of angiogenic and anti-angiogenic factors in endothelial cells. *Eur.J.Physiol.*, 2001; 441(6): R129

Zakrzewicz, A.; Secomb, T. W.; Pries, A. R.: Angioadaptation: Keeping the vascular system in shape. *News Physiol.Sci.*, 2002; 17: 197-201

Zelzer, E.; Levy, Y.; Kahana, C.; Shilo, B. Z.; Rubinstein, M.; Cohen, B.: Insulin induces transcription of target genes through the hypoxia-inducible factor HIF-1 α / ARNT. *EMBO.J.*, 1998; 17: 5085-5094

Zhou, A.; Egginton, S.; Hudlicka, O.; Brown, M. D.: Internal division of capillaries in rat skeletal muscle in response to chronic vasodilator treatment with alpha1-antagonist prazosin. *Cell.Tissue.Res.*, 1998; 293(2):293-303