

Imaging Molecular and Trace Element Changes in Scrapie-Infected Nervous Tissue with a Time Course Study



Dissertation

zur Erlangung des akademischen Grades des Doktors der
Naturwissenschaften (Dr. rer. nat.)

eingereicht im Fachbereich Biologie, Chemie, Pharmazie
der Freien Universität Berlin

vorgelegt von

Ariane Kretlow
aus Berlin

Mai, 2007

1. Gutachter: Prof. Dr. Dieter Naumann
2. Gutachter: Prof. Dr. Ronald Gust

Disputation am 04. Juli 2007

CONTENTS

Index of Figures	iv
Index of Tables	vi
1. Introduction	1
1.1 Literature overview	1
1.1.1 Protein misfolding diseases – General Aspects	1
1.1.2 Transmissible spongiform encephalopathies	1
1.1.2.1 Human prion diseases	2
1.1.2.2 Animal prion diseases	5
1.1.2.3 The prion protein: PrP ^C and PrP ^{Sc}	9
1.1.2.4 Conversion of PrP ^C to PrP ^{Sc}	11
1.1.2.5 Prion strains	12
1.1.2.6 Metals in prion diseases	14
1.1.2.7 The scrapie-263K hamster model	15
1.1.3 Reovirus T3C9	17
1.2 The vertebrate nervous system	18
1.2.1 The central nervous system	19
1.2.2 The peripheral nervous system	19
1.3 FTIR microspectroscopy	20
1.3.1 Basic principles of IR spectroscopy	20
1.3.2 The FTIR spectrometer	22
1.3.3 Advantages of FTIR versus dispersive IR spectrometer	24
1.3.4 FTIR microspectroscopy of biological tissue	25
1.4 X-Ray fluorescence microprobe	29
1.5 Previous FTIRMS studies on scrapie infected nervous tissue	31
1.5.1 FTIR-based detection of pathological changes in scrapie-infected hamsters	31
1.5.2 FTIR microspectroscopy of dorsal root ganglia utilizing a synchrotron IR light source	34
1.6 Aims	36
2. Materials and Methods	39
2.1 Scrapie	39
2.1.1 TSE strains and animal infection	39
2.1.2 Sample preparation	40
2.2 Reovirus	41
2.2.1 Reovirus T3C9	41
2.2.2 Mouse fibroblasts L929	41
2.2.3 Medium	42
2.2.4 Cultivation of Reovirus T3C9	42
2.2.5 Transmission Electron Microscopy	43
2.2.6 Animal infection	43
2.2.7 Sample preparation	44
2.3 Histology	44
2.3.1 Hematoxylin-Eosin staining	44
2.3.2 Cresyl violet staining	45
2.3.3 GFAP - Staining for Astrocytes	45

2.3.4	F4/80 - Staining for Microglia	47
2.3.5	Immunohistochemistry	48
2.3.6	Western Blot	50
2.4	FTIR microspectroscopy	54
2.4.1	DRG of 263K scrapie-infected hamsters	54
2.4.2	DRG of terminally diseased 263K and ME7 infected hamsters	55
2.4.3	DMNV of 263K infected hamsters at 70 dpi	56
2.4.4	DMNV of Reovirus T3C9 infected hamsters	57
2.5	XRF Microprobe	57
2.6	Data Analysis	58
2.6.1	protein/lipid ratio	58
2.6.2	β -sheet content and distribution	59
2.6.3	α -helix content and distribution	59
2.6.4	relative protein content	60
2.6.5	β -sheet / α -helix ratio and distribution	60
2.6.6	'fingerprint' (predominantly phosphates and carbohydrates)	60
2.6.7	lipid composition	60
2.6.8	Cluster Analysis	61
2.6.9	Statistical Analysis	61
2.6.10	Derivative spectra	61
2.6.11	XRF microprobe	61
3.	Results	63
3.1	Spectral changes in DRG of 263K scrapie-infected hamsters	64
3.2	DMNV of 263K infected hamsters at 70 dpi	71
3.3	DRG of terminally diseased 263K and ME7 infected hamsters	77
3.4	DMNV of Reovirus T3C9 infected hamsters	85
3.5	Comparison of Reovirus to Scrapie induced spectral changes	89
3.6	XRF Microprobe pathogenesis study	93
4.	Discussion	99
4.1	Spectral and molecular changes during scrapie pathogenesis	99
4.2	Onset of scrapie induced changes detected in the DMNV	105
4.3	Specificity of scrapie induced spectral changes	107
4.4	Alterations in the DMNV of reovirus infected hamster	110
4.5	Distribution and content of trace metals in scrapie	113
4.5.1	Changes in the total metal content in DRG neurons during scrapie pathogenesis	113
4.5.2	Subcellular distribution and content of Cu, Zn, Fe, Ca, P and Mn	115
4.6	Limitations and Possible error sources	118
4.6.1	Experimental design	118
4.6.2	Analysis	119
5.	Conclusions	121
6.	Abstract	125
7.	Zusammenfassung	129
8.	References	133

List of Abbreviations	159
List of Publications	161
Acknowledgements	163

INDEX OF FIGURES

Figure 1.1:	Secondary structure of PrP ^C and PrP ^{Sc}	11
Figure 1.2:	Simplified illustration of the two proposed hypotheses for prion conversion	12
Figure 1.3:	PrP27-30 2nd derivative spectra in the amide sensitive spectral ranges	13
Figure 1.4:	Initial spread of the 263K agent after oral uptake	16
Figure 1.5:	Structure of Orthoreovirus	18
Figure 1.6:	Illustration of the different vibrational modes	21
Figure 1.7:	Morse and harmonic oscillator potential	22
Figure 1.8:	Schematic assembly of a spectrometer	23
Figure 1.9:	IR spectrum of a biological sample	25
Figure 1.10:	Principles of XRF	30
Figure 1.11:	Typical XRF microprobe spectrum	31
Figure 1.12:	FTIR-based detection of pathological changes in scrapie-infected hamsters	35
Figure 2.2.5:	TEM photomicrographs of Reovirus T3C9 infected L929 mouse fibroblasts and sucrose gradients	43
Figure 2.3.3:	GFAP staining of a Reovirus T3C9 and a control hamster	47
Figure 2.3.5:	3F4 stained sections of DRG of 263K, ME7 and control hamster	49
Figure 2.3.6:	Western Blot Analysis	53
Figure 2.6.2:	Curve fitting result of a spectrum derived from a terminally infected animal	59
Figure 3.1.1:	Mean values + average deviation for the β -sheet, α -helical and relative protein content	65
Figure 3.1.2:	Photomicrographs and corresponding FTIRM images of the β -sheet distribution for control and infected ganglia at different time points	67
Figure 3.1.3:	Photomicrographs of the antibody 3F4 stained adjacent sections	68
Figure 3.1.4:	Comparison of the results from the FTIRMS of infected animals with 3F4 stained section of the same tissue	69
Figure 3.1.5:	Correlation of β -sheet to α -helix content in scrapie-infected DRG at 100 dpi and terminal stage	69
Figure 3.1.6:	Mean values + average deviation for the CH_3/CH_2 ratio and results of the carbohydrate/phosphate region	71
Figure 3.2.1:	3F4 immunostained adjacent sections of the DMNV of an infected and a control animal	72
Figure 3.2.2:	Dendrogram and average spectra shown as a result of the Cluster Analysis performed on the detailed measurement of an infected animal at 70 dpi in the range $1300 - 1000 \text{ cm}^{-1}$	73
Figure 3.2.3:	Dendrogram and average spectra shown as a result of the Cluster Analysis of normalized first derivative spectra of the DMNV of 16 infected and 8 control animals at 70 dpi in the range $1100 - 1000 \text{ cm}^{-1}$	74
Figure 3.2.4:	First derivative average spectra of scrapie-infected and control animals at 70 dpi, 90 dpi, 120 dpi and at the terminal stage of disease	75
Figure 3.2.5:	Photomicrographs, β -sheet distribution and lipid composition of two control animals and three 70 dpi scrapie infected animals	76

Figure 3.2.6: Results of the analysis of the total protein content, the carbohydrate/phosphate region, the β -sheet content and the lipid composition of five 70 dpi infected and three control animals	77
Figure 3.3.1: β -sheet to α -helix ratio of two control animals, four 263K and two ME7 infected animals and β -sheet to α -helix ratio of ME7 infected DRG in relation to the position of the ganglion	79
Figure 3.3.2: β -sheet to α -helix ratio of one ME7, one 263K infected animal and one control animal	79
Figure 3.3.3: β -sheet to α -helix ratio and average spectra of one ME7, one 263K infected animal and one control animal	81
Figure 3.3.4: Original and 2nd derivative average spectra of two 263K infected and two ME7 infected animals	82
Figure 3.3.5: 1st derivative and original spectra of averages from control, 263K and ME7 infected animals	83
Figure 3.3.6: Analysis of the phosphate/carbohydrate and the lipid region of control, 263K and ME7 infected animals	84
Figure 3.4.1: Cresyl violet stained photomicrograph and protein/lipid ratio of the Medulla oblongata of a 10 day old control hamster	86
Figure 3.4.2: Dendrogram as a result of a Cluster Analysis and vector normalized first derivatives of one animal in the region 1300–1000 cm ⁻¹	86
Figure 3.4.3: Dendrogram as a result of a Cluster Analysis on 1st derivative average spectra in the range 1100–1000 cm ⁻¹ of the DMNV from seven reovirus infected and six age matched control animals and vector normalized first derivatives in the region 1300–1000 cm ⁻¹	87
Figure 3.4.4: 2 nd derivative spectra of the DMNV of reovirus infected and control animals in the region 3050 – 2800 cm ⁻¹	88
Figure 3.4.5: Statistical analysis of the lipid composition, fingerprint region and total protein content of six control and seven infected hamsters	89
Figure 3.5.1: 1 st derivative spectra of the DMNV of reovirus infected, age-matched control, terminal scrapie infected and age-matched control in the region 1300 – 1000 cm ⁻¹	90
Figure 3.5.2: Original spectra of the DMNV of reovirus infected, age-matched control, terminal scrapie infected and age-matched control, 70 dpi scrapie infected and age-matched control in the region 1300 – 1000 cm ⁻¹	91
Figure 3.5.3: Dendrogram as a result of a cluster analysis on vector-normalized 1 st derivative average spectra of control and infected animals at different ages	92
Figure 3.6.1: Cu, Zn, Fe, Ca, Mn, K and P content compared to control over the course of the disease	95
Figure 3.6.2: Cu, Zn, Fe, Ca, and Mn content and distribution of a 70 dpi, a 100 dpi, a fcs, a terminal scrapie infected and a terminal control calculated from coarse scans	96
Figure 3.6.3: Cu, Zn, Fe, Ca, P and Mn distribution of one control and one terminal scrapie-infected animal calculated from the fine scans	98

INDEX OF TABLES

Table 1.1:	Overview of the human and animal prion diseases	9
Table 1.2:	IR band assignment in biological molecules	27
Table 1.3:	Assignment of IR absorption bands to different secondary structures of proteins	29
Table 2.1:	List of official name, acronym, strain, time point and tissue of the animals used in this thesis	40

Index of Tables
