

6. Literaturverzeichnis

- Aarts, N., Metz, M., Holub, E., Staskawicz, B. J., Daniels, M. J. & Parker, J. E. (1998) Different requirements for *EDS1* and *NDR1* by disease resistance genes define at least two *R* gene-mediated signaling pathways in *Arabidopsis*. Proc. Natl. Acad. Sci. USA **95**, 10306-10311
- Anderson, P. A., Lawrence, G. J., Morrish, B. C., Ayliffe, M. A., Finnegan, E. J. & Ellis, J. G. (1997) Inactivation of the flax rust resistance gene *M* associated with loss of a repeated unit within the leucine-rich repeat coding region. Plant Cell **9**, 641-651
- 'The Arabidopsis Genome Initiative' (2000) Analysis of the genome sequence of the flowering plant *Arabidopsis thaliana*. Nature **408**, 796-815
- Arbeiter, A. (1997) Charakterisierung von YACs zur Erstellung eines YAC-Contigs über das *RPB1*-Gen von *Arabidopsis thaliana* (L.) Heynh.. Diplomarbeit, Inst. für Angewandte Genetik, Freie Universität Berlin
- Arbeiter, A., Kobelt, P., Sacristán, M.D., Luerßen, H. & Siemens, J. (2001) The interaction of the obligate biotrophic pathogen *Plasmodiophora brassicae* with *Arabidopsis thaliana*. 12th International Conference on Arabidopsis Research, June 2001, Madison, Wisconsin, USA
- Ayers, G. W. & Lelacheur, K. E. (1972) Genetics of resistance in rutabaga to races of *Plasmodiophora brassicae*. Can. J. Plant Sci. **52**, 897-900
- Bent, A. F. (1996) Plant disease resistance genes: Function meets structure. Plant Cell **8** (10), 1757-1771
- Botella, M. A., Parker, J. E., Frost, L. N., Bittnereddy, P. D., Beynon, J. L., Daniels, M. J., Holub, E. B. & Jones, J. D. G. (1998) Three genes of the *Arabidopsis RPP1* complex resistance locus recognize distinct *Peronospora parasitica* avirulence determinants. Plant Cell **10** (11), 1847-1860
- Breyne, P., Rombaut, D., Van Gysel, A., Van Montagu, M. & Gerats, T. (1999) AFLP analysis of genetic diversity within and between *Arabidopsis thaliana* ecotypes. Mol. Gen. Genet. **261**, (4-5), 627-634
- Cao, H., Li, X. & Dong, X. (1998) Generation of broad-spectrum disease resistance by overexpression of an essential regulatory gene in systemic acquired resistance. Proc. Natl. Acad. Sci. USA **95**, 6531-6536
- Cavell, A. C., Lydiate, D. J., Parkin, I. A. P., Dean, C. & Trick, M. (1998) Collinearity between a 30-centimorgan segment of *Arabidopsis thaliana* chromosome 4 and duplicated regions within the *Brassica napus* genome. Genome **41**, 62-69
- Chang, C., Bowman, J. L., DeJohn, A. W., Lander, E. S. & Meyerowitz, E. M. (1988) Restriction fragment length polymorphism linkage map for *Arabidopsis thaliana*. Proc. Natl. Acad. Sci. **85**, 6856-6860
- Choi, S., Creelman, R.A., Mullet, J.E. & Wing, R. (1995) Construction and characterisation of a bacterial artificial chromosome library of *Arabidopsis thaliana*. Plant Mol. Biol. Rep. **13**, 124-128
- Cooke, R., Raynal, M., Laudie, M., Grellet, F., Delseny, M., Morris, P. C., Guerrier, D., Giraudat, J., Quigley, F., Clabault, G., Li, Y. F., Mache, R., Krivitzky, M., Gy, I. J. J., Kreis, M., Lecharny, A., Parmentier, Y., Marbach, J., Fleck, J., Clement, B., Philipp, G., Herve, C., Bardet, C., Tremousaygue, D., Lescure, B., Lacomme, C., Roby, D., Jourjon, M.-F., Chabrier, P., Charpentier, J.-L., Desprez, T., Amselem, J., Chiapello, H. & Höfte, H. (1996) Further progress towards a catalogue of all *Arabidopsis* genes: analysis of a set of 5000 non-redundant ESTs. Plant J. **9** (1), 101-124

- Creusot, F., Fouilloux, E., Dron, M., Lafleur, J., Picard, G., Billault, A., Le Paslier, D., Cohen, D., Chabouté, M.-E., Durr, A., Fleck, J., Gigot, C., Camilleri, C., Bellini, C., Caboche, M. and Bouchez, D. (1995) The CIC library: a large insert YAC library for genome mapping in *Arabidopsis thaliana*. *Plant J.* **8** (5), 763-770
- Diederichsen, E. (1992) Kombination verschiedener Resistzen gegenüber *Plasmodiophora brassicae* Wor. in resynthetisierten Formen von amphidiploiden *Brassica*-Arten. Dissertation, Inst. für Angewandte Genetik, Freie Universität Berlin
- Diederichsen, E. & Sacristán, M. D. (1996) Disease response of resynthesized *Brassica napus* L. lines carrying different combinations of resistance to *Plasmodiophora brassicae* Wor.. *Plant Breedings* **115**, 5-10
- Dekhuijzen, H. M. & Overeem, J. C. (1971) The role of cytokinins in clubroot formation. *Physiol. Pl. Path.* **1**, 151-161
- Dekhuijzen, H. M. (1979) Electron microscopic studies on the root hairs and cortex of a susceptible and a resistant variety of *Brassica campestris* infected with *Plasmodiophora brassicae*. *Neth. J. Pl. Path.* **85**, 1-17
- Delaney, T. P. (1997) Genetic dissection of acquired resistance to disease. *Plant Physiol.* **113**, 5-12
- Delseny, M., Cooke, R., Comella, P., Wu, H. J., Raynal, M. & Grellet, F. (1997) The *Arabidopsis thaliana* genome project [review]. *Comptes Rendus de l' Academie des Sciences de la Vie-Life Sciences*. **320** (8), 589-599
- Dixon, G. R. (1981) Vegetable crop diseases. MacMillan Publishers, London, p 137
- Dixon, M. S., Jones, D. A., Keddie, J. S., Thomas, C. M., Harrison, K. & Jones, J. D. G. (1996) The tomato *Cf-2* disease resistance locus comprises two functional genes encoding leucine-rich repeat proteins. *Cell* **84**, 451-459
- Doke, N., Miura, L. M., Park, H.-J., Noritake, T., Yoshioka, H. & Kawakita, K. (1996) The oxidative burst protects plants against pathogen attack: mechanism and role as an emergency signal for plant bio-defence – a review. *Gene* **179**, 45-51
- Ecker, J.R. (1990) PFGE and YAC analysis of the *Arabidopsis* genome. *Methods: A Companion to Methods in Enzymology* **1** (2), 186-194
- Edwards, K., Johnstone, C. & Thompson, C. (1991) A simple and rapid method for the preparation of plant genomic DNA for PCR analysis. *Nucleic Acids Res.* **19** (6), 1349
- Ellis, J., Dodds, P., Pryor, T. (2000) Structure, function and evolution of plant disease resistance genes. *Curr. Opin. Plant Biol.* **3**, 278-284
- Erschadi, S., Haberer, G., Schöniger, M. & Torres-Ruiz, R. A. (2000) Estimating genetic diversity of *Arabidopsis thaliana* ecotypes with amplified fragment length polymorphisms (AFLP). *Theor. Appl. Genet.* **100**, 633-640
- Fan, Q. Q. & Petes, T. D. (1996) Relationship between nuclease-hypersensitive sites and meiotic recombination hot spot activity at the *HIS4* locus of *Saccharomyces cerevisiae*. *Mol. Cell. Biol.* **5**, 2037-2043
- Fantl, W. J., Johnson, D. E. & Williams, L. T. (1993) Signalling by receptor tyrosine kinases. *Annual. Rev. Biochem.* **62**, 453-481
- Faris, J. D., Haen, K. M. & Gill, B. S. (2000) Saturation mapping of a gene-rich recombination hot spot region in wheat. *Genetics* **154**, 823-835
- Fatokun, C. A., Menancio-Hautea, D. I., Danesh, D. & Young, N. D. (1992) Evidence for orthologous seed weight genes in cowpea and mung bean based on RFLP mapping. *Genetics* **132**, 841-846
- Feinberg, A. P. & Vogelstein, B. (1983) A technique for radiolabeling DNA restriction endonuclease fragments to high specific activity. *Anal. Biochem.* **132**, 6-13
- Flor, H. H. (1956) The complementary genetic systems in flax and flax rust. *Adv. in Genet.* **8**, 29-54

- Fuchs, H. & Sacristán, M. D. (1996) Identification of a gene in *Arabidopsis thaliana* controlling resistance to clubroot (*Plasmodiophora brassicae*) and characterization of the resistance response. *Mol. Plant-Microbe Interact.* **9** (2), 91-97
- Gabriel, D. W. & Rolfe, B. G. (1990) Working models of specific recognition in plant-microbe interactions. *Annu. Rev. Phytopathol.* **28**, 365-391
- Gebhardt, C., Ritter, E., Barone, A., Debener, T., Walkemeier, B., Schachtschabel, U., Kaufmann, H., Thompson, R. D., Bonierbale, M. W., Ganal, M., Tanksley, S. D. & Salamini, F. (1991) RFLP maps of potato and their alignment with the homoeologous tomato genome. *Theor. Appl. Genet.* **83**, 49-57
- Gibson, S. I. & Somerville, C. (1992) Chromosome walking in *Arabidopsis thaliana* using yeast artificial chromosomes. In: *Methods in Arabidopsis research* (Koncz, C., Chua, N. H., and Shell, J., Hrsg.) World Scientific Publishing Co. Pte. Ltd., Singapor, pp 119-143
- Gilchrist, D. G. (1998) Programmed cell death in plant disease: The purpose and promise of cellular suicide. *Annu. Rev. Phytopathol.* **36**, 393-414
- Gill, K. S., Gill, B. S., Endo, T. R. & Boiko, E. V. (1996a) Identification and high-density mapping of gene-rich regions in chromosome group 5 of wheat. *Genetics* **143**, 1001-1012
- Gill, K. S., Gill, B. S., Endo, T. R. & Taylor, T. (1996b) Identification and high-density mapping of gene-rich regions in chromosome group 1 of wheat. *144*, 1883-1891
- Glazebrook, J. (1999) Genes controlling expression of defense responses in *Arabidopsis*. *Curr. Opin. Plant Biol.* **2**, 280-286
- Grant, M. R., McDowell, J. M., Sharpe, A. G., Torres Zabala, M., Lydiate, D. J. & Dangl, J. L. (1998) Independent deletions of a pathogen-resistance gene in *Brassica* and *Arabidopsis*. *Proc. Natl. Acad. Sci. USA* **95**, 15843-15848
- Grill, E. and Somerville, C. (1991) Construction and characterization of a yeast artificial chromosome library of *Arabidopsis* which is suitable for chromosome walking. *Mol. Gen. Genet.* **226**, 484-490
- Grsic, S., Neuhaus, K., Sauerteig, S. & Ludwig-Müller, J. (1997) The role of nitrilase during symptom development of the clubroot disease. *Pl. Physiol. Suppl.* **114**, 27
- Grsic, S., Kirchheim, B., Pieper, K., Fritsch, M., Hilgenberg, W. & Ludwig-Müller, J. (1998) Auxin biosynthesis in clubroot diseased Chinese cabbage plants and induction by jasmonic acid. *Physiol. Plant.* **105**, 521-531
- Hammond-Kosack, K. E., & Jones, J. D. G. (1996) Resistance gene-dependent plant defense responses. *Plant Cell* **8** (10), 1773-1791
- Hammond-Kosack, K. E., & Jones, J. D. G. (2000) Responses to plant pathogens. In: Buchanan, B., Grussem, W., Jones, R., Eds. *Biochemistry & Molecular Biology of Plants*. American Society of Plant Physiologists, 1102-1156
- Hardtke, C. S., Muller, J. & Berleth, T. (1996) Genetic similarity among *Arabidopsis thaliana* ecotypes estimated by DNA sequence comparison. *Plant Mol. Biol.* **32** (5), 915-922
- Hauge, B. M., Hanley, S. M., Cartinhour, S., Cherry, J. M., Goodman, H. M.; Koornneef, M., Stam, P.; Chang, C., Kemplin, S., Medrano, L. & Meyerowitz, E. M. (1993) An integrated genetic/RFLP map of the *Arabidopsis thaliana* genome. *Plant J.* **3** (5), 745-754
- Höfte, H., Desprez, T., Amselem, J., Chiapello, H., Caboche, M., Moisan, A., Jourjon, M.-F., Charpentier, J.-L., Berthomieu, P., Guerrier, D., Giraudat, J., Quigley, F., Thomas, F., Yu, D.-Y., Mache, R., Raynal, M., Cooke, R., Grellet, F., Delseny, M., Parmentier, Y., De Marcillac, G., Gigot, C., Fleck, J., Phillips, G., Axelos, M., Bardet, C., Tremousaygue, D. & Lescure, B. (1993) An inventory of 1152 expressed sequence tags obtained by partial sequencing of cDNAs from *Arabidopsis thaliana*. *Plant J.* **4** (6), 1051-1061
- Hunt, M. D., Neuenschwander, U. H., Delaney, T. P., Weymann, K. B., Friedrich, L. B.,

- Lawton, K. A., Steiner, H.-Y. & Ryals, J. A. (1996) Recent advances in systemic acquired resistance research – a review. *Gene* **179**, 89-95
- Hunter, T. (1991) Protein kinase classification. *Methods in Enzymology* **200**, 3-37
- Hutcheson, S. W. (1998) Current concepts of active defense in plants. *Annu. Rev. Phytopathol.* **36**, 59-90
- James, R. V. & Williams, P. H. (1980) Clubroot resistance and linkage in *Brassica campestris*. *Phytopathol.* **70** (8), 776-779
- Jarvis, P., Lister, C., Szabo, V. & Dean, C. (1994) integration of CAPS markers into the RFLP map generated using recombinant inbred lines of *Arabidopsis thaliana*. *Plant Mol. Biol.* **24**, 685-687
- Johnston, T. D. (1968) Club root in *Brassica*: a standard inoculation technique and the specification of races. *Plant Path.* **17**, 184-187
- Jordan, T. (1999) Kartierung molekularer Marker in der *RPB1*-Region. Diplomarbeit, Inst. für Angewandte Genetik, Freie Universität Berlin
- King, G., Nienhuis, J. & Hussey, C. (1993) Genetic diversity among ecotypes of *Arabidopsis thaliana* estimated by analysis of restriction fragment length polymorphisms. *Theor. Appl. Genet.* **86**, 1028-1032
- Kobelt, P. (2000) Die Verbreitung von sekundären Plasmodien von *Plasmodiophora brassicae* (Wor.) im Wurzelgewebe von *Arabidopsis thaliana* nach immunhistologischer Markierung des plasmoidalen Zytoskeletts Dissertation, Inst. für Angewandte Genetik, Freie Universität Berlin
- Kobelt, P., Siemens, J. & Sacristán, M.D. (2000) Histological characterisation of the incompatible interaction between *Arabidopsis thaliana* and the obligate biotrophic pathogen *Plasmodiophora brassicae*. *Mycol. Res.* **104**, 220-225
- Koch, E., Cox, R. & Williams, P. H. (1991) Infection of *Arabidopsis thaliana* by *Plasmodiophora brassicae*. *J. Phytopathol.* **132**, 99-104
- Köhn, C. (1997) Feinkartierung des Resistenzgens *RPB1* von *Arabidopsis thaliana* (L.) Heynh. gegen *Plasmodiophora brassicae* Wor. mit Hilfe molekularer Marker als Basis für eine kartengestützte Klonierung. Dissertation, Institut für Angewandte Genetik, Freie Universität Berlin
- Koornneef, M. (1994) *Arabidopsis* genetics. In: "Arabidopsis" (Meyerowitz, E. M. and Somerville, C. R., eds.) Cold Spring Harbor Laboratory Press, New York, USA, pp 89-120
- Kosambi, D. D. (1944) The estimation of map distances from recombination values. *Ann. Eugen.* **12**, 172-175
- Kroll, T. K., Lacy, G. H. & Moore, L. D. (1983) A quantitative description of the colonization of susceptible and resistant radish plants by *Plasmodiophora brassicae*. *Phytopath. Z.* **108**, 97-105
- King, G., Nienhuis, J. & Hussey, C. (1993) Genetic similarity among ecotypes of *Arabidopsis thaliana* estimated by analysis of restriction fragment length polymorphisms. *Theor. Appl. Genet.* **86**, 1028-1032
- Kowalski, S. P., Lan, T.-H., Feldmann, K. A. & Paterson, A. H. (1994) Comparative mapping of *Arabidopsis thaliana* and *Brassica oleracea* chromosomes reveals islands of conserved organization. *Genetics* **138**, 499-510
- Kurata, N., Moore, G., Nagamura, Y., Foote, T., Yano, M., Minobe, Y. & Gale, M. (1994) Conservation of genome structure between rice and wheat. *Bio/Technology* **12**, 276-278
- Lagercrantz, U., Putterill, J., Coupland, G. & Lydiate, D. J. (1996) Comparative mapping in *Arabidopsis* and *Brassica*, fine scale genome collinearity and congruence of genes controlling flowering time. *Plant J.* **9**, 13-20
- Laurens, F. & Thomas, G. (1993) Inheritance of resistance to clubroot (*Plasmodiophora brassicae* Wor.) in kale (*Brassica oleracea* ssp. *acephala*). *Hereditas* **119**, 253-262

- Leister, R. T., Ausubel, F. M. & Katagiri, F. (1996) Molecular recognition of pathogen attack occurs inside of plant cells in plant disease resistance specified by the *Arabidopsis* genes *RPS2* and *RPM1*. Proc. Natl. Acad. Sci. USA **93** (26), 15497-15502
- Liang, P. & Pardee, A. B. (1992) Differential display of eucaryotic messenger RNA by means of the polymerase chain reaction. Science **257**, 967-971
- Lichten, M. & Goldman, A. S. H. (1995) Meiotic recombination hotspots. Annu. Rev. Genet. **29**, 423-444
- Lister, C. & Dean, C. (1993) Recombinant inbred lines for mapping RFLP and phenotypic markers in *Arabidopsis thaliana*. Plant J. **4** (4), 745-750
- Liu, Y.-G., Mitsukawa, N., Vazquez-Tello, A. & Whittier, R. (1995) Generation of a high-quality P1 library of *Arabidopsis* suitable for chromosome walking. Plant J. **7** (2), 351-358
- Liu, Y.-G., Mitsukawa, N., Lister, C., Dean, C. & Whittier, R. F. (1996) Isolation and mapping of a new set of 129 RFLP markers in *Arabidopsis thaliana* using recombinant inbred lines. Plant J. **10** (4), 733-736
- Loh, Y.-T. & Martin, G. B. (1995) The disease-resistance gene *Pto* and the fenthion-sensitivity gene *Fen* encode closely related functional protein kinases. Proc. Natl. Acad. Sci. USA **92**, 4181-4184
- Ludwig-Müller, J., Bendel, U., Thermann, P., Ruppel, M., Epstein, E. & Hilgenberg, W. (1993) Concentrations of indole-3-acetic acid in plants of tolerant and susceptible varieties of Chinese cabbage infected with *Plasmodiophora brassicae* Woron.. New Phytol. **125**, 763-769
- Ludwig-Müller, J. (1999) *Plasmodiophora brassicae*, the causal agent of clubroot disease: a review on molecular and biochemical events in pathogenesis. Journal of Plant Diseases and Protection **106** (2), 109-127
- Marra, M., Kucaba, T., Sekhon, M., Hillier, L., Martienssen, R., Chinwalla, A., Crockett, J., Fedele, J., Grover, H., Gund, C., McCombie, W. R., McDonald, K., McPherson, J., Mudd, N., Parnell, L., Schein, J., Seim, R., Shelby, P., Waterston, R. & Wilson, R. (1999) A map for sequence analysis of the *Arabidopsis thaliana* genome. Nature Genet. **22**, 265-270
- Martin, G. B., Brommonschenkel, S. H., Chunwongse, J., Frary, A., Gan, M. W., Spivey, R., Wu, T., Earle, E. D. & Tanksley, S. (1993) Map-based cloning of a protein kinase gene conferring disease resistance in tomato. Science **262**, 1432-1436
- Martin, G. B., Frary, A., Wu, T., Brommonschenkel, S. H., Chunwongse, J., Earle, E. D. & Tanksley, S. (1994) A member of the tomato *Pto* family confers sensitivity to fenthion resulting in rapid cell death. Plant Cell **6**, 1543-1552
- Martin, G. B. (1999) Functional analysis of plant disease resistance genes and their downstream effectors. Curr. Opin. Plant Biol. **2**, 273-279
- Meyer, K., Leube, M. P. & Grill, E. (1994) A protein phosphatase 2C involved in ABA signal transduction in *Arabidopsis thaliana*. Science **264**, 1452-1455
- Meyerowitz, E. M. (1994) Structure and organization of the *Arabidopsis thaliana* nuclear genome. In: "Arabidopsis" (Meyerowitz, E. M. and Somerville, C. R., eds.) Cold Spring Harbor Laboratory Press, New York, USA, pp 21-36
- Morgner, M. (1995) Mikroskopische Untersuchungen der Reaktionen empfindlicher und resistenter Genotypen verschiedener *Brassica*-Arten gegenüber *Plasmodiophora brassicae* (Wor.). Dissertation, Institut für Angewandte Genetik, Freie Universität Berlin

- Mozo, T., Fischer, S., Shizuya, H. & Altmann, T. (1998) Construction and characterisation of the IGF *Arabidopsis* BAC library. Mol. Gen. Genet. **258**, 562-570
- Mozo, T., Dewar, K., Dunn, P., Ecker, J. R., Fischer, S., Kloska, S., Lehrach, H., Marra, M., Martienssen, R., Meier-Ewert, S. & Altmann, T. (1999) A complete BAC-based physical map of the *Arabidopsis thaliana* genome. Nature Genet. **22**, 271-275
- Müller, P. & Hilgenberg, W. (1986) Isomers of zeatin and zeatin riboside in clubroot tissue: evidence for trans-zeatin biosynthesis by *Plasmochiphora brassicae*. Physiologia Pl. **66**, 245-250
- Nam, H.-G., Giraudat, J., den Boer, B., Moonan, F., Loos, W. D. B., Hauge, B. M. & Goodman, H. M. (1989) Restriction fragment length polymorphism linkage map of *Arabidopsis thaliana*. Plant Cell **1**, 699-705
- Newman, T., De Bruijn, F. J., Green, P., Keegstra, K., Kende, H., McIntosh, L., Ohlrogge, J., Raikhel, N., Somerville, S., Thomashow, M., Retzel, E., & Somerville, C. (1994) Genes galore – a summary of methods for accessing results from large-scale partial sequencing of anonymous *Arabidopsis* cDNA clones. Plant Physiol. **106** (4), 1241-1255
- Ochman, H., Gerber, A. S. & Hartl, D. L. (1988) Genetic applications of an inverse polymerase chain reaction. Genetics **120**, 621-623
- Ohta, K., Shibata, T. & Nicolas, A. (1994) Changes in chromatin structure at recombination initiation sites during yeast meiosis. EMBO J. **13**, 5754-5763
- Oldroyd, G. E. D. & Staskawicz, B. J. (1998) Genetically engineered broad-spectrum disease resistance in tomato. Proc. Natl. Acad. Sci. USA **95** (17), 10300-10305
- Osborn, T. C., Kole, C., Parkin, I. A. P., Sharpe, A. G., Kuiper, M., Lydiate, D. J. & Trick, M. (1997) Comparison of flowering time genes in *Brassica rapa*, *B. napus* and *Arabidopsis thaliana*. Genetics **146**, 1123-1129
- Parker, J. E., Coleman, M. J., Szabo, V., Frost, L. N., Schmidt, R., Vanderbiezen, E. A., Moores T., Dean, C., Daniels, M. J. & Jones, J. D. G. (1997) The *Arabidopsis* Downy Mildew resistance gene *RPP5* shares similarity to the toll and interleukin-1 receptors with *N* and *L6*. Plant Cell **9** (6), 879-894
- Parniske, M., Wulff, B. B. H., Bonnema, G., Thomas, C. M., Jones, D. A. & Jones, J. D. G. (1999) Homologues of the *Cf-9* disease resistance gene (*Hcr9s*) are present at multiple loci on the short arm of tomato chromosome 1. Mol. Plant-Microbe Interact. **12** (2), 93-102
- Penninckx, I., Eggemont, K., Terras, F., Thomma, B., Samblanx, G., Buchala, A., Metraux, J.-P., Manners, J. & Broekaert, W. (1996) Pathogen-induced systemic activation of a plant defensin gene in *Arabidopsis* follows a salicylic acid-independent pathway. Plant Cell **8**, 2309-2323
- Penninckx, I. A. M. A., Thomma, B. P. H. J., Buchala, A., Metraux, J. & Broekaert, W. (1998) Concomitant activation of jasmonate and ethylene response pathways is required for induction of a plant defensin gene in *Arabidopsis*. Plant Cell **10**, 2103-2113
- Petes, T. D. (2001) Meiotic recombination hot spots and cold spots. Nature Reviews Genetics **2**, 360-369
- Rausch, T., Butcher, D. N. & Hilgenberg, W. (1981) Nitrilase activity in clubroot diseased plants. Physiologia Pl. **52**, 467- 470
- Reuber, T. L. & Ausubel, F. M. (1996) Isolation of *Arabidopsis* genes that differentiate between resistance responses mediated by the *RPS2* and *RPM1* disease resistance genes. Plant Cell **8**, 241-249
- Riley, J., Butler, R., Ogilvie, D., Finniear, R., Jenner, D., Powell, S., Anand, R., Smith, J.C. & Markham, A.F. (1990) A novel, rapid method for the isolation of terminal sequences from yeast artificial chromosome (YAC) clones. Nucleic Acids Res. **18** (10), 2887-2890

- Ritter, C. & Dangl, J. L. (1996) Interference between two specific pathogen recognition events mediated by distinct plant disease resistance genes. *Plant Cell* **8** (2), 251-257
- Rogers, S. O. & Bendich, A. J. (1985) Extraction of DNA from milligram amounts of fresh, herbarium and mummified plant tissues. *Plant Mol. Biol.* **5**, 69-76
- Sacristán, M. D. & Hoffmann, F. (1979) Direct infection of embryogenic tissue cultures of haploid *B. napus* with resting spores of *Plasmodiophora brassicae*. *Theor. Appl. Genet.* **54**, 129-132
- Sacristán, M. D., Gerdemann-Knörck, M. & Schieder, O. (1989) Incorporation of hygromycin resistance in *Brassica nigra* and its transfer to *B. napus* through asymmetric protoplast fusion. *Theor. Appl. Genet.* **78**, 194-200
- Sadowski, J., & Quiros, C. F. (1998) Organization of an *Arabidopsis thaliana* gene cluster on chromosome 4 including the *RPS2* gene, in the *Brassica nigra* genome. *Theor. Appl. Genet.* **96**, 468-474
- Salmeron, J. M., Barker, S. J., Carland, F. M., Mehta, A. Y. & Staskawicz, B. J. (1994) Tomato mutants altered in bacterial disease resistance provide evidence for a new locus controlling pathogen recognition. *Plant Cell* **6**, 511-520
- Salmeron, J. M., Oldroyd, G. E. D., Rommens, C. M. T., Scofield, S. R., Kim, H.-S., Lavelle, D. T., Dahlbeck, D. & Staskawicz, B. J. (1996) Tomato *Prf* is a member of the leucine-rich repeat class of plant disease resistance genes and lies embedded within the *Pto* kinase gene cluster. *Cell* **86**, 123-133
- Sanger, F., Nicklen, S. & Coulson, A. R. (1977) DNA sequencing with chain-terminating inhibitors. *Proc. Natl. Acad. Sci. USA* **75**, 5463-5467
- Scheffler, J. A., Sharpe, A. G., Schmidt, H., Sperling, P., Parkin, I.A. P., Lydiate, D. J. & Heinz, E. (1997) Desaturase multigene families of *Brassica napus* arose through genome duplication. *Theor. Appl. Genet.* **94**, 583-591
- Schmidt, R., Cnops, G. Bancroft, I. & Dean, C. (1992) Construction of an overlapping YAC library of the *Arabidopsis thaliana* genome. *Aust. J. Plant Physiol.* **19**, 341-351
- Schmidt, R., West, J., Love, K., Lenehan, Z., Lister, C., Thompson, H., Bouchez, D. & Dean, C. (1995) Physical map and organization of *Arabidopsis thaliana* chromosome 4. *Science* **270**, 480-483
- Scofield, S. R., Tobias, C. M., Rathjen, J. P., Chang, J. H., Lavelle, D. T., Michelmore, R. W. & Staskawicz, B. J. (1996) Molecular basis of gene-for-gene specificity in bacterial speck disease of tomato. *Science* **274** (5295), 2063-2065
- Shiroishi, T., Sagai, T. & Moriwaki, K. (1993) Hotspots of meiotic recombination in the mouse major histocompatibility complex. *Genetica* **88**, 187-196
- Southern, E. M. (1975) Detection of specific sequence among DNA fragments separated by gelelectrophoresis. *Journal of Molecular Biology* **98**, 503-517
- Staskawicz, B. J. Ausubel, F. M. Baker, B. J., Ellis, J. G. & Jones, J. D. G. (1995) Molecular genetics of plant disease resistance. *Science* **268**, 661-667
- Sun, T.-P., Straus, D. & Ausubel, F. M. (1992) Cloning *Arabidopsis* genes by genomic subtraction. In: Methods in *Arabidopsis* research. (Koncz, C., Chua, N.-H. & Schell, J. eds.) World Scientific Publishing Co. Pte. Ltd., Singapur, pp331-341
- Sillito, D., Parkin, I. A. P., Mayerhofer, R., Lydiate, D. J. & Good, A. G. (2000) *Arabidopsis thaliana*: A source of candidate disease resistance genes for *Brassica napus*. *Genome* **43**, 452-460
- Tang, X. Y., Xie, M. T., Kim, Y. J., Zhou, J. M., Klessig, D. F. & Martin, G. D. (1999) Overexpression of *Pto* activates defense responses and confers broad resistance. *Plant Cell* **11**, 15-29

- Tanksley, S. D., Ganal, M. W., Prince, J. P., de Vicente, M. C., Bonierbale, M. W., Broun, P., Fulton, T. M., Giovannoni, J. J., Grandillo, S. & Martin, G. B. (1992) High density molecular linkage maps of the tomato and potato genomes. *Genetics* **132**, 1141-1160
- Theologis, A., Ecker, J. R., Palm, C. J., Federspiel, N. A., Kaul, S., White, O., Alonso, J., Altafi, H., Araujo, R., Bowman, C. L., Brooks, S. Y., Buehler, E., Chan, A., Chao, Q., Chen, H., Cheuk, R. F., Chin, C., W., Chung, M. K., Conn, L., Conway, A. B., Conway, A. R., Creasy, T. H., Dewar, K., Dunn, P., Etgu, P. et al. (2000) Sequence and analysis of chromosome 1 of the Plant *Arabidopsis thaliana*. *Nature* **408**, 816-820
- Thilmony, R.L., Chen, Z., Bressan, R. A. & Martin, G. B. (1995) Expression of the tomato *Pto* gene in tobacco enhances resistance to *Pseudomonas syringae* pv. tabaci expressing *avrPto*. *Plant Cell* **7** (10), 1529-1236
- Thomas, C. M., Dixon, M. S., Parniske, M., Golstein, C. & Jones, J. D. G. (1998) Genetic and molecular analysis of tomato *Cf* genes for resistance to *Cladosporium fulvum*. *Philosophical Transactions of the Royal Society of London - Series B: Biological Sciences* **353** (1374), 1413-1424
- Thomma, B. P. H. J., Eggermont, K., Penninckx, I.A. M. A., Mauch-Mani, B., Vogelsang, R., Cammue, B. P. A. & Broekaert, W. F. (1998) Separate jasmonate-dependent and salicylate-dependent defense-response pathways in *Arabidopsis* are essential for resistance to distinct microbial pathogens. *Proc. Natl. Acad. Sci. USA* **95**, 15107-15111
- Ullrich, H. Lättig, K. Brennicke, A. & Knoop, V. (1997) Mitochondrial DNA variations and nuclear RFLPs reflect different genetic similarities among 23 *Arabidopsis thaliana* ecotypes. *Plant Mol. Biol.* **33**, 37-45
- Voorrips, R. E., Jongerius, M. C. & Kanne, H. J. (1997) Mapping of two genes for resistance to clubroot (*Plasmodiophora brassicae*) in a population of doubledhaploid lines of *Brassica oleracea* by means of RFLP and AFLP markers. *Theor. Appl. Genet.* **94**, 75-82
- Wagenblatt, B. (1992) Kreuzungsexperimente zur Vererbung der Kohlhernieresistenz bei *Brassica nigra*. Diplomarbeit, Inst. für Angewandte Genetik, Freie Universität Berlin
- Wagner, A. (1998) The fate of duplicated genes: loss or new function? *BioEssays* **20**, 785-788
- Walsh, J. B. (1995) How often do duplicated genes evolve new function? *Genetics* **139**, 421-428
- Ward, E. R. & Jen, G. C. (1990) Isolation of single-copy sequence clones from a yeast artificial chromosome library of randomly-sheared *Arabidopsis thaliana* DNA. *Plant Mol. Biol.* **14**, 561-568
- Weisaeth, G. (1961) Zur Züchtung hernieresistenten Weißkohls II. Prüfung auf Resistenz und Leistung von Inzuchtlinien aus mitteleuropäischen resistenten Herkünften. *Z. Pflanzenz.* **46**, 20-45
- Whitham, S. Mc Cormick, S. & Baker, B. (1996) The *N*-Gene of tobacco confers resistance to tobacco mosaic virus in transgenic tomato. *Proc. Natl. Acad. Sci. USA* **93**, 8776-8781
- Williams, J. G. K., Kubelik, A. R., Livak, K. J., Rafalski, J. A. & Tingey, J. V. (1990) DNA polymorphisms amplified by arbitrary primers are useful as genetic markers. *Nucleic Acids Res.* **18** (22), 6531-6535
- Wit, F. & Van de Weg, M. (1964) Clubroot resistance in turnips (*Brassica campestris* L.). *Euphytica* **13**, 9-18
- Wu, T. C. & Lichten, M. (1995) Factors that affect the location and frequency of meiosis-induced double-strand breaks in *Saccharomyces cerevisiae*. *Genetics* **140**, 55-66
- Yu, I., Parker, J. & Bent, A. F. (1998) Gene-for-gene disease resistance without the hypersensitive response in *Arabidopsis dnd1* mutant. *Proc. Natl. Acad. Sci. USA* **95** (13), 7819-7824