

7 Referenzen

- Altschul, S. F., Gish, W., Miller, W., Myers, E. W., und Lipman, D. J. (1990) Basic local alignment search tool. *J.Mol.Biol.* 215: 403-410
- Ambagala, A. P., Solheim, J. C., und Srikuamran, S. (2005) Viral interference with MHC class I antigen presentation pathway: The battle continues. *Vet.Immunol.Immunopathol.* 107: 1-15
- Andersson, M., Freland, S., Johansson, M. H., Wallin, R., Sandberg, J. K., Chambers, B. J., Christensson, B., Lendahl, U., Lemieux, S. et. al. (1998) MHC class I mosaic mice reveal insights into control of Ly49C inhibitory receptor expression in NK cells. *J.Immunol.* 161: 6475-6479
- Arakawa, H. und Buerstedde, J. M. (2004) Immunoglobulin gene conversion: insights from bursal B cells and the DT40 cell line. *Dev.Dyn.* 229: 458-464
- Baker, K. E. und Parker, R. (2004) Nonsense-mediated mRNA decay: terminating erroneous gene expression. *Curr.Opin.Cell Biol.* 16: 293-299
- Bakker, A. B., Wu, J., Phillips, J. H., und Lanier, L. L. (2000) NK cell activation: distinct stimulatory pathways counterbalancing inhibitory signals. *Hum.Immunol.* 61: 18-27
- Barclay, A. N. (1999) Ig-like domains: evolution from simple interaction molecules to sophisticated antigen recognition. *Proc.Natl.Acad.Sci.U.S.A* 96: 14672-14674
- Barnes, P. D. und Grundy, J. E. (1992) Down-regulation of the class I HLA heterodimer and beta 2-microglobulin on the surface of cells infected with cytomegalovirus. *J.Gen.Viro.* 73 (Pt 9): 2395-2403
- Barten, R., Torkar, M., Haude, A., Trowsdale, J., und Wilson, M. J. (2001) Divergent and convergent evolution of NK-cell receptors. *Trends Immunol.* 22: 52-57
- Beersma, M. F., Bijlmakers, M. J., und Ploegh, H. L. (1993) Human cytomegalovirus down-regulates HLA class I expression by reducing the stability of class I H chains. *J.Immunol.* 151: 4455-4464
- Bernardi, G. (2000) Isochores and the evolutionary genomics of vertebrates. *Gene* 241: 3-17
- Biassoni, R., Cantoni, C., Pende, D., Sivori, S., Parolini, S., Vitale, M., Bottino, C., und Moretta, A. (2001) Human natural killer cell receptors and co-receptors. *Immunol.Rev.* 181: 203-214
- Biassoni, R., Pessino, A., Bottino, C., Pende, D., Moretta, L., und Moretta, A. (1999) The murine homologue of the human NKp46, a triggering receptor involved in the induction of natural cytotoxicity. *Eur.J.Immunol.* 29: 1014-1020
- Binstadt, B. A., Brumbaugh, K. M., und Leibson, P. J. (1997) Signal transduction by human NK cell MHC-recognition receptors. *Immunol.Rev.* 155: 197-203

- Bjorkman, P. J. und Parham, P. (1990) Structure, function, and diversity of class I major histocompatibility complex molecules. *Annu.Rev.Biochem.* 59: 253-288
- Blery, M., Olcese, L., und Vivier, E. (2000) Early signaling via inhibitory and activating NK receptors. *Hum.Immunol.* 61: 51-64
- Borges, L. und Cosman, D. (2000) LIRs/ILTs/MIRs, inhibitory and stimulatory Ig-superfamily receptors expressed in myeloid and lymphoid cells. *Cytokine Growth Factor Rev.* 11: 209-217
- Borges, L., Hsu, M. L., Fanger, N., Kubin, M., und Cosman, D. (1997) A family of human lymphoid and myeloid Ig-like receptors, some of which bind to MHC class I molecules. *J.Immunol.* 159: 5192-5196
- Boyington, J. C., Motyka, S. A., Schuck, P., Brooks, A. G., und Sun, P. D. (2000) Crystal structure of an NK cell immunoglobulin-like receptor in complex with its class I MHC ligand. *Nature* 405: 537-543
- Campbell, K. S., Yusa, S., Kikuchi-Maki, A., und Catina, T. L. (2004) NKp44 triggers NK cell activation through DAP12 association that is not influenced by a putative cytoplasmic inhibitory sequence. *J.Immunol.* 172: 899-906
- Church, G. M. und Gilbert, W. (1984) Genomic sequencing. *Proc.Natl.Acad.Sci.U.S.A* 81: 1991-1995
- Clemetson, J. M., Polgar, J., Magnenat, E., Wells, T. N., und Clemetson, K. J. (1999) The platelet collagen receptor glycoprotein VI is a member of the immunoglobulin superfamily closely related to Fc α R and the natural killer receptors. *J.Biol.Chem.* 274: 29019-29024
- Colonna, M., Nakajima, H., und Cella, M. (2000) A family of inhibitory and activating Ig-like receptors that modulate function of lymphoid and myeloid cells. *Semin.Immunol.* 12: 121-127
- Colonna, M., Navarro, F., Bellon, T., Llano, M., Garcia, P., Samaridis, J., Angman, L., Cella, M., und Lopez-Botet, M. (1997) A common inhibitory receptor for major histocompatibility complex class I molecules on human lymphoid and myelomonocytic cells. *J.Exp.Med.* 186: 1809-1818
- Colonna, M. und Samaridis, J. (1995) Cloning of immunoglobulin-superfamily members associated with HLA-C and HLA-B recognition by human natural killer cells. *Science* 268: 405-408
- Corpet, F. (1988) Multiple sequence alignment with hierarchical clustering. *Nucleic Acids Res.* 16: 10881-10890
- Cosman, D., Fanger, N., und Borges, L. (1999) Human cytomegalovirus, MHC class I and inhibitory signalling receptors: more questions than answers. *Immunol.Rev.* 168: 177-185
- Cosman, D., Fanger, N., Borges, L., Kubin, M., Chin, W., Peterson, L., und Hsu, M. L. (1997) A novel immunoglobulin superfamily receptor for cellular and viral MHC class I molecules. *Immunity*. 7: 273-282
- Costello, R. T., Sivori, S., Marcenaro, E., Lafage-Pochitaloff, M., Mozziconacci, M. J., Reviron, D., Gastaut, J. A., Pende, D., Olive, D. et. al. (2002) Defective expression and function

of natural killer cell-triggering receptors in patients with acute myeloid leukemia. *Blood* 99: 3661-3667

Dawkins, R., Leelayuwat, C., Gaudieri, S., Tay, G., Hui, J., Cattley, S., Martinez, P., und Kulski, J. (1999) Genomics of the major histocompatibility complex: haplotypes, duplication, retroviruses and disease. *Immunol Rev* 167: 275-304

de Wit, T. P., Morton, H. C., Capel, P. J., und van De Winkel, J. G. (1995) Structure of the gene for the human myeloid IgA Fc receptor (CD89). *J.Immunol.* 155: 1203-1209

DeLano, W. L. (2002) The PyMOL Molecular Graphics System (2002) on World Wide Web <http://www.pymol.org>.

Dennis, G., Jr., Kubagawa, H., und Cooper, M. D. (2000) Paired Ig-like receptor homologs in birds and mammals share a common ancestor with mammalian Fc receptors. *Proc.Natl.Acad.Sci.U.S.A* 97: 13245-13250

Dennis, G., Jr., Stephan, R. P., Kubagawa, H., und Cooper, M. D. (1999) Characterization of paired Ig-like receptors in rats. *J.Immunol.* 163: 6371-6377

Djeu, J. Y., Jiang, K., und Wei, S. (2002) A view to a kill: signals triggering cytotoxicity. *Clin.Cancer Res.* 8: 636-640

Döhring, C., Samaridis, J., und Colonna, M. (1996) Alternatively spliced forms of human killer inhibitory receptors. *Immunogenetics* 44: 227-230

du Pasquier L. (2001) The immune system of invertebrates and vertebrates. *Comp Biochem.Physiol B Biochem.Mol.Biol.* 129: 1-15

Ehlers, A., Beck, S., Forbes, S. A., Trowsdale, J., Volz, A., Younger, R., und Ziegler, A. (2000) MHC-linked olfactory receptor loci exhibit polymorphism and contribute to extended HLA/OR-haplotypes. *Genome Res.* 10: 1968-1978

Ezumi, Y., Uchiyama, T., und Takayama, H. (2000a) Molecular cloning, genomic structure, chromosomal localization, and alternative splice forms of the platelet collagen receptor glycoprotein VI. *Biochem.Biophys.Res.Commun.* 277: 27-36

Ezumi, Y., Uchiyama, T., und Takayama, H. (2000b) Molecular cloning, genomic structure, chromosomal localization, and alternative splice forms of the platelet collagen receptor glycoprotein VI. *Biochem.Biophys.Res.Commun.* 277: 27-36

Falco, M., Cantoni, C., Bottino, C., Moretta, A., und Biassoni, R. (1999) Identification of the rat homologue of the human NKp46 triggering receptor. *Immunol.Lett.* 68: 411-414

Fan, Q. R., Long, E. O., und Wiley, D. C. (2000) A disulfide-linked natural killer cell receptor dimer has higher affinity for HLA-C than wild-type monomer. *Eur.J.Immunol.* 30: 2692-2697

Faure, M. und Long, E. O. (2002) KIR2DL4 (CD158d), an NK cell-activating receptor with inhibitory potential. *J.Immunol.* 168: 6208-6214

Feinberg, A. P. und Vogelstein, B. (1984) "A technique for radiolabeling DNA restriction endonuclease fragments to high specific activity". Addendum. *Anal.Biochem.* 137: 266-267

- Galtier, N., Gouy, M., und Gautier, C. (1996) SEAVIEW and PHYLO_WIN: two graphic tools for sequence alignment and molecular phylogeny. *Comput.Appl.Biosci.* 12: 543-548
- Guethlein, L. A., Flodin, L. R., Adams, E. J., und Parham, P. (2002) NK cell receptors of the orangutan (*Pongo pygmaeus*): a pivotal species for tracking the coevolution of killer cell Ig-like receptors with MHC- C. *J.Immunol.* 169: 220-229
- Guex, N. und Peitsch, M. C. (1997) SWISS-MODEL and the Swiss-PdbViewer: an environment for comparative protein modeling. *Electrophoresis* 18: 2714-2723
- Hansasuta, P., Dong, T., Thananchai, H., Weekes, M., Willberg, C., Aldemir, H., Rowland-Jones, S., und Braud, V. M. (2004) Recognition of HLA-A3 and HLA-A11 by KIR3DL2 is peptide-specific. *Eur.J.Immunol.* 34: 1673-1679
- Hao, L. und Nei, M. (2004) Genomic organization and evolutionary analysis of Ly49 genes encoding the rodent natural killer cell receptors: rapid evolution by repeated gene duplication. *Immunogenetics* 56: 343-354
- Harpaz, Y. und Chothia, C. (1994) Many of the immunoglobulin superfamily domains in cell adhesion molecules and surface receptors belong to a new structural set which is close to that containing variable domains. *J.Mol.Biol.* 238: 528-539
- Hieter, P. A., Hollis, G. F., Korsmeyer, S. J., Waldmann, T. A., und Leder, P. (1981) Clustered arrangement of immunoglobulin lambda constant region genes in man. *Nature* 294: 536-540
- Hillier, L. W., Miller, W., Birney, E., Warren, W., Hardison, R. C., Ponting, C. P., Bork, P., Burt, D. W., Groenen, M. A. et. al. (2004) Sequence and comparative analysis of the chicken genome provide unique perspectives on vertebrate evolution. *Nature* 432: 695-716
- Hoelsbrekken, S. E., Nylenne, O., Saether, P. C., Slettedal, I. O., Ryan, J. C., Fossum, S., und Dissen, E. (2003) Cutting edge: molecular cloning of a killer cell Ig-like receptor in the mouse and rat. *J.Immunol.* 170: 2259-2263
- Honjo, T., Nakai, S., Nishida, Y., Kataoka, T., Yamawaki-Kataoka, Y., Takahashi, N., Obata, M., Shimizu, A., Yaoita, Y. et. al. (1981) Rearrangements of immunoglobulin genes during differentiation and evolution. *Immunol.Rev.* 59: 33-67
- Kaufman, J., Milne, S., Gobel, T. W., Walker, B. A., Jacob, J. P., Auffray, C., Zoorob, R., und Beck, S. (1999a) The chicken B locus is a minimal essential major histocompatibility complex. *Nature* 401: 923-925
- Kaufman, J., Milne, S., Gobel, T. W., Walker, B. A., Jacob, J. P., Auffray, C., Zoorob, R., und Beck, S. (1999b) The chicken B locus is a minimal essential major histocompatibility complex. *Nature* 401: 923-925
- Kehrel, B., Wierwille, S., Clemetson, K. J., Anders, O., Steiner, M., Knight, C. G., Farndale, R. W., Okuma, M., und Barnes, M. J. (1998) Glycoprotein VI is a major collagen receptor for platelet activation: it recognizes the platelet-activating quaternary structure of collagen, whereas CD36, glycoprotein IIb/IIIa, and von Willebrand factor do not. *Blood* 91: 491-499
- Kinet, J. P. und Launay, P. (2000) Fc alpha/microR: single member or first born in the family? *Nat.Immunol.* 1: 371-372

- Kollnberger, S., Bird, L. A., Roddis, M., Hacquard-Bouder, C., Kubagawa, H., Bodmer, H. C., Breban, M., McMichael, A. J., und Bowness, P. (2004a) HLA-B27 heavy chain homodimers are expressed in HLA-B27 transgenic rodent models of spondyloarthritis and are ligands for paired Ig-like receptors. *J.Immunol.* 173: 1699-1710
- Kollnberger, S., Bird, L. A., Roddis, M., Hacquard-Bouder, C., Kubagawa, H., Bodmer, H. C., Breban, M., McMichael, A. J., und Bowness, P. (2004b) HLA-B27 heavy chain homodimers are expressed in HLA-B27 transgenic rodent models of spondyloarthritis and are ligands for paired Ig-like receptors. *J.Immunol.* 173: 1699-1710
- Kubagawa, H., Burrows, P. D., und Cooper, M. D. (1997) A novel pair of immunoglobulin-like receptors expressed by B cells and myeloid cells. *Proc.Natl.Acad.Sci.U.S.A* 94: 5261-5266
- Lander, E. S., Linton, L. M., Birren, B., Nusbaum, C., Zody, M. C., Baldwin, J., Devon, K., Dewar, K., Doyle, M. et. al. (2001) Initial sequencing and analysis of the human genome. *Nature* 409: 860-921
- Lanier, L. L. (1998) Follow the leader: NK cell receptors for classical and nonclassical MHC class I. *Cell* 92: 705-707
- Lanier, L. L., Corliss, B. C., Wu, J., Leong, C., und Phillips, J. H. (1998) Immunoreceptor DAP12 bearing a tyrosine-based activation motif is involved in activating NK cells. *Nature* 391: 703-707
- Laun, K., Coggill, P., Palmer, S., Sims, S., Ning, Z., Ragoussis, J., Volpi, E., Wilson, N., Beck, S. et. al. (2006) A Maximally Expanded Leukocyte Receptor Complex in the Chicken Contrasts its Minimal Essential Major Histocompatibility Complex (zur Publikation eingereicht).
- Lebbink, R. J., de Ruiter, T., Verbrugge, A., Bril, W. S., und Meynard, L. (2004) The mouse homologue of the leukocyte-associated Ig-like receptor-1 is an inhibitory receptor that recruits Src homology region 2-containing protein tyrosine phosphatase (SHP)-2, but not SHP-1. *J.Immunol.* 172: 5535-5543
- Letunic, I., Goodstadt, L., Dickens, N. J., Doerks, T., Schultz, J., Mott, R., Ciccarelli, F., Copley, R. R., Ponting, C. P. et. al. (2002) Recent improvements to the SMART domain-based sequence annotation resource. *Nucleic Acids Res.* 30: 242-244
- Liang, S., Baibakov, B., und Horuzsko, A. (2002) HLA-G inhibits the functions of murine dendritic cells via the PIR-B immune inhibitory receptor. *Eur.J.Immunol.* 32: 2418-2426
- Litman, G. W., Anderson, M. K., und Rast, J. P. (1999) Evolution of antigen binding receptors. *Annu.Rev.Immunol.* 17: 109-147
- Ljunggren, H. G. und Karre, K. (1990) In search of the 'missing self': MHC molecules and NK cell recognition. *Immunol.Today* 11: 237-244
- Lodden, M. B. und Lanier, L. L. (2005) Viral modulation of NK cell immunity. *Nat.Rev.Microbiol.* 3: 59-69

- Long, E. O., Barber, D. F., Burshtyn, D. N., Faure, M., Peterson, M., Rajagopalan, S., Renard, V., Sandusky, M., Stebbins, C. C. et. al. (2001) Inhibition of natural killer cell activation signals by killer cell immunoglobulin-like receptors (CD158). *Immunol.Rev.* 181: 223-233
- Martin, A. M., Kulski, J. K., Witt, C., Pontarotti, P., und Christiansen, F. T. (2002) Leukocyte Ig-like receptor complex (LRC) in mice and men. *Trends Immunol.* 23: 81-88
- Maruoka, T., Nagata, T., und Kasahara, M. (2004) Identification of the rat IgA Fc receptor encoded in the leukocyte receptor complex. *Immunogenetics* 55: 712-716
- McQueen, K. L. und Parham, P. (2002) Variable receptors controlling activation and inhibition of NK cells. *Curr.Opin.Immunol.* 14: 615-621
- McQueen, K. L., Wilhelm, B. T., Harden, K. D., und Mager, D. L. (2002) Evolution of NK receptors: a single Ly49 and multiple KIR genes in the cow. *Eur.J.Immunol.* 32: 810-817
- Meyaard, L., Adema, G. J., Chang, C., Woollatt, E., Sutherland, G. R., Lanier, L. L., und Phillips, J. H. (1997) LAIR-1, a novel inhibitory receptor expressed on human mononuclear leukocytes. *Immunity*. 7: 283-290
- Meyaard, L., van der Vuurst de Vries AR, de Ruiter, T., Lanier, L. L., Phillips, J. H., und Clevers, H. (2001) The epithelial cellular adhesion molecule (Ep-CAM) is a ligand for the leukocyte-associated immunoglobulin-like receptor (LAIR). *J.Exp.Med.* 194: 107-112
- Monteiro, R. C. und van De Winkel, J. G. (2003) IgA Fc receptors. *Annu.Rev.Immunol.* 21: 177-204
- Moretta, A., Bottino, C., Vitale, M., Pende, D., Cantoni, C., Mingari, M. C., Biassoni, R., und Moretta, L. (2001) Activating receptors and coreceptors involved in human natural killer cell-mediated cytotoxicity. *Annu.Rev.Immunol.* 19: 197-223
- Moretta, A. und Moretta, L. (1997) HLA class I specific inhibitory receptors. *Curr.Opin.Immunol.* 9: 694-701
- Moretta, A., Sivori, S., Vitale, M., Pende, D., Morelli, L., Augugliaro, R., Bottino, C., und Moretta, L. (1995) Existence of both inhibitory (p58) and activatory (p50) receptors for HLA-C molecules in human natural killer cells. *J.Exp.Med.* 182: 875-884
- Morton, H. C., Pleass, R. J., Storset, A. K., Dissen, E., Williams, J. L., Brandtzaeg, P., und Woof, J. M. (2004) Cloning and characterization of an immunoglobulin A Fc receptor from cattle. *Immunology* 111: 204-211
- Mullis, K., Falloona, F., Scharf, S., Saiki, R., Horn, G., und Erlich, H. (1992) Specific enzymatic amplification of DNA in vitro: the polymerase chain reaction. 1986. *Biotechnology* 24: 17-27
- Nei, M. und Gojobori, T. (1986) Simple methods for estimating the numbers of synonymous and nonsynonymous nucleotide substitutions. *Mol.Biol.Evol.* 3: 418-426
- Nei, M. und Rooney, A. P. (2005) Concerted and Birth-and-Death Evolution of Multigene Families. *Annu.Rev.Genet.*

- Nieswandt, B., Bergmeier, W., Schulte, V., Rackebrandt, K., Gessner, J. E., und Zirngibl, H. (2000) Expression and function of the mouse collagen receptor glycoprotein VI is strictly dependent on its association with the FcRgamma chain. *J.Biol.Chem.* 275: 23998-24002
- Nikolaidis, N., Klein, J., und Nei, M. (2005a) Origin and evolution of the Ig-like domains present in mammalian leukocyte receptors: insights from chicken, frog, and fish homologues. *Immunogenetics* 57: 151-157
- Nikolaidis, N., Makalowska, I., Chalkia, D., Makalowski, W., Klein, J., und Nei, M. (2005b) Origin and evolution of the chicken leukocyte receptor complex. *Proc.Natl.Acad.Sci.U.S.A* 102: 4057-4062
- Ohno, S. (1970) Evolution by gene duplication. New York: Springer.
- Ota, T. und Nei, M. (1994) Variance and covariances of the numbers of synonymous and nonsynonymous substitutions per site. *Mol.Biol.Evol.* 11: 613-619
- Pessino, A., Sivori, S., Bottino, C., Malaspina, A., Morelli, L., Moretta, L., Biassoni, R., und Moretta, A. (1998) Molecular cloning of NKp46: a novel member of the immunoglobulin superfamily involved in triggering of natural cytotoxicity. *J.Exp.Med.* 188: 953-960
- Rajagopalan, S., Fu, J., und Long, E. O. (2001) Cutting edge: induction of IFN-gamma production but not cytotoxicity by the killer cell Ig-like receptor KIR2DL4 (CD158d) in resting NK cells. *J.Immunol.* 167: 1877-1881
- Rajalingam, R., Parham, P., und Abi-Rached, L. (2004) Domain shuffling has been the main mechanism forming new hominoid killer cell Ig-like receptors. *J.Immunol.* 172: 356-369
- Ravetch, J. V. und Lanier, L. L. (2000) Immune inhibitory receptors. *Science* 290: 84-89
- Samaridis, J. und Colonna, M. (1997) Cloning of novel immunoglobulin superfamily receptors expressed on human myeloid and lymphoid cells: structural evidence for new stimulatory and inhibitory pathways. *Eur.J.Immunol.* 27: 660-665
- Schultz, J., Milpetz, F., Bork, P., und Ponting, C. P. (1998) SMART, a simple modular architecture research tool: identification of signaling domains. *Proc.Natl.Acad.Sci.U.S.A* 95: 5857-5864
- Scott, P. und Trinchieri, G. (1995) The role of natural killer cells in host-parasite interactions. *Curr.Opin.Immunol.* 7: 34-40
- Sharma, J. M. und Okazaki, W. (1981) Natural killer cell activity in chickens: target cell analysis and effect of antithymocyte serum on effector cells. *Infect.Immun.* 31: 1078-1085
- Shiina, T., Shimizu, S., Hosomichi, K., Kohara, S., Watanabe, S., Hanzawa, K., Beck, S., Kulski, J. K., und Inoko, H. (2004) Comparative genomic analysis of two avian (quail and chicken) MHC regions. *J.Immunol.* 172: 6751-6763
- Shiroishi, M., Tsumoto, K., Amano, K., Shirakihara, Y., Colonna, M., Braud, V. M., Allan, D. S., Makadzange, A., Rowland-Jones, S. et. al. (2003) Human inhibitory receptors Ig-like transcript 2 (ILT2) and ILT4 compete with CD8 for MHC class I binding and bind preferentially to HLA-G. *Proc.Natl.Acad.Sci.U.S.A* 100: 8856-8861

- Smit, A., Hubley, R., und Green, P. (2004) RepeatMasker Open-3.0.1996-2004 <http://www.repeatmasker.org>
- Songyang, Z., Shoelson, S. E., Chaudhuri, M., Gish, G., Pawson, T., Haser, W. G., King, F., Roberts, T., Ratnofsky, S. et. al. (1993) SH2 domains recognize specific phosphopeptide sequences. *Cell* 72: 767-778
- Sonnhammer, E. L. und Durbin, R. (1995) A dot-matrix program with dynamic threshold control suited for genomic DNA and protein sequence analysis. *Gene* 167: GC1-10
- Southern, E. M. (1975) Detection of specific sequences among DNA fragments separated by gel electrophoresis. *J.Mol.Biol.* 98: 503-517
- Stewart-Jones, G. B., di Gleria, K., Kollnberger, S., McMichael, A. J., Jones, E. Y., und Bowness, P. (2005) Crystal structures and KIR3DL1 recognition of three immunodominant viral peptides complexed to HLA-B*2705. *Eur.J.Immunol.* 35: 341-351
- Storset, A. K., Kulberg, S., Berg, I., Boysen, P., Hope, J. C., und Dissen, E. (2004) NKp46 defines a subset of bovine leukocytes with natural killer cell characteristics. *Eur.J.Immunol.* 34: 669-676
- Storset, A. K., Slettedal, I. O., Williams, J. L., Law, A., und Dissen, E. (2003) Natural killer cell receptors in cattle: a bovine killer cell immunoglobulin-like receptor multigene family contains members with divergent signaling motifs. *Eur.J.Immunol.* 33: 980-990
- Suto, Y., Maenaka, K., Yabe, T., Hirai, M., Tokunaga, K., Tadok, K., und Juji, T. (1996) Chromosomal localization of the human natural killer cell class I receptor family genes to 19q13.4 by fluorescence in situ hybridization. *Genomics* 35: 270-272
- Takahashi, T., Yawata, M., Raudsepp, T., Lear, T. L., Chowdhary, B. P., Antczak, D. F., und Kasahara, M. (2004) Natural killer cell receptors in the horse: evidence for the existence of multiple transcribed LY49 genes. *Eur.J.Immunol.* 34: 773-784
- Tay, C. H., Szomolanyi-Tsuda, E., und Welsh, R. M. (1998) Control of infections by NK cells. *Curr.Top.Microbiol.Immunol.* 230: 193-220
- Tomasello, E., Blery, M., Vely, E., und Vivier, E. (2000) Signaling pathways engaged by NK cell receptors: double concerto for activating receptors, inhibitory receptors and NK cells. *Semin.Immunol.* 12: 139-147
- Trowsdale, J. (2001) Genetic and functional relationships between MHC and NK receptor genes. *Immunity*. 15: 363-374
- Trowsdale, J., Barten, R., Haude, A., Stewart, C. A., Beck, S., und Wilson, M. J. (2001) The genomic context of natural killer receptor extended gene families. *Immunol.Rev.* 181: 20-38
- Ujike, A., Takeda, K., Nakamura, A., Ebihara, S., Akiyama, K., und Takai, T. (2002) Impaired dendritic cell maturation and increased T(H)2 responses in PIR-B(-/-) mice. *Nat.Immunol.* 3: 542-548
- Unkeless, J. C. und Jin, J. (1997) Inhibitory receptors, ITIM sequences and phosphatases. *Curr.Opin.Immunol.* 9: 338-343

- Vankayalapati, R., Wizel, B., Weis, S. E., Safi, H., Lakey, D. L., Mandelboim, O., Samten, B., Porgador, A., und Barnes, P. F. (2002) The NKp46 receptor contributes to NK cell lysis of mononuclear phagocytes infected with an intracellular bacterium. *J.Immunol.* 168: 3451-3457
- Viertlboeck, B. C., Crooijmans, R. P., Groenen, M. A., und Gobel, T. W. (2004) Chicken Ig-like receptor B2, a member of a multigene family, is mainly expressed on B lymphocytes, recruits both Src homology 2 domain containing protein tyrosine phosphatase (SHP)-1 and SHP-2, and inhibits proliferation. *J.Immunol.* 173: 7385-7393
- Vilches, C., Pando, M. J., und Parham, P. (2000) Genes encoding human killer-cell Ig-like receptors with D1 and D2 extracellular domains all contain untranslated pseudoexons encoding a third Ig-like domain. *Immunogenetics* 51: 639-646
- Vitale, M., Bottino, C., Sivori, S., Sanseverino, L., Castriconi, R., Marcenaro, E., Augugliaro, R., Moretta, L., und Moretta, A. (1998) NKp44, a novel triggering surface molecule specifically expressed by activated natural killer cells, is involved in non-major histocompatibility complex-restricted tumor cell lysis. *J.Exp.Med.* 187: 2065-2072
- Vitale, M., Castriconi, R., Parolini, S., Pende, D., Hsu, M. L., Moretta, L., Cosman, D., und Moretta, A. (1999) The leukocyte Ig-like receptor (LIR)-1 for the cytomegalovirus UL18 protein displays a broad specificity for different HLA class I alleles: analysis of LIR-1 + NK cell clones. *Int Immunol* 11: 29-35
- Volz, A., Wende, H., Laun, K., und Ziegler, A. (2001) Genesis of the ILT/LIR/MIR clusters within the human leukocyte receptor complex. *Immunol.Rev.* 181: 39-51
- Wagtmann, N., Rojo, S., Eichler, E., Mohrenweiser, H., und Long, E. O. (1997) A new human gene complex encoding the killer cell inhibitory receptors and related monocyte/macrophage receptors. *Curr.Biol.* 7: 615-618
- Wende H. (2004) Genetische Charakterisierung des „Leukocyte Receptor Complex“ und Entwicklung einer Methode zum Nachweis seiner Produkte im Einzelzellmaßstab. Dissertation Universität Potsdam (<http://pub.ub.uni-potsdam.de/2004meta/0029/door.htm>)
- Wende, H., Colonna, M., Ziegler, A., und Volz, A. (1999) Organization of the leukocyte receptor cluster (LRC) on human chromosome 19q13.4. *Mamm.Genome* 10: 154-160
- Wende, H., Volz, A., und Ziegler, A. (2000) Extensive gene duplications and a large inversion characterize the human leukocyte receptor cluster. *Immunogenetics* 51: 703-713
- Williams AF (1987) A year in the life of the immunoglobulin superfamily. *Immunology Today* 8: 298-303
- Willcox, B. E., Thomas, L. M., und Bjorkman, P. J. (2003a) Crystal structure of HLA-A2 bound to LIR-1, a host and viral major histocompatibility complex receptor. *Nat.Immunol.* 4: 913-919
- Willcox, B. E., Thomas, L. M., und Bjorkman, P. J. (2003b) Crystal structure of HLA-A2 bound to LIR-1, a host and viral major histocompatibility complex receptor. *Nat.Immunol.* 4: 913-919
- Williams, A. F. und Barclay, A. N. (1988) The immunoglobulin superfamily--domains for cell surface recognition. *Annu.Rev.Immunol.* 6: 381-405

- Wilson, M. J., Torkar, M., Haude, A., Milne, S., Jones, T., Sheer, D., Beck, S., und Trowsdale, J. (2000) Plasticity in the organization and sequences of human KIR/ILT gene families. Proc.Natl.Acad.Sci.U.S.A 97: 4778-4783
- Wilson, M. J., Torkar, M., und Trowsdale, J. (1997) Genomic organization of a human killer cell inhibitory receptor gene. *Tissue Antigens* 49: 574-579
- Wu, J., Cherwinski, H., Spies, T., Phillips, J. H., und Lanier, L. L. (2000) DAP10 and DAP12 form distinct, but functionally cooperative, receptor complexes in natural killer cells. *J.Exp.Med.* 192: 1059-1068
- Wu, T. T. und Kabat, E. A. (1970) An analysis of the sequences of the variable regions of Bence Jones proteins and myeloma light chains and their implications for antibody complementarity. *J.Exp.Med.* 132: 211-250
- Yamashita, I., Nagata, T., Tada, T., und Nakayama, T. (1993) CD69 cell surface expression identifies developing thymocytes which audition for T cell antigen receptor-mediated positive selection. *Int.Immunol.* 5: 1139-1150
- Yusa, S., Catina, T. L., und Campbell, K. S. (2002) SHP-1- and phosphotyrosine-independent inhibitory signaling by a killer cell Ig-like receptor cytoplasmic domain in human NK cells. *J.Immunol.* 168: 5047-5057
- Zekarias, B., Ter Huurne, A. A., Landman, W. J., Rebel, J. M., Pol, J. M., und Gruys, E. (2002) Immunological basis of differences in disease resistance in the chicken. *Vet.Res.* 33: 109-125