

## 8 Bibliographie

- Applequist, S.E., Selg, M., Raman, C. und Jack, H.M. (1997) Cloning and characterization of HUPF1, a human homolog of the *Saccharomyces cerevisiae* nonsense mRNA-reducing UPF1 protein. *Nucleic Acids Res.*, **25**, 814-821.
- Batt, D.B., Luo, Y. und Carmichael, G.G. (1994) Polyadenylation and transcription termination in gene constructs containing multiple tandem polyadenylation signals. *Nucleic Acids Res.*, **22**, 2811-6.
- Beelman, C.A. und Parker, R. (1995) Degradation of mRNA in eukaryotes. *Cell*, **81**, 179-183.
- Belgrader, P., Cheng, J. und Maquat, L.E. (1993) Evidence to implicate translation by ribosomes in the mechanism by which nonsense codons reduce the nuclear level of human triosephosphate isomerase mRNA. *Proc. Natl. Acad. Sci. U. S. A.*, **90**, 482-486.
- Bhattacharya, A., Czaplinski, K., Trifillis, P., He, F., Jacobson, A. und Peltz, S.W. (2000) Characterization of the biochemical properties of the human Upf1 gene product that is involved in nonsense-mediated mRNA decay. *RNA*, **6**, 1226-1235.
- Brocke, K.S., Neu-Yilik, G., Gehring, N.H., Hentze, M.W. und Kulozik, A.E. (2002) The human intronless melanocortin 4-receptor gene is NMD insensitive. *Hum. Mol. Genet.*, **11**, 331-335.
- Brouwer, R., Allmang, C., Raijmakers, R., van Aarssen, Y., Egberts, W.V., Petfalski, E., van Venrooij, W.J., Tollervey, D. und Pruijn, G.J. (2001) Three novel components of the human exosome. *J. Biol. Chem.*, **276**, 6177-6184.
- Burkard, K.T. und Butler, J.S. (2000) A nuclear 3'-5' exonuclease involved in mRNA degradation interacts with Poly(A) polymerase and the hnRNA protein Npl3p. *Mol Cell Biol*, **20**, 604-16.
- Caceres, J.F., Sreaton, G.R. und Krainer, A.R. (1998) A specific subset of SR proteins shuttles continuously between the nucleus and the cytoplasm. *Genes Dev.*, **12**, 55-66.

- Cai, Z., Gorin, A., Frederick, R., Ye, X., Hu, W., Majumdar, A., Kettani, A. und Patel, D.J. (1998) Solution structure of P22 transcriptional antitermination N peptide-boxB RNA complex. *Nature Struct. Biol.*, **5**, 203-212.
- Carter, M.S., Doskow, J., Morris, P., Li, S., Nhim, R.P., Sandstedt, S. und Wilkinson, M.F. (1995) A regulatory mechanism that detects premature nonsense codons in T-cell receptor transcripts in vivo is reversed by protein synthesis inhibitors in vitro. *J. Biol. Chem.*, **270**, 28995-29003.
- Carter, M.S., Li, S. und Wilkinson, M.F. (1996) A splicing-dependent regulatory mechanism that detects translation signals. *EMBO J.*, **15**, 5965-5975.
- Cattaneo, M., Chantarangkul, V., Taioli, E., Santos, J.H. und Tagliabue, L. (1999) The G20210A mutation of the prothrombin gene in patients with previous first episodes of deep-vein thrombosis: prevalence and association with factor V G1691A, methylenetetrahydrofolate reductase C677T and plasma prothrombin levels. *Thromb Res.*, **93**, 1-8.
- Cazzola, M. und Skoda, R.C. (2000) Translational pathophysiology: a novel molecular mechanism of human disease. *Blood*, **95**, 3280-3288.
- Chang, J.C. und Kan, Y.W. (1979) beta 0 thalassemia, a nonsense mutation in man. *Proc. Natl. Acad. Sci. U. S. A.*, **76**, 2886-2889.
- Chen, C.Y., Gherzi, R., Ong, S.E., Chan, E.L., Raijmakers, R., Pruijn, G.J., Stoecklin, G., Moroni, C., Mann, M. und Karin, M. (2001) AU Binding Proteins Recruit the Exosome to Degrade ARE-Containing mRNAs. *Cell*, **107**, 451-464.
- Chen, C.Y. und Shyu, A.B. (1995) AU-rich elements: characterization and importance in mRNA degradation. *Trends Biochem. Sci.*, **20**, 465-470.
- Chen, F., MacDonald, C.C. und Wilusz, J. (1995) Cleavage site determinants in the mammalian polyadenylation signal. *Nucleic Acids Res.*, **23**, 2614-20.
- Chou, Z.F., Chen, F. und Wilusz, J. (1994) Sequence and position requirements for uridylate-rich downstream elements of polyadenylation signals. *Nucleic Acids Res.*, **22**, 2525-31.
- Cilley, C.D. und Williamson, J.R. (1997) Analysis of bacteriophage N protein and peptide binding to boxB RNA using polyacrylamide gel coelectrophoresis (PACE). *RNA*, **3**, 57-67.

- Colgan, D.F. und Manley, J.L. (1997) Mechanism and regulation of mRNA polyadenylation. *Genes Dev.*, **11**, 2755-66.
- Coller, J.M., Gray, N.K. und Wickens, M.P. (1998) mRNA stabilization by poly(A) binding protein is independent of poly(A) and requires translation. *Genes Dev.*, **12**, 3226-3235.
- Conne, B., Stutz, A. und Vassalli, J.D. (2000) The 3' untranslated region of messenger RNA: A molecular 'hotspot' for pathology? *Nat Med.*, **6**, 637-41.
- Cui, Y., Dinman, J.D. und Peltz, S.W. (1996) Mof4-1 is an allele of the UPF1/IFS2 gene which affects both mRNA turnover and -1 ribosomal frameshifting efficiency. *EMBO J.*, **15**, 5726-5736.
- Cui, Y., Hagan, K.W., Zhang, S. und Peltz, S.W. (1995) Identification and characterization of genes that are required for the accelerated degradation of mRNAs containing a premature translational termination codon. *Genes Dev.*, **9**, 423-436.
- Culbertson, M.R. (1999) RNA surveillance. Unforeseen consequences for gene expression, inherited genetic disorders and cancer. *Trends Genet.*, **15**, 74-80.
- Czaplinski, K., Ruiz-Echevarria, M.J., Gonzalez, C.I. und Peltz, S.W. (1999) Should we kill the messenger? The role of the surveillance complex in translation termination and mRNA turnover. *BioEssays*, **21**, 685-696.
- Czaplinski, K., Ruiz-Echevarria, M.J., Paushkin, S.V., Han, X., Weng, Y., Perlick, H.A., Dietz, H.C., Ter-Avanesyan, M.D. und Peltz, S.W. (1998) The surveillance complex interacts with the translation release factors to enhance termination and degrade aberrant mRNAs. *Genes Dev.*, **12**, 1665-1677.
- Czaplinski, K., Weng, Y., Hagan, K.W. und Peltz, S.W. (1995) Purification and characterization of the Upf1 protein: a factor involved in translation and mRNA degradation. *RNA*, **1**, 610-623.
- Danckwardt, S., Neu-Yilik, G., Thermann, R., Frede, U., Hentze, M.W. und Kulozik, A.E. (2002) Abnormally spliced beta-globin mRNAs: a single point mutation generates transcripts sensitive and insensitive to nonsense-mediated mRNA decay. *Blood*, **99**, 1811-1816.

- Dang, O.D., Vindigni, A. und Di Cera, E. (1995) An allosteric switch controls the procoagulant and anticoagulant activities of thrombin. *Proc. Natl. Acad. Sci. U. S. A.*, **92**, 5977-5981.
- De Gregorio, E., Baron, J., Preiss, T. und Hentze, M.W. (2001) Tethered-function analysis reveals that eIF4E can recruit ribosomes independent of its binding to the cap structure. *RNA*, **7**, 106-113.
- De Gregorio, E., Preiss, T. und Hentze, M.W. (1999) Translation driven by an eIF4G core domain in vivo. *EMBO J.*, **18**, 4865-4874.
- De Stefano, V., Finazzi, G. und Mannucci, P.M. (1996) Inherited thrombophilia: pathogenesis, clinical syndromes, and management. *Blood*, **87**, 3531-3544.
- Dehlin, E., Wormington, M., Korner, C.G. und Wahle, E. (2000) Cap-dependent deadenylation of mRNA. *EMBO J.*, **19**, 1079-1086.
- Denning, G., Jamieson, L., Maquat, L.E., Thompson, E.A. und Fields, A.P. (2001) Cloning of a novel phosphatidylinositol kinase-related kinase: characterization of the human SMG-1 RNA surveillance protein. *J. Biol. Chem.*, **276**, 22709-22714.
- Dietz, H.C. und Hamosh, A. (1996) Nonstop treatment of cystic fibrosis. *Nature Med.*, **2**, 608-609.
- Dietz, H.C., McIntosh, I., Sakai, L.Y., Corson, G.M., Chalberg, S.C., Pyeritz, R.E. und Francomano, C.A. (1993) Four novel FBN1 mutations: significance for mutant transcript level and EGF-like domain calcium binding in the pathogenesis of Marfan syndrome. *Genomics*, **17**, 468-475.
- Dominski, Z. und Marzluff, W.F. (1999) Formation of the 3' end of histone mRNA. *Gene*, **239**, 1-14.
- Dominski, Z., Zheng, L.X., Sanchez, R. und Marzluff, W.F. (1999) Stem-loop binding protein facilitates 3'-end formation by stabilizing U7 snRNP binding to histone pre-mRNA. *Mol. Cell. Biol.*, **19**, 3561-3570.
- Dreyfuss, G., Matunis, M.J., Pinol-Roma, S. und Burd, C.G. (1993) hnRNP proteins and the biogenesis of mRNA. *Annu. Rev. Biochem.*, **62**, 289-321.

- Forch, P., Puig, O., Kedersha, N., Martinez, C., Granneman, S., Seraphin, B., Anderson, P. und Valcarcel, J. (2000) The apoptosis-promoting factor TIA-1 is a regulator of alternative pre-mRNA splicing. *Mol. Cell*, **6**, 1089-1098.
- Frischmeyer, P.A. und Dietz, H.C. (1999) Nonsense-mediated mRNA decay in health and disease. *Hum. Mol. Genet.*, **8**, 1893-1900.
- Gao, M., Fritz, D.T., Ford, L.P. und Wilusz, J. (2000) Interaction between a poly(A)-specific ribonuclease and the 5' cap influences mRNA deadenylation rates in vitro. *Mol. Cell*, **5**, 479-488.
- Gatfield, D., Le Hir, H., Schmitt, C., Braun, I.C., Kocher, T., Wilm, M. und Izaurralde, E. (2001) The DExH/D box protein HEL/UAP56 is essential for mRNA nuclear export in Drosophila. *Curr. Biol.*, **11**, 1716-1721.
- Ghilardi, N., Wiestner, A. und Skoda, R.C. (1998) Thrombopoietin production is inhibited by a translational mechanism. *Blood*, **92**, 4023-4030.
- Gonzalez, C.I., Ruiz-Echevarria, M.J., Vasudevan, S., Henry, M.F. und Peltz, S.W. (2000) The yeast hnRNP-like protein Hrp1/Nab4 marks a transcript for nonsense-mediated mRNA decay. *Mol. Cell*, **5**, 489-499.
- Graveley, B.R. (2001) Alternative splicing: increasing diversity in the proteomic world. *Trends Genet.*, **17**, 100-107.
- Gray, N.K., Coller, J.M., Dickson, K.S. und Wickens, M. (2000) Multiple portions of poly(A)-binding protein stimulate translation in vivo. *EMBO J.*, **19**, 4723-4733.
- Gray, N.K. und Wickens, M. (1998) Control of translation initiation in animals. *Annu. Rev. Cell Dev. Biol.*, **14**, 399-458.
- Grosset, C., Chen, C.Y., Xu, N., Sonenberg, N., Jacquemin-Sablon, H. und Shyu, A.B. (2000) A mechanism for translationally coupled mRNA turnover: interaction between the poly(A) tail and a c-fos RNA coding determinant via a protein complex. *Cell*, **103**, 29-40.
- Hagemeier, C. (1996) Site-directed mutagenesis using a uracil-containing phagemid template. *Methods Mol. Biol.*, **57**, 45-54.

- Hall, G.W. und Thein, S. (1994) Nonsense codon mutations in the terminal exon of the beta-globin gene are not associated with a reduction in beta-mRNA accumulation: a mechanism for the phenotype of dominant beta-thalassemia. *Blood*, **83**, 2031-2037.
- Hanahan, D., Jessee, J. und Bloom, F.R. (1991) Plasmid transformation of *Escherichia coli* and other bacteria. *Methods Enzymol.*, **204**, 63-113.
- He, F., Peltz, S.W., Donahue, J.L., Rosbash, M. und Jacobson, A. (1993) Stabilization and ribosome association of unspliced pre-mRNAs in a yeast upf1- mutant. *Proc. Natl. Acad. Sci. U.S.A.*, **90**, 7034-7038.
- Hentze, M.W. (2001) Protein synthesis. Believe it or not-translation in the nucleus. *Science*, **293**, 1058-1059.
- Hentze, M.W., Caughman, S.W., Rouault, T.A., Barriocanal, J.G., Dancis, A., Harford, J.B. und Klausner, R.D. (1987a) Identification of the iron-responsive element for the translational regulation of human ferritin mRNA. *Science*, **238**, 1570-1573.
- Hentze, M.W. und Kuhn, L.C. (1996) Molecular control of vertebrate iron metabolism: mRNA-based regulatory circuits operated by iron, nitric oxide, and oxidative stress. *Proc. Natl. Acad. Sci. U. S. A.*, **93**, 8175-8182.
- Hentze, M.W. und Kulozik, A.E. (1999) A perfect message: RNA surveillance and nonsense-mediated decay. *Cell*, **96**, 307-310.
- Hentze, M.W., Rouault, T.A., Caughman, S.W., Dancis, A., Harford, J.B. und Klausner, R.D. (1987b) A cis-acting element is necessary and sufficient for translational regulation of human ferritin expression in response to iron. *Proc. Natl. Acad. Sci. U. S. A.*, **84**, 6730-6734.
- Howard, M., Frizzel, R.A. und Bedwell, D.M. (1996) Aminoglycoside antibiotics restore CFTR function by overcoming premature stop mutations. *Nature Med.*, **2**, 467-469.
- Iborra, F.J., Jackson, D.A. und Cook, P.R. (2001) Coupled transcription and translation within nuclei of mammalian cells. *Science*, **293**, 1139-1142.
- International Human Genome Sequencing Consortium (2001) Initial sequencing and analysis of the human genome. *Nature*, **409**, 860-921.

- Johnstone, O. und Lasko, P. (2001) Translational regulation and RNA localization in Drosophila oocytes and embryos. *Annu. Rev. Genet.*, **35**, 365-406.
- Kataoka, N., Diem, M.D., Kim, V.N., Yong, J. und Dreyfuss, G. (2001) Magoh, a human homolog of Drosophila mago nashi protein, is a component of the splicing-dependent exon-exon junction complex. *EMBO J.*, **20**, 6424-6433.
- Kataoka, N., Yong, J., Kim, V.N., Velazquez, F., Perkinson, R.A., Wang, F. und Dreyfuss, G. (2000) Pre-mRNA splicing imprints mRNA in the nucleus with a novel RNA-binding protein that persists in the cytoplasm. *Mol. Cell*, **6**, 673-682.
- Kim, V.N. und Dreyfuss, G. (2001) Nuclear mRNA binding proteins couple pre-mRNA splicing and post-splicing events. *Mol. Cells*, **12**, 1-10.
- Kim, V.N., Kataoka, N. und Dreyfuss, G. (2001a) Role of the nonsense-mediated decay factor hUpf3 in the splicing-dependent exon-exon junction complex. *Science*, **293**, 1832-1836.
- Kim, V.N., Yong, J., Kataoka, N., Abel, L., Diem, M.D. und Dreyfuss, G. (2001b) The Y14 protein communicates to the cytoplasm the position of exon-exon junctions. *EMBO J.*, **20**, 2062-2068.
- Koster, T., Rosendaal, F.R., de Ronde, H., Briet, E., Vandebroucke, J.P. und Bertina, R.M. (1993) Venous thrombosis due to poor anticoagulant response to activated protein C: Leiden Thrombophilia Study. *Lancet*, **342**, 1503-1506.
- Kugler, W., Enssle, J., Hentze, M.W. und Kulozik, A.E. (1995) Nuclear degradation of nonsense mutated beta-globin mRNA: a post-transcriptional mechanism to protect heterozygotes from severe clinical manifestations of beta-thalassemia? *Nucleic Acids Res*, **23**, 413-8.
- Lane, D.A. und Grant, P.J. (2000) Role of hemostatic gene polymorphisms in venous and arterial thrombotic disease. *Blood*, **95**, 1517-32.
- Le Hir, H., Gatfield, D., Izaurralde, E. und Moore, M.J. (2001) The exon-exon junction complex provides a binding platform for factors involved in mRNA export and nonsense-mediated mRNA decay. *EMBO J.*, **20**, 4987-4997.

- Le Hir, H., Izaurrealde, E., Maquat, L.E. und Moore, M.J. (2000a) The spliceosome deposits multiple proteins 20-24 nucleotides upstream of mRNA exon-exon junctions. *EMBO J.*, **19**, 6860-6869.
- Le Hir, H., Moore, M.J. und Maquat, L.E. (2000b) Pre-mRNA splicing alters mRNP composition: evidence for stable association of proteins at exon-exon junctions. *Genes Dev.*, **14**, 1098-1108.
- Leeds, P., Peltz, S.W., Jacobson, A. und Culbertson, M.R. (1991) The product of the yeast UPF1 gene is required for rapid turnover of mRNAs containing a premature translational termination codon. *Genes Dev.*, **5**, 2303-2314.
- Legault, P., Li, J., Mogridge, J., Kay, L.E. und Greenblatt, J. (1998) NMR structure of the bacteriophage lambda N peptide/boxB RNA complex: recognition of a GNRA fold by an arginine-rich motif. *Cell*, **93**, 289-299.
- Levine, B.J., Chodchoy, N., Marzluff, W.F. und Skoulchi, A.I. (1987) Coupling of replication type histone mRNA levels to DNA synthesis requires the stem-loop sequence at the 3' end of the mRNA. *Proc. Natl. Acad. Sci. U. S. A.*, **84**, 6189-6193.
- Long, R.M., Gu, W., Lorimer, E., Singer, R.H. und Chartrand, P. (2000) She2p is a novel RNA-binding protein that recruits the Myo4p-She3p complex to ASH1 mRNA. *EMBO J.*, **19**, 6592-6601.
- Luo, M.J. und Reed, R. (1999) Splicing is required for rapid and efficient mRNA export in metazoans. *Proc. Natl. Acad. Sci. U. S. A.*, **96**, 14937-14942.
- Luo, M.L., Zhou, Z., Magni, K., Christoforides, C., Rappaport, J., Mann, M. und Reed, R. (2001) Pre-mRNA splicing and mRNA export linked by direct interactions between UAP56 and Aly. *Nature*, **413**, 644-647.
- Lykke-Andersen, J. (2001) mRNA quality control: Marking the message for life or death. *Curr. Biol.*, **11**, R88-R91.
- Lykke-Andersen, J., Shu, M.D. und Steitz, J.A. (2000) Human Upf proteins target an mRNA for nonsense-mediated decay when bound downstream of a termination codon. *Cell*, **103**, 1121-1131.
- Lykke-Andersen, J., Shu, M.D. und Steitz, J.A. (2001) Communication of the position of exon-exon junctions to the mRNA surveillance machinery by the protein RNPS1. *Science*, **293**, 1836-1839.

- Makris, M., Preston, F.E., Beauchamp, N.J., Cooper, P.C., Daly, M.E., Hampton, K.K., Bayliss, P., Peake, I.R. und Miller, G.J. (1997) Co-inheritance of the 20210A allele of the prothrombin gene increases the risk of thrombosis in subjects with familial thrombophilia. *Thromb Haemost*, **78**, 1426-9.
- Maquat, L.E. (1995) When cells stop making sense: effects of nonsense codons on RNA metabolism in vertebrate cells. *RNA*, **1**, 453-465.
- Maquat, L.E. (2001) The power of point mutations. *Nature Genet.*, **27**, 5-6.
- Maquat, L.E. und Li, X. (2001) Mammalian heat shock p70 and histone H4 transcripts, which derive from naturally intronless genes, are immune to nonsense-mediated decay. *RNA*, **7**, 445-456.
- Martin, F., Schaller, A., Eglite, S., Schumperli, D. und Muller, B. (1997) The gene for histone RNA hairpin binding protein is located on human chromosome 4 and encodes a novel type of RNA binding protein. *EMBO J.*, **16**, 769-778.
- Martinelli, I. (2001) Risk factors in venous thromboembolism. *Thromb. Haemost.*, **86**, 395-403.
- McGarvey, T., Rosonina, E., McCracken, S., Li, Q., Arnaout, R., Mientjes, E., Nickerson, J.A., Awrey, D., Greenblatt, J., Grosveld, G. und Blencowe, B.J. (2000) The acute myeloid leukemia-associated protein, DEK, forms a splicing-dependent interaction with exon-product complexes. *J. Cell. Biol.*, **150**, 309-320.
- Mendell, J.T. und Dietz, H.C. (2001) When the message goes awry: disease-producing mutations that influence mRNA content and performance. *Cell*, **107**, 411-414.
- Mendell, J.T., Medghalchi, S.M., Lake, R.G., Noensie, E.N. und Dietz, H.C. (2000) Novel Upf2p orthologues suggest a functional link between translation initiation and nonsense surveillance complexes. *Mol. Cell. Biol.*, **20**, 8944-8957.
- Minvielle-Sebastia, L. und Keller, W. (1999) mRNA polyadenylation and its coupling to other RNA processing reactions and to transcription. *Curr Opin Cell Biol*, **11**, 352-7.

- Mitrovich, Q.M. und Anderson, P. (2000) Unproductively spliced ribosomal protein mRNAs are natural targets of mRNA surveillance in *C. elegans*. *Genes Dev.*, **14**, 2173-2184.
- Mogridge, J., Mah, T.F. und Greenblatt, J. (1995) A protein-RNA interaction network facilitates the template-independent cooperative assembly on RNA polymerase of a stable antitermination complex containing the lambda N protein. *Genes Dev.*, **9**, 2831-2845.
- Muhlrad, D. und Parker, R. (1994) Premature translational termination triggers mRNA decapping. *Nature*, **370**, 578-581.
- Muhlrad, D. und Parker, R. (1999) Aberrant mRNAs with extended 3' UTRs are substrates for rapid degradation by mRNA surveillance. *RNA*, **5**, 1299-307.
- Muller, B. und Schumperli, D. (1997) The U7 snRNP and the hairpin binding protein: Key players in histone mRNA metabolism. *Semin. Cell. Dev. Biol.*, **8**, 567-576.
- Neu-Yilik, G., Gehring, N.H., Thermann, R., Frede, U., Hentze, M.W. und Kulozik, A.E. (2001) Splicing and 3' end formation in the definition of nonsense-mediated decay-competent human beta-globin mRNPs. *EMBO J.*, **20**, 532-540.
- Ostareck, D.H., Ostareck-Lederer, A., Wilm, M., Thiele, B.J., Mann, M. und Hentze, M.W. (1997) mRNA silencing in erythroid differentiation: hnRNP K and hnRNP E1 regulate 15-lipoxygenase translation from the 3' end. *Cell*, **89**, 597-606.
- Ostareck-Lederer, A., Ostareck, D.H. und Hentze, M.W. (1998) Cytoplasmic regulatory functions of the KH-domain proteins hnRNPs K and E1/E2. *Trends Biochem. Sci.*, **23**, 409-411.
- Pandey, N.B., Chodchoy, N., Liu, T.J. und Marzluff, W.F. (1990) Introns in histone genes alter the distribution of 3' ends. *Nucleic Acids Res.*, **18**, 3161-3170.
- Perlick, H.A., Medghalchi, S.M., Spencer, F.A., Kendzior, R.J. und Dietz, H.C. (1996) Mammalian orthologues of a yeast regulator of nonsense transcript stability. *Proc. Natl. Acad. Sci. U.S.A.*, **93**, 10928-10932.

- Poort, S.R., Rosendaal, F.R., Reitsma, P.H. und Bertina, R.M. (1996) A common genetic variation in the 3'-untranslated region of the prothrombin gene is associated with elevated plasma prothrombin levels and an increase in venous thrombosis. *Blood*, **88**, 3698-703.
- Preiss, T. und Hentze, M.W. (1998) Dual function of the messenger RNA cap structure in poly(A)-tail-promoted translation in yeast. *Nature*, **392**, 516-520.
- Ross, J. und Pizarro, A. (1983) Human beta and delta globin messenger RNAs turn over at different rates. *J Mol Biol*, **167**, 607-17.
- Ruiz-Echevarria, M.J., Czaplinski, K. und Peltz, S.W. (1996) Making sense of nonsense in yeast. *Trends Biochem. Sci.*, **21**, 433-438.
- Ruiz-Echevarria, M.J., Gonzalez, C.I. und Peltz, S.W. (1998a) Identifying the right stop: determining how the surveillance complex recognizes and degrades an aberrant mRNA. *EMBO J.*, **17**, 575-589.
- Ruiz-Echevarria, M.J., Yasenchak, J.M., Han, X., Dinman, J.D. und Peltz, S.W. (1998b) The upf3 protein is a component of the surveillance complex that monitors both translation and mRNA turnover and affects viral propagation. *Proc. Natl. Acad. Sci. U.S.A.*, **95**, 8721-8726.
- Russel, M., Kidd, S. und Kelley, M.R. (1986) An improved filamentous helper phage for generating single-stranded plasmid DNA. *Gene*, **45**, 333-338.
- Salles, F.J., Richards, W.G. und Strickland, S. (1999) Assaying the polyadenylation state of mRNAs. *Methods*, **17**, 38-45.
- Schell, T., Kulozik, A.E. und Hentze, M.W. (2002) Integration of splicing, transport and translation to achieve mRNA quality control by the nonsense-mediated decay pathway. *Genome Biol.*, **3**, 1006.1-1006.6.
- Schwabe, G.C., Tinschert, S., Buschow, C., Meinecke, P., Wolff, G., Gillessen-Kaesbach, G., Oldridge, M., Wilkie, A.O., Komec, R. und Mundlos, S. (2000) Distinct mutations in the receptor tyrosine kinase gene ROR2 cause brachydactyly type B. *Am. J. Hum. Genet.*, **67**, 822-831.

- Serin, G., Gersappe, A., Black, J.D., Aronoff, R. und Maquat, L.E. (2001) Identification and characterization of human orthologues to *Saccharomyces cerevisiae* Upf2 protein and Upf3 protein (*Caenorhabditis elegans* SMG-4). *Mol. Cell. Biol.*, **21**, 209-223.
- Sheets, M.D., Ogg, S.C. und Wickens, M.P. (1990) Point mutations in AAUAAA and the poly (A) addition site: effects on the accuracy and efficiency of cleavage and polyadenylation in vitro. *Nucleic Acids Res.*, **18**, 5799-5805.
- Shirley, R.L., Lelivelt, M.J., Schenkman, L.R., Dahlseid, J.N. und Culbertson, M.R. (1998) A factor required for nonsense-mediated mRNA decay in yeast is exported from the nucleus to the cytoplasm by a nuclear export signal sequence. *J. Cell Sci.*, **111**, 3129-3143.
- Shyu, A.B. und Wilkinson, M.F. (2000) The double lives of shuttling mRNA binding proteins. *Cell*, **102**, 135-138.
- Siomi, H. und Dreyfuss, G. (1997) RNA-binding proteins as regulators of gene expression. *Curr. Opin. Genet. Dev.*, **7**, 345-353.
- Sun, X., Perlick, H.A., Dietz, H.C. und Maquat, L.E. (1998) A mutated human homologue to yeast Upf1 protein has a dominant-negative effect on the decay of nonsense-containing mRNAs in mammalian cells. *Proc. Natl. Acad. Sci. U.S.A.*, **95**, 10009-10014.
- Sureau, A., Gattoni, R., Dooghe, Y., und Stevenin, J.J.S. (2001) SC35 autoregulates its expression by promoting splicing events that destabilize its mRNAs. *EMBO J.*, **20**, 1785-1796.
- Tarun, S.Z. und Sachs, A.B. (1995) A common function for mRNA 5' and 3' ends in translation initiation in yeast. *Genes Dev.*, **9**, 2997-3007.
- Thein, S.L., Hesketh, C., Taylor, P., Temperley, I.J., Hutchinson, R.M., Old, J.M., Wood, W.G., Clegg, J.B. und Weatherall, D.J. (1990) Molecular basis for dominantly inherited inclusion body beta-thalassemia. *Proc. Natl. Acad. Sci. U. S. A.*, **87**, 3924-3928.
- Thermann, R., Neu-Yilik, G., Deters, A., Frede, U., Wehr, K., Hagemeier, C., Hentze, M.W. und Kulozik, A.E. (1998) Binary specification of nonsense codons by splicing and cytoplasmic translation. *EMBO J.*, **17**, 3484-94.

- Tosetto, A., Missiaglia, E., Frezzato, M. und Rodeghiero, F. (1999) The VITA project: prothrombin G20210A mutation and venous thromboembolism in the general population. *Thromb Haemost*, **82**, 1395-8.
- van Hoof, A. und Parker, R. (1999) The exosome: a proteasome for RNA? *Cell*, **99**, 347-50.
- Venter, J.C., Adams, M.D., Myers, E.W., Li, P.W., Mural, R.J., Sutton, G.G., Smith, H.O., Yandell, M., Evans, C.A., Holt, R.A., Gocayne, J.D., Amanatides, P., Ballew, R.M. und Huson, D.H. (2001) The Sequence of the Human Genome. *Science*, **291**, 1304-1351.
- Vieira, J. und Messing, J. (1987) Production of single-stranded plasmid DNA. *Methods Enzymol.*, **153**, 3-11.
- Wahle, E. und Ruegsegger, U. (1999) 3'-End processing of pre-mRNA in eukaryotes. *FEMS Microbiol Rev*, **23**, 277-95.
- Wang, Z.F., Whitfield, M.L., Ingledue 3rd, T.C., Dominski, Z. und Marzluff, W.F. (1996) The protein that binds the 3' end of histone mRNA: a novel RNA-binding protein required for histone pre-mRNA processing. *Genes Dev.*, **10**, 3028-3040.
- Weng, Y., Czaplinski, K. und Peltz, S.W. (1996a) Genetic and biochemical characterization of mutations in the ATPase and helicase regions of the Upf1 protein. *Mol. Cell. Biol.*, **16**, 5477-5490.
- Weng, Y., Czaplinski, K. und Peltz, S.W. (1996b) Identification and characterization of mutations in the UPF1 gene that affect nonsense suppression and the formation of the Upf protein complex but not mRNA turnover. *Mol. Cell. Biol.*, **16**, 5491-5506.
- Weng, Y., Czaplinski, K. und Peltz, S.W. (1998) ATP is a cofactor of the Upf1 protein that modulates its translation termination and RNA binding activities. *RNA*, **4**, 205-214.
- Whitfield, M.L., Zheng, L.X., Baldwin, A., Ohta, T., Hurt, M.M. und Marzluff, W.F. (2000) Stem-loop binding protein, the protein that binds the 3' end of histone mRNA, is cell cycle regulated by both translational and posttranslational mechanisms. *Mol. Cell. Biol.*, **20**, 4188-4198.

- Wilkinson, M.F. und Shyu, A.B. (2001) Multifunctional regulatory proteins that control gene expression in both the nucleus and the cytoplasm. *Bioessays*, **23**, 775-787.
- Wilusz, C.J., Gao, M., Jones, C.L., Wilusz, J. und Peltz, S.W. (2001a) Poly(A)-binding proteins regulate both mRNA deadenylation and decapping in yeast cytoplasmic extracts. *RNA*, **7**, 1416-1424.
- Wilusz, C.J., Wormington, M. und Peltz, S.W. (2001b) The cap-to-tail guide to mRNA turnover. *Nat. Rev. Mol. Cell. Biol.*, **2**, 237-246.
- Wylenzek, M., Geisen, C., Staphenhorst, L., Wielckens, K. und Klingler, K.R. (2001) A novel point mutation in the 3' region of the prothrombin gene at position 20221 in a Lebanese/Syrian family. *Thromb. Haemost.*, **85**, 943-944.
- Xu, N., Chen, C.Y. und Shyu, A.B. (1997) Modulation of the fate of cytoplasmic mRNA by AU-rich elements: key sequence features controlling mRNA deadenylation and decay. *Mol. Cell. Biol.*, **17**, 4611-4621.
- Yamashita, A., Ohnishi, T., Kashima, I., Taya, Y. und Ohno, S. (2001) Human SMG-1, a novel phosphatidylinositol 3-kinase-related protein kinase, associates with components of the mRNA surveillance complex and is involved in the regulation of nonsense-mediated mRNA decay. *Genes Dev.*, **15**, 2215-2228.
- Zambetti, G., Stein, J. und Stein, G. (1990) Role of messenger RNA subcellular localization in the posttranscriptional regulation of human histone gene expression. *J. Cell. Physiol.*, **144**, 175-182.
- Zhang, J. und Maquat, L.E. (1996) Evidence that the decay of nucleus-associated nonsense mRNA for human triosephosphate isomerase involves nonsense codon recognition after splicing. *RNA*, **2**, 235-243.
- Zhang, J., Sun, X., Qian, Y., LaDuca, J.P. und Maquat, L.E. (1998a) At least one intron is required for the nonsense-mediated decay of triosephosphate isomerase mRNA: a possible link between nuclear splicing and cytoplasmic translation. *Mol. Cell. Biol.*, **18**, 5272-5283.

- Zhang, J., Sun, X., Qian, Y. und Maquat, L.E. (1998b) Intron function in the nonsense-mediated decay of beta-globin mRNA: indications that pre-mRNA splicing in the nucleus can influence mRNA translation in the cytoplasm. *RNA*, **4**, 801-815.
- Zhang, S., Ruiz-Echevarria, M.J., Quan, Y. und Peltz, S.W. (1995) Identification and characterization of a sequence motif involved in nonsense-mediated mRNA decay. *Mol. Cell. Biol.*, **15**, 2231-2244.
- Zhao, J., Hyman, L. und Moore, C. (1999) Formation of mRNA 3' ends in eukaryotes: mechanism, regulation, and interrelationships with other steps in mRNA synthesis. *Microbiol Mol Biol Rev*, **63**, 404-45.
- Zhou, Z., Luo, M.J., Straesser, K., Katahira, J., Hurt, E. und Reed, R. (2000) The protein Aly links pre-messenger-RNA splicing to nuclear export in metazoans. *Nature*, **407**, 401-405.