

6. Literatur

Altschul, S.F., Madden, T.L., Schaffer, A.A., Zhang, J., Zhang Z., Miller, W., and Lipman, D.J. (1997). Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res.* **25:** 3389-3402

Barut B., and Zon, L. (2000). Realizing the potential of zebrafish as a model for human disease. *Physiol Genomics* **2:** 49-51. Review.

Brownlie,A.,Donovan,A.,Pratt,S.J.,Paw,B.H.,Oates,A.C.,Brugnara,C.,Witkowska,H.E.,Sassa, S., and Zon,L. (1998). Positional cloning of the zebrafish sauternes gene: a model for congenital sideroblastic anaemia. *Nature Genetics* **20:** 244-250

Chen, J. and Fishman, M. (1996). Zebrafish tinman homolog demarcates the heart field and initiates myocardial differentiation. *Development* **122:** 3809-3816

Chesi, M.P., Bergsagel, L., Shonukan, O.O., Martelli, M.L., Brents, L.A., Chen, T., Schrock, T., Ried, T., and Kuehl W.M. (1998). Frequent dysregulation of the c-maf proto-oncogene at 16q23 by translocation to an Ig locus in multiple myeloma. *Blood* **91:** 4457-63

Clark, M.D., Hennig, S., Herwig, R., Clifton, S.W., Marra, M.A., Lehrach, H., Johnson, S.L., Group tW, and WU-GSCnEST Group (2001). An oligonucleotide fingerprint normalized and expressed sequence tag characterized zebrafish cDNA library. *Genome Res.* **11:** 1594-602

Classen, M. et al., (1998) Innere Medizin, Urban & Schwarzenberg, München

Cooke, J., Moens, C., Roth, L., Durbin, L., Shiomi, K., Brennan, C., Kimmel, C., Wilson, S., and Holder, N. (2001). Eph signalling functions downstream of Val to regulate cell sorting and boundary formation in the caudal hindbrain. *Development* **128:** 571-580

Development. (1996). **123**

Dick, A., Hild, M., Bauer, H., Imai, Y., Maifeld, H., Schier, A.F., Talbot, W.S., Bouwmeester, T., and Hammerschmidt, M. (2000). Essential role of Bmp7 (*snailhouse*) and its prodomain in dorsoventral patterning of the zebrafish embryo. *Development* **127**: 343–354.

Dickmeis, T., Mourrain, P., Saint-Etienne, L., Fischer, N., Aanstad, P., Clark, M., Strahle, U., and Rosa, F. (2001). A crucial component of the endoderm formation pathway, CASANOVA, is encoded by a novel sox-related gene. *Genes Dev.* **15**: 1487-1492

Driever, W., Solnica-Krezel, L., Schier, A.F., Neuhauss, S.C.F., Malicki, J., Stemple, D.L., Stainier, D.Y.R., Zwartkruis, F., Abdelilah, S., and Rangini, Z. et al. (1996). A genetic screen for mutations affecting embryogenesis in zebrafish. *Development* **123**: 37–46.

Drummond, I.A., Majumdar, A., Hentschel, H., Elger, M., Solnica-Krezel, L., Schier, A.F., Neuhauss, S.C., Stemple, D.L., Zwartkruis, F., Rangini, Z., Driever, W., and Fishman, M.C. (1998). Early development of the zebrafish pronephros and analysis of mutations affecting pronephric function. *Development* **125**: 4655-4667

Eisen, J.S. (1996). Zebrafish make a big splash. *Cell* **87** :969-977

Ekker, S. (2000). Morphants: a new systematic vertebrate functional genomics approach. *Yeast* **17**: 302-306. Review.

Gawantka, V., Pollet, N., Delius, H., Vingron, M., Pfister, R., Nitsch, R., Blumenstock, C., and Niehrs, C. (1998). Gene expression screening in Xenopus identifies molecular pathways, predicts gene function and provides a global view of embryonic patterning. *Mech. Dev.* **77**: 95-141

Gilbert, S.F. (1997). *Developmental Biology*, Sinauer Associates, Sunderland Massachusetts

Gritsman, K., Zhang, J., Cheng, S., Heckscher, E., Talbot, W.S., and Schier, AF. (1999). The EGF–CFC protein one-eyed pin-head is essential for nodal signaling. *Cell* **97**: 121–132.

Haffter, P., Granato, M., Brand, M., Mullins, M.C., Hammerschmidt, M., Kane, D.A., Odenthal, J., van Eeden, F.J.M., Jiang, Y.-J., and Heisenberg, C.-P. et al. (1996). The identification of genes with unique and essential functions in the development of the zebrafish, *Danio rerio*. *Development* **123**: 1–36.

Hammerschmidt, M., Pelegri, F., Mullins, M.C., Kane, D.A., van Eeden, F.J., Granato, M., Brand, M., Furutani-Seiki, M., Haffter, P., Heisenberg, C.P., Jiang, Y.J., Kelsh, R.N., Odenthal, J., Warga, R.M., and Nusslein-Volhard, C. (1996). dino and mercedes, two genes regulating dorsal development in the zebrafish embryo. *Development* **123**: 95–102

Hild, M., Dick, A., Rauch, G.J., Meier, A., Bouwmeester, T., Haffter, P., and Hammer-schmidt, M. (1999). The *smad5* mutation *somitabun* blocks Bmp2b signaling during early dorso-ventral patterning of the zebrafish embryo. *Development* **126**: 2149–2159.

Ingham, P.W. (1997). Zebrafish genetics and its implications for understanding vertebrate development. *Hum. Mol. Genet.* **6**: 1755–1760

Jowett, T., and Lettice, L. (1994). Whole-mount *in situ* hybridizations on zebrafish embryos using a mixture of digoxigenin- and fluorescein-labelled probes. *Trends Genet* **10**: 73–74

Kerckaert, J.P., Deweindt, C., Tilly, H., Quief, S., Lecocq, G., and Bastard, C. (1993). LAZ3, a novel zinc-finger encoding gene, is disrupted by recurring chromosome 3q27 translocations in human lymphomas. *Nature Genetics* **5**: 66–70

Kikuchi, Y., Agathon, A., Alexander, J., Thisse, C., Waldron, S., Yelon, D., Thisse, B., and Stainier, D.Y. (2001). casanova encodes a novel Sox-related protein necessary and sufficient for early endoderm formation in zebrafish. *Genes Dev.* **15**: 1493-1505

Kimmel, C. B., Ballard, W. W., Kimmel, S. R., Ullmann, B. and Schilling, T. (1995). Stages of embryonic development of the zebrafish. *Dev. Dyn.* **203**: 253-310

Kishimoto, Y., Lee, K-H., Zon, L., Hammerschmidt, M., and Schulte-Merker, S. (1997). The molecular nature of zebrafish *swirl*: BMP2 function is essential during early dorsoventral patterning. *Development* **124**: 4457–4466

Majumdar, A., Lun, K., Brand, M., and Drummond, I.A. (2000). Zebrafish no isthmus reveals a role for pax2.1 in tubule differentiation and patterning events in the pronephric primordia. *Development* **127**: 2089-2098

Metscher, B.D., and Ahlberg, P.E. (1999). Zebrafish in context: uses of a laboratory model in comparative studies. *Dev. Biol.* **210**: 1-14

Moens, C.B., Cordes, S.P., Giorgianni, M.W., Barsh, G.S., and Kimmel, C.B. (1998). Equivalence in the genetic control of hindbrain segmentation in fish and mouse. *Development* **125**:381–391.

Mullins, M.C., and Nüsslein-Volhard, C. (1993). Mutational approaches to studying embryonic pattern formation in the zebrafish. *Curr. Opin. Genet. Dev.* **3**: 648-654

Mullins, M.C. and Rosenthal, A. (1999). Development of noradrenergic neurons in the zebrafish hindbrain requires BMP, FGF8, and the homeodomain protein soulless / Phox2a. *Neuron* **24**: 555–566.

Neidhardt, L., Gasca, S., Wertz, K., Obermayr, F., Worpenberg, S., Lehrach, H., and Herrmann, B.G. (2000). Large-scale screen for genes controlling mammalian embryogenesis, using high-throughput gene expression analysis in mouse embryos. *Mech. Dev.* **98**: 77-94

Nguyen, V.H., Schmid, B., Trout, J., Connors, S.A., Ekker, M., and Mullins, M.C. (1998). Ventral and lateral regions of the zebrafish gastrula, including the neural crest progenitors, are established by a *bmp2b / swirl* pathway of genes. *Dev. Biol.* **199**: 93–110.

Nüsslein-Volhard, C., and Wieschaus, E. (1980). Mutations affecting segment number and polarity in Drosophila. *Nature* **287**: 795-801

Orkin, S.H. (1995) Regulation of globin gene expression in erythroid cells. *Eur. J. Biochem.* **231**: 271-281

Postlethwait, J.H. and Talbot, W.S. (1997). Zebrafish genomics: From mutants to genes. *Trends Genet.* **13**: 183–190.

Ransom, D.G., Haffter, P., Odenthal, J., Brownlie, A., Vogelsang, E., Kelsh, R.N., Brand, M., van Eeden, F.J., Furutani-Seiki, M., Granato, M., Hammerschmidt, M., Heisenberg, C.P., Jiang, Y.J., Kane, D.A., Mullins, M.C., and Nusslein-Volhard, C. (1996). Characterization of zebrafish mutants with defects in embryonic hematopoiesis. *Development* **123**: 311-319

Reifers, F., Bohli, H., Walsh, E.C., Crossley, P.H., Stainier, D.Y., and Brand, M. (1998). Fgf8 is mutated in zebrafish *acerebellar* (*ace*) mutants and is required for maintenance of midbrain-hindbrain boundary development and somitogenesis. *Development* **125**: 2381–2395.

Reiter, J.F., Alexander, J., Rodaway, A., Yelon, D., Patient, R., Holder, N., and Stainier, D.Y. (1999). Gata5 is required for the development of the heart and endoderm in zebrafish. *Genes and Dev.* **13**: 2983–2995.

Sambrook, J., Fritsch, E.F., and Maniatis, T. (1989). Molecular Cloning, A Laboratory Manual: Cold Spring Harbor Laboratory Press

Schier, A.F., and Talbot, W.S. (2001). Nodal signaling and the zebrafish organizer. *Int. J. Dev. Biol.* **45**: 289-297

Schmid, B., Furthauer, M., Connors, S.A., Trout, J., Thisse, C., and Mullins, M.C. (2000). Equivalent genetic roles for *bmp7 / snailhouse* and *bmp2b / swirl* in dorsoventral pattern formation. *Development* **127**: 957–967.

Schulte-Merker, S., van Eeden, F.J., Halpern, M.E., Kimmel, C.B., and Nüsslein-Volhard, C. (1994). *no tail (ntl)* is the zebrafish homologue of the mouse *T (Brachyury)* gene. *Development* **120**: 1009–1015.

Schulte-Merker, S., Lee, K.J., McMahon, A.P., and Hammerschmidt, M. (1997). The zebrafish organizer requires *chordino*. *Nature* **387**: 862–863.

Shepherd, P.R., and Kahn, B.B. (1997). Glucose transporters and insulin action-implications for insulin resistance and diabetes mellitus. *N. Engl. J. Med.* **337**: 248-257.

Solnica-Krezel, L., Schier, A.F., and Driever, W. (1994). Efficient recovery of ENU-induced mutations from the zebrafish germline. *Genetics* **136**: 1401-1420

Stainier, D., Weinstein, B., Detrich, H., Zon, L., and Fishman, M. (1995). *cloche*, an early acting zebrafish gene, is required by both the endothelial and haematopoietic lineages. *Development* **121**: 3141-3150

Streisinger, G., Walker, C., Dower, N., Knauber, D., and Singer, F. (1981). Production of clones of homozygous diploid zebrafish (*Brachydanio rerio*). *Nature* **291**: 293–296

Torra, R., Badenas, C., Darnell, A., Bru, C., Escorsell, A., and Estivill, X. (1997). Autosomal dominant polycystic kidney disease with anticipation and Caroli's disease associated with a PKD1 mutation. *Kidney. Int.* **52**: 33-38

Vogel, G. (2000). Zebrafish earns its stripes in genetic screens. *Science* **288**: 1160-1161

Wang, H., Long, Q., Marty, S.D., Sassa, S., and Lin, S. (1998). A zebrafish model for hepatoerythropoietic porphyria. *Nat. Genet.* **20**: 239–243.

Weinstein, B.M., Stemple, D.L., Driever, W., and Fishman, M.C. (1995). *Gridlock*, a localized heritable vascular patterning defect in the zebrafish. *Nat. Med.* **1**: 1143–1147.

Westerfield, M. (1993). The Zebrafish Book. University of Oregon Press, Eugene, Oregon

Yelon, D., Ticho, B., Halpern, M.E., Ruvinsky, I., Ho, R.K., Silver, L.M., and Stainier, D.Y. (2000). The bHLH transcription factor hand2 plays parallel roles in zebrafish heart and pectoral fin development. *Development* **127**: 2573-2582

Zhang, J., Talbot, W.S., and Schier, A.F. (1998). Positional cloning identifies zebrafish *one-eyed pinhead* as a permissive EGF-related ligand required during gastrulation. *Cell* **92**: 241–251.

Zhao, Z., Cao, Y., Li, M., Meng, A. (2001). Double-stranded RNA injection produces nonspecific defects in zebrafish. *Dev. Biol.* **229**: 215-223

Zhong, T.P., Rosenberg, M., Mohideen, M.A.P.K., Weinstein, B., and Fishman, M.C. (2000). *Gridlock*, an HLH gene required for assembly for the aorta in zebrafish. *Science* **287**: 1820–1824.

Zon, L. (1999). Zebrafish: a new model for human disease. *Genome Res.* **9**: 99–100