

6 Literaturverzeichnis

- Aberle, H., Bauer, A., Stappert, J., Kispert, A., Kemler, R. (1997)**
-Catenin is targeted for the ubiquitin-proteasome pathway. *EMBO J.* **16**: 3797-3804
- Aghib, D.F. und McCrea, P.D. (1995)**
The E-cadherin complex contains the src substrate p120. *Exp Cell Res.* **218**: 359-369
- Angres, B., Müller, A.H.J., Kellermann, J., Hausen, P. (1991)**
Differential expression of two cadherins in *Xenopus laevis*. *Development* **111**: 829-844
- Aota, S., Nagai, T., Yamada, K.M. (1991)**
Characterization of regions of fibronectin beside the Arginine-Glycin-Aspartic acid sequence required for adhesive function of the cell binding domain using site-directed mutagenesis. *J. Biol. Chem.* **266**: 15938-15943
- Arendt, D., Nübler-Jung, K. (1997)**
Dorsal or ventral: similarities in fate maps and gastrulation patterns in annelids, arthropods and chordates. *Mech. Dev.* **61**: 7-21
- Artinger, M., Blitz, I., Inoue, K., Tran, U., Cho, K.W.Y. (1997)**
Interaction of goosecoid and brachyury in *Xenopus* mesoderm patterning. *Mech. Dev.* **65**: 187-196
- Bahnot, P., Brink, M., Samos, C.H., Hsieh, J.-C., Wang, Y., Macke, J.P., Andrew, D., Nathans, J., Nusse, R. (1996)**
A new member of the frizzled family from *Drosophila* functions as a Wingless receptor. *Nature* **382**: 225-230
- Axelrod, J.D., Matsuno, K., Artavanis-Tsakonas, S., Perrimon, N. (1996)**
Interaction between wingless and notch signaling pathways mediated by dishevelled. *Science* **271**: 1826-1832
- Basler, K. und Hafen, E. (1991)** Specification of cell fate in the developing eye of *Drosophila*. *Bioessays* **13**: 621-631
- Behrens, J., Birchmeier, W., Goodman, S.L., Imhof, B.A. (1985)**
Dissociation of Madin-Darby canine kidney epithelial cells by the monoclonal antibody anti-Arc-1: mechanistic aspects and identification of the antigen as a component related to uvomorulin. *J. Cell Biol.* **101**: 1307-1315
- Behrens, J., Mareel, M.M., van Roy, F., Birchmeier, W. (1989)**
Dissecting tumor cell invasion: epithelial cells acquire invasive properties after the loss of uvomorulin-mediated cell-cell adhesion. *J. Cell Biol.* **108**: 2435-2447
- Behrens, J., Frixen, U., Schipper, J., Weidner, M., Birchmeier, W. (1992)**
Cell adhesion in invasion and metastasis. *Sem. Cell Biol.* **3**: 169-178

- Behrens, J., von Kries, J.P., Kühl, M., Bruhn, L., Wedlich, D., Grosschedl, R., Birchmeier, W.** (1996) Functional interaction of -catenin with the transcription factor LEF-1. *Nature* **382**: 638-642
- Berndorff, D., Gessner, R., Kreft, B., Schnoy, N., Lajous-Petter, A.-M., Loch, N., Reutter, W., Hortsch, M., Tauber, R.** (1994) Liver-intestine cadherin: cloning and characterization of a novel Ca^{2+} -dependent cell adhesion molecule expressed in liver and intestine. *J. Cell Biol.* **125**: 1353-1369
- Bisbee, C., Baker, M., Wilson, A., Hadji-Azimi, I., Fischberg, M.** (1977) Albumin phylogenie for clawed frogs (*Xenopus*). *Science* **195**: 785-787
- Blaschuk, O.W., Sullivan, R., David, S., Pouliot, Y.** (1990) Identification of a cadherin cell adhesion recognition sequence. *Dev. Biol.* **139**: 227-229
- Boller, K., Vestweber, D., Kemler, R.** (1985) Cell adhesion molecule uvomorulin is located in the intermediate junctions of adult intestinal epithelial cells. *J. Cell Biol.* **100**: 327-332
- Boucaut, J.C. und Darribère, T.** (1983) Presence of fibronectin during early embryogenesis in the amphibian *Pleurodeles waltlii*. *Cell Differ.* **12**: 77-83
- Boucaut, J.C., Darribère, T., Poole, T.J., Aoyama, H., Yamada, K.M., Thiery, J.P.** (1984) Biologically active synthetic peptides as probes for embryonic development: A competitive peptide inhibitor of fibronectin function inhibits gastrulation in amphibian embryos and neural crest cell migration in avian embryos. *J. Cell Biol.* **99**: 1822-1830
- Brannon, M., Gomperts, M., Sumoy, L., Moon, R.T., Kimelman, D.** (1997) A beta-catenin/Xtcf-3 complex bid to the siamois promoter to regulate dorsal axis specification in *Xenopus*. *Genes Dev.* **11**: 2359-2370
- Broders, F., Girault, J.M., Simmeneau, L., Suzuki, S., Thiery, J.P.** (1993) Sequence and distribution of *Xenopus laevis* E-cadherin transcripts. *Cell Adh. Comm.* **1**: 265-277
- Broders, F. und Thiery, J.P.** (1995) Contribution of cadherins to directional cell migration and histogenesis in *Xenopus* embryos. *Cell Adh. Comm.* **3**: 419-440
- Burridge, K., Fath, K., Kelly, T., Nuckolls, G., Turner, C.** (1988) Focal adhesions: transmembrane junctions between the extracellular matrix and the cytoskeleton. *Annu. Rev. Cell Biol.* **4**: 487-525
- Butz, S. und Kemler, R.** (1994) Distinct cadherin-catenin complexes in Ca^{2+} -dependent cell-cell adhesion. *FEBS Lett.* **355**: 195-200
- Buxton, R.S., Magee, A.I.** (1992) Structure and interactions of desmosomal and other cadherins. *Sem. Cell Biol.* **3**: 157-167

- Carnac, G., Kodjabachian, L., Gurdon, J.B., Lemaire, P.** (1996)
The homeobox gene *siamois* is a target of the wnt dorsalisation pathway and triggers organiser activity in the absence of mesoderm. *Development* **122**: 3055-3065
- Cepek, K.L., Shaw, S.K., Parker, C.M., Russel, G.J., Morrow, J.S., Rimm, D.L., Brenner, M.B.** (1994)
Adhesion between epithelial cells and T-lymphocytes mediated by E-cadherin and integrin. *Nature* **372**: 190-193
- Chen, P., Gupta, K., Wells, A.** (1994)
Cell movement elicited by epidermal growth factor receptor requires kinase and autoprophosphorylation but is separable from mitogenesis. *J. Cell Biol.* **124**: 547-555
- Choi, Y.S., Sehgal, R., McCrea, P., Gumbiner, B.** (1990)
A cadherin-like protein in eggs and cleaving embryos of *Xenopus laevis* is expressed in oocytes in response to progesterone. *J. Cell Biol.* **110**: 1575-1582
- Choi, Y.S. und Gumbiner, B.** (1989)
Expression of cell adhesion molecule E-cadherin in *Xeopus* begins at gastrulation and predominates in the ectoderm. *J. Cell Biol.* **108**: 2449-2458
- Clark, E:A., Brugge, J.S.** (1995)
Integrins and signal transduction pathways: the road taken. *Science* **268**: 233-239
- Cole, G.J., Loewy, A., Cross, N.V., Akeson, R., Glaser, L.** (1986)
Topographic localization of the heparin-binding domain of the neural cell adhesion molecule N-CAM. *J. Cell Biol.* **103**, 1739-1744
- Cook, D., Fry, M.J., Hughes, K., Sumathipala, R., Woodgett, J.R., Dale, T.C.** (1996) Wingless inactivates glycogen synthase kinase-3 via an intracellular signalling pathway which involves a protein kinase C. *EMBO J.* **15**: 45326-45336
- Cui, Y., Brown, J.D., Moon, R.T., Christian, J.L.** (1995)
Xwnt-8b: A maternally expressed *Xenopus* wnt gene with a potential role in establishing the dorsoventral axis. *Development* **121**: 2177-2186
- Cunningham, B.A., Hemperly, J.J., Murray, B.A., Prediger, E.A., Brackenbury, R., Edelman, G.M.** (1987)
Neural cell adhesion molecule: structure, immunoglobulin-like domains, cell surface modulation, and alternative RNA splicing. *Science* **236**: 799-806
- Damsky, C.H., Richa, J., Solter, D., Knudsen, K., Buck, C.A.** (1983)
Identification and purification of a cell surface glycoprotein mediating intercellular adhesion in embryonic and adult tissues. *Cell* **34**: 455-466
- Danen, E.H., Aota, S., van Kraats, A.A., Yamada, K.M., Ruiter, D.J., van Muijen, G.N.P.** (1995) Requirement for the synergy site for cell adhesion to fibronectin depends on the activation of integrin β_1 . *J. Biol. Chem.* **270**: 21612-21618

- Daniel, J.M. und Reynolds, A.B. (1997)**
Tyrosine phosphorylation and cadherin/catenin function.
BioEssays 19: 883-891
- Darribère, T., Guida, K., Larjava, H., Johnson, K.E., Yamada, K.M., Thiery, J.-P., Boucaut, J.-C. (1990)** In vivo analysis of γ_1 subunit function in fibronectin matrix assembly. *J. Cell Biol.* 110: 1813-1823
- DeMarais, A.A. und Moon, R.T. (1992)** The armadillo homologs β -catenin and plakoglobin are differentially expressed during early development of *Xenopus laevis*. *Dev. Biol.* 153: 337-346
- DeRobertis, E.M., Sasai, Y. (1996)**
A common plan for dorsoventral patterning in Bilateria. *Nature* 380: 37-40
- DeSimone, D.W. (1997)**
Regulation of Integrin Adhesive Function in Early Development.
3rd. International Colloquium on Cellular Signal Recognition and Transduction, Abstract: 51
- Detrick, R.J., Dickey, D., Kintner, C.R. (1990)**
The effects of N-cadherin misexpression on morphogenesis in *Xenopus* embryos. *Neuron* 4: 493-506
- Dinshaw, J.P. und Gumbiner, B.M. (1995)**
Zipping together a cell adhesion interface. *Nature* 374: 306-307
- Donalies, M., Cramer, M., Ringwald, M., Starzinski-Powitz, A. (1991)**
Expression of a novel member of the cadherin multigene family correlates with differentiation of skeletal muscle cells. *Proc. Natl. Acad. Sci. USA* 88: 8024-8028
- Dosch, R., Gawantka, V., Delius, H., Blumenstock, C., Niehrs, C. (1997)**
BMP-4 acts as a morphogen in dorsoventral mesoderm patterning in *Xenopus*. *Development* 124: 2325-2334
- Du, S.J., Purcell, S., Christian, J.L., McGrew, L.L., Moon, R.T. (1995)**
Identification of distinct classes and functional domains of Wnts through expression of wild-type and chimeric proteins in *Xenopus* embryos. *Mol. Cell Biol.* 15: 2625-2634
- Duband, J., Dufour, S., Hatta, K., Takeichi, M., Edelman, G.M., Thiery, J.P. (1987)** Adhesion molecules during somitogenesis in the avian embryo. *J. Cell Biol.* 127: 521-535
- Dufuor, S., Saint-Jeannet, J.P., Broders, F., Wedlich, D., Thiery, J.P. (1994)** Differential perturbations in the morphogenesis of anterior structures induced by overexpression of truncated XB- and N-cadherins in *Xenopus* embryos. *J. Cell Biol.* 127: 521-535
- Edelman, G.M., Thiery, J.P. (1985)**
The cell in contact: adhesions and junctions as morphogenetic determinants.
John Wiley & Sons, New York

- Edelman, G.M. (1988)**
Morphoregulatory molecules. *Biochemistry* **27**: 3533-3543
- Espeseth, A., Johnson, E., Kintner, C. (1992)**
Xenopus F-cadherin, a novel member of the cadherin family of cell adhesion molecules, is expressed at boundaries in the neural tube. *Mol. Cell Neurosci.* **6**: 199-211
- Fagotto, F., Funayama, N., Glück, U., Gumbiner, B.M. (1996)**
Binding to cadherins antagonizes the signaling activity of -catenin during axis formation. *J. Cell Biol.* **132**: 1105-1114
- Fagotto, F., Guger, K., Gumbiner, B.M. (1997)**
Induction of the primary dorsalizing center in Xenopus by the Wnt/GSK/ -catenin signaling pathway, but not by Vg1, activin or noggin. *Development* **124**: 453-460
- Fainsod, A., Deißler, K., Yelin, R., Marom, K., Epstein, M., Pillemer, G., Steinbeisser, H., Blum, M. (1997)**
The dorsalizing and neural inducing gene follistatin is an antagonist of BMP-4. *Mech. Dev.* **63**: 39-50
- Faull, R.J., Kovach, N.L., Harlan, J.M., Ginsberg, M.H. (1993)**
Affinity modulation of integrin $\alpha_5\beta_1$: regulation of the functional response by soluble fibronectin. *J. Cell Biol.* **121**: 155-162
- Finnemann, S., Kühl, M., Otto, G., Wedlich, D. (1995)**
Cadherin transfection of Xenopus XTC cells downregulates expression of substrate adhesion molecules. *Mol. Cell Biol.* **15**: 5082-5091
- Finnemann, S., Mitrik, I., Hess, M., Otto, G., Wedlich, D. (1997)**
Uncoupling of XB/U-cadherin-catenin complex formation from its function in cell-cell adhesion. *Journal Biol. Chem.* **272**: 11856-11862
- Friedlander, D.R., Mege, R.-M., Cunningham, B.A., Edelmann, G.M. (1989)**
Cell sorting-out is modulated by both the specificity and amount of different cell adhesion molecules (CAMs) expressed on cell surfaces. *Proc. Natl. Acad. Sci. USA* **86**: 7043-7047
- Friedle, H., Rastegar, S., Paul, H., Kaufmann, E., Knöchel, W. (1998)**
Xvent-1 mediates BMP-4-induced suppression of the dorsal-lip-specific early response gene XFD-1' in Xenopus embryos. *EMBO J.* **17**: 2298-2307
- Funayama, N., Fagotto, F., McCrea, P., Gumbiner, B. (1995)** Embryonic axis induction by the armadillo repeat domain of -catenin: Evidence for intracellular signalling. *J. Cell Biol.* **128**: 959-968
- Gallin, W.J., Sorkin, B.C., Edelman, G.M., Cunningham, B.A. (1987)**
Sequence analysis of a cDNA clone encoding the liver cell adhesion molecule L-CAM. *Proc. Natl. Acad. Sci. USA* **84**: 2808-2812
- Geiger, B., Volberg, T., Sabany, Y., Volk, T. (1990)**
In Edelman, G.M., Cunningham, B.A., Thiery, J.P.: Morhoregulatory molecules, John Wiley & Sons, New York: 57-79

- Geiger, B. und Ayalon, O. (1992)**
Cadherins. Annu. Rev. Cell Biol. **8**: 307-332
- Gentry, L. und Rohrschneider, L. (1984)**
Common features of the yes and src gene products defined by peptide-specific antibodies.
Cell Diff. Dev. **31**: 647-75
- Gerhardt, J., Danilchik, M., Doniach, T., Roberts, S., Rowning, B., Stewart, R. (1989)** Cortical rotation of the Xenopus egg: consequences for the anteroposterior pattern of embryonic development. *Development Supplement* **107**: 37-52
- Gilbert, S. F. (1988)**
Developmental Biology, *Sinauer Associates, Inc.*, 2nd ed., pp. 247-249
- Ginsberg, D., DeSimone, D., Geiger, B. (1991)**
Expression of a novel cadherin (EP-cadherin) in unfertilized eggs and early embryos.
Development **111**: 315-325
- Giudice, G. (1962)**
Restitution of whole larvae from disaggregated cells of sea urchin embryos.
Dev. Biol. **5**: 402-411
- Gomperts, M., Garcia-Castro, M., Wylie, C., Heasman, J. (1994)**
Interactions between primordial germ cells play a role in their migration in mouse embryos.
Development **120**: 135-141
- Gradl, D., Kühl, M., Wedlich, D. (1997)** Catenin regulates the crosstalk between cell-cell and cell-matrix adhesion in XTC-cells. *3rd. International Colloquium on Cellular Signal Recognition and Transduction, Abstract*: 42
- Graff, J.M. (1997)**
Embryonic patterning: to BMP or not to BMP, that is the question. *Cell* **89**: 171-174
- Gumbiner, B., Simons, K. (1986)**
A functional assay for proteins involved in establishing an epithelial occluding barrier: identification of a uvomorulin-like polypeptide. *J. Cell Biol.* **102**: 457-468
- Gurdon, J.B. (1987)**
Embryonic induction-molecular prospects. *Development* **99**: 285-306
- Gurdon, J.B., Lemaire, P., Kato, K. (1993)**
Community Effects and Related Phenomena in Development. *Cell* **75**: 831-834
- Hamaguchi, M., Matsuyoshi, M., Onishi, Y., Gotoh, B., Takeichi, M., Nagai, Y. (1993)** p60v-src causes tyrosine phosphorylation and inactivation of the N-cadherin-catenin cell adhesion system. *EMBO J.* **12**: 307-314
- Hatta, K., Nose, A., Nagafuchi, A., Takeichi, M. (1988)**
Cloning and expression of cDNA encoding a neural calcium-dependent cell adhesion molecule: its identity in the cadherin gene family. *J. Cell Biol.* **106**: 873-881

- He, X., Saint-Jeannet, J.P., Woodgett, J.R., Varmus, H.E., Dawid, I.B. (1995)
Glycogen synthase kinase-3 and dorsoventral patterning in Xenopus embryos.
Nature **374**: 617-622
- Heasman, J., Crawford, A., Goldstone, K., Garder-Hamrick, P., Gumbiner, B.,
McCrea, P., Kintner, C., Noro, C.Y., Wylie, C. (1994)
Overexpression of cadherin and underexpression of -catenin inhibit dorsal mesoderm
induction in early Xenopus embryos. *Cell* **79**: 791-803
- Heasman, J., Ginsberg, D., Geiger, B., Goldstone, K., Pratt, T., Yoshida-Noro,
C., Wylie, C. (1994a) A functional test for maternally inherited cadherin in Xenopus
shows its importance in cell adhesion at the blastula stage. *Development* **120**: 49-57
- Herrenknecht, K., Ozawa, M., Eckerskorn, C., Lottspeich, F., Lenter, M.,
Kemler, R. (1991) The uvomorulin-anchorage protein catenin is a vinculin homologue.
Proc. Natl. Acad. Sci. USA **88**: 9156-9160
- Herzberg, F., Wildermuth, V., Wedlich, D. (1991)
Expression of Xbcd, a novel cadherin, during oogenesis and early development of Xenopus.
Mech. Dev. **35**: 33-42
- Hinck, L., Nähke, I., Papkoff, J., Nelson, W. (1994)
Dynamics of cadherin/catenin complex formation: Novel protein interaction and pathways of
complex assembly. *J. Cell Biol.* **125**: 1327-1340
- Hirano, S., Nose, A., Hatta, K., Kawakami, A., Takeichi, M. (1987)
Calcium-dependent cell cell adhesion molecules (cadherins): subclass specificities and
possible involvements of actin bundles. *J. Cell Biol.* **105**: 2501-2510
- Hirano, S., Kimoto, N., Shimoyama, Y., Hirohashi, S., Takeichi, M. (1992)
Identification of a neural -catenin as a key regulator of cadherin function and multicellular
organization. *Cell* **70**: 293-301
- Hodivala, K.J. und Watt, F.M. (1994)
Evidence That Cadherins Play a Role in the Downregulation of Integrin Expression That
Occurs during Keratinocyte Terminal Differentiation. *J. Cell Biol.* **124**: 589-600
- Holt, C.E., Lemaire, P., Gurdon, J.B. (1994)
Cadherin-mediated cell interactions are necessary for the activation of MyoD in Xenopus
mesoderm. *Proc. Natl. Acad. Sci. USA* **91**: 10844-10848
- Hopwood, W.D., Pluck, A., Gurdon, J.B. (1989)
MyoD expression in the forming somites is an early response to mesoderm induction in
Xenopus embryos. *EMBO J.* **8**: 3409-3417
- Hoschützky, H., Aberle, H., Kemler, R. (1994)
-Catenin mediates the interaction of the cadherin-catenin complex with epidermal growth
factor receptor. *J. Cell Biol.* **127**: 1375-1380

- Howard, J.E., Hirst, E.M.A., Smith, J.C. (1992)**
Are α_1 integrins involved in Xenopus gastrulation? *Mech. Dev.* **38**: 109-120
- Huttenlocher, A., Sandborg, R.R., Horwitz, A.F. (1995)**
Adhesion in cell migration. *Curr. Opin. Cell Biol.* **7**: 697-706
- Hyafil, F., Babinet, C., Jacob, F. (1981)**
Cell-cell interactions in early embryogenesis: a molecular approach to the role of calcium. *Cell* **26**: 447-454
- Hynes, R.O. (1992)**
Integrins: Versatility, modulation and signalling in cell adhesion. *Cell* **69**: 11-25
- Inuzuka, H., Miyatani, S., Takeichi, M. (1991)**
R-Cadherin: a novel Ca^{2+} -dependent cell-cell adhesion molecule expressed in the retina. *Neuron* **7**: 69-79
- Isaacs, H.V., Pownall, M.E., Slack, J.M.W. (1994)**
eFGF regulates Xbra expression during Xenopus gastrulation. *EMBO J.* **13**: 4469-4481
- Iwamoto, T., Taniguchi, M., Asai, N., Ohkusu, K., Nakashima, I., Takahashi, M. (1993)** cDNA cloning of mouse ret proto-oncogene and its sequence similarity to the cadherin superfamily. *Oncogene* **8**: 1087-1091
- Jaffe, S.H., Friedlander, D.R., Matsuzaki, F., Crossin, K.L., Cunningham, B.A., Edelman, G.M. (1990)** Differential effects of the cytoplasmic domains of cell adhesion molecules on cell aggregation and sorting out. *Proc. Natl. Acad. Sci. USA* **87**: 3589-3593
- Joos, T.O., Whittaker, C.A., Meng, F., DeSimone, D.W., Gnau, V., Hausen, P. (1995)** Integrin α_5 during early development of Xenopus laevis. *Mech. Dev.* **50**: 187-199
- Karecla, P.I., Bowden, S.J., Green, S.J., Kilshaw, P.J. (1995)**
Recognition of E-cadherin on epithelial cells by the mucosal T cell integrin $\alpha_M\beta_2$ ($\alpha_E\beta_7$). *Eur. J. Immunol.* **25**: 852-856
- Karnovsky, A. und Klymkowsky, M.W. (1995)**
Anterior axis duplication in Xenopus induced by the over-expression of the cadherin-binding protein plakoglobin. *Proc. Natl. Acad. Sci. USA* **92**: 4522-4526
- Keller, R. (1986)**
The cellular basis of amphibian gastrulation. *Developmental Biology*, vol. 2 (ed. L.W. Browder), Plenum Press New York: 241-323
- Keller, R.E., Danilchik, M., Gimlich, R., Shih, J. (1985)**
The function of convergent extension during gastrulation of Xenopus laevis. *J. Embryol. Exp. Morph.* **89** Supplement: 185-209
- Keller, R. und Jansa, S. (1992)**
Xenopus Gastrulation Without a Blastocoel Roof. *Dev. Dyn.* **195**: 162-176

- Kemler, R., Ozawa, M., Ringwald, M. (1989)**
Calcium-dependent cell adhesion molecules. *Curr. Op. Cell Biol.* **1**: 892-897
- Kimura, Y., Matsunami, H., Inoue, T., Shimamura, K., Uchida, N., Ueno, T., Miyazaki, T., Takeichi, M. (1995)** Cadherin-11 expressed in association with mesenchymal morphogenesis in head, somite, and limb bud of early mouse embryos. *Dev. Biol.* **169**: 347-358
- Kinch, M.S., Clark, G.J. Der, Burridge, K. (1995)**
Tyrosine-phosphorylation regulates the adhesions of Ras-transformed breast epithelia. *J. Cell Biol.* **130**: 461-471
- Kintner, C. (1992)**
Regulation of embryonic cell adhesion by the cadherin cytoplasmic domain. *Cell* **69**: 225-236
- Kirschner, M., Newport, J., Gerhart, J. (1985)**
The timing of early developmental events in Xenopus. *Trends Genet.* **1**, 41-47
- Klingensmith, J., Nusse, R., Perrimon, N. (1994)**
The Drosophila segment polarity gene *dishevelled* encodes a novel protein required for response to the wingless signal. *Genes & Dev.* **8**: 118-130
- Knudsen, K.A. und Wheelock, M.J. (1992)** Plakoglobin, or a 83-kD homologue distinct from -catenin, interacts with E-cadherin and N-cadherin. *J. Cell Biol.* **118**: 671-679
- Koch, P.J., Goldschmidt, M.D., Zimbelmann, R., Troyanovsky, R., Franke, W.W. (1992)** Complexity and expression patterns of the desmosomal cadherins. *Proc. Natl. Acad. Sci. USA* **89**: 353-357
- Kofron, M., Spagnuolo, A., Klymkowsky, M., Wylie, C., Heasman, J. (1997)**
The roles of maternal -catenin and plakoglobin in the early Xenopus embryo. *Development* **124**: 1553-1560
- Koster, J.G., Eizema, K., Peterson-Maduro, L.J., Stegman, B.I., Destree, O.H. (1996)** Analysis of Wnt/Engrailed Signaling in Xenopus Embryos Using Biolistics. *Dev. Biol.* **173**: 348-352
- Kühl, M., Finnemann., S., Binder, O., Wedlich, D. (1996)**
Dominant negative expression of a cytoplasmically deleted mutant of XB/U-cadherin disturbs mesoderm migration during gastrulation in *Xenopus laevis*. *Mech. Dev.* **54**: 71-82
- Kühl, M. und Wedlich, D. (1995)**
XB/U-Cadherin mRNA contains cytoplasmic polyadenylation elements and is polyadenylated during oocyte maturation in *Xenopus laevis*. *Biochim. Biophys. Acta* **1262**:95-98
- Kühl, M. und Wedlich, D. (1996)**
Xenopus cadherins: sorting out types and functions in embryogenesis. *Dev. Dyn.* **207**: 121-134

Ku, M. und Melton, D.A. (1993)
Xwnt-11: A maternally expressed Xenopus wnt gene. *Development* 119: 1161-1173

Lampugnani, M.G., Corada, M., Caveda, L., Breviario, F., Ayalon, O., Geiger, B., Dejana, E. (1995) The Molecular Organisation of Endothelial Cell to Cell Junctions: Differential Association of Plakoglobin, -catenin, and -catenin with Vascular Endothelial Cadherin (VE-cadherin). *J. Cell Biol.* 129: 203-217

Larabell, C.A., Torres, M., Rowning, B.A., Yost, C., Miller, J.R., Wu, M., Kimelman, D., Moon, R.T. (1997)
Establishment of the dorso-ventral axis in Xenopus embryos presaged by early asymmetries in -catenin that are modulated by the Wnt signaling pathway.
Development 122: 1281-12849

Larue, L., Antos, C., Butz, S., Huber, O., Delmas, V., Dominis, M., Kemler, R. (1996) A role for cadherins in tissue formation. *Development* 122: 3185-3194

Laurent, M.N., Blitz, I.L., Hashimoto, C., Rothbacher, U., Cho, K.W. (1997)
The Xenopus homeobox gene twin mediates Wnt induction of goosecoid in establishment of Spemann's organizer. *Development* 124: 4905-4916

Lee, G., Hynes, R.O., Kirshner, M. (1984) Temporal and spatial regulation of fibronectin in early Xenopus development. *Cell* 36:729-740

Lee, G. und Gumbiner, B. (1995)
Disruption of gastrulation movements in Xenopus by a dominant negative mutant of C-cadherin. *Dev. Biol.* 171: 363-373

Levine, E., Gumbiner, B., Thiery, J.P. (1991)
The distribution of E-cadherin during Xenopus laevis development.
Development 111: 159-169

Levine, E., Hyun Lee, C., Kintner, C., Gumbiner, B.M. (1994)
Selective disruption of E-cadherin function in early Xenopus embryos by a dominant negative mutant. *Development* 120: 901-909

Lewis, J.E., Wahl III, J.K., Sass, K.M., Jensen, P.J., Johnson, K.R., Wheelock, M.J. (1997) Cross-Talk between Adherens Junctions and Desmosomes Depends on Plakoglobin. *J. Cell Biol.* 136: 919-934

Lewis, J.M., Schwartz, M.A. (1995)
Mapping in vivo associations of cytoplasmic proteins with integrin α_1 cytoplasmic domain mutants. *J. Cell Sci.* 6: 151-160

Liaw, C.W., Cannon, C., Power, M.D., Kiboneka, P.K., Rubin, L.L. (1990)
Identification and cloning of two species of cadherins in bovine endothelial cells.
EMBO J. 9: 2701-2708

Mangold, O. (1933)
Über die Induktionsfähigkeit verschiedener Bezirke der Neurula von Urodelen.
Naturwissenschaften 21: 761-766

- Mansouri, A., Spurr, N., Goodfellow, P.B., Kemler, R. (1988)**
Characterization and chromosomal localization of the gene encoding the human cell adhesion molecule uvomorulin. *Differentiation* **38**: 67-71
- Matsumoto, K., Matsumoto, K., Nakamura, T., Kramer, R.H. (1994)**
Hepatocyte growth factor/scatter factor induces tyrosine phosphorylation of focal adhesion kinase (p125FAK) and promotes migration and invasion by oral squamous cell carcinoma cells. *J. Biol. Chem.* **269**: 31807-31813
- Matsunami, H. (1993)**
Cell binding specificity of mouse R-cadherin and chromosomal mapping of the gene. *J. Cell Sci.* **106**: 401-409
- McCrea, P. und Gumbiner, B. (1991)**
Purification of a 92-kD cytoplasmic protein tightly associated with the cell-cell adhesion molecule E-cadherin (uvomorulin). Characterization and extractability of the protein complex from the cell cytostructure. *J. Biol. Chem.* **266**: 4514-4520
- McCrea, P., Brieher, W., Gumbiner, B. (1993)** Induction of a secondary axis in Xenopus by antibodies to -Catenin. *J. Cell Biol.* **123**: 477-484
- McNeill, H., Ozawa, M., Kemler, R., Nelson, W.J. (1990)**
Novel function of the cell adhesion molecule uvomorulin as an inducer of cell surface polarity. *Cell* **62**: 309-316
- Merriam, J.M., Rubenstein, A.B., Klymkowsky, M.W. (1997)**
Cytoplasmically Anchored Plakoglobin Induces a WNT-like Phenotype in Xenopus. *Dev. Biol.* **185**: 67-81
- Miller, J.R. und Moon, R.T. (1997)**
Analysis of the Signaling Activities of Localization Mutants of -Catenin during Axis Specification in Xenopus. *J. Cell Biol.* **139**: 229-243
- Miyatani, S., Shimamura, K., Hatta, M., Nagafuchi, A., Nose, A., Matsunaga, M., Hatta, K., Takeichi, M. (1989)** Neural cadherin: role in selective cell-cell adhesion. *Science* **245**: 631-635
- Moscona, A. (1961)**
Rotation-mediated histogenetic aggregation of dissociated cells: a quantifiable approach to cell interaction in vitro. *Exp. Cell Res.* **22**: 455-475
- Molenaar, M., van de Wetering, M., Oosterwegel, M., Peterson-Maduro, J., Godsave, S., Korinek, V., Roose, J., Destree, O., Clevers, H. (1996)**
Xtcf-3 transcription factor mediates -catenin-induced axis formation in Xenopus embryos. *Cell* **86**: 391-399
- Müller, A.H.J., Angres, B., Hausen, P. (1992)**
U-Cadherin in Xenopus oogenesis and oocyte maturation. *Development* **114**: 533-543

- Müller, A.H.J., Kühl, M., Finnemann, S., van der Poel, S., Schneider, S., Hausen, P., Wedlich, D. (1994) Xenopus cadherins: the maternal pool comprises distinguishable members of the family. *Mech. Dev.* **47**: 213-223
- Müller, A.H.J. und Hausen, P. (1995)
Epithelial cell polarity in Xenopus development. *Dev. Dyn.* **202**: 405-420
- Mullins, M., Hammerschmidt, M., Haffter, P., Nüsslein-Vollhard, C. (1993)
Mutagenesis and screen for embryonic mutants in the zebrafish, *Brachydanio rerio*.
Tagung der Gesellschaft für Entwicklungsbiologie, München, Abstract
- Munemitsu, S., Souza, B., Müller, O., Albert, I., Rubinfeld, B., Polakis, P. (1994)
(1994) The APC gene product associates with microtubules in vivo and promotes their assembly in vitro. *Cancer Research* **54**: 3676-3681
- Näthke, I., Hinck, L., Swedlow, J., Papkoff, J., Nelson, W. (1994)
Defining interaction and distributions of cadherin and catenin complexes in polarized epithelial cells. *J. Cell Biol.* **125**: 1341-1352
- Nagafuchi, A., Shirayoshi, Y., Okazaki, K., Yasuda, K., Takeichi, M. (1987)
Transformation of cell adhesion properties by exogenously introduced E-cadherin cDNA. *Nature* **329**: 341-343
- Nagafuchi, A. und Takeichi, M. (1988)
Cell binding function of E-cadherin is regulated by the cytoplasmic domain.
EMBO J. **7**: 3679-3784
- Nagafuchi, A., Takeichi, M., Tsukita, S. (1991)
The 102 kD cadherin associated protein: similarity to vinculin and posttranscriptional regulation of expression. *Cell* **65**: 849-857
- Nakatsuji, N. und Johnson, K.E. (1983) Comparative study of extracellular fibrils on the ectodermal layer of gastrulae of five amphibian species. *J. Cell Sci.* **59**: 61-70
- Napolitano, E.W., Venstrom, K., Wheeler, E.F., Reichardt, L.F. (1991)
Molecular cloning and characterization of B-cadherin, a novel chick cadherin.
J. Cell Biol. **113**: 893-905
- Niehrs, C., Keller, R., Cho, K.W.Y., De Robertis, E.M. (1993)
The Homeboxgene goosecoid Controls Cell Migration in Xenopus Embryos. *Cell* **72**: 491-503
- Nieuwkoop, P.D. und Faber, J. (1967)
Normal table of *Xenopus laevis*. (Daudin), Amsterdam, North Holland Publishing Co.
- Nose, A., Nagafuchi, A., Takeichi, M. (1987)
Isolation of placental cDNA: identification of a novel gene family for cell-cell adhesion molecules. *EMBO J.* **6**: 3655-3661
- Nose, A., Tsuji, K., Takeichi, M. (1988)
Expressed recombinant cadherins mediate cell sorting in model systems.
Cell **54**: 993-1001

Nose, A., Tsuji, K., Takeichi, M. (1990)

Localization of specificity determining sites in cadherin cell adhesion molecules.
Cell **61**: 147-155

Nusse, R. und Varmus, H.E. (1992)

Wnt genes. *Cell* **69**: 1073-1087

Ozawa, M., Baribault, H., Kemler, R. (1989)

The cytoplasmic domain of the cell adhesion molecule uvomorulin associates with three independent proteins structurally related in different species. *EMBO J.* **8**: 1711-1717

Ozawa, M. und Kemler, R. (1990)

Correct proteolytic cleavage is required for the cell adhesive function of uvomorulin.
J. Cell Biol. **111**: 1645-1650

Ozawa, M., Engel, J., Kemler, R. (1990a)

Single amino acid substitutions in one Ca^{2+} binding site of uvomorulin abolish the adhesive function. *Cell* **63**: 1033-1038

Ozawa, M., Ringwald, M., Kemler, R. (1990b)

Uvomorulin-catenin complex formation is regulated by a specific domain in the cytoplasmic region of the cell adhesion molecule. *Proc. Natl. Acad. Sci. USA* **87**: 4246-4250

Ozawa, M., Hoschützky, H., Herrenknecht, K., Kemler, R. (1991)

A possible new adhesive site in the cell adhesion molecule uvomorulin.

Mech. Dev. **33**: 49-56

Ozawa, M. und Kemler, R. (1992)

Molecular organization of the uvomorulin-catenin complex. *J. Cell Biol.* **116**: 989-996

Overduin, M., Harvey, T.S., Bagby, S., Tong, K.I., Yau, P., Takeichi, M.,

Ikura, M. (1995) Solution structure of the epithelial cadherin domain responsible for selective cell adhesion. *Science* **267**: 386-389

Papkoff, J., Rubinfeld, B., Schryver, B., Polakis, P. (1996)

Wnt-1 regulates free pools of catenins and stabilizes APC-catenin complexes.

Mol. Cell Biol. **16**: 2128-2134

Peifer, M., McCrea, P., Green, K., Wieschaus, E., Gumbiner, B. (1992)

The vertebrate adhesive junction proteins beta-catenin and plakoglobin and the Drosophila segment polarity gene armadillo form a multigene family with similar properties.

J. Cell Biol. **118**: 681-691

Peyrieras, N., Bause, E., Legler, G., Vasilov, R., Claesson, L. (1983)

Effects of the glucosidase inhibitors nojirimycin and deoxynojirimycin on the biosynthesis of membrane and secretory glycoproteins. *EMBO J.* **2**: 823-832

Peyrieras, N., Louvard, D., Jaco, F. (1985)

Characterization of antigens recognized by monoclonal and polyclonal antibodies directed against uvomorulin. *Proc. Natl. Acad. Sci. USA* **82**: 8067-8071

- Piccolo, S., Sasai, Y., Lu, B., De Robertis, E.M. (1996)**
Dorsoventral patterning in Xenopus: inhibition of ventral signals by direct binding of chordin to BMP-4. *Cell* **86**: 589-598
- Pierce, S.B. und Kimelman, D. (1995)**
Regulation of Speman organizer formation by the intracellular kinase Xgsk-3. *Development* **121**: 755-765
- Pokutta, S., Herrenknecht, K., Kemler, R., Engel, J. (1994)**
Conformational changes of the recombinant extracellular domain of E-cadherin upon calcium binding. *Eur. J. Biochem.* **223**: 1019-1026
- Ramos, J.W. und DeSimone, D.W. (1996)** Xenopus embryonic cell adhesion to fibronectin: Position-specific activation of RGD/synergy-site-dependent migratory behavior at gastrulation. *J. Cell Biol.* **134**: 227-240
- Ramos, J.W., Whittaker, C.A., DeSimone, D.W. (1996)**
Integrin-dependent adhesive activity is spatially controlled by inductive signals at gastrulation. *Development* **122**: 2873-2883
- Ransom, D.G., Hens, M.D., DeSimone, D.W. (1993)**
Integrin Expression in Early Amphibian Embryos: cDNA Cloning and Characterization of Xenopus ₁, ₂, ₃, and ₆ Subunits. *Dev.Biol.* **160**:265-275
- Rantsch, B., Dours, M.T. (1989)**
Selective expression of a novel cadherin in the pathways of the developing motor and commissural axons. *Soc. Neurosci. Abstr.* **15**: 382-386
- Rantsch, B., Dours-Zimmermann, M.T. (1991)**
T-Cadherin, a novel cell adhesion molecule inthe nervous system lacks the conserved cytoplasmic region. *Neuron* **7**: 391-402
- Represa, J. und Slack, J.M.W. (1989)**
Mesoderm induction by the mesoderm of Xenopus neurulae. *Int. J. Dev. Biol.* **33**, 397-401
- Reyes, A.A., Akeson, R., Brezina, L., Cole, G.J. (1990)**
Structural requirements for neural cell adhesion molecule-heparin interaction. *Cell Int.* **1**:567-576
- Reynolds, A.B., Daniel, J., McCrea, P.D., Wheelock, M.J., Wu, J., Zhang, Z. (1994)** Identification of a new catenin: The tyrosine kinase substrate p120^{CAS} associates with E-cadherin complexes. *Mol. Cell Biol.* **14**: 8333-8342
- Ringwald, M., Schuh, R., Vestweber, D., Eistetter, H., Lottspeich, F., Engel, J., Dölz, R., Jähnig, F., Epplen, J., Mayer, S., Müller, C., Kemler, R. (1987)**
The structure of cell adhesion molecule uvomorulin. Insights into the molecular mechanism of Ca²⁺-dependent cell adhesion. *EMBO J.* **6**: 3647-3653
- Rowning, B.A., Wells, J., Wu, M., Gerhart, J.C., Moon, R.T., Larabell, C.A. (1997)** Microtubule mediated transport of organelles and location of -catenin to future dorsal side of Xenopus eggs. *Proc. Natl. Acad. Sci. USA* **94**: 1224-1229

Rupp, R.A.W. und Weintraub, H. (1991) Ubiquitous MyoD transcription at the mid blastula transition precedes induction-dependent MyoD expression in presumptive mesoderm of *Xenopus laevis*. *Cell* **65**: 927-937

Sanger, F. und Coulson, A.R. (1977) A sequencing with chain-terminating inhibitors. *Proc. Natl. Acad. Sci. USA* **74**: 5463-5467

Sano, K., Tanihara, H., Heimark, R.L., Obata, S., Davidson, M., St. John, T., Taketani, S., Suzuki, S. (1993) Protocadherins: a large family of cadherin-related molecules in central nervous system. *EMBO J.* **12**: 2249-2256

Sasai, Y., Lu, B., Steinbeisser, H., Geissert, D., Gont, L.K., De Robertis, E.M. (1994) Xenopus chordin: a novel dorsalizing factor activated by organizer-specific homeobox genes. *Cell* **70**: 779-790

Sastry, S.K. und Horwitz, A.F. (1993) Integrin cytoplasmic domains: mediators of cytoskeletal linkages and extra- and intracellular initiated transmembrane signalling. *Curr. Opin. Cell Biol.* **5**: 819-831

Schuh, R., Vestweber, D., Riede, L., Ringwald, M., Rosenberg, U.B., Jaeckle, H., Kemler, R. (1986) Molecular cloning of the mouse cell adhesion molecule uvomorulin: cDNA contains a B1-related sequence. *Proc. Natl. Acad. Sci. USA* **83**: 1364-1368

Shapiro, L., Fannon, A.M., Kwong, P.D., Thomson, A., Lehmann, M.S., Grübel, G., Legrand, J.-F., Als-Nielsen, J., Colman, D.R., Hendrickson, W.A. (1995) Structural basis of cell-cell adhesion by cadherins. *Nature* **374**: 327-337

Shibamoto, S.M., Hayakawa, M., Takeuchi, K., Hori, T., Miyazawa, K., Kitamura, N., Johnson, K.R., Wheelock, M.J., Matsuyoshi, N., Takeichi, M., Ito, F. (1995) Association of p120, a tyrosine kinase substrate, with E-cadherin/catenin complexes. *J. Cell Biol.* **128**: 949-957

Shimoyama, Y., Yoshida, T., Terada, M., Shimosato, Y., Abe, O., Hirohashi, S. (1989) Molecular cloning of a human Ca^{2+} -dependent cell-cell adhesion molecule homologous to mouse placental cadherin: its low expression in human placental tissues. *J. Cell Biol.* **109**: 1787-1794

Shirayoshi, Y., Nose, A., Iwasaki, K., Takeichi, M. (1986) N-linked oligosaccharides are not involved in the function of cell-cell binding glycoprotein E-cadherin. *Cell Struct. Funct.* **11**: 245-252

Simcha, I., Shtutman, M., Salomon, D., Zhurinsky, J., Sadot, E., Geiger, B., Ben-Ze'ev, A. (1998) Differential Nuclear Translocation and Transactivation Potential of -Catenin and Plakoglobin. *J. Cell Biol.* **141**: 1433-1448

Simmoneau, L., Broders, F., Thiery, J.P. (1992) N-Cadherin transcripts in *Xenopus laevis* from early tailbud to tadpole. *Dev. Dyn.* **194**: 247-260

- Sive, H.** (1993)
The frog prince-ss: A molecular formula for dorsoventral patterning in *Xenopus*.
Genes & Dev. 7: 1-12
- Slack, J.M.W.** (1991)
From Egg to Embryo. *Cambridge University Press*, Cambridge: 93-110
- Slack, J.M.W.** (1994)
Inducing factors in *Xenopus* early embryos. *Curr. Biol.* 4: 116-126
- Slack, J.M.W.** (1996)
High hops of transgenic frogs. *Nature* 383:765-766
- Smith, W.C., Harland, R.M.** (1992)
Expression cloning of noggin, a new dorsalizing factor localized to the Spemann organizer in *Xenopus* embryos. *Cell* 70: 829-840
- Smith, J.C., Symes, K., Hynes, R.O., DeSimone, D.** (1990) Mesoderm induction and the control of gastrulation in *Xenopus laevis*: The roles of fibronectin and integrins.
Development 108: 229-238
- Sokol, S.Y.** (1993) Mesoderm formation in *Xenopus* ectodermal explants overexpressing Xwnt-8: evidence for a cooperating signal reaching the animal pole by gastrulation.
Development 118: 1335-1342
- Sokol, S.Y.** (1994) The pregastrula establishment of gene expression pattern in *Xenopus* embryos: Requirements for local cell interactions and for protein synthesis.
Dev. Biol. 166: 782-788
- Sokol, S.Y. und Melton, D.A.** (1992)
Interaction of Wnt and Activin in Dorsal Mesoderm Induction in *Xenopus*.
Dev. Biol. 154: 348-355
- Sokol, S.Y., Klingensmith, J., Perrimon, N., Itoh, K.** (1995)
Dorsalizing and neuralizing properties of Xdsh, a maternally expressed *Xenopus* homolog of dishevelled. *Development* 121: 1637-1647
- Spemann, H.** (1938)
Embryonic Development and Induction. *Yale university Press*, New Haven: 214
- Spemann, H. und Mangold, H.** (1924)
Über Induktion von Embryonalanlagen durch Implantation artfremder Organisatoren.
Roux´s Archiv 100: 599-638
- Spiegel, M. und Spiegel, E.S.** (1975)
Reaggregation of dissociated embryonic sea urchin cells. *Am. Zool.* 15: 583-606
- Staddon, J.M., Smales, C., Schulze, C., Esch, F.S., Rubin, L.L.** (1995)
p120, a p120-related protein (p100), and the cadherin/catenin complex.
J. Cell Biol. 130: 369-381

Stappert, J. und Kemler, R. (1994)

A short core region of E-cadherin is essential for catenin binding and is highly phosphorylated. *Cell Adhes. Commun.* 2: 319-327

Steinberg, M.S. und Poole, T.J. (1982)

Cellular adhesive differentials as determinants of morphogenetic movements and organ segregation. S. Subtelny and P.B. Green (eds.), Developmental Order: Its Origin and Regulation. Alan R. Liss, New York: 351-378

Suzuki, S.T. (1996)

Structural and functional diversity of cadherin superfamily: are new members of cadherin superfamily involved in signal transduction pathway? *J. Cell Bioch.* 61: 531-542

Takeichi, M. (1977)

Functional correlations between cell adhesive properties and some cell surface proteins. *J. Cell Biol.* 75: 464-474

Takeichi, M. (1987)

Cadherins: a molecular family essential for selective cell-cell adhesion and animal morphogenesis. *Trends Gen.* 3: 213-217

Takeichi, M. (1988)

The cadherins: cell-cell adhesion molecules controlling animal morphogenesis. *Development* 102: 639-655

Takeichi, M. (1990)

Cadherins: a molecular family important in selective cell-cell adhesion. *Annu. Rev. Biochem.* 59: 237-252

Takeichi, M. (1995)

Morphogenetic roles of classical cadherins. *Curr. Opin. Cell Biol.* 7: 619-627

Tanahara, H., Sano, K., Heimark, R.L., St. John, T., Suzuki, S. (1994)

Cloning of five human cadherins clarifies characteristic features of cadherin extracellular domain and provides further evidence for two structurally different types of cadherins. *Cell Adh. Comm.* 2: 15-26

Tashiro, K., Too, O., Nakamura, H., Koga, C., Ito, Y., Hikasa, H., Shiokawa, K. (1995) Cloning and expression of cDNA for a novel Xenopus cadherin (XmN-cadherin), expressed maternally and later neural-specifically in embryogenesis.

Mech. Dev. 54: 161-171

Theisen, H., Purcell, J., Bennet, M., Kansagara, D., Marsh, J.L. (1994)

Dishevelled is required during wingless signaling to establish both cell polarity and cell identity. *Development* 120: 347-360

Thiery, J.P., Delouvee, A., Gallin, W.J., Cunningham, B.A., Edelman, G.M.

(1984) Ontogenetic expression of the cell adhesion molecule L-CAM is found in epithelia derived from the three germ layers. *Dev. Biol.* 102: 61-78

- Tooi, O., Fujii, G., Tashiro, K., Shiokawa, K.** (1994)
Molecular cloning of cDNA for XTCAD-1, a novel Xenopus cadherin, and its expression in adult tissue and embryos of *Xenopus laevis*. *Biochim. Biophys. Acta* **1219**: 121-128
- Torres, M., Yang-Snyder, J.A., Purcell, S.M., DeMarais, A.A., McGrew, L.L., Moon, R.T.** (1996) Activities of the Wnt-1 class of secreted signaling factors are antagonized by the Wnt-5a class and by a dominant negative cadherin in early *Xenopus* development. *J. Cell Biol.* **113**: 867-879
- Townes, P. und Holtfreter, J.** (1955)
Directed movements and selective adhesion of embryonic amphibian cells. *J. Exp. Zool.* **128**: 53-120
- Trinkaus, J.** (1963)
The cellular basis of Fundulus epiboly. *Dev. Biol.* **7**: 513-532
- Tsukita, S., Olshi, K., Akiyama, T., Yamanashi, Y., Yamamoto, T.** (1991)
Specific proto-oncogenic tyrosine kinases of src family are enriched in cell-to-cell adherens junctions where the level of tyrosine phosphorylation is elevated. *J. Cell Biol.* **113**: 867-879
- Vestweber, D., Gossler, A., Boller, K., Kemler, R.** (1987)
Expression and distribution of cell adhesion molecule uvomorulin in mouse preimplantation embryos. *Dev. Biol.* **124**: 452-456
- Volk, T. und Geiger, B.** (1984)
A 135-kD membrane protein of intercellular adherens junctions. *EMBO J.* **3**: 2249-2260
- Volk, T. und Geiger, B.** (1986)
A-CAM: A 135 kD receptor of intracellular adherens junction: immunoelectron microscopy, localization and biochemical studies. *J. Cell Biol.* **103**: 1441-1450
- Volk, T., Cohen, O., Geiger, B.** (1987)
Formation of heterotypic adherens-type junctions between L-CAM containing liver cells and A-CAM containing lens cells. *Exp. Cell Res.* **191**: 186-193
- Wagner, G.** (1995)
E-Cadherin: a distant member of the immunoglobulin superfamily. *Science* **267**: 342
- Wahl, J.K., Sacco, P.A., McGranahan-Sadler, T.M., Sauppé, L.M., Wheelock, M.J., Johnson, K.R.** (1996)
Plakoglobin domains that define its association with the desmosomal cadherins and the classical cadherins: identification of unique and shared domains. *J. Cell Sci.* **109**: 1143-1154
- Wang, S., Krinks, M., Lin, K., Luyten, F.P., Moos Jr., M.** (1997)
Frzb, a secreted protein expressed in the Spemann organizer, binds and inhibits Wnt-8. *Cell* **88**: 757-766

- Watabe, M., Nagafuchi, A., Tsukita, S., Takeichi, M.** (1994)
Induction of polarized cell-cell association and retardation of growth by activation of the E-cadherin-catenin adhesion system in a dispersed carcinoma line.
J. Cell Biol. **127**: 247-256
- Wenstrom, S., Siegbahn, A., Yokote, K., Arvidsson, A.K., Heldin, C.H., Mori, S., Claesson-Welsh, S.** (1994)
Membrane ruffling and chemotaxis transduced by the PDGF -receptor require the binding site for phosphatidylinositol 3' kinase. *Oncogene* **9**: 651-660
- Weston, M.J.D., Kato, K., Gurdon, J.B.** (1994)
A community effect is required for amphibian notochord differentiation.
Roux's Arch. Dev. Biol. **203**: 250-253
- Wheelock, M.J.** (1990)
Catenin association with E-cadherin changes with the state of polarity of HT-29 cells.
Exp. Cell Res. **191**: 186-193
- Wheelock, M.J. und Knudsen, K.A.** (1991)
N-Cadherin associated proteins in chicken muscle. *Differentiation* **46**: 35-42
- Whittaker, C.A., DeSimone, D.W.** (1993) Integrin a subunit mRNAs are differentially expressed in early Xenopus embryos. *Development* **117**: 1239-1249
- Wieslander, L.** (1979) A simple method to recover high molecular weight RNA and DNA after elektrophoretic separation in low gelling temperature agarose gels.
Analytical Biochem. **98**: 305-309
- Williams, E.J., Furness, J., Walsh, F.S., Doherty, P.** (1994)
Activation of the FGF receptor underlies neurite outgrowth stimulated by L1, N-CAM, and N-Cadherin. *Neuron* **13**: 583-594
- Wilson, P.A. und Melton, D.A.** (1994)
Mesodermal patterning by an inducer gradient depends on secondary cell-cell communication. *Curr. Biol.* **4**: 676-686
- Winkelbauer, R.** (1989)
A reinvestigation of the role of fibronectin in amphibian gastrulation.
J. Cell Biol. **107**: 815
- Winkelbauer, R., Selchow, A., Nagel, M., Angres, B.** (1992)
Cell Interaction and Its Role in Mesoderm Cell Migration During Xenopus Gastrulation.
Dev. Dyn. **195**: 290-302
- Winkelbauer, R., Selchow, A., Nagel, M., Angres, B.** (1992)
Cell interaction and its role in mesoderm cell migration during Xenopus gastrulation.
Dev. Dynam. **195**: 290-302
- Winkelbauer, R. und Keller, R.E.** (1996)
Fibronectin, Mesoderm Migration, and Gastrulation in Xenopus. *Dev. Biol.* **177**: 413-426

Wylie, C., Kofron, M., Payne, C., Anderson, R., Hosobuchi, M., Joseph, E., Heasman, J. (1996) Maternal β -catenin establishes a 'dorsal signal' in early Xenopus embryos. *Development* **122**: 2987-2996

Yap, S., Niessen, C.M., Gumbiner, B.M. (1998)
The Juxtamembrane Region of the Cadherin Cytoplasmic Tail Supports Lateral Clustering, Adhesive Strengthening, and Interaction wih p120^{ctn}.
J. Cell Biol. **141**: 779-789

Yost, H.J. (1992) Regulation of vertebrate left-right asymmetries by extracellular matrix. *Nature* **357**: 158-161

Yost, C., Torres, M., Miller, J.R., Huang, E., Kimelman, D., Moon, R.T.
(1996) The axis-inducing activity, stability, and subcellular distribution of γ -catenin is regulated in Xenopus embryos by glycogen synthase kinase 3. *Genes & Dev.* **10**: 1443-1454

Zeng, L., Fagotto, F., Zhang, T., Hsu, W., Vasicek, T.J., Perry III, W.L., Lee, J.J., Tilghman, S.M., Gumbiner, B.M., Costantini, F. (1997)
The mouse fused locus encodes axin, an inhibitor of the Wnt signaling pathway that regulates embryonic axis formation. *Cell* **90**: 181-192

Zimmermann, L.B., De Jesus-Escobar, J.E., Harland, R.M. (1996)
The Spemann organizer signal noggin binds and inactivates bone morphogenetic protein-4.
Cell **86**: 599-606