

V LITERATURVERZEICHNIS

Abrahamson SL, Wu HM, Williams RE, Der K, Ottah N, Little R, Gazzano-Santoro H, Theofan G, Bauer R, Leigh S, Orme A, Horwitz AH, Carroll SF, Dedrick RL.: Biochemical characterization of recombinant fusions of lipopolysaccharide binding protein and bactericidal/permeability-increasing protein. Implications in biological activity. J Biol Chem. 1997 Jan 24; 272(4): 2149-55.

Akashi S, Ogata H, Kirikae F, Kirikae T, Kawasaki K, Nishijima M, Shimazu R, Nagai Y, Fukudome K, Kimoto M, Miyake K.: Regulatory roles for CD14 and phosphatidylinositol in the signaling via toll-like receptor 4-MD-2. Biochem Biophys Res Commun. 2000 Feb 5;268(1):172-7.

Alberti C, Brun-Buisson C, Burchardi H, Martin C, Goodman S, Artigas A, Sicignano A, Palazzo M, Moreno R, Boulme R, Lepage E, Le Gall R.: Epidemiology of sepsis and infection in ICU patients from an international multicentre cohort study. Intensive Care Med. 2002 Feb;28(2):108-21. Epub 2001 Dec 04. Erratum in: Intensive Care Med 2002 Apr;28(4):525-6.

Alexander C, Rietschel ET.: Bacterial lipopolysaccharides and innate immunity. J Endotoxin Res 2001; (3): 67-202. Review

Au-Young J, Fielding CJ.: Synthesis and secretion of wild-type and mutant human plasma cholesteryl ester transfer protein in baculovirus-transfected insect cells: the carboxyl-terminal region is required for both lipoprotein binding and catalysis of transfer. Proc Natl Acad Sci U S A. 1992 May 1;89(9):4094-8.

Baenkler, H.W. (Hrsg): *Innere Medizin*. Hippokrates Verlag im Thieme-Verlag, Stuttgart 1999

Bazil V, Strominger JL.: Shedding as a mechanism of down-modulation of CD14 on stimulated human monocytes. *J Immunol* 1991 Sep 1;147(5):1567-74

Beamer LJ, Carroll SF, Eisenberg D.: Crystal structure of human BPI and two bound phospholipids at 2.4 angstrom resolution. *Science*. 1997 Jun 20;276(5320):1861-4.

Beamer LJ, Carroll SF, Eisenberg D.: The three-dimensional structure of human bactericidal/permeability-increasing protein: implications for understanding protein-lipopolysaccharide interactions. *Biochem Pharmacol*. 1999 Feb 1;57(3):225-9. Review.

Beamer, L.J., Carroll, S.F., Eisenberg, D.: The BPI/LBP family of proteins: a structural analysis of conserved regions. *Protein Sci* 1998; 7/4: 906-914

Beutler B, Hoebe K, Du X, Janssen E, Georgel P, Tabeta K.: Lps2 and signal transduction in sepsis: at the intersection of host responses to bacteria and viruses. *Scand J Infect Dis*. 2003;35(9):563-7. Review.

Beutler B.: Tlr4: central component of the sole mammalian LPS sensor. *Curr Opin Immunol*. 2000 Feb;12(1):20-6. Review.

Bone RC.: The pathogenesis of sepsis. *Ann Intern Med* 1991 Sep 15;115(6):457-69

Brun-Buisson C, Doyon F, Carlet J, Dellamonica P, Gouin F, Lepoutre A, Mercier JC, Offenstadt G, Regnier B.: Incidence, risk factors, and outcome of severe sepsis and septic shock in adults. A multicenter prospective study in intensive care units. French ICU Group for Severe Sepsis. *JAMA*. 1995 Sep 27;274(12):968-74.

Cavaillon JM, Adib-Conquy M, Fitting C, Adrie C, Payen D.: Cytokine cascade in sepsis. *Scand J Infect Dis*. 2003;35(9):535-44. Review.

da Silva Correia J, Soldau K, Christen U, Tobias PS, Ulevitch RJ.: Lipopolysaccharide is in close proximity to each of the proteins in its membrane receptor complex. transfer from CD14 to TLR4 and MD-2. J Biol Chem. 2001 Jun 15;276(24):21129-35.

Elsbach P. The bactericidal/permeability-increasing protein (BPI) in antibacterial host defense. J Leukoc Biol. 1998 Jul;64(1):14-8. Review.

Fink, P.C.: Septischer Schock und Endotoxinämie. In: Peter, H.-H., Pichler, W.J. (Hrsg.): *Klinische Immunologie*. 2. Auflage, Urban & Schwarzenberg, München 1996

Frey EA, Miller DS, Jahr TG, Sundan A, Bazil V, Espevik T, Finlay BB, Wright SD: Soluble CD14 participates in the response of cells to lipopolysaccharide. J Exp Med. 1992 Dec 1;176(6):1665-71.

Friedman G, Silva E, Vincent JL.: Has the mortality of septic shock changed with time. Crit Care Med. 1998 Dec;26(12):2078-86. Review.

Gardlund B, Sjolín J, Nilsson A, Roll M, Wickerts CJ, Wretling B.: Plasma levels of cytokines in primary septic shock in humans: correlation with disease severity. J Infect Dis. 1995 Jul;172(1):296-301.

Glauser MP, Zanetti G, Baumgartner JD, Cohen J.: Septic shock: pathogenesis. Lancet. 1991 Sep 21;338(8769):732-6. Review.

Grube BJ, Cochane CG, Ye RD, Green CE, McPhail ME, Ulevitch RJ, Tobias PS.: Lipopolysaccharide binding protein expression in primary human hepatocytes and HepG2 hepatoma cells. J Biol Chem. 1994 Mar 18;269(11):8477-82.

Gupta D, Kirkland TN, Viriyakosol S, Dziarski R: CD14 is a cell-activating receptor for bacterial peptidoglycan. *J Biol Chem*. 1996 Sep 20;271(38):23310-6.

Gutsmann T, Muller M, Carroll SF, MacKenzie RC, Wiese A, Seydel U.: Dual role of lipopolysaccharide (LPS)-binding protein in neutralization of LPS and enhancement of LPS-induced activation of mononuclear cells. *Infect Immun*. 2001 Nov;69(11):6942-50.

Hailman E, Albers JJ, Wolfbauer G, Tu AY, Wright SD.: Neutralization and transfer of lipopolysaccharide by phospholipid transfer protein. *J Biol Chem* 1996 May 24;271(21):12172-8

Hailman E, Lichenstein HS, Wurfel MM, Miller DS, Johnson DA, Kelley M, Busse LA, Zukowski MM, Wright SD.: Lipopolysaccharide (LPS)-binding protein accelerates the binding of LPS to CD14. *J Exp Med*. 1994 Jan 1;179(1):269-77.

Hailman E, Vasselon T, Kelley M, Busse LA, Hu MC, Lichenstein HS, Detmers PA, Wright SD.: Stimulation of macrophages and neutrophils by complexes of lipopolysaccharide and soluble CD14. *J Immunol* 1996 Jun 1; 156(11): 4384-90

Hamann L, Schumann RR, Flad HD, Brade L, Rietschel ET, Ulmer AJ.: Binding of lipopolysaccharide (LPS) to CHO cells does not correlate with LPS-induced NF-kappaB activation. *Eur J Immunol* 2000 Jan; 30(1): 211-6

Han J, Mathison JC, Ulevitch RJ, Tobias PS.: Lipopolysaccharide (LPS) binding protein, truncated at Ile-197, binds LPS but does not transfer LPS to CD14. *J Biol Chem*. 1994 Mar 18; 269(11): 8172-5.

Haziot A, Ferrero E, Kontgen F, Hijiya N, Yamamoto S, Silver J, Stewart CL, Goyert SM: Resistance to endotoxin shock and reduced dissemination of gram-negative bacteria in CD14-deficient mice. *Immunity*. 1996 Apr;4(4):407-14.

Haziot A, Rong GW, Lin XY, Silver J, Goyert SM: Recombinant soluble CD14 prevents mortality in mice treated with endotoxin (lipopolysaccharide). *J Immunol.* 1995 Jun 15;154(12):6529-32

Heine H, Rietschel ET, Ulmer AJ.: The biology of endotoxin. *Mol Biotechnol* 2001 Nov; 19(3): 279-96. Review

Heumann D, Gallay P, Barras C, Zaech P, Ulevitch RJ, Tobias PS, Glauser MP, Baumgartner JD.: Control of lipopolysaccharide (LPS) binding and LPS-induced tumor necrosis factor secretion in human peripheral blood monocytes. *J Immunol* 1992 Jun 1;148(11):3505-12

Hubacek JA, Buchler C, Aslanidis C, Schmitz G.: The genomic organization of the genes for human lipopolysaccharide binding protein (LBP) and bactericidal permeability increasing protein (BPI) is highly conserved. *Biochem Biophys Res Commun.* 1997 Jul 18;236(2):427-30.

Huuskonen J, Wohlfahrt G, Jauhiainen M, Ehnholm C, Telemann O, Olkkonen VM.: Structure and phospholipid transfer activity of human PLTP: analysis by molecular modeling and site-directed mutagenesis. *J Lipid Res.* 1999 Jun;40(6):1123-30.

Jack RS, Fan X, Bernheiden M, Rune G, Ehlers M, Weber A, Kirsch G, Mentel R, Furll B, Freudenberg M, Schmitz G, Stelter F, Schutt C: Lipopolysaccharide-binding protein is required to combat a murine gram-negative bacterial infection. *Nature* 1997 Oct 16; 389(6652): 742-5

Janeway CA Jr.: Introduction: The role of innate immunity in the adaptive immune response. *Seminars In Immunology* 1998, Vol 10: 329-350

Jiang, X., Bruce, C., Cocke, T., Wang, S., Boguski, M., Tall, R.T.: Point mutagenesis of positively charged amino acids of cholesteryl ester transfer protein: conserved residues within the lipid transfer/lipopolysaccharide binding protein gene family essential for function. *Biochemistry* 1995; 34: 7258-7263

Johnston RB Jr.: Current concepts: immunology. Monocytes and macrophages. *N Engl J Med.* 1988 Mar 24;318(12):747-52. Review.

Juan TS, Hailman E, Kelley MJ, Busse LA, Davy E, Empig CJ, Narhi LO, Wright SD, Lichenstein HS: Identification of a lipopolysaccharide binding domain in CD14 between amino acids 57 and 64. *J Biol Chem.* 1995 Mar 10;270(10):5219-24.

Juan TS, Kelley MJ, Johnson DA, Busse LA, Hailman E, Wright SD, Lichenstein HS: Soluble CD14 truncated at amino acid 152 binds lipopolysaccharide (LPS) and enables cellular response to LPS. *J Biol Chem.* 1995 Jan 20;270(3):1382-7.

Juan TS, Hailman E, Kelley MJ, Wright SD, Lichenstein HS: Identification of a domain in soluble CD14 essential for lipopolysaccharide (LPS) signaling but not LPS binding. *J Biol Chem.* 1995 Jul 21;270(29):17237-42.

Kirschning CJ, Au-Young J, Lamping N, Reuter D, Pfeil D, Seilhamer JJ, Schumann RR.: Similar organization of the lipopolysaccharide-binding protein (LBP) and phospholipid transfer protein (PLTP) genes suggests a common gene family of lipid-binding proteins. *Genomics* 1997 Dec 15;46(3):416-25

Kirschning CJ, Unbehaun A, Fiedler G, Hallatschek W, Lamping N, Pfeil D, Schumann RR.: The transcriptional activation pattern of lipopolysaccharide binding protein (LBP) involving transcription factors AP-1 and C/EBP beta. *Immunobiology* 1997 Dec;198(1-3):124-35

Kirschning C, Unbehauen A, Lamping N, Pfeil D, Herrmann F, Schumann RR.: Control of transcriptional activation of the lipopolysaccharide binding protein (LBP) gene by proinflammatory cytokines. *Cytokines Cell Mol Ther* 1997 Mar;3(1):59-62

Kopp EB, Medzhitov R.: The Toll-receptor family and control of innate immunity. *Curr Opin Immunol*. 1999 Feb;11(1):13-8. Review.

Krutzik SR, Sieling PA, Modlin RL.: The role of Toll-like receptors in host defense against microbial infection. *Curr Opin Immunol*. 2001 Feb;13(1):104-8.

Lamping N, Dettmer R, Schroder NW, Pfeil D, Hallatschek W, Burger R, Schumann RR.: LPS-binding protein protects mice from septic shock caused by LPS or gram-negative bacteria. *J Clin Invest*. 1998 May 15;101(10):2065-71.

Lamping N, Hoess A, Yu B, Park TC, Kirschning CJ, Pfeil D, Reuter D, Wright SD, Herrmann F, Schumann RR.: Effects of site-directed mutagenesis of basic residues (Arg 94, Lys 95, Lys 99) of lipopolysaccharide (LPS)-binding protein on binding and transfer of LPS and subsequent immune cell activation. *J Immunol*. 1996 Nov 15;157(10):4648-56.

Lee JD, Kato K, Tobias PS, Kirkland TN, Ulevitch RJ.: Transfection of CD14 into 70Z/3 cells dramatically enhances the sensitivity to complexes of lipopolysaccharide (LPS) and LPS binding protein. *J Exp Med* 1992 Jun 1;175(6):1697-705

Loeffler, G.: Leber. In: Löffler, G., Petrides, P.E. (Hrsg.): *Biochemie und Pathobiochemie*. 5. Auflage, Springer-Verlag, Heidelberg 1997

Lynn WA, Liu Y, Golenbock DT.: Neither CD14 nor serum is absolutely necessary for activation of mononuclear phagocytes by bacterial lipopolysaccharide. *Infect Immun* 1993 Oct;61(10):4452-61

Martin GS, Mannino DM, Eaton S, Moss M.: The epidemiology of sepsis in the United States from 1979 through 2000. *N Engl J Med.* 2003 Apr 17;348(16):1546-54.

Martin TR, Mathison JC, Tobias PS, Leturcq DJ, Moriarty AM, Maunder RJ, Ulevitch RJ.: Lipopolysaccharide binding protein enhances the responsiveness of alveolar macrophages to bacterial lipopolysaccharide. Implications for cytokine production in normal and injured lungs. *J Clin Invest* 1992 Dec;90(6):2209-19

Mathison JC, Tobias PS, Wolfson E, Ulevitch RJ.: Plasma lipopolysaccharide (LPS)-binding protein. A key component in macrophage recognition of gram-negative LPS. *J Immunol.* 1992 Jul 1; 149(1): 200-6.

McCallum KL, Schoenhals G, Laakso D, Clarke B, Whitfield C.: A high-molecular-weight fraction of smooth lipopolysaccharide in *Klebsiella* serotype O1:K20 contains a unique O-antigen epitope and determines resistance to nonspecific serum killing. *Infect Immun.* 1989 Dec;57(12):3816-22.

Medzhitov R, Janeway CA Jr.: Innate immunity: the virtues of a nonclonal system of recognition. *Cell* 1997 Oct 31;91(3):295-8

Medzhitov R, Janeway CA Jr.: Innate immune recognition: mechanisms and pathways. *Immunol Rev* 2000 Feb;173:89-97

Medzhitov R, Janeway CA Jr.: Decoding the patterns of self and nonself by the innate immune system. *Science.* 2002 Apr 12;296(5566):298-300.

Nathan CF.: Secretory products of macrophages. *J Clin Invest.* 1987 Feb;79(2):319-26. Review.

Onishi HR, Pelak BA, Gerckens LS, Silver LL, Kahan FM, Chen MH, Patchett AA, Galloway SM, Hyland SA, Anderson MS, Raetz CR.: Antibacterial agents that inhibit lipid A biosynthesis. *Science* 1996 Nov 8;274(5289):980-2

Opal SM.: Interactions between coagulation and inflammation. *Scand J Infect Dis.* 2003;35(9):545-54. Review.

Park CT, Wright SD.: Fibrinogen is a component of a novel lipoprotein particle: factor H-related protein (FHRP)-associated lipoprotein particle (FALP). *Blood.* 2000 Jan 1;95(1):198-204.

Park CT, Wright SD.: Plasma lipopolysaccharide-binding protein is found associated with a particle containing apolipoprotein A-I, phospholipid, and factor H-related proteins. *J Biol Chem.* 1996 Jul 26;271(30):18054-60.

Poltorak A, He X, Smirnova I, Liu MY, Van Huffel C, Du X, Birdwell D, Alejos E, Silva M, Galanos C, Freudenberg M, Ricciardi-Castagnoli P, Layton B, Beutler B.: Defective LPS signaling in C3H/HeJ and C57BL/10ScCr mice: mutations in Tlr4 gene. *Science.* 1998 Dec 11;282(5396):2085-8

Pugin J, Heumann ID, Tomasz A, Kravchenko VV, Akamatsu Y, Nishijima M, Glauser MP, Tobias PS, Ulevitch RJ: CD14 is a pattern recognition receptor. *Immunity.* 1994 Sep;1(6):509-16.

Pugin J, Schurer-Maly CC, Leturcq D, Moriarty A, Ulevitch RJ, Tobias PS.: Lipopolysaccharide activation of human endothelial and epithelial cells is mediated by lipopolysaccharide-binding protein and soluble CD14. *Proc Natl Acad Sci U S A* 1993 Apr 1;90(7):2744-8

Ramadori G, Meyer zum Buschenfelde KH, Tobias PS, Mathison JC, Ulevitch RJ.: Biosynthesis of lipopolysaccharide-binding protein in rabbit hepatocytes. Pathobiology. 1990; 58(2): 89-94.

Remick DG.: Applied molecular biology of sepsis. J Crit Care. 1995 Dec;10(4):198-212. Review.

Rietschel ET, Brade H, Brade L, Kaca W, Kawahara K, Lindner B, Luderitz T, Tomita T, Schade U, Seydel U, et al.: Newer aspects of the chemical structure and biological activity of bacterial endotoxins. Prog Clin Biol Res 1985;189:31-51

Rietschel ET, Brade L, Brandenburg K, Flad HD, de Jong-Leuveninck J, Kawahara K, Lindner B, Loppnow H, Luderitz T, Schade U, et al.: Chemical structure and biologic activity of bacterial and synthetic lipid A. Rev Infect Dis. 1987 Sep-Oct; 9 Suppl 5: S527-36. Review.

Rietschel ET, Brade H, Holst O, Brade L, Muller-Loennies S, Mamat U, Zahringer U, Beckmann F, Seydel U, Brandenburg K, Ulmer AJ, Mattern T, Heine H, Schletter J, Loppnow H, Schonbeck U, Flad HD, Hauschildt S, Schade UF, Di Padova F, Kusumoto S, Schumann RR.: Bacterial endotoxin: Chemical constitution, biological recognition, host response, and immunological detoxification. Curr Top Microbiol Immunol. 1996;216:39-81. Review.

Roitt, I., Brostoff, J., Male, D. (Hrsg.): *Immunology*. 5.Auflage, Mosby, London 1998

Le Roy D, Di Padova F, Adachi Y, Glauser MP, Calandra T, Heumann D.: Critical role of lipopolysaccharide-binding protein and CD14 in immune responses against gram-negative bacteria. J Immunol. 2001 Sep 1;167(5):2759-65.

Sands KE, Bates DW, Lanken PN, Graman PS, Hibberd PL, Kahn KL, Parsonnet J, Panzer R, Orav EJ, Snyderman DR.: Epidemiology of sepsis syndrome in 8 academic medical centers. Academic Medical Center Consortium Sepsis Project Working Group. JAMA. 1997 Jul 16;278(3):234-40.

Schroemm, A.B., Brandenburg, K., Rietschel, E.Th., Flad, H.-D., Carroll, S.F., Seydel, U.: Lipopolysaccharide-binding protein mediates CD14-independent intercalation of lipopolysaccharide into phospholipid membranes. FEBS Letters 1996; 399: 267-271

Schroemm AB, Brandenburg K, Loppnow H, Moran AP, Koch MH, Rietschel ET, Seydel U.: Biological activities of lipopolysaccharides are determined by the shape of their lipid A portion. Eur J Biochem 2000 Apr; 267(7): 2008-13

Schumann RR, Belka C, Reuter D, Lamping N, Kirschning CJ, Weber JR, Pfeil D.: Lipopolysaccharide activates caspase-1 (interleukin-1-converting enzyme) in cultured monocytic and endothelial cells. Blood 1998 Jan 15;91(2):577-84

Schumann RR.: Function of lipopolysaccharide (LPS)-binding protein (LBP) and CD14, the receptor for LPS/LBP complexes: a short review. Res Immunol. 1992 Jan;143(1):11-5. Review.

Schumann RR, Kirschning CJ, Unbehauen A, Aberle HP, Knope HP, Lamping N, Ulevitch RJ, Herrmann F.: The lipopolysaccharide-binding protein is a secretory class 1 acute-phase protein whose gene is transcriptionally activated by APRF/STAT/3 and other cytokine-inducible nuclear proteins. Mol Cell Biol 1996 Jul;16(7):3490-503

Schumann RR, Lamping N, Hoess A.: Interchangeable endotoxin-binding domains in proteins with opposite lipopolysaccharide-dependent activities. J Immunol 1997 Dec 1;159(11):5599-605

Schumann RR, Latz E: Lipopolysaccharide-Binding Protein. In: Jack RS (ed): *CD14 in the Inflammatory Response*. Chem Immunol. Basel, Karger, 2000, vol 74, pp 42-60

Schumann RR, Leong SR, Flagg GW, Gray PW, Wright SD, Mathison JC, Tobias PS, Ulevitch RJ.: Structure and function of lipopolysaccharide binding protein. *Science*. 1990 Sep 21; 249(4975): 1429-31.

Schumann RR, Rietschel ET, Loppnow H.: The role of CD14 and lipopolysaccharide-binding protein (LBP) in the activation of different cell types by endotoxin. *Med Microbiol Immunol (Berl)*. 1994 Dec;183(6):279-97. Review.

Schumann RR, Zweigner J.: A novel acute-phase marker: lipopolysaccharide binding protein (LBP). *Clin Chem Lab Med*. 1999 Mar;37(3):271-4. Review.

Schütt C, Schumann R.: The endotoxin receptor CD14. *Immun Infekt* 1993 Apr;21(2):36-40

Shah, P.M.: Infektionskrankheiten. In: *Innere Medizin*. Greten, H. (Hrsg.), Thieme Verlag, 2001

Shimazu R, Akashi S, Ogata H, Nagai Y, Fukudome K, Miyake K, Kimoto M.: MD-2, a molecule that confers lipopolysaccharide responsiveness on Toll-like receptor 4. *Exp Med*. 1999 Jun 7;189(11):1777-82.

Silverman, M.H., Ostro, M.J.: Bacterial endotoxin in human disease. Xoma, Berkeley 1999

Somerville JE Jr, Cassiano L, Bainbridge B, Cunningham MD, Darveau RP.: A novel *Escherichia coli* lipid A mutant that produces an antiinflammatory lipopolysaccharide. *J Clin Invest.* 1996 Jan 15;97(2):359-65.

Stelter F, Witt S, Furl B, Jack RS, Hartung T, Schutt C: Different efficacy of soluble CD14 treatment in high- and low-dose LPS models. *Eur J Clin Invest.* 1998 Mar;28(3):205-13.

Sugiyama T, Wright SD.: Soluble CD14 mediates efflux of phospholipids from cells. *J Immunol.* 2001 Jan 15;166(2):826-31

Tapping RI, Akashi S, Miyake K, Godowski PJ, Tobias PS.: Toll-like receptor 4, but not toll-like receptor 2, is a signaling receptor for *Escherichia* and *Salmonella* lipopolysaccharides. *J Immunol.* 2000 Nov 15;165(10):5780-7.

Theofan G, Horwitz AH, Williams RE, Liu PS, Chan I, Birr C, Carroll SF, Meszaros K, Parent JB, Kasler H, et al.: An amino-terminal fragment of human lipopolysaccharide-binding protein retains lipid A binding but not CD14-stimulatory activity.. *J Immunol.* 1994 Apr 1; 152(7): 3623-9.

Tobias PS, Mathison J, Mintz D, Lee JD, Kravchenko V, Kato K, Pugin J, Ulevitch RJ.: Participation of lipopolysaccharide-binding protein in lipopolysaccharide-dependent macrophage activation. *Am J Respir Cell Mol Biol.* 1992 Sep; 7(3): 239-45. Review.

Tobias PS, Mathison JC, Ulevitch RJ.: A family of lipopolysaccharide binding proteins involved in responses to gram-negative sepsis. *J Biol Chem.* 1988 Sep 25;263(27):13479-81.

Tobias PS, Soldau K, Iovine NM, Elsbach P, Weiss J.: Lipopolysaccharide (LPS)-binding proteins BPI and LBP form different types of complexes with LPS. *J Biol Chem.* 1997 Jul 25; 272(30): 18682-5.

Tobias PS, Soldau K, Gegner JA, Mintz D, Ulevitch RJ.: Lipopolysaccharide binding protein-mediated complexation of lipopolysaccharide with soluble CD14.. *J Biol Chem.* 1995 May 5; 270(18): 10482-8.

Tobias, P.S., Soldau, K., Ulevitch, R.J.: Identification of a lipid binding site in the acute phase reactant lipopolysaccharide binding protein. *J Biol Chem* 1989; 264/18: 10867-10871

Tobias PS, Soldau K, Ulevitch RJ.: Isolation of a lipopolysaccharide-binding acute phase reactant from rabbit serum. *J Exp Med* 1986 Sep 1;164(3):777-93

Triantafilou M, Triantafilou K.: Lipopolysaccharide recognition: CD14, TLRs and the LPS-activation cluster. *Trends Immunol.* 2002 Jun;23(6):301-4. Review.

Ulevitch RJ, Tobias PS.: Recognition of gram-negative bacteria and endotoxin by the innate immune system. *Curr Opin Immunol* 1999 Feb;11(1):19-22

Vasselon T, Hailman E, Thieringer R, Detmers PA.: Internalization of monomeric lipopolysaccharide occurs after transfer out of cell surface CD14. *J Exp Med.* 1999 Aug 16;190(4):509-21.

Viriyakosol S, Kirkland TN: The N-terminal half of membrane CD14 is a functional cellular lipopolysaccharide receptor. *Infect Immun.* 1996 Feb;64(2):653-6.

Viriyakosol S, Tobias PS, Kitchens RL, Kirkland TN.: MD-2 binds to bacterial lipopolysaccharide. *J Biol Chem.* 2001 Oct 12;276(41):38044-51.

Vreugdenhil AC, Dentener MA, Snoek AM, Greve JW, Buurman WA.: Lipopolysaccharide binding protein and serum amyloid A secretion by human intestinal epithelial cells during the acute phase response. *J Immunol.* 1999 Sep 1; 163(5): 2792-8.

Wang PY, Kitchens RL, Munford RS.: Phosphatidylinositides bind to plasma membrane CD14 and can prevent monocyte activation by bacterial lipopolysaccharide. *J Biol Chem.* 1998 Sep 18; 273(38): 24309-13.

Weidemann B, Brade H, Rietschel ET, Dziarski R, Bazil V, Kusumoto S, Flad HD, Ulmer AJ.: Soluble peptidoglycan-induced monokine production can be blocked by anti-CD14 monoclonal antibodies and by lipid A partial structures. *Infect Immun.* 1994 Nov;62(11):4709-15.

Wright SD, Ramos RA, Tobias PS, Ulevitch RJ, Mathison JC.: CD14, a receptor for complexes of lipopolysaccharide (LPS) and LPS binding protein. *Science* 1990 Sep 21;249(4975):1431-3

Wright, S.D., Sugiyama, T.: Soluble CD14 mediates efflux of phospholipids from cells. *J Immunol* 2001; *166*: 826-831

Wright, S.D.: Innate recognition of microbial lipids. In: *Inflammation: Basic Principles and Clinical Correlates*. 3. Auflage, herausgegeben von John I. Gallin und Ralph Snyderman, Lippincott Williams & Wilkins, Philadelphia 1999

Wurfel MM, Hailman E, Wright SD.: Soluble CD14 acts as a shuttle in the neutralization of lipopolysaccharide (LPS) by LPS-binding protein and reconstituted high density lipoprotein. *J Exp Med.* 1995 May 1;181(5):1743-54.

Wurfel MM, Kunitake ST, Lichenstein H, Kane JP, Wright SD.: Lipopolysaccharide (LPS)-binding protein is carried on lipoproteins and acts as a cofactor in the neutralization of LPS. *J Exp Med.* 1994 Sep 1; 180(3): 1025-35.

Wurfel MM, Monks BG, Ingalls RR, Dedrick RL, Delude R, Zhou D, Lamping N, Schumann RR, Thieringer R, Fenton MJ, Wright SD, Golenbock D.: Targeted deletion of the lipopolysaccharide (LPS)-binding protein gene leads to profound suppression of LPS responses ex vivo, whereas in vivo responses remain intact. *J Exp Med* 1997 Dec 15; 186(12): 2051-6

Wurfel MM, Wright SD.: Lipopolysaccharide-binding protein and soluble CD14 transfer lipopolysaccharide to phospholipid bilayers: preferential interaction with particular classes of lipid. *J Immunol.* 1997 Apr 15;158(8):3925-34.

Yang KK, Dorner BG, Merkel U, Ryffel B, Schutt C, Golenbock D, Freeman MW, Jack RS.: Neutrophil Influx in Response to a Peritoneal Infection with Salmonella Is Delayed in Lipopolysaccharide-Binding Protein or CD14-Deficient Mice. *J Immunol.* 2002 Oct 15; 169(8): 4475-4480.

Yu, B., Hailman, E., Wright, S.D.: Lipopolysaccharide binding protein and soluble CD14 catalyze exchange of phospholipids. *J Clin Invest* 1997; 99/2: 315-324

Yu, B., Wright, S.D.: Catalytic Properties of lipopolysaccharide (LPS) binding protein. *J Biol Chem* 1996; 271/8: 4100-4105

Zweigner J, Gramm HJ, Singer OC, Wegscheider K, Schumann RR.: High concentrations of lipopolysaccharide-binding protein in serum of patients with severe sepsis or septic shock inhibit the lipopolysaccharide response in human monocytes. *Blood* 2001 Dec 15;98(13):3800-8