

6. REFERENCES

- Adler, A., J., Scheller, A., Robins, D., M. (1993). The stringency and magnitude of androgen-specific gene activation are combinatorial functions of receptor and nonreceptor binding site sequences. *Mol Cell Biol* **13**: 6326-6335
- Abate-Shen, C. (2002). Deregulated homeobox gene expression in cancer: cause or consequence? *Nature Cancer Reviews* **2**: 777-785
- Abdel-Hafiz, H., Takimoto, G. S., Tung, L., Horwitz, K. B. (2002). The inhibitory function in human progesterone receptor N termini binds SUMO-1 protein to regulate autoinhibition and transrepression. *J Biol Chem* **277**: 33950-33956
- Barbulescu, K., Geserick, C., Schüttke, I., Schleuning, W. D., Haendler, B. (2001). New androgen response elements in the murine Pem promoter mediate selective transactivation. *Mol Endocrinol* **15**: 1803-1816
- Bevan C., L., Hoare, S., Claessens, F., Heery, D., M., Parker, M., G. (1999). The AF1 and AF2 domains of the androgen receptor interact with distinct regions of SRC1. *Mol Cell Biol* **19**: 8383-8392
- Beverdam, A., Brouwer, A., Reijnen, M., Korving, J., Meijlink, F. (2001). Severe nasal clefting and abnormal embryonic apoptosis in *Alx3/Alx4* double mutant mice. *Development* **128**: 3975-3986
- Biggin, M., D. and McGinnis W. (1997). Regulation of segmentation and segmental identity of *Drosophila* homeoproteins: the role of DNA binding in functional activity and specificity. *Development* **124**: 4425-4433.
- Birk, O., S., Casiano, D., E., Wassif, C., A., Cogliati, T., Zhao, L., Zhao, Y., Grinberg, A., Huang, S., Kreidberg, J., A., Parker, K., L., Porter, F., D., Westphal, H. (2000). The LIM homeobox gene *Lhx9* is essential for mouse gonad formation. *Nature* **403**: 909-913
- Boulikas, T. (1994). Putative nuclear localization signals (NLS) in protein transcription factors. *J Cell Biochem* **55**: 32-58
- Branford, W., W., Zhao, G.-Q., Valerius, M., T., Weinstein, M., Birkenmeier, E., H., Rowe, L., B. and Potter, S., S. (1997). *Spx1*, a novel X-linked homeobox gene expressed during spermatogenesis. *Mech Dev* **65**: 87-98.
- Breathnach, R. and Chambon, P. (1981). Organization and expression of eucaryotic split genes coding for proteins. *Annu Rev Biochem* **50**: 349-383
- Celis, L., Claessens, F., Peeters, B., Heyns, W., Verhoeven, G., Rombauts, W. (1993). Proteins interacting with an androgen-responsive unit in the C3(1) gene intron. *Mol Cell Endocrinol* **94**: 165-72
- Chang, C., Saltzman, A., Yeh, S., Young, W., Keller, E., Lee, H., J., Wang, C., Mizokami, A. (1995). Androgen receptor: an overview. *Crit Rev Eukaryot Gene Expr* **5**: 97-125
- Chen, S., Wang, J., Yu, G., Liu, W. and Pearce, D. (1997). Androgen and glucocorticoid receptor heterodimer formation. A possible mechanism for mutual inhibition of transcriptional activity. *J Biol Chem* **272**: 14087-14092

- Claessens, F., Alen, P., Devos, A., Peeters, B., Verhoeven, G., Rombauts, W. (1996). The androgen-specific probasin response element 2 interacts differentially with androgen and glucocorticoid receptors. *J Biol Chem* **271**: 19013-19016
- Claessens, F., Verrijdt, G., Schoenmakers, E., Haelens, A., Peeters, B., Verhoeven, G., Rombauts W. (2001). Selective DNA binding by the androgen receptor as a mechanism for hormone-specific gene regulation. *J Steroid Biochem Mol Biol* **76**: 23-30
- Cleutjens C., B., Steketee, K., van Eekelen C., C., van der Korput J., A., Brinkmann A., O., Trapman J. (1997). Both androgen receptor and glucocorticoid receptor are able to induce prostate-specific antigen expression, but differ in their growth-stimulating properties of LNCaP cells. *Endocrinology* **138**: 5293-300.
- Cleutjens, C., B., van Eekelen, C., C., van der Korput H., A., Brinkmann A., O., Trapman J. (1996). Two androgen response regions cooperate in steroid hormone regulated activity of the prostate-specific antigen promoter. *J Biol Chem* **271**: 6379-6388
- Conaway, R., C., Brower, C., S., Conaway, J., W. (2002). Emerging roles of ubiquitin in transcription regulation. *Science* **296**: 1254 - 1258
- Couette, B., Le Ricousse, S., Fortin, D., Rafestin-Oblin, M., E., Richard-Foy, H. (1994). The establishment of the long terminal repeat of the mouse mammary tumor virus into CV-1 cells allows a functional analysis of steroid receptors. *Biochim Biophys Acta* **1219**: 607-612
- Courey, A., J. (2001). Cooperativity in transcriptional control. *Curr Biology* **10**: 250–252
- Cramer P. (2002). Multisubunit RNA polymerases. *Curr Opin Struct Biol* **12**: 89–97
- Dasen, J., S., Barbera, J., P., Herman, T., S., Connell, S., O., Olson, L., Ju, B., Tollkuhn, J., Baek, S., H., Rose, D., W., Rosenfeld, M., G. (2001). Temporal regulation of a paired-like homeodomain repressor/TLE corepressor complex and a related activator is required for pituitary organogenesis. *Genes Dev* **15**: 3193-3207
- Dotzlaw, H., Moehren, U., Mink, S., Cato, A.C., Iñiguez-Lluhí, J.A., Baniahmad, A. (2002). The amino terminus of the human AR is target for corepressor action and antihormone agonism. *Mol Endocrinol* **16**: 661-673
- Feng, W., Ribeiro, R, C., J., Wagner, R., L., Nguyen, H., Apriletti, J., W., Fletterick, R., J., Baxter, J., D., Kushner, P., J., West, B., L. (1998). Hormone-dependent coactivator binding to a hydrophobic cleft on nuclear receptors. *Science* **280**: 1747-1749
- Fey, Y. and Hugues, T., E. (2000). Nuclear trafficking of photoreceptor protein crx: the targeting sequence and pathologic implications. *Invest. Ophthalmol Vis Sci* **41**: 2849-2856
- Fohn, L., E. and Behringer, R., R. (2001). *ESX1L*, a novel X chromosome-linked human homeobox gene expressed in the placenta and testis. *Genomics* **74**: 105-108
- Freedman, L., P. (1992). Anatomy of the steroid receptor zinc finger region. *Endocrine Rev.* **13**: 129-145
- Freedman, L., P. (1999). Multimeric coactivator complexes for steroid/nuclear receptors. *Trends in Endocrinology and Metabolism* **10**: 403-407
- Fromental-Ramin, C., Warot, X., Lakkaraju, S., Favier, B., Haack, H., Birling, C., Dierich, A., Dollé, P., Chambon, P. (1996). Specific and redundant functions of the paralogous *Hoxa-9* and *Hoxd-9* genes in forelimb and axial skeleton patterning. *Development* **122**: 461-472

- Galliot, B., de Vargas, C., Miller, D. (1999). Evolution of homeobox genes: Q₅₀ paired-like genes founded the paired class. *Dev Genes Evol* **209**: 186-197
- Gehring, W., J., Affolter, M., Bürglin, T. (1994a). Homeodomain proteins. *Ann Rev Biochem* **63**: 487-526
- Gehring, W., J., Quian, Y., Q., Billeter, M., Furukubo-Tokunaga, K., Schier, A., F., Resendez-Perez, D., Affolter, M., Otting, G., Wüthrich, K. (1994b). Homeodomain-DNA recognition. *Cell* **78**: 211-223
- Gelmann, E., P. (2002). Molecular biology of the androgen receptor. *J Clinical Oncology* **20**: 3001-3015
- Gendron, R., L., Paradis, H., Hsieh-Li, H., M., Lee, D., W., Potter, S., S., Markoff, E. (1997). Abnormal uterine stromal and glandular function associated with maternal reproductive defects in *Hoxa11* null mice. *Biol Reprod* **56**: 1097-1105
- Gross, M., Liu, B., Tan, J., French, F., S., Carey, M., Shuai, K. (2001). Distinct effects of PIAS proteins on androgen-mediated gene activation in prostate cancer cells. *Oncogene* **20**: 3880-3887
- Haendler, B. (2002). Androgen-selective gene regulation in the prostate. *Biomed Pharmacother* **56**: 78-83
- Han Y., J., Lee, Y., H., Chun, J., Y. (2000). Identification and characterization of *Psx-2*, a novel member of the *Psx* (placenta-specific homeobox) family. *Gene* **241**: 149-155
- Han, Y., J., Park, A., R., Sung, D., Y. and Chun, J., Y. (1998). *Psx*, a novel murine homeobox gene expressed in placenta. *Gene* **207**: 159-166
- Hard, T., Dahlman, K., Carlstedt-Duke, J., Gustafsson, J., A., Rigler, R. (1990). Cooperativity and specificity in the interactions between DNA and the glucocorticoid receptor DNA-binding domain. *Biochemistry* **29**: 5358-5364
- He B., Kemppainen J. A., Voegel J. J., Gronemeyer H., Wilson E. M. (1999). Activation Function 2 in the Human Androgen Receptor Ligand Binding Domain Mediates Interdomain Communication with the NH₂-terminal Domain . *J Biol Chem* **274**: 37219-37225
- Heinemeyer, T., Wingender, E., Reuter, I., Hermjakob, H., Kel, A., E., Kel, O., V., Ignatieva, E., V., Ananko, E., A., Podkolodnaya, O., A., Kolpakov, F., A., Podkolodny, N., L., Kolchanov, N., A. (1998). Databases on transcriptional regulation: TRANSFAC, TRRD, and COMPEL. *Nucleic Acids Res* **26**: 362-367
- Heinlein, C., A. and Chang, C. (2002). The roles of androgen receptor and androgen-binding proteins in nongenomic androgen actions. *Endocrine Rev* **23**: 175-200
- Ho, K., C., Marschke, K., B., Tan, J., Power, S., G., Wilson, E., M., French, F., S. (1993). A complex response element in intron 1 of the androgen-regulated 20-kDa protein gene displays cell type-dependent androgen receptor specificity. *J Biol Chem* **268**: 27226-27235
- Hughes, I., A., Lim, H., N., Martin, H., Mongan, N., P., Dovey, L., Ahmed, S., F., Hawkins, J., R. (2001). Developmental aspect of androgen action. *Mol Cell Endo* **185**: 33-41
- Iñiguez-Lluhí, J., A. and Pearce, D. (2000). A common motif within the negative regulatory regions of multiple factors inhibits their transcriptional synergy. *Mol Cell Biol* **20**: 6040-6050

- Jans, D., A., Xiao, C., Y. and Lam, M., H., C. (2000). Nuclear targeting signal recognition: a key control point in nuclear transport? *BioEssays* **22**: 532-544
- Kaspar, F., Klocker, H., Denninger, A., and Cato, A. C. B. (1993). A mutant androgen receptor from patients with Reifenstein syndrome: identification of the function of a conserved alanine residue in the D box of steroid receptors. *Mol Cell Biol* **13**: 7850-7858
- Kasper, S., Rennie, P., S., Bruchovsky, N., Sheppard, P., C., Cheng, H., Lin, L., Shiu, R., P., Snoek, R., Matusik, R., J. (1994). Cooperative binding of androgen receptor to two DNA sequences is required for androgen induction of the probasin gene. *J Biol Chem* **269**: 31763-31769
- Kokontis, J., M. and Liao, S. (1999). Molecular action of androgen in the normal and neoplastic prostate. *Vitam Horm* **55**: 219-307
- Kotaja, N., Aittomäki, S., Silvennoinen, O., Palvimo, J. J., Jänne, O. A. (2000). ARIP3 (androgen receptor-interacting protein 3) and other PIAS (protein inhibitor of activated STAT) proteins differ in their ability to modulate steroid receptor-dependent transcriptional activation. *Mol Endocrinol* **14**: 1986-2000
- Kotaja, N., Karvonen, U., Jänne, O., A., Palvimo, J., J. (2002). PIAS proteins modulate transcription factors by functioning as SUMO-1 ligases. *Mol Cell Biol* **22**: 5222-5234
- Krauss, C., M., Turksoy, R., N., Atkins, L., McLaughlin, C., Brown, L., G., Page, D., C. (1987). Familial premature ovarian failure due to an interstitial deletion of the long arm of the X chromosome. *N England J Med* **317**: 125-131
- La Spada, A., R., Wilson E., M., Lubahn, D., B., Harding, A., E., Fishbeck, K., H. (1991). AR gene mutation in X-linked spinal and bulbar muscular atrophy. *Nature* **352**: 77-79
- Latchman, D., S. (1998). Eukaryotic transcription factors. 3rd edition. *Academic Press, San Diego*.
- Lefstin, J., A., and Yamamoto, K., R. (1998). Allosteric effects of DNA on transcriptional regulators. *Nature* **392**: 885-888
- Leo, C. and Chen, J., D., (2000). The SRC family of nuclear receptor coactivators. *Gene* **245**: 1-11
- Li, Y. and Behringer, R., R. (1998). *Esx1* is an X-chromosome-imprinted regulator of placental development and fetal growth. *Nature Gen* **20**: 309-311
- Li, Y., Lemaire, P. and Behringer, R., R. (1997). *Esx1*, a novel X chromosome-linked homeobox gene expressed in mouse extraembryonic tissues and male germ cells. *Dev Biol* **188**: 85-95
- Lin, H., Wang, L., Hu, Y., Altuwajiri, S., Chang, C. (2002). Phosphorylation-dependent ubiquitylation and degradation of androgen receptor by Akt require Mdm2 E3 ligase. *EMBO J* **21**: 4037-4048.
- Lindsey J., S., and Wilkinson, M., F. (1996). Pem: a testosterone- and LH-regulated homeobox gene expressed in mouse Sertoli cells and epididymis. *Dev Biol* **179**: 471-484
- List, H., J., Lozano, C., Lu, J., Danielsen, M., Wellstein, A., Riegel, A., T. (1999). Comparison of chromatin remodeling and transcriptional activation of the mouse mammary tumor virus promoter by the androgen and glucocorticoid receptor. *Exp Cell Res* **250**: 414-22

- Liu, W., Wang, J., Yu, G., Pearce, D. (1996). Steroid receptor transcriptional synergy is potentiated by disruption of the DNA-binding domain dimer interface. *Mol Endocrinol* **10**: 1399-1406
- Lo, K. and Smale, S., T. (1996). Generality of a functional initiator consensus sequence. *Gene* **182**: 13-22
- Loven, M., A., Likhite, V., S., Choi, I., Nardulli, A., M. (2001). Estrogen response elements alter coactivator recruitment through allosteric modulation of estrogen receptor beta conformation. *J Biol Chem* **276**: 45282-45288
- Lu, S., Jenster, G., Epner, D., E. (2000). Androgen induction of cyclin-dependent kinase inhibitor p21 gene: role of androgen receptor and transcription factor Sp1 complex. *Mol Endocrinol* **14**: 753-60
- Lucas, P., C., and Granner, D., K. (1992). Hormone response domains in gene transcription. *Annu Rev Biochem* **61**: 1131-1173
- Luisi, B., F., Xu, W., X., Otwinowski, Z., Freedman, L., P., Yamamoto, K., R., Sigler, P., B. (1991). Crystallographic analysis of the interaction of the glucocorticoid receptor with DNA. *Nature* **352**: 497-505
- MacLean, H., E., Warne, G., L., Zajac, J., D. (1997). Localization of functional domains in the androgen receptor. *J Steroid Biochem Mol Biol* **62**: 233-242;
- Maiti, S., Doskow, J., Li, S., Nhim, R., P., Lindsey, J., S., Wilkinson, M., F. (1996a). The *Pem* homeobox gene. Androgen-dependent and -independent promoters and tissue-specific alternative RNA splicing. *J Biol Chem* **271**: 17536-17546
- Maiti, S., Doskow, J., Sutton, K., Nhim, R., P., Lawlor, D., A., Levan, K., Lindsey, J., S., Wilkinson, M., F. (1996b). The *Pem* homeobox gene: rapid evolution of the homeodomain, X chromosomal localization, and expression in reproductive tissues. *Genomics* **34**: 304-316
- Maiti, S., Meistrich, M., L., Wilson, G., Shetty, G., Marcelli, M., McPhaul, M., J., Morris, P., L. and Wilkinson, M., F. (2001). Irradiation selectively inhibits expression from the androgen-dependent *Pem* homeobox gene promoter in Sertoli cells. *Endocrinology* **142**: 1567-1577
- Mangelsdorf D., J., Thummel C., Beato M. et al. (1995). The nuclear receptor family: The second decade. *Cell* **83**: 835-839
- Mann, R., S. (1995). The specificity of homeotic gene function. *Bio Essays* **17**: 855-863.
- Matias, P., M., Carrondo M., A., Coelho, R., Thomaz, M., Zhao, X., Y., Wegg, A., Crusius, K., Egner, U., Donner, P. (2002). Structural basis for the glucocorticoid response in a mutant human androgen receptor (AR(ccr)) derived from an androgen-independent prostate cancer. *J Med Chem* **45**:1 439-1446
- Matias, P., M., Donner, P., Coelho R., Thomaz, M., Peixoto, C., Macedo, S., Otto, N., Joschko, S., Scholz, P., Wegg, A., Bäsler, S., Schäfer, M., Egner, U., Carrondo, M., A. (2000). Structural evidence for ligand specificity in the binding domain of the human androgen receptor. *J Biol Chem* **275**: 26164-26171
- McDonnell, D., P. and Norris, J., D. (2002). Connections and regulation of the human estrogen receptor. *Science* **296**: 1642-1644

- Miller, D., J., Hayward, D., C., Reece-Hoyes, J., S., Scholten, I., Catmull, J., Gehring, W., J., Callaerts, P., Larsen, J., E., Ball, E., E. (2000). *Pax* gene diversity in the basal cnidarian *Acropora millepora* (Cnidaria, Anthozoa): implications for the evolution of the *Pax* gene family. *Proc Natl Acad Sci U S A* **97**: 4475-4480
- Moede, T., Leibiger, B., Pour, H., G., Berggren, P.-O., Leibiger, I., B. (1999). Identification of a nuclear localization signal, RRMKWKK, in the homeodomain transcription factor PDX-1. *FEBS Lett* **461**: 229-234
- Muller, J., M., Isele, U., Metzger, E., Rempel, A., Moser, M., Pscherer, A., Breyer, T., Holubarsch, C., Buettner, R., Schüle, R. (2000). FHL2, a novel tissue-specific coactivator of the androgen receptor. *EMBO J* **19**:359-69
- Nelson, C., C., Hendy, S., C., Shukin, R., J., Cheng, H., Bruchovsky, N., Koop, B., F., Rennie, P., S. (1999). Determinants of DNA sequence specificity of the androgen, progesterone, and glucocorticoid receptors: evidence for differential steroid receptor response elements. *Mol Endocrinol* **13**: 2090-2107
- Nelson, S., A., and Robins, D., M. (1997). Regulatory capacity of an androgen-specific enhancer of the mouse *Slp* gene in transgenic mice. *Mol Cell Endocrinol* **133**:89-97
- Nishida, T. and Yasuda, H. (2002). PIAS1 and PIASx function as SUMO-E3 ligases toward androgen receptor, and repress androgen receptor-dependent transcription. *J Biol Chem* **277**: 41311-41317
- Norris, R., A. and Kern, M., J. (2001). Identification of domains mediating transcription activation, repression, and inhibition in the paired-related homeobox protein, Prx2 (S8). *DNA Cell Biol* **20**: 89-99
- Nussinov, R. (1990). Sequence signals in eukaryotic upstream regions. *Crit Rev Biochem Mol Biol* **25**: 185-224
- Ostendorff, H., P., Peirano, R., I., Peters, M., A., Schluter, A., Bossenz, M., Scheffner, M., Bach, I. (2002). Ubiquitination-dependent cofactor exchange on LIM homeodomain transcription factors. *Nature* **416**: 99–103
- Pinsonneault, J., Florence, B., Vaessin, H., McGinnis, W. (1997). A model for extradenticle function as a switch that changes HOX proteins from repressor to activators. *EMBO J* **16**: 2032-2042
- Pitman, J., L., Lin, T., P., Kleeman, J., E., Erickson, G., F., McLeod, C., L. (1998). Normal reproductive and macrophage function in *Pem* homeobox-deficient mice. *Dev Biol* **202**: 196-214
- Poukka, H., Aarnisalo, P., Karvonen, U., Palvimo, J., J., Jänne, O., A. (1999). Ubc9 interacts with the androgen receptor and activates receptor-dependent transcription. *J Biol Chem* **274**: 19441-19446
- Poukka, H., Karvonen, U., Jänne, O., A. and Palvimo, J., J. (2000). Covalent modification of the androgen receptor by small ubiquitin-like modifier 1 (SUMO-1). *Proc Natl Acad Sci U S A* **97**: 14145-14150
- Quarmby, V., E., Yarbrough, W., G., Lubahn, D., B., French, F., S., Wilson, E., M. (1990). Autologous down-regulation of androgen receptor messenger ribonucleic acid. *Mol Endocrinol* **4**: 22-28

- Rao, M., K., Wayne, C., M., Wilkinson, M., F. (2002). Pem homeobox gene regulatory sequences that direct androgen-dependent developmentally regulated gene expression in different subregions of the epididymis. *J Biol Chem* **277**: 48771-48778
- Rao, E., Weiss, B., Fukami, M., Rump, A., Niesler, B., Mertz, A., Muroya, K., Binder, G., Kirsch, S., Winkelmann, M., Nordsiek, G., Heinrich, U., Breuning, M., H., Ranke, M., B., Rosenthal, A., Ogata, T., Rappold, G., A. (1997). Pseudoautosomal deletions encompassing a novel homeobox gene cause growth failure in idiopathic short stature and Turner syndrome. *Nat Genet* **16**: 54-62
- Reichardt, M., H., Kaestner, K., H., Tuckermann J., Kretz, O., Wessely, O., Bock, R., Gass, P., Schmid, W., Herrlich, P., Angel, P., Schütz, G. (1998). DNA binding of the glucocorticoid receptor is not essential for survival. *Cell* **93**: 531 –541
- Rennie, P., S., Bruchofsky, N., Leco, K. J., Sheppard, P., C., McQueen, S., A., Cheng, H., Snoek, R., Hamel, A., Bock, M. E., MacDonald, B., S., Nickel, B., E., Chang, C., Liao, S., Cattinin, P., A., and Matusik, R., J. (1993). Characterization of two cis-acting DNA elements involved in the androgen regulation of the probasin gene. *Mol Endocrinol* **7**: 23-36
- Rini B., I., Small E., J. (2002). Prostate Cancer Update. *Curr Opin Oncol* **14**: 286 – 291
- Roche, P., J., Hoare, S., A., Parker, M., G. (1992). A consensus DNA-binding site for the androgen receptor. *Mol Endocrinol* **6**: 2229-2235
- Rundlett S., E., Miesfeld, R., L. (1995). Quantitative differences in androgen and glucocorticoid receptor DNA binding properties contribute to receptor-selective transcriptional regulation. *Mol Cell Endocrinol* **109**: 1-10
- Sack, J., S., Kish K., F., Wang, C., Attar, R., M., Kiefer, S., E., An, Y., Wu, G., Y., Scheffler, J., E., Salvati, M., E., Krystek, S., R., Jr, Weinmann R., Einspahr, H., M. (2001). Crystallographic structures of the ligand-binding domains of the androgen receptor and its T877A mutant complexed with the natural agonist dihydrotestosterone. *Proc Natl Acad Sci USA* **98**: 4904 - 4909
- Sambrook, J., Fritsch, E., Maniatis, T. (1989). Molecular Cloning, A laboratory manual. *Cold Spring Harbour Laboratory Press, Cold Spring Harbor, New York*.
- Sasaki, A. W., Doskow, J., MacLeod, C. L., Rogers, M. B., Gudas, L. J. and Wilkinson, M. F. (1991). The oncofetal gene *Pem* encodes a homeobox and is regulated in primordial and pre-muscle stem cells. *Mech Dev* **34**: 155-164
- Satokata, I., Benson, G., Maas, R. (1995). Sexually dimorphic sterility phenotypes in *Hoxa10*-deficient mice. *Nature* **374**: 460-463
- Scheller, A., Hughes, E., Golden, K., L., Robins, D., M. (1998). Multiple receptor domains interact to permit, or restrict, androgen-specific gene activation. *J Biol Chem* **273**: 24216-24222
- Schoenmakers, E., Alen, P., Verrijdt, G., Peeters, B., Verhoeven, G., Rombauts, W., Claessens, F. (1999). Differential DNA binding by the androgen and glucocorticoid receptors involves the second Zn-finger and a C-terminal extension of the DNA-binding domains. *Biochem J* **341**: 515-521
- Schwabe, J., W., Chapman, L. and Rhodes, D. (1995). The oestrogen receptor recognizes an imperfectly palindromic response element through an alternative side-chain conformation. *Structure* **3**: 201-213

- Schwidetzky, U., Schleuning, W., D., Haendler, B. (1997). Isolation and characterization of the androgen-dependent mouse cysteine-rich secretory protein-1 (CRISP-1) gene. *Biochem J* **321**: 325-332
- Shan, L., X., Rodriguez, M., C., Janne, O., A. (1990). Regulation of androgen receptor protein and mRNA concentrations by androgens in rat ventral prostate and seminal vesicles and in human hepatoma cells. *Mol Endocrinol* **11**: 1636-46
- Shiau, A., K., Bastard, D., Loria, P., M., Cheng, L., Kushner, P., J., Agard, D., A., Green, G., L. (1998). The structural basis of estrogen receptor/coactivator recognition and the antagonism of this interaction by tamoxifen. *Cell* **95**: 927-937
- Simental, J., A., Sar, M., Lane, M., V., French, F., S., Wilson, E., M. (1991). Transcriptional activation and nuclear targeting signals of the human androgen receptor. *J Biol Chem* **266**: 510-518.
- Smale, S., T. (1997). Transcriptional initiation from TATA-less promoters within eukaryotic protein-coding genes. *Biochim Biophys Acta* **1351**: 73-88
- Sutton, K., A., Maiti, S., Tribley, W., A., Lindsey, J., S., Meistrich, M., L., Bucana, C., D., Sanborn, B., M., Joseph, D., R., Griswold, M., D., Cornwall, G., A., Wilkinson, M., F. (1998). Androgen regulation of the Pcm homeodomain gene in mice and rat Sertoli and epididymal cells. *J Androl* **19**: 21-30
- Swanson, W., J., Clark, A., G., Waldrip-Dail, H., M., Wolfner, M., F., Aquadro, C., F. (2001). Evolutionary EST analysis identifies rapidly evolving male reproductive proteins in *Drosophila*. *Proc Natl Acad Sci USA* **98**: 7375-7379
- Szapary, D., Xu, M., Simon, S., S. Jr. (1996). Induction properties of a transiently transfected glucocorticoid-responsive gene vary with glucocorticoid receptor concentration. *J Biol Chem* **271**: 30576-30582
- Takasaki, N., Mclsaac, R., Dean, J. (2000). *Gpbox (Psx2)*, a homeobox gene preferentially expressed in female germ cells at the onset of sexual dimorphism in mice. *Dev. Biol.* **223**: 181-193
- Takasaki, N., Rankin, T. and Dean, J. (2001). Normal gonadal development in mice lacking GPBOX, a homeobox protein expressed in germ cells at the onset of sexual dimorphism. *Mol Cell Biol* **21**: 8197-8202
- Takehita, A., Yen, P., M., Ikeda, M., Cardona, G., R., Liu, Y., Koibuchi, N., Norwitz, E., R., Chin, W., W. (1998). Thyroid hormone response elements differentially modulate the interactions of thyroid hormone receptors with two receptor binding domains in the steroid receptor coactivator-1. *J Biol Chem* **273**: 21554-21562
- Tanaka, M., Schinke, M., Liao, H., S., Yamasaki, N., Izumo, S. (2001). Nkx2.5 and Nkx2.6, homologs of drosophila tinman, are required for development of the pharynx. *Mol Cell Biol* **21**: 4391-4398
- Tian, S., Poukka, H., Palvimo, J., J., Jänne, O., A. (2002). SUMO-1 modification of the glucocorticoid receptor. *Biochem J* **367**: 907-911
- Torres, M., Gomez-Pardo, E., Dressler, G. R., Gruss, P. (1995). Pax-2 controls multiple steps of urogenital development. *Development* **121**: 4057-4065

- Treisman, J., Harris, E., Wilson, D., Desplan, C. (1992). The homeodomain: a new face for the helix-turn-helix? *BioEssays* **14**: 145-150
- Tsai, M., J., and O'Malley, B., W. (1994). Molecular mechanisms of action of steroid/thyroid receptor superfamily members. *Ann Rev Biochem* **63**: 451-486
- Verrijdt, G., Schauwaers, K., Haelens, A., Rombauts, W., Claessens, F. (2002). Functional interplay between two response elements with distinct binding characteristics dictates androgen specificity of the mouse sex-limited protein enhancer *J Biol Chem* **227**: 35191-35201
- Verrijdt, G., Schoenmakers, E., Alen, P., Haelens, A., Peeters, B., Rombauts, W., Claessens, F. (1999). Androgen specificity of a response unit upstream of the human secretory component gene is mediated by differential receptor binding to an essential androgen response element. *Mol Endocrinol* **13**:1558-1570
- Visakorpi, T., Hyytinen, E., Koivisto, P., Tanner, M., Keinanen, R., Palmberg, C., Palotie, A., Tammela, T., Isola, J., Kallioniemi, O., P. (1995). In vivo amplification of the androgen receptor gene and progression of human prostate cancer. *Nat Genet* **9**: 401-406
- Wilson, T., M. and Moore, J., T. (2002). Genomics versus orphan nuclear receptors – A half-time report. *Mol Endo* **16**: 1135-1144
- Wimmer, Z., Zhu, X., X., Rouillard, J., M., Ambros, P., F., Lamb, B. J., Kuick, R., Eckart, M., Weinhausl, A., Fonatsch, C. and Hanash, S. M. (2002). Combined restriction landmark genomic scanning and virtual genome scans identify a novel human homeobox gene, ALX3, that is hypermethylated in neuroblastoma. *Genes Chromosomes Cancer* **33**: 285-294
- Wood, J., R., Likhite, V., S., Loven, M., A., Nardulli, A., M. (2001). Allosteric modulation of estrogen receptor conformation by different estrogen response elements. *Mol Endocrinol* **15**: 1114-1126
- Woychik, N., A., and Hampsey, M. (2002). The RNA polymerase II machinery: structure illuminates function. *Cell* **108**: 453 - 463
- Wuyts, W., Cleiren, E., Homfray, T., Rasore-Quartino, A., Vanhoenacker, F., Van Hul, W. (2000). The ALX4 homeobox gene is mutated in patients with ossification defects of the skull (foramina parietalia permagna, OMIM 168500). *J Med Gene* **37**: 916-920
- Xu, L., C., Thali, M., Schaffner, W. (1991). Upstream box/TATA box order is the major determinant of the direction of transcription. *Nucleic Acids Res* **19** : 6699-6704
- Yan, Y.-T., Stein, S., M., Ding, J., Shen, M., M., Abate-Shen, C. (2000). A novel PF/PN motif inhibits nuclear localization and DNA binding activity of the ESX1 homeoprotein. *Mol Cell Biol* **20**: 661-671
- Yeh, S. and Chang, C. (1996). Cloning and characterization of a specific coactivator, ARA70, for the androgen receptor in human prostate cells. *Proc Natl Acad Sci U S A* **93**: 5517-21
- Yeh, S., Kang, H., Y., Miyamoto, H., Nishimura, K., Chang, H., C., Ting, H., J., Rahman, M., Lin, H., K., Fujimoto, N., Hu, Y., C., Mizokami, A., Huang, K., E., Chang C. (1999). Differential induction of androgen receptor transactivation by different androgen receptor coactivators in human prostate cancer DU145 cells. *Endocrine* **11**: 195–202

Zellmer, E., Zhang, Z., Greco, D., Rhodes, J., Cassel, S., Lewis, E. J. (1995). A homeodomain protein selectively expressed in noradrenergic tissue regulates transcription of neurotransmitter biosynthetic genes. *J Neurosci* **15**: 8109-8120

Zhao, X., Y., Malloy, P., J., Krishnan, A., V., Swami, S., Navone N., M., Peehl, D., M., Feldman, D. (2000). Glucocorticoids can promote androgen-independent growth of prostate cancer cells through a mutated androgen receptor. *Nat Med* **6**: 703-706