

References

- Agler, H. L., Evans, J., Colecraft, H. M., & Yue, D. T. (2003). Custom distinctions in the interaction of G-protein β subunits with N-type ($\text{Ca}_V2.2$) versus P/Q-type ($\text{Ca}_V2.1$) calcium channels. *J. Gen. Physiol.*, **121**, 495–510.
- Ahn, J.-Y., Rong, R., Liu, X., & Ye, K. (2004). PIKE/nuclear PI 3-kinase signaling mediates the antiapoptotic actions of NGF in the nucleus. *EMBO J.*, **23**, 3995–4006.
- Alessi, D. R., Andjelkovic, M., Caudwell, B., Cron, P., Morrice, N., Cohen, P., & Hemmings, B. A. (1996). Mechanism of activation of protein kinase B by insulin and IGF-1. *EMBO J.*, **15**, 6541–6551.
- Ali, K., Bilancio, A., Thomas, M., Pearce, W., Gilfillan, A. M., Tkaczyk, C., Kuehn, N., Gray, A., Giddings, J., Peskett, E., Fox, R., Bruce, I., Walker, C., Sawyer, C., Okkenhaug, K., Finan, P., & Vanhaesebroeck, B. (2004). Essential role for the p110 δ phosphoinositide 3-kinase in the allergic response. *Nature*, **431**, 1007–1011.
- Allam, A. & Marshall, A. J. (2005). Role of the adaptor proteins Bam32, TAPP1 and TAPP2 in lymphocyte activation. *Immunol. Lett.*, **97**, 7–17.
- Altschul, S. F., Gish, W., Miller, W., Myers, E. W., & Lipman, D. J. (1990). Basic local alignment search tool. *J. Mol. Biol.*, **215**, 403–410.
- Anderson, K. E., Lipp, P., Bootman, M., Ridley, S. H., Coadwell, J., Rönstrand, L., Lennartsson, J., Holmes, A. B., Painter, G. F., Thuring, J., Lim, Z.-Y., Erdjument-Bromage, H., Grewal, A., Tempst, P., Stephens, L. R., & Hawkins, P. T. (2000). DAPP1 undergoes a PI 3-kinase-dependent cycle of plasma-membrane recruitment and endocytosis upon cell stimulation. *Curr. Biol.*, **10**, 1403–1412.
- Arcaro, A. & Wymann, M. P. (1993). Wortmannin is a potent phosphatidylinositol 3-kinase inhibitor: the role of phosphatidylinositol 3,4,5-triphosphate in neutrophil responses. *Biophys. J.*, **296**, 297–301.
- Arcaro, A., Zvelebil, M. J., Wallasch, C., Ullrich, A., Waterfield, M. D., & Domin, J. (2000). Class II phosphoinositide 3-kinases are downstream targets of activated polypeptide growth factor receptors. *Mol. Cell. Biol.*, **20**, 3817–3830.

REFERENCES

- Arkin, M. R. & Wells, J. A. (2004). Small-molecule inhibitors of protein–protein interactions: Progressing towards the dream. *Nat. Rev. Drug Disc.*, **3**, 301–317.
- Bacqueville, D., Déléris, P., Mendre, C., Pieraggi, M.-T., Chap, H., Guillon, G., Perret, B., & Breton-Douillon, M. (2001). Characterization of a G protein-activated phosphoinositide 3-kinase in vascular smooth muscle cell nuclei. *J. Biol. Chem.*, **276**, 22170–22176.
- Baier, R., Bondeva, T., Klinger, R., Bondev, A., & Wetzker, R. (1999). Retinoic acid induces selective expression of phosphoinositide 3-kinase γ in myelomonocytic U937 cells. *Cell. Growth Differ.*, **10**, 447–456.
- Beeton, C. A., Chance, E. M., Foukas, L. C., & Shepherd, P. R. (2000). Comparison of the kinetic properties of the lipid- and protein-kinase activities of the p110 α and p110 β catalytic subunits of class-Ia phosphoinositide 3-kinases. *Biochem. J.*, **350**, 353–359.
- Bers, D. M. (2002). Cardiac excitation–contraction coupling. *Nature*, **415**, 198–205.
- Bhaskar, P. T. & Hay, N. (2007). The two TORCs and Akt. *Dev. Cell*, **12**, 487–502.
- Biondi, R. M., Kieloch, A., A. Currie, R., Deak, M., & Alessi, D. R. (2001). The PIF-binding pocket in PDK1 is essential for activation of S6K and SGK, but not PKB. *EMBO J.*, **20**, 4380–4390.
- Blomberg, N., Baraldi, E., Nilges, M., & Saraste, M. (1999). The PH superfold: a structural scaffold for multiple functions. *Trends Biochem. Sci.*, **24**, 441–445.
- Bogan, A. A. & Thorn, K. S. (1998). Anatomy of hot spots in protein interfaces. *J. Mol. Biol.*, **280**, 1–9.
- Bokoch, G. M., Vlahos, C. J., Wang, Y., Knaus, U. G., & Traynor-Kaplan, A. E. (1996). Rac GTPase interacts specifically with phosphatidylinositol 3-kinase. *Biochem. J.*, **315**, 775–779.
- Bonacci, T. M., Mathews, J. L., Yuan, C., Lehmann, D. M., Malik, S., Wu, D., Font, J. L., Bidlack, J. M., & Smrcka, A. V. (2006). Differential targeting of G $\beta\gamma$ -subunit signaling with small molecules. *Science*, **312**, 443–446.
- Bondev, A., Rubio, I., & Wetzker, R. (1999). Differential regulation of lipid and protein kinase activities of phosphoinositide 3-kinase γ *in vitro*. *Biol. Chem.*, **380**, 1337–1340.
- Bondeva, T., Pirola, L., Bulgarelli-Leva, G., Rubio, I., Wetzker, R., & Wymann, M. P. (1998). Bifurcation of lipid and protein kinase signals of PI3K γ to the protein kinases PKB and MAPK. *Science*, **282**, 293–296.
- Boylan, J. M. & Gruppuso, P. A. (2002). Insulin receptor substrate-1 is present in hepatocyte nuclei from intact rats. *Endocrinology*, **143**, 4178–4183.

- Brock, C. (2002). *Untersuchungen zur Regulation von Klasse I Phosphoinositid-3-Kinasen in vivo*. Inaugural-Dissertation, Fachbereich Biologie, Chemie, Pharmazie, Freie Universität Berlin.
- Brock, C., Schaefer, M., Reusch, H. P., Czupalla, C., Michalke, M., Spicher, K., Schultz, G., & Nürnberg, B. (2003). Roles of G $\beta\gamma$ in membrane recruitment and activation of p110 γ /p101 phosphoinositide 3-kinase γ . *J. Cell Biol.*, **160**, 89–99.
- Brown, R. A., Domin, J., Arcaro, A., Waterfield, M. D., & Shepherd, P. R. (1999). Insulin activates the α isoform of class II phosphoinositide 3-kinase. *J. Biol. Chem.*, **274**, 14529–14532.
- Brugnera, E., Haney, L., Grimsley, C., Lu, M., Walk, S. F., Tosello-Trampont, A.-C., Macara, I. G., Madhani, H., Fink, G. R., & Ravichandran, K. S. (2002). Unconventional Rac-GEF activity is mediated through the Dock180–ELMO complex. *Nat. Cell Biol.*, **4**, 574–582.
- Brummelkamp, T. R., Bernards, R., & Agami, R. (2002). A system for stable expression of short interfering RNAs in mammalian cells. *Science*, **296**, 550–553.
- Brunet, A., Bonni, A., Zigmond, M. J., Lin, M. Z., Juo, P., Hu, L. S., Anderson, M. J., Arden, K. C., Blenis, J., & Greenberg, M. E. (1999). Akt promotes cell survival by phosphorylating and inhibiting a forkhead transcription factor. *Cell*, **96**, 857–868.
- Burgering, B. M. & Kops, G. J. (2002). Cell cycle and death control: Long live forkheads. *Trends Biochem. Sci.*, **27**, 352–360.
- Byfield, M. P., Murray, J. T., & Backer, J. M. (2005). hVps34 is a nutrient-regulated lipid kinase required for activation of p70 S6 kinase. *J. Biol. Chem.*, **280**, 33076–33082.
- Cabrera-Vera, T. M., Vanhauwe, J., Thomas, T. O., Medkova, M., Preininger, A., Mazzoni, M. R., & Hamm, H. E. (2003). Insights into G protein structure, function, and regulation. *Endocr. Rev.*, **24**, 765–781.
- Camps, M., Rückle, T., Ji, H., Ardisson, V., Rintelen, F., Shaw, J., Ferrandi, C., Chabert, C., Gillieron, C., Francon, B., Martin, T., Gretener, D., Perrin, D., Leroy, D., Vitte, P. A., Hirsch, E., Wymann, M. P., Cirillo, R., Schwarz, M. K., & Rommel, C. (2005). Blockade of PI3K γ suppresses joint inflammation and damage in mouse models of rheumatoid arthritis. *Nat. Med.*, **11**, 936–943.
- Cantor, C. R. & Schimmel, P. R. (1980). *Biophysical Chemistry, Part II: Techniques for the Study of Biological Structure and Function*. W. H. Freeman and Company, San Francisco.
- Cardone, M. H., Roy, N., Stennicke, H. R., Salvesen, G. S., Franke, T. F., Stanbridge, E., Frisch, S., & Reed, J. C. (1998). Regulation of cell death protease caspase-9 by phosphorylation. *Science*, **282**, 1318–1321.

REFERENCES

- Carnevale, K. A. & Cathcart, M. K. (2001). Calcium-independent phospholipase A₂ is required for human monocyte chemotaxis to monocyte chemoattractant protein 1. *J. Immunol.*, **167**, 3414–3421.
- Casamayor, A., Morrice, N. A., & Alessi, D. R. (1999). Phosphorylation of Ser-241 is essential for the activity of 3-phosphoinositide-dependent protein kinase-1: identification of five sites of phosphorylation *in vivo*. *Biochem. J.*, **342**, 287–292.
- Chardin, P., Paris, S., Antonny, B., Robineau, S., Béraud-Dufour, S., Jackson, C. L., & Chabre, M. (1996). A human exchange factor for ARF contains Sec7- and pleckstrin-homology domains. *Nature*, **384**, 481–484.
- Charest, P. G. & Firtel, R. A. (2006). Feedback signaling controls leading-edge formation during chemotaxis. *Curr. Opin. Genet. Dev.*, **16**, 339–347.
- Charest, P. G. & Firtel, R. A. (2007). Big roles for small GTPases in the control of directed cell movement. *Biochem. J.*, **401**, 377–390.
- Chen, J., DeVivo, M., Dingus, J., Harry, A., Li, J., Sui, J., Carty, D. J., Blank, J. L., Exton, J. H., Stoffel, R. H., Inglese, J., Lefkowitz, R. J., Logothetis, D. E., Hildebrandt, J. D., & Iyengar, R. (1995). A region of adenylyl cyclase 2 critical for regulation by G protein $\beta\gamma$ subunits. *Science*, **268**, 1166–1169.
- Chen, L., Iijima, M., Tang, M., Landree, M. A., Huang, Y. E., Xiong, Y., Iglesias, P. A., & Devreotes, P. N. (2007). PLA₂ and PI3K/PTEN pathways act in parallel to mediate chemotaxis. *Dev. Cell*, **12**, 603–614.
- Chen, Q., Powell, D. W., Rane, M. J., Singh, S., Butt, W., Klein, J. B., & McLeish, K. R. (2003). Akt phosphorylates p47^{phox} and mediates respiratory burst activity in human neutrophils. *J. Immunol.*, **170**, 5302–5308.
- Choi, Y. H., Park, S., Hockman, S., Zmuda-Trzebiatowska, E., Svennelid, F., Haluzik, M., Gavrilova, O., Ahmad, F., Pepin, L., Napolitano, M., Taira, M., Sundler, F., Holst, L. S., Degerman, E., & Manganiello, V. C. (2006). Alterations in regulation of energy homeostasis in cyclic nucleotide phosphodiesterase 3B-null mice. *J. Clin. Invest.*, **116**, 3240–3251.
- Christoforidis, S., Miaczynska, M., Ashman, K., Wilm, M., Zhao, L., Yip, S.-C., Waterfield, M. D., Backer, J. M., & Zerial, M. (1999). Phosphatidylinositol-3-OH kinases are Rab5 effectors. *Nat. Cell Biol.*, **1**, 249–252.
- Clackson, T. & Wells, J. A. (1995). A hot spot of binding energy in a hormone-receptor interface. *Science*, **267**, 383–386.

- Clapham, D. & Neer, E. (1999). G protein $\beta\gamma$ subunits. *Annu. Rev. Pharmacol. Toxicol.*, **37**, 167–203.
- Claycomb, W. C., Lanson, Jr., N. A., Stallworth, B. S., Egeland, D. B., Delcarpio, J. B., Bahinsky, A., & Izzo, Jr., N. J. (1998). HL-1 cells: A cardiac muscle cell line that contracts and retains phenotypic characteristics of the adult cardiomyocyte. *Proc. Natl. Acad. Sci. U.S.A.*, **95**, 2979–2984.
- Crackower, M. A., Oudit, G. Y., Kozieradzki, I., Sarao, R., Sun, H., Sasaki, T., Hirsch, E., Suzuki, A., Shioi, T., Irie-Sasaki, J., Sah, R., Cheng, H.-Y. M., Rybin, V. O., Lembo, G., Fratta, L., Oliveira-dos-Santos, A. J., Benovic, J. L., Kahn, C. R., Izumo, S., Steinberg, S. F., Wymann, M. P., Backx, P. H., & Penninger, J. M. (2002). Regulation of myocardial contractility and cell size by distinct PI3K-PTEN signaling pathways. *Cell*, **110**, 737–749.
- Cross, D. A., Alessi, D. R., Cohen, P., Andjelkovich, M., & Hemmings, B. A. (1995). Inhibition of glycogen synthase kinase-3 by insulin mediated by protein kinase B. *Nature*, **378**, 785–789.
- Cunha-Melo, J. R., Gonzaga, H. M. S., Ali, H., Huang, F. L., Huang, K.-P., & Beaven, M. A. (1989). Studies of protein kinase C in the rat basophilic leukemia (RBL-2H3) cell reveal that antigen-induced signals are not mimicked by the actions of phorbol myristate acetate and Ca^{2+} ionophore. *J. Immunol.*, **143**, 2617–2625.
- Czupalla, C., Culo, M., Müller, E. C., Brock, C., Reusch, H. P., Spicher, K., Krause, E., & Nürnberg, B. (2003). Identification and characterization of the autophosphorylation sites of phosphoinositide 3-kinase isoforms β and γ . *J. Biol. Chem.*, **278**, 11536–11545.
- Côté, J.-F., Motoyama, A. B., Bush, J. A., & Vuori, K. (2005). A novel and evolutionarily conserved PtdIns(3,4,5) P_3 -binding domain is necessary for DOCK180 signalling. *Nat. Cell Biol.*, **7**, 797–807.
- Datta, S. R., Dudek, H., Tao, X., Masters, S., Fu, H., Gotoh, Y., & Greenberg, M. E. (1997). Akt phosphorylation of BAD couples survival signals to the cell-intrinsic death machinery. *Cell*, **91**, 231–241.
- Davis, T. L., Bonacci, T. M., Sprang, S. R., & Smrcka, A. V. (2005). Structural and molecular characterization of a preferred protein interaction surface on G protein $\beta\gamma$ subunits. *Biochemistry*, **44**, 10593–10604.
- Degerman, E., Belfrage, P., & Manganiello, V. C. (1997). Structure, localization, and regulation of cGMP-inhibited phosphodiesterase (PDE3). *J. Biol. Chem.*, **272**, 6823–6826.
- Del Prete, A., Vermi, W., Dander, E., Otero, K., Barberis, L., Luini, W., Bernasconi, S., Sironi, M., Santoro, A., Garlanda, C., Facchetti, F., Wymann, M. P., Vecchi, A., Hirsch, E., Mantovani,

REFERENCES

- A., & Sozzani, S. (2004). Defective dendritic cell migration and activation of adaptive immunity in PI3K γ -deficient mice. *EMBO J.*, **23**, 3505–3515.
- Dhand, R., Hara, K., Hiles, I., Bax, B., Gout, I., Panayotou, G., Fry, M. J., Yonezawa, K., Kasuga, M., & Waterfield, M. D. (1994a). PI 3-kinase: Structural and functional analysis of intersubunit interactions. *EMBO J.*, **13**, 551–521.
- Dhand, R., Hiles, I., Panayotou, G., Roche, S., Fry, M., Gout, I., Totty, N. F., Truong, O., Vicendo, P., Yonezawa, K., Kasuga, M., Courtneidge, S. A., & Waterfield, M. D. (1994b). PI 3-kinase is a dual specificity enzyme: autoregulation by an intrinsic protein-serine kinase activity. *EMBO J.*, **13**, 522–533.
- Doble, B. W. & Woodgett, J. R. (2003). GSK-3: tricks of the trade for a multi-tasking kinase. *J. Cell Sci.*, **116**, 1175–1186.
- Dodge, K. L., Khouangsathiene, S., Kapiloff, M. S., Mouton, R., Hill, E. V., Houslay, M. D., Langeberg, L. K., & Scott, J. D. (2001). mAKAP assembles a protein kinase A/PDE4 phosphodiesterase cAMP signaling module. *EMBO J.*, **20**, 1921–1930.
- Domin, J., Gaidarov, I., Smith, M. E., Keen, J. H., & Waterfield, M. D. (2000). The class II phosphoinositide 3-kinase PI3K-C2 α is concentrated in the trans-Golgi network and present in clathrin-coated vesicles. *J. Biol. Chem.*, **275**, 11943–11950.
- Dorner, B. G., Scheffold, A., Rolph, M. S., Hüser, M. B., Kaufmann, S. H. E., Radbruch, A., Flesch, I. E. A., & Kroczeck, R. A. (2002). MIP-1 α , MIP-1 β , RANTES, and ATAC/lymphotactin function together with IFN- γ as type 1 cytokines. *Proc. Natl. Acad. Sci. U.S.A.*, **99**, 6181–6186.
- Doukas, J., Wrasidlo, W., Noronha, G., Dneprovskaja, E., Fine, R., Weis, S., Hood, J., Soll, A. D. R., & Cheresch, D. (2006). Phosphoinositide 3-kinase γ/δ inhibition limits infarct size after myocardial ischemia/reperfusion injury. *Proc. Natl. Acad. Sci. U.S.A.*, **103**, 19866–19871.
- D'Souza-Schorey, C. & Chavrier, P. (2006). ARF proteins: roles in membrane traffic and beyond. *Nat. Rev. Mol. Cell Biol.*, **7**, 347–358.
- Déléris, P., Gayral, S., & Breton-Douillon, M. (2006). Nuclear PtdIns(3,4,5)P $_3$ signaling: an ongoing story. *J. Cell. Biochem.*, **98**, 469–485.
- Elbashir, S. M., Harborth, J., Lendeckel, W., Yalcin, A., Weber, K., & Tuschl, T. (2001a). Duplexes of 21-nucleotide RNAs mediate RNA interference in cultured mammalian cells. *Nature*, **411**, 494–498.

- Elbashir, S. M., Martinez, J., Patkanioswka, A., Lendeckel, W., & Tuschl, T. (2001b). Functional anatomy of siRNAs for mediating efficient RNAi in *Drosophila melanogaster* embryo lysate. *EMBO J.*, **20**, 6877–6888.
- Ellson, C. D., Andrews, S., Stephens, L. R., & Hawkins, P. T. (2002). The PX domain: a new phosphoinositide-binding module. *J. Cell Sci.*, **115**, 1099–1105.
- Engelman, J. A., Luo, J., & Cantley, L. C. (2006). The evolution of phosphatidylinositol 3-kinases as regulators of growth and metabolism. *Nat. Rev. Genet.*, **7**, 606–619.
- Falasca, M. & Mafucci, T. (2007). Role of class II phosphoinositide 3-kinase in cell signalling. *Biochem. Soc. Trans.*, **35**, 211–214.
- Fan, Q.-W., Knight, Z. A., Goldenberg, D. D., Yu, W., Mostov, K. E., Stokoe, D., Shokat, K. M., & Weiss, W. A. (2006). A dual PI3 kinase/mTOR inhibitor reveals emergent efficacy in glioma. *Cancer Cell*, **9**, 341–349.
- Ferguson, G. J., Milne, L., Kulkarni, S., Sasaki, T., Walker, S., Andrews, S., Crabbe, T., Finan, P., Jones, G., Jackson, S., Camps, M., Rommel, C., Wymann, M., Hirsch, E., Hawkins, P., & Stephens, L. (2007). PI(3)K γ has an important context-dependent role in neutrophil chemokinesis. *Nat. Cell Biol.*, **9**, 86–91.
- Finkelstein, L. D. & Schwartzberg, P. L. (2004). Tec kinases: shaping T-cell activation through actin. *Trends Cell Biol.*, **14**, 443–451.
- Fire, A., Xu, S., Montgomery, M. K., Kostas, S. A., Driver, S. E., & Mello, C. C. (1998). Potent and specific genetic interference by double-stranded RNA in *Caenorhabditis elegans*. *Nature*, **391**, 806–811.
- Fleming, I. N., Gray, A., & Downes, C. P. (2000). Regulation of the Rac1-specific exchange factor Tiam1 involves both phosphoinositide 3-kinase-dependent and -independent components. *Biochem. J.*, **351**, 173–182.
- Fogg, V. C., Azpiazu, I., Linder, M. E., Smrcka, A., Scarlata, S., & Gautam, N. (2001). Role of the γ subunit prenyl moiety in G protein $\beta\gamma$ complex interaction with phospholipase C β . *J. Biol. Chem.*, **276**, 41797–41802.
- Ford, C. E., Skiba, N. P., Bae, H., Daaka, Y., Reuveny, E., Shekter, L. R., Rosal, R., Weng, G., Yang, C.-S., Iyengar, R., Miller, R. J., Jan, L. Y., Lefkowitz, R. J., & Hamm, H. E. (1998). Molecular basis for interactions of G protein $\beta\gamma$ subunits with effectors. *Science*, **280**, 1271–1274.
- Foukas, L. C. & Shepherd, P. R. (2004). eIF4E binding protein 1 and H-Ras are novel substrates for the protein kinase activity of class-I phosphoinositide 3-kinase. *Biochem. Biophys. Res. Comm.*, **319**, 541–549.

REFERENCES

- Foukas, L. C., Beeton, C. A., Jensen, J., Phillips, W. A., & Shepherd, P. R. (2004). Regulation of phosphoinositide 3-kinase by its intrinsic serine kinase activity in vivo. *Mol. Cell. Biol.*, **24**, 966–975.
- Frank, R. (1992). Spot synthesis: an easy technique for the positionally addressable, parallel chemical synthesis. *Tetrahedron*, **48**, 9217–9232.
- Funamoto, S., Meili, R., Lee, S., Perry, L., & Firtel, R. A. (2002). Spatial and temporal regulation of 3-phosphoinositides by PI 3-kinase and PTEN mediates chemotaxis. *Cell*, **109**, 611–623.
- Förster, T. (1948). Zwischenmolekulare Energiewanderung und Fluoreszenz. *Ann. Physik*, **2**, 55–67.
- Gaidarov, I., Smith, M. E., Domin, J., & Keen, J. H. (2001). The class II phosphoinositide 3-kinase PI3K-C2 α is activated by clathrin and regulates clathrin-mediated membrane trafficking. *Mol. Cell*, **7**, 443–449.
- Gaidarov, I., Chen, Q., Falck, J. R., Reddy, K. K., & Keen, J. H. (1996). A functional phosphatidylinositol 3,4,5-trisphosphate/phosphoinositide binding domain in the clathrin adaptor AP-2 α subunit. *J. Biol. Chem.*, **271**, 20922–20929.
- Gaidarov, I. & Keen, J. H. (1999). Phosphoinositide–AP-2 interactions required for targeting to plasma membrane clathrin-coated pits. *J. Cell Biol.*, **146**, 755–764.
- Geering, B., Cutillas, P. R., & Vanhaesebroeck, B. (2007). Regulation of class IA PI3Ks: is there a role for monomeric PI3K subunits?. *Biochem. Soc. Trans.*, **35**, 199–203.
- Gibson, S. K. & Gilman, A. G. (2005). G α and G β subunits both define selectivity of G protein activation by α 2-adrenergic receptors. *Proc. Natl. Acad. Sci. U.S.A.*, **103**, 212–217.
- Gilfillan, A. M. & Tkaczyk, C. (2006). Integrated signalling pathways for mast-cell activation. *Nat. Rev. Immunol.*, **6**, 218–230.
- Gillham, H., Golding, M. C., Pepperkok, R., & Gullick, W. J. (1999). Intracellular movement of green fluorescent protein-tagged phosphatidylinositol 3-kinase in response to growth factor receptor signaling. *J. Cell Biol.*, **146**, 869–880.
- Gimm, O., Perren, A., Weng, L.-P., Marsh, D. J., Yeh, J. J., Ziebold, U., Gil, E., Hinze, R., Delbridge, L., Lees, J. A., Mutter, G. L., Robinson, B. G., Komminoth, P., Dralle, H., & Eng, C. (2000). Differential nuclear and cytoplasmic expression of PTEN in normal thyroid tissue, and benign and malignant epithelial thyroid tumors. *Am. J. Pathol.*, **156**, 1693–1700.
- Gish, W. & States, D. J. (1993). Identification of protein coding regions by database similarity search. *Nat. Genet.*, **3**, 266–272.

- Gonnet, G. H., Cohen, M. A., & Benner, S. A. (1992). Exhaustive matching of the entire protein sequence database. *Science*, **256**, 1443–1445.
- Gouet, P., Robert, X., & Courcelle, E. (2003). ESPript/ENDscript: extracting and rendering sequence and 3D information from atomic structures of proteins. *Nucleic Acids Res.*, **31**, 3320–3323.
- Gray, A., Van Der Kaay, J., & Downes, C. P. (1999). The pleckstrin homology domains of protein kinase B and GRP1 (general receptor for phosphoinositides-1) are sensitive and selective probes for the cellular detection of phosphatidylinositol 3,4-bisphosphate and/or phosphatidylinositol 3,4,5-trisphosphate *in vivo*. *Biochem. J.*, **344**, 929–936.
- Grynkiewicz, G., Poenie, M., & Tsien, R. Y. (1985). A new generation of Ca^{2+} indicators with greatly improved fluorescence properties. *J. Biol. Chem.*, **260**, 3440–3450.
- Gu, H. & Neel, B. G. (2003). The 'Gab' in signal transduction. *Trends Cell Biol.*, **13**, 122–130.
- Gu, H., Saito, K., Klamann, L. D., Shen, J., Fleming, T., Wang, Y., Pratt, J. C., Lin, G., Lim, B., Kinet, J.-P., & Neel, B. G. (2001). Essential role for Gab2 in the allergic response. *Nature*, **412**, 186–190.
- Guertin, D. A., Stevens, D. M., Thoreen, C. C., Burds, A. A., Kalaany, N. Y., Moffat, J., Brown, M., Fitzgerald, K. J., & Sabatini, D. M. (2006). Ablation in mice of the mTORC components *raptor*, *riCTOR*, or *mLst8* reveals that mTORC2 is required for signaling to Akt-FOXO and PKC α , but not S6K1. *Dev. Cell*, **11**, 859–871.
- Gukovsky, I., Cheng, J. H., Nam, K. J., Lee, O. T., Lugea, A., Fischer, L., Penninger, J. M., Pandol, S. J., & Gukovskaya, A. S. (2004). Phosphatidylinositol 3-kinase γ regulates key pathologic responses to cholecystokinin in pancreatic acinar cells. *Gastroenterology*, **126**, 554–566.
- Haar, E. V., Lee, S.-I., Bandhakavi, S., Griffin, T. J., & Kim, D.-H. (2007). Insulin signalling to mTOR mediated by the Akt/PKB substrate PRAS40. *Nat. Cell Biol.*, **9**, 316–323.
- Hamilton, A. J. & Baulcombe, D. C. (1999). A species of small antisense RNA in posttranscriptional gene silencing in plants. *Science*, **286**, 950–952.
- Hamm, H. E. (1998). The many faces of G protein signaling. *J. Biol. Chem.*, **273**, 669–672.
- Han, A., Saijo, K., Mecklenbräuer, I., Tarakhovskiy, A., & Nussenzweig, M. C. (2003). Bam32 links the B cell receptor to ERK and JNK and mediates B cell proliferation but not survival. *Immunity*, **19**, 621–632.

REFERENCES

- Han, J., Luby-Phelps, H., Das, B., Shu, X., Xia, Y., Mosteller, R. D., Krishna, U. M., Falck, J. R., White, M. A., & Broek, D. (1998). Role of substrates and products of PI 3-kinase in regulating activation of Rac-related guanosine triphosphatases by Vav. *Science*, **279**, 558–560.
- Hannigan, M., Zhan, L., Li, Z., Ai, Y., Wu, D., & Huang, C. K. (2002). Neutrophils lacking phosphoinositide 3-kinase γ show loss of directionality during N-formyl-Met-Leu-Phe-induced chemotaxis. *Proc. Natl. Acad. Sci. U.S.A.*, **99**, 3603–3608.
- Hannon, G. J. (2002). RNA interference. *Nature*, **418**, 244–250.
- Hawkins, P. T., Anderson, K. E., Davidson, K., & Stephens, L. R. (2006). Signalling through class I PI3Ks in mammalian cells. *Biochem. Soc. Trans.*, **34**, 647–662.
- Heim, R. & Tsien, R. Y. (1996). Engineering green fluorescent protein for improved brightness, longer wavelengths and fluorescence energy transfer. *Curr. Biol.*, **6**, 178–182.
- Hellwig, N., Albrecht, N., Harteneck, C., Schultz, G., & Schaefer, M. (2005). Homo- and heteromeric assembly of TRPV channel subunits. *J. Cell Sci.*, **118**, 917–928.
- Hellwig, N., Plant, T. D., Janson, W., Schäfer, M., Schultz, G., & Schaefer, M. (2004). TRPV1 acts as proton channel to induce acidification in nociceptive neurons. *J. Biol. Chem.*, **279**, 34553–34561.
- Henikoff, S. & Henikoff, J. G. (1992). Amino acid substitution matrices from protein blocks. *Proc. Natl. Acad. Sci. U.S.A.*, **89**, 10915–10919.
- Hennessy, B. T., Smith, D. L., Ram, P. T., Lu, Y., & Mills, G. B. (2005). Exploiting the PI3K/Akt pathway for cancer drug discovery. *Nat. Rev. Drug Disc.*, **4**, 988–1004.
- Hiles, I. D., Otsu, M., Volinia, S., Fry, M. J., Gout, I., Dhand, R., Panayotou, G., Riuz-Larrea, F., Thompson, A., Totty, N. F., Hsuan, J. J., Courtneidge, S. A., Parker, P. J., & Waterfield, M. D. (1992). Phosphatidylinositol 3-kinase: Structure and expression of the 110 kd catalytic subunit. *Cell*, **70**, 419–429.
- Hirsch, E., Bosco, O., Tropel, P., Laffargue, M., Calvez, R., Altruda, F., Wymann, M. P., & Montecuccio, G. (2001). Resistance to thromboembolism in PI3K γ -deficient mice. *FASEB J.*, **15**, 2019–2021.
- Hirsch, E., Katanaev, V. L., Garlanda, C., Azzolino, O., Pirola, L., Silengo, L., Sozzani, S., Mantovani, A., Altruda, F., & Wymann, M. P. (2000). Central role for G protein-coupled phosphoinositide 3-kinase γ in inflammation. *Science*, **287**, 1049–1053.
- Hofmann, T., Schaefer, M., Schultz, G., & Gudermann, T. (2002). Subunit composition of mammalian transient receptor potential channels in living cells. *Proc. Natl. Acad. Sci. U.S.A.*, **99**, 7461–7466.

- Holz, M. K., Ballif, B. A., Gygi, S. P., & Blenis, J. (2005). mTOR and S6K1 mediate assembly of the translation preinitiation complex through dynamic protein interchange and ordered phosphorylation events. *Cell*, **123**, 569–580.
- Hresko, R. C. & Mueckler, M. (2005). mTOR·RICTOR is the Ser⁴⁷³ kinase for Akt/protein kinase B in 3T3-L1 adipocytes. *J. Biol. Chem.*, **280**, 40406–40416.
- Hu, P., Mondino, A., Skolnik, E. Y., & Schlessinger, J. (1993). Cloning of a novel, ubiquitously expressed human phosphatidylinositol 3-kinase and identification of its binding site on p85. *Mol. Cell. Biol.*, **13**, 7677–7688.
- Huber, M., Helgason, C. D., Damen, J. E., Liu, L., Humphries, R. K., & Krystal, G. (1998). The src homology 2-containing inositol phosphatase (SHIP) is the gatekeeper of mast cell degranulation. *Proc. Natl. Acad. Sci. U.S.A.*, **95**, 11330–11335.
- Hunter, T. (1995). When is a lipid kinase not a lipid kinase? When it is a protein kinase. *Cell*, **83**, 1–4.
- Hurley, J. H. & Misra, S. (2000). Signaling and subcellular targeting by membrane-binding domains. *Annu. Rev. Biophys. Biomol. Struct.*, **29**, 49–79.
- Iijima, M. & Devreotes, P. (2002). Tumor suppressor PTEN mediates sensing of chemoattractant gradients. *Cell*, **109**, 599–610.
- Ingham, R. J., Santos, L., Dang-Lawson, M., Holgado-Madruga, M., Dudek, P., Maroun, C. R., Wong, A. J., Matsuuchi, L., & Gold, M. R. (2001). The Gab1 docking protein links the B cell antigen receptor to the phosphatidylinositol 3-kinase/Akt signaling pathway and to the SHP2 tyrosine phosphatase. *J. Biol. Chem.*, **276**, 12257–12265.
- Inglese, J., Luttrell, L. M., Iniguez-Lluhi, J. A., Touhara, K., Koch, W. J., & Lefkowitz, R. J. (1994). Functionally active targeting domain of the β -adrenergic receptor kinase: An inhibitor of $G_{\beta\gamma}$ -mediated stimulation of type II adenylyl cyclase. *Proc. Natl. Acad. Sci. U.S.A.*, **91**, 3637–3641.
- Inoki, K., Li, Y., Zhu, T., Wu, J., & Guan, K.-L. (2002). TSC2 is phosphorylated and inhibited by Akt and suppresses mTOR signalling. *Nat. Cell Biol.*, **4**, 648–657.
- Irvine, R. F. (2003). Nuclear lipid signalling. *Nat. Rev. Mol. Cell Biol.*, **4**, 1–12.
- Iwase, M., Bishop, S. P., Uechi, M., Vatner, D. E., Shannon, R. P., Kudej, R. K., Wight, D. C., Wagner, T. E., Ishikawa, Y., Homcy, C. J., & Vatner, S. F. (1996). Adverse effects of chronic endogenous sympathetic drive induced by cardiac G_s overexpression. *Circ. Res.*, **78**, 517–524.

REFERENCES

- Jacinto, E., Facchinetti, V., Liu, D., Soto, N., Wei, S., Jung, S. Y., Huang, Q., Qin, J., & Su, B. (2006). SIN1/MIP1 maintains rictor-mTOR complex integrity and regulates Akt phosphorylation and substrate specificity. *Cell*, **127**, 125–137.
- Jackson, S. P., Schoenwaelder, S. M., Goncalves, I., Nesbitt, W. S., Yap, C. L., Wright, C. E., Kenche, V., Anderson, K. E., Dopheide, S. M., Yuan, Y., Sturgeon, S. A., Prabakaran, H., Thompson, P. E., Smith, G. D., Shepherd, P. R., Daniele, N., Kulkarni, S., Abbott, B., Sylik, D., Jones, C., Lu, L., Giuliano, S., Hugan, S. C., Angus, J. A., Robertson, A. D., & Salem, H. H. (2005). PI 3-kinase p110 β : a new target for antithrombotic therapy. *Nat. Med.*, **11**, 507–514.
- Jackson, T. R., Kearns, B. G., & Theibert, A. B. (2000). Cytohesins and centaurins: mediators of PI 3-kinase-regulated Arf signaling. *Trends Biochem. Sci.*, **25**, 489–495.
- Jiménez, C., Hernández, C., Pimentel, B., & Carrera, A. C. (2002). The p85 regulatory subunit controls sequential activation of phosphoinositide 3-kinase by Tyr kinases and Ras. *J. Biol. Chem.*, **277**, 41556–41562.
- Jones, G. E., Prigmore, E., Calvez, R., Hogan, C., Dunn, G. A., Hirsch, E., Wymann, M. P., & Ridley, A. J. (2003). Requirement for PI 3-kinase γ in macrophage migration to MCP-1 and CSF-1. *Exp. Cell Res.*, **290**, 120–131.
- Kalkbrenner, F., Degtiar, V. E., Schenker, M., Brendel, S., Zobel, A., Hescheler, J., Wittig, B., & Schultz, G. (1995). Subunit composition of G_o proteins functionally coupling galanin receptors to voltage-gated calcium channels. *EMBO J.*, **14**, 4728–4737.
- Kapeller, R., Prasad, K. V., Janssen, O., Hou, W., Schaffhausen, B. S., Rudd, C. E., & Cantley, L. C. (1994). Identification of two SH3-binding motifs in the regulatory subunit of phosphatidylinositol 3-kinase. *J. Biol. Chem.*, **269**, 1927–1933.
- Katakami, Y., Kaibuchi, K., Sawamura, M., Takai, Y., & Nishizuka, Y. (1984). Synergistic action of protein kinase C and calcium for histamine release from rat peritoneal mast cells. *Biochem. Biophys. Res. Comm.*, **121**, 573–578.
- Kenworthy, A. K. (2001). Imaging protein-protein interactions using fluorescence resonance energy transfer. *Methods*, **24**, 289–296.
- Kerchner, K. R., Clay, R. L., McCleery, G., Watson, N., McIntire, W. E., Myung, C.-S., & Garrison, J. C. (2004). Differential sensitivity of phosphatidylinositol 3-kinase p110 γ to isoforms of G protein $\beta\gamma$ dimers. *J. Biol. Chem.*, **279**, 44554–44562.
- Kerfant, B.-G., Rose, R. A., Sun, H., & Backx, P. H. (2006). Phosphoinositide 3-kinase γ regulates cardiac contractility by locally controlling cyclic adenosine monophosphate levels. *Trends Cardiovasc Med.*, **16**, 250–256.

- Keskin, O., Ma, B., & Nussinov, R. (2005). Hot regions in protein-protein interactions: The organization and contribution of structurally conserved hot spot residues. *J. Mol. Biol.*, **345**, 1281–1294.
- Kim, S. J. (1998). Insulin rapidly induces nuclear translocation of PI3-kinase in HepG2 cells. *Biochem. Mol. Biol. Int.*, **46**, 187–196.
- Klarlund, J. K., Guilherme, A., Holik, J. J., Virbasius, J. V., Chawla, A., & Czech, M. P. (1997). Signaling by phosphoinositide-3,4,5-trisphosphate through proteins containing pleckstrin and Sec7 homology domains. *Science*, **275**, 1927–1930.
- Klarlund, J. K., Rameh, L. E., Cantley, L. C., Buxton, J. M., Holik, J. J., Sakelis, C., Patki, V., Corvera, S., & Czech, M. P. (1998). Regulation of GRP1-catalyzed ADP ribosylation factor guanine nucleotide exchange by phosphatidylinositol-3,4,5-trisphosphate. *J. Biol. Chem.*, **273**, 1859–1862.
- Kleuss, C., Scherübl, H., Hescheler, J., Schultz, G., & Wittig, B. (1992). Different β -subunits determine G-protein interaction with transmembrane receptors. *Nature*, **358**, 424–426.
- Kleuss, C., Scherübl, H., Hescheler, J., Schultz, G., & Wittig, B. (1993). Selectivity in signal transduction determined by γ subunits of heterotrimeric G proteins. *Science*, **259**, 832–834.
- Klippel, A., Escobedo, J. A., Fantl, W. J., & Williams, L. T. (1992). The C-terminal SH2 domain of p85 accounts for the high affinity and specificity of the binding of phosphatidylinositol 3-kinase to platelet-derived growth factor β receptor. *Mol. Cell. Biol.*, **12**, 1451–1459.
- Klippel, A., Escobedo, J. A., Hu, Q., & Williams, L. T. (1993). A region of the 85-kilodalton (kDa) subunit of phosphatidylinositol 3-kinase binds the 110-kDa catalytic subunit *in vivo*. *Mol. Cell. Biol.*, **13**, 5560–5566.
- Knight, Z. A., Gonzalez, B., Feldman, M. E., Zunder, E. R., Goldenberg, D. D., Williams, O., Loewith, R., Stokoe, D., Balla, A., Toth, B., Balla, T., Weiss, W. A., Williams, R. L., & Shokat, K. M. (2006). A pharmacological map of the PI3-K family defines a role for p110 α in insulin signaling. *Cell*, **125**, 733–747.
- Kolanus, W., Nagel, W., Schiller, B., Zeitlmann, L., Godar, S., Stockinger, H., & Seed, B. (1996). α L β 2 integrin/LFA-1 binding to ICAM-1 induced by cytohesin-1, a cytoplasmic regulatory molecule. *Cell*, **86**, 233–242.
- Kondo, Y., Shen, L., Yan, P. S., Huang, T. H.-M., & Issa, J.-P. J. (2004). Chromatin immunoprecipitation microarrays for identification of genes silenced by histone H3 lysine 9 methylation. *Proc. Natl. Acad. Sci. U.S.A.*, **101**, 7398–7403.

REFERENCES

- Koyasu, S. (2003). The role of PI3K in immune cells. *Nat. Immunol.*, **4**, 313–319.
- Kozak, M. (1987). At least six nucleotides preceding the AUG initiator codon enhance translation in mammalian cells. *J. Mol. Biol.*, **196**, 947–950.
- Kramer, A. & Schneider-Mergener, J. (1998). Synthesis and screening of peptide libraries on continuous cellulose membrane supports. *Methods Mol. Biol.*, **87**, 25–39.
- Krugmann, S., Anderson, K. E., Ridley, S. H., Risso, N., McGregor, A., Coadwell, J., Davidson, K., Eguinoa, A., Ellson, C. D., Lipp, P., Manifava, M., Ktistakis, N., Painter, G., Thuring, J. W., Cooper, M. A., Lim, Z.-Y., Holmes, A. B., Dove, S. K., Michell, R. H., Grewal, A., Nazarian, A., Erdjument-Bromage, H., Tempst, P., Stephens, L. R., & Hawkins, P. T. (2002a). Identification of ARAP3, a novel PI3K effector regulating both Arf and Rho GTPases, by selective capture on phosphoinositide affinity matrices. *Mol. Cell*, **9**, 95–108.
- Krugmann, S., Cooper, M. A., Williams, D. H., Hawkins, P. T., & Stephens, L. R. (2002b). Mechanism of the regulation of type IB phosphoinositide 3OH-kinase by G-protein $\beta\gamma$ subunits. *Biochem. J.*, **362**, 725–731.
- Krugmann, S., Hawkins, P. T., Pryor, N., & Braselmann, S. (1999). Characterizing the interactions between the two subunits of the p101/p110 γ phosphoinositide 3-kinase and their role in the activation of this enzyme by G $\beta\gamma$ subunits. *J. Biol. Chem.*, **274**, 17152–17158.
- Krystal, G. (2000). Lipid phosphatases in the immune system. *Semin. Immunol.*, **12**, 397–403.
- Kular, G., Loubtchenkov, M., Swigart, P., Whatmore, J., Ball, A., Cockcroft, S., & Wetzker, R. (1997). Co-operation of phosphatidylinositol transfer protein with phosphoinositide 3-kinase γ in the formylmethionylleucylphenylalanine-dependent production of phosphatidylinositol 3,4,5-trisphosphate in human neutrophils. *Biochem. J.*, **325**, 299–301.
- Kurosu, H., Maehama, T., Okada, T., Yamamoto, T., Hoshino, S.-I., Fukui, Y., Ui, M., Hazeki, O., & Katada, T. (1997). Heterodimeric phosphoinositide 3-kinase consisting of p85 and p110 β is synergistically activated by the $\beta\gamma$ subunits of G proteins and phosphotyrosyl peptide. *J. Biol. Chem.*, **272**, 24252–24256.
- Lachyankar, M. B., Sultana, N., Schonhoff, C. M., Mitra, P., Poluha, W., Lambert, S., Quesenberry, P. J., Litofsky, N. S., Recht, L. D., Nabi, R., Miller, S. J., Ohta, S., Neel, B. G., & Ross, A. H. (2000). A role for nuclear PTEN in neuronal differentiation. *J. Neurosci.*, **20**, 1404–1413.
- Laemmli, U. K. (1970). Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature*, **227**, 680–685.

- Laffargue, M., Calvez, R., Finan, P., Trifilieff, A., Barbier, M., Altruda, F., Hirsch, E., & Wymann, M. P. (2002). Phosphoinositide 3-kinase γ is an essential amplifier of mast cell function. *Immunity*, **16**, 441–451.
- Lam, K., Carpenter, C. L., Ruderman, N. B., Friel, J. C., & Kelly, K. L. (1994). The phosphatidylinositol 3-kinase serine kinase phosphorylates IRS-1. *J. Biol. Chem.*, **269**, 20648–20652.
- Lambert, J. M., Lambert, Q. T., Reuther, G. W., Malliri, A., Siderovski, D. P., Sondek, J., Collard, J. G., & Der, C. J. (2002). Tiam1 mediates Ras activation of Rac by a PI(3)K-independent mechanism. *Nat. Cell Biol.*, **4**, 621–625.
- Leevers, S. J., Weinkove, D., MacDougall, L. K., Hafen, E., & Waterfield, M. D. (1996). The *Drosophila* phosphoinositide 3-kinase Dp110 promotes cell growth. *EMBO J.*, **15**, 6584–6594.
- Lei, Q., Jones, M. B., Talley, E. M., Schrier, A. D., McIntire, W. E., Garrison, J. C., & Bayliss, D. A. (2000). Activation and inhibition of G protein-coupled inwardly rectifying potassium (Kir3) channels by G protein $\beta\gamma$ subunits. *Proc. Natl. Acad. Sci. U.S.A.*, **97**, 9771–9776.
- Lemmon, M. A. (2004). Pleckstrin homology domains: not just for phosphoinositides. *Biochem. Soc. Trans.*, **32**, 707–711.
- Lemmon, M. A. & Ferguson, K. M. (2000). Signal-dependent membrane targeting by pleckstrin homology (PH) domains. *Biochem. J.*, **350**, 1–18.
- Lenz, J. C., Reusch, H. P., Albrecht, N., Schultz, G., & Schaefer, M. (2002). Ca^{2+} -controlled competitive diacylglycerol binding of protein kinase C isoenzymes in living cells. *J. Cell Biol.*, **159**, 291–302.
- Leopoldt, D., Hanck, T., Exner, T., Maier, U., Wetzker, R., & Nürnberg, B. (1998). $G\beta\gamma$ stimulates phosphoinositide 3-kinase- γ by direct interaction with two domains of the catalytic p110 subunit. *J. Biol. Chem.*, **273**, 7024–7029.
- Leslie, N. R. & Downes, C. P. (2002). PTEN: The down side of PI 3-kinase signalling. *Cell. Signal.*, **14**, 285–295.
- Li, Y., Sternweis, P. M., Charnecki, S., Smith, T. F., Gilman, A. G., Neer, E. J., & Kozasa, T. (1998). Sites for $G\alpha$ binding on the G protein β subunit overlap with sites for regulation of phospholipase $C\beta$ and adenylyl cyclase. *J. Biol. Chem.*, **273**, 16265–16272.
- Li, Z., Jiang, H., Xie, W., Zhang, Z., Smrcka, A. V., & Wu, D. (2000). Roles of PLC- β 2 and - β 3 and PI3K γ in chemoattractant-mediated signal transduction. *Science*, **287**, 1046–1049.

REFERENCES

- Lim, W. K., Myung, C.-S., Garrison, J. C., & Neubig, R. R. (2001). Receptor-G protein γ specificity: γ_{11} shows unique potency for A₁ adenosine and 5-HT_{1A} receptors. *Biochemistry*, **40**, 10532–10541.
- Lindmo, K. & Stenmark, H. (2006). Regulation of membrane traffic by phosphoinositide 3-kinases. *J. Cell Sci.*, **119**, 605–614.
- Liu, H. & Maurice, D. H. (1998). Expression of cyclic GMP-inhibited phosphodiesterases 3A and 3B (PDE3A and PDE3B) in rat tissues: Differential subcellular localization and regulated expression by cyclic AMP. *Br. J. Pharmacol.*, **125**, 1501–1510.
- Locati, M., Lamorte, G., Luini, W., Introna, M., Bernasconi, S., Mantovani, A., & Sozzani, S. (1996). Inhibition of monocyte chemotaxis to C-C chemokines by antisense oligonucleotide for cytosolic phospholipase A₂. *J. Biol. Chem.*, **271**, 6010–6016.
- Lodowski, D. T., Pitcher, J. A., Capel, W. D., Lefkowitz, R. J., & Tesmer, J. J. G. (2003). Keeping G proteins at bay: A complex between G protein-coupled receptor kinase 2 and G $\beta\gamma$. *Science*, **300**, 1256–1262.
- Lowry, W. E. & Huang, X. Y. (2002). G protein $\beta\gamma$ subunits act on the catalytic domain to stimulate Bruton's agammaglobulinemia tyrosine kinase. *J. Biol. Chem.*, **277**, 1488–1492.
- Lu, P.-J., Hsu, A.-L., Wang, D.-S., Yan, H. Y., Yin, H. L., & Chen, C.-S. (1998). Phosphoinositide 3-kinase in rat liver nuclei. *Biochemistry*, **37**, 5738–5745.
- Luckow, V. A., Lee, S. C., Barry, G. F., & Olins, P. O. (1993). Efficient generation of infectious recombinant baculoviruses by site-specific transposon-mediated insertion of foreign genes into a baculovirus genome propagated in *Escherichia coli*. *J. Virol.*, **67**, 4566–4579.
- Luckow, V. A. & Summers, M. D. (1988). Trends in the development of baculovirus expression vectors. *Bio/Technology*, **6**, 47–55.
- Lupia, E., Goffi, A., Giuli, P. D., Azzolino, O., Bosco, O., Patrucco, E., Vivaldo, M. C., Ricca, M., Wymann, M. P., Hirsch, E., Montrucchio, G., & Emanuelli, G. (2004). Ablation of phosphoinositide 3-kinase- γ reduces the severity of acute pancreatitis. *Am. J. Pathol.*, **165**, 2003–2011.
- Löhning, M., Hutloff, A., Kallinich, T., Mages, H. W., Bonhagen, K., Radbruch, A., Hamelmann, E., & Kroczek, R. A. (2003). Expression of ICOS in vivo defines CD4⁺ effector T cells with high inflammatory potential and a strong bias for secretion of interleukin 10. *J. Exp. Med.*, **197**, 181–193.

- Ma, B., Elkayam, T., Wolfson, H., & Nussinov, R. (2003). Protein-protein interactions: Structurally conserved residues distinguish between binding sites and exposed protein surfaces. *Proc. Natl. Acad. Sci. U.S.A.*, **100**, 5772–5777.
- MacDonald, P. E., Joseph, J. W., Yau, D., Diao, J., Asghar, Z., Dai, F., Oudit, G. Y., Patel, M. M., Backx, P. H., & Wheeler, M. B. (2004). Impaired glucose-stimulated insulin secretion, enhanced intraperitoneal insulin tolerance, and increased β -cell mass in mice lacking the p110 γ isoform of phosphoinositide 3-kinase. *Endocrinology*, **145**, 4078–4083.
- Macrez-Leprêtre, N., Kalkbrenner, F., Morel, J.-L., Schultz, G., & Mironneau, J. (1997). G protein heterotrimer $G\alpha_{13}\beta_1\gamma_3$ couples the angiotensin AT_{1A} receptor to increases in cytoplasmic Ca^{2+} in rat portal vein myocytes. *J. Biol. Chem.*, **272**, 10095–10102.
- Mafucci, T., Cooke, F. T., Foster, F. M., Traer, C. J., Fry, M. J., & Falasca, M. (2005). Class II phosphoinositide 3-kinase defines a novel signaling pathway in cell migration. *J. Cell Biol.*, **169**, 789–799.
- Mahadevan, D., Thanki, N., Singh, J., McPhie, P., Zangrilli, D., Wang, L.-M., Guerrero, C., LeVine, III, H., Humblet, C., Saldanha, J., Gutkind, J. S., & Najmabadi-Haske, T. (1995). Structural studies on the PH domains of Dbl, Sos1, IRS-1, and β ARK1 and their differential binding to $G\beta\gamma$ subunits. *Biochemistry*, **34**, 9111–9117.
- Maier, U., Babich, A., & Nürnberg, B. (1999). Roles of non-catalytic subunits in $G\beta\gamma$ -induced activation of class I phosphoinositide 3-kinase isoforms β and γ . *J. Biol. Chem.*, **274**, 29311–29317.
- Manning, B. D., Tee, A. R., Logsdon, M. N., Blenis, J., & Cantley, L. C. (2002). Identification of the tuberous sclerosis complex-2 tumor suppressor gene product tuberlin as a target of the phosphoinositide 3-kinase/Akt pathway. *Mol. Cell*, **10**, 151–162.
- Martelli, A. M., Borgatti, P., Bortul, R., Manfredini, M., Massari, L., Capitani, S., & Neri, L. M. (2000). Phosphatidylinositol 3-kinase translocates to the nucleus of osteoblast-like MC3T3-E1 cells in response to insulin-like growth factor I and platelet-derived growth factor but not to the proapoptotic cytokine tumor necrosis factor α . *J. Bone Miner. Res.*, **15**, 1716–1730.
- Maurice, D. H., Palmer, D., Tilley, D. G., Dunkerley, H. A., Netherton, S. J., Raymond, D. R., Elbatarny, H. S., & Jimmo, S. L. (2003). Cyclic nucleotide phosphodiesterase activity, expression, and targeting in cells of the cardiovascular system. *Mol. Pharmacol.*, **64**, 533–546.
- Mayeenuddin, L. H., McIntire, W. E., & Garrison, J. C. (2006). Differential sensitivity of P-Rex1 to isoforms of G protein $\beta\gamma$ dimers. *J. Biol. Chem.*, **281**, 1913–1920.

REFERENCES

- Mayo, L. D. & Donner, D. B. (2002). The PTEN, Mdm2, p53 tumor suppressor–oncoprotein network. *Trends Biochem. Sci.*, **27**, 462–467.
- McIntire, W. E., MacCleery, G., & Garrison, J. C. (2001). The G protein β subunit is a determinant in the coupling of G_s to the β_1 -adrenergic and A2a adenosine receptors. *J. Biol. Chem.*, **276**, 15801–15809.
- McMillan, K., Adler, M., Auld, D. S., Baldwin, J. J., Blasko, E., Browne, L. J., Chelsky, D., Davey, D., Dolle, R. E., Eagen, K. A., Erickson, S., Feldman, R. I., Glaser, C. B., Mallari, C., Morrissey, M. M., Ohlmeyer, M. H. J., Pan, G., Parkinson, J. F., Phillips, G. B., Polokoff, M. A., Sigal, N. H., Vergona, R., Whitlow, M., Young, T. A., & Devlin, J. J. (2000). Allosteric inhibitors of inducible nitric oxide synthase dimerization discovered via combinatorial chemistry. *Proc. Natl. Acad. Sci. U.S.A.*, **97**, 1506–1511.
- Merrifield, C. J., Moss, S. E., Ballestrem, C., Imhof, B. A., Giese, G., Wunderlich, I., & Almers, W. (1999). Endocytic vesicles move at the tips of actin tails in cultured mast cells. *Nat. Cell Biol.*, **1**, 72–74.
- Metjian, A., Roll, R. L., Ma, A. D., & Abrams, C. S. (1999). Agonists cause nuclear translocation of phosphatidylinositol 3-kinase γ . *J. Biol. Chem.*, **274**, 27943–27947.
- Meunier, F. A., Osborne, S. L., Hammond, G. R., Cooke, F. T., Parker, P. J., Domin, J., & Schiavo, G. (2005). Phosphatidylinositol 3-kinase C2 α is essential for ATP-dependent priming of neurosecretory granule exocytosis. *Mol. Biol. Cell*, **16**, 4841–4851.
- Milburn, C. C., Deak, M., Kelly, S. M., Price, N. C., Alessi, D. R., & van Aalten, D. M. F. (2003). Binding of phosphatidylinositol 3,4,5-trisphosphate to the pleckstrin homology domain of protein kinase B induces a conformational change. *Biochem. J.*, **375**, 531–538.
- Miura, K., Jacques, K. M., Stauffer, S., Kubosaki, A., Zhu, K., Hirsch, D. S., Resau, J., Zheng, Y., & Randazzo, P. A. (2002). ARAP1: A point of convergence for Arf and Rho signaling. *Mol. Cell*, **9**, 109–119.
- Moon, K. D., Post, C. B., Durden, D. L., Zhou, Q., De, P., Harrison, M. L., & Geahlen, R. L. (2005). Molecular basis for a direct interaction between the Syk protein-tyrosine kinase and phosphoinositide 3-kinase. *J. Biol. Chem.*, **280**, 1543–1551.
- Mora, A., Komander, D., van Aalten, D. M., & Alessi, D. R. (2004). PDK1, the master regulator of AGC kinase signal transduction. *Semin. Cell Dev. Biol.*, **15**, 161–170.
- Morel, J. L., Macrez-Leprêtre, N., & Mironneau, J. (1996). Angiotensin II-activated Ca^{2+} entry-induced release of Ca^{2+} from intracellular stores in rat portal vein myocytes. *Br. J. Pharmacol.*, **118**, 73–78.

- Morishita, R., Ueda, H., Kato, K., & Asano, T. (1998). Identification of two forms of the γ subunit of G protein, γ_{10} and γ_{11} , in bovine lung and their tissue distribution in the rat. *FEBS Lett.*, **428**, 85–88.
- Morris, J. Z., Tissenbaum, H. A., & Ruvkun, G. (1996). A phosphatidylinositol-3-OH kinase family member regulating longevity and diapause in *Caenorhabditis elegans*. *Nature*, **382**, 536–539.
- Myers, M. P., Pass, I., Batty, I. H., Van der Kaay, J., Stolarov, J. P., Hemmings, B. A., Wigler, M. H., Downes, C. P., & Tonks, N. K. (1998). The lipid phosphatase activity of PTEN is critical for its tumor suppressor function. *Proc. Natl. Acad. Sci. U.S.A.*, **95**, 13513–13518.
- Naccache, P. H., Levasseur, S., Lachance, G., Chakravarti, S., Bourgoin, S. G., & McGoll, S. R. (2000). Stimulation of human neutrophils by chemotactic factors is associated with the activation of phosphatidylinositol 3-kinase γ . *J. Biol. Chem.*, **275**, 23636–23641.
- Naga Prasad, S. V., Barak, L. S., Rapacciuolo, A., Caron, M. G., & Rockman, H. A. (2001). Agonist-dependent recruitment of phosphoinositide 3-kinase to the membrane by β -adrenergic receptor kinase 1. *J. Biol. Chem.*, **276**, 18953–18959.
- Naga Prasad, S. V., Esposito, G., Mao, L., Koch, W. J., & Rockman, H. A. (2000). $G\beta\gamma$ -dependent phosphoinositide 3-kinase activation in hearts with *in vivo* pressure overload hypertrophy. *J. Biol. Chem.*, **275**, 4693–4698.
- Naga Prasad, S. V., Jayatilleke, A., Madamanchi, A., & Rockman, H. A. (2005). Protein kinase activity of phosphoinositide 3-kinase regulates β -adrenergic receptor endocytosis. *Nat. Cell Biol.*, **7**, 785–796.
- Naga Prasad, S. V., Laporte, S. A., Chamberlain, D., Caron, M. G., Barak, L., & Rockman, H. A. (2002). Phosphoinositide 3-kinase regulates β_2 -adrenergic receptor endocytosis by AP-2 recruitment to the receptor/ β -arrestin complex. *J. Cell Biol.*, **158**, 563–575.
- Nakai, K. & Horton, P. (1999). PSORT: a program for detecting sorting signals in proteins and predicting their subcellular localization. *Trends Biochem. Sci.*, **24**, 34–36.
- Napoli, C. A., Lemieux, C., & Jorgensen, R. (1990). Introduction of a chimeric chalcone synthase gene in *Petunia* results in reversible cosuppression of homologous genes *in trans*. *Plant Cell*, **2**, 279–289.
- Neduva, V. & Russell, R. B. (2006). Peptides mediating interaction networks: new leads at last. *Curr. Opin. Biotechnol.*, **17**, 465–471.

REFERENCES

- Nienaber, J. J., Tachibana, H., Naga Prasad, S. V., Esposito, G., Wu, D., Mao, L., & Rockman, H. A. (2003). Inhibition of receptor-localized PI3K preserves cardiac β -adrenergic receptor function and ameliorates pressure overload heart failure. *J. Clin. Invest.*, **112**, 1067–1079.
- Nishio, M., Watanabe, K.-I., Sasaki, J., Taya, C., Takasuga, S., Iizuka, R., Balla, T., Yamazaki, M., Watanabe, H., Itoh, R., Kuroda, S., Horie, Y., Förster, I., Mak, T. W., Yonekawa, H., Penninger, J. M., Kanaho, Y., Suzuki, A., & Sasaki, T. (2007). Control of cell polarity and motility by the PtdIns(3,4,5)P₃ phosphatase SHIP1. *Nat. Cell Biol.*, **9**, 36–44.
- Nobukuni, T., Joaquin, M., Roccio, M., Dann, S. G., Kim, S. Y., Gulati, P., Byfield, M. P., Backer, J. M., Bos, F. N. J. L., Zwartkruis, F. J. T., & Thomas, G. (2005). Amino acids mediate mTOR/raptor signaling through activation of class 3 phosphatidylinositol 3OH-kinase. *Proc. Natl. Acad. Sci. U.S.A.*, **102**, 14238–14243.
- Ogasawara, M., Kim, S.-C., Adamik, R., Togawa, A., Ferrans, V. J., Takeda, K., Kirby, M., Moss, J., & Vaughan, M. (2000). Similarities in function and gene structure of cytohesin-4 and cytohesin-1, guanine nucleotide-exchange proteins for ADP-ribosylation factors. *J. Biol. Chem.*, **275**, 3221–3230.
- Okkenhaug, K., Ali, K., & Vanhaesebroeck, B. (2007). Antigen receptor signalling: a distinctive role for the p110 δ isoform of PI3K. *Trends Immunol.*, **28**, 80–87.
- Oliver, J. M., Seagrave, J., Stump, R. F., Pfeiffer, J. R., & Deanin, G. G. (1988). Signal transduction and cellular response in RBL-2H3 mast cells. *Prog. Allergy*, **42**, 185–245.
- Oltersdorf, T., Elmore, S. W., Shoemaker, A. R., Armstrong, R. C., Augeri, D. J., Belli, B. A., Bruncko, M., Deckwerth, T. L., Dinges, J., Hajduk, P. J., Joseph, M. K., Kitada, S., Korsmeyer, S. J., Kunzer, A. R., Letai, A., Li, C., Mitten, M. J., Nettesheim, D. G., Ng, S., Nimmer, P. M., O'Connor, J. M., Oleksijew, A., Petros, A. M., Reed, J. C., Shen, W., Tahir, S. K., Thompson, C. B., Tomaselli, K. J., Wang, B., Wendt, M. D., Zhang, H., Fesik, S. W., & Rosenberg, S. H. (2005). An inhibitor of Bcl-2 family proteins induces regression of solid tumours. *Nature*, **435**, 677–681.
- Orme, M. H., Alrubaie, S., Bradley, G. L., Walker, C. D., & Levers, S. J. (2006). Input from Ras is required for maximal PI(3)K signalling in *Drosophila*. *Nat. Cell Biol.*, **8**, 1298–1302.
- Oudit, G. A., Sun, H., Kerfant, B.-G., Crackower, M. A., Penninger, J. M., & Backx, P. H. (2004). The role of phosphoinositide 3-kinase and PTEN in cardiovascular physiology and disease. *J. Mol. Cell. Cardiol.*, **37**, 449–471.
- Oudit, G. Y., Crackower, M. A., Eriksson, U., Sarao, R., Kozieradzki, I., Sasaki, T., Irie-Sasaki, J., Gidrewicz, D., Rybin, V. O., Wada, T., Steinberg, S. F., Backx, P. H., & Penninger,

- J. M. (2003). Phosphoinositide 3-kinase γ -deficient mice are protected from isoproterenol-induced heart failure. *Circulation*, **108**, 2147–2152.
- Ozawa, K., Szallasi, Z., Kazanietz, M. G., Blumberg, P. M., Mischak, H., Mushinski, J. F., & Beaven, M. A. (1993). Ca^{2+} -dependent and Ca^{2+} -independent isozymes of protein kinase C mediate exocytosis in antigen-stimulated rat basophilic RBL-2H3 mast cells. *J. Biol. Chem.*, **268**, 1749–1756.
- Ozes, O. N., Mayo, L. D., Gustin, J. A., Pfeffer, S. R., Pfeffer, L. M., & Donner, D. B. (1999). NF- κ B activation by tumour necrosis factor requires the Akt serine-threonine kinase. *Nature*, **401**, 82–85.
- Pacold, M. E., Suire, S., Perisic, O., Lara-Gonzalez, L., Davis, C. T., Walker, E. H., Hawkins, P. T., Stephens, L. R., Eccleston, J. F., & Williams, R. L. (2000). Crystal structure and functional analysis of Ras binding to its effector phosphoinositide 3-kinase γ . *Cell*, **103**, 931–943.
- Panchenko, M. P., Saxena, K., Li, Y., Charnecki, S., Sternweis, P. M., Smith, T. F., Gilman, A. G., Kozasa, T., & Neer, E. J. (1998). Sites important for PLC β_2 activation by the G protein $\beta\gamma$ subunit map to the sides of the β propeller structure. *J. Biol. Chem.*, **273**, 28298–28304.
- Parekh, D. B., Ziegler, W., & Parker, P. J. (2000). Multiple pathways control protein kinase C phosphorylation. *EMBO J.*, **19**, 496–503.
- Patrucco, E., Notte, A., Barberis, L., Selvetella, G., Maffei, A., Brancaccio, M., Marengo, S., Russo, G., Azzolino, O., Rybalkin, S. D., Silengo, L., Altruda, F., Wetzker, R., Wymann, M. P., Lembo, G., & Hirsch, E. (2004). PI3K γ modulates the cardiac response to chronic pressure overload by distinct kinase-dependent and -independent effects. *Cell*, **118**, 375–387.
- Pawley, J. B. (Ed.). (1995). *Handbook of Biological Confocal Microscopy* (2nd edition). Plenum Press, New York.
- Pearce, G., Angeli, V., Randolph, G. J., Junt, T., von Andrian, U., Schnittler, H.-J., & Jessberger, R. (2006). Signaling protein SWAP-70 is required for efficient B cell homing to lymphoid organs. *Nat. Immunol.*, **7**, 827–834.
- Penn, R. B., Pronin, A. N., & Benovic, J. L. (2000). Regulation of G protein-coupled receptor kinases. *Trends Cardiovasc Med.*, **10**, 81–89.
- Perisic, O., Wilson, M. I., Karathanassis, D., Bravo, J., Pacold, M. E., Ellson, C. D., Hawkins, P. T., Stephens, L., & Williams, R. L. (2004). The role of phosphoinositides and phosphorylation in regulation of NADPH oxidase. *Adv. Enzyme Regul.*, **44**, 279–298.

REFERENCES

- Perrino, C., Naga Prasad, S. V., Schroder, J. N., Hata, J. A., Milano, C., & Rockman, H. A. (2005). Restoration of β -adrenergic receptor signaling and contractile function in heart failure by disruption of the β ARK1/phosphoinositide 3-kinase complex. *Circulation*, **111**, 2597–2587.
- Pinho, V., Souza, D. G., Barsante, M. M., Hamer, F. P., De Freitas, M. S., Rossi, A. G., & Teixeira, M. M. (2005). Phosphoinositide-3 kinases critically regulate the recruitment and survival of eosinophils in vivo: importance for the resolution of allergic inflammation. *J. Leukoc. Biol.*, **77**, 800–810.
- Pitcher, J. A., Freedman, N. J., & Lefkowitz, R. J. (1998). G protein-coupled receptor kinases. *Annu. Rev. Biochem.*, **67**, 653–692.
- Pleiman, C. M., Hertz, W. M., & Cambier, J. C. (1994). Activation of phosphatidylinositol-3' kinase by Src-family kinase SH3 binding to the p85 subunit. *Science*, **263**, 1609–1612.
- Pomel, V., Klicic, J., Covini, D., Church, D. D., Shaw, J. P., Roulin, K., Burgat-Charvillon, F., Valognes, D., Camps, M., Chabert, C., Gillieron, C., Françon, B., Perrin, D., Leroy, D., Gretener, D., Nichols, A., Vitte, P. A., Carboni, S., Rommel, C., Schwarz, M. K., & Rückle, T. (2006). Furan-2-ylmethylene thiazolidinediones as novel, potent, and selective inhibitors of phosphoinositide 3-kinase γ . *J. Med. Chem.*, **49**, 3857–3871.
- Puri, K. D., Doggett, T. A., Huang, C.-Y., Douangpanya, J., Hayflick, J. S., Turner, M., Penninger, J., & Diacovo, T. G. (2005). The role of endothelial PI3K γ activity in neutrophil trafficking. *Blood*, **106**, 150–157.
- Quignard, J.-F., Mironneau, J., Carricaburu, V., Fournier, B., Babich, A., Nürnberg, B., Mironneau, C., & Macrez, N. (2001). Phosphoinositide 3-kinase γ mediates angiotensin II-induced stimulation of L-type calcium channels in vascular myocytes. *J. Biol. Chem.*, **276**, 32545–32551.
- Raftopoulou, M. & Hall, A. (2004). Cell migration: Rho GTPases lead the way. *Dev. Biol.*, **265**, 23–32.
- Ramkumar, V., Styles, G. L., Beaven, M. A., & Ali, H. (1993). The A₃ adenosine receptor is the unique adenosine receptor which facilitates release of allergic mediators in mast cells. *J. Biol. Chem.*, **268**, 16887–16890.
- Randazzo, P. A. & Hirsch, D. S. (2004). Arf GAPs: multifunctional proteins that regulate membrane traffic and actin remodelling. *Cell. Signal.*, **16**, 401–413.
- Reif, K., Okkenhaug, K., Sasaki, T., Penninger, J. M., Vanhaesebroeck, B., & Cyster, J. G. (2004). Cutting edge: Differential roles for phosphoinositide 3-kinases, p110 γ and p110 δ , in lymphocyte chemotaxis and homing. *J. Immunol.*, **173**, 2236–2240.

- Reineke, U., Volkmer-Engert, R., & Schneider-Mergener, J. (2001). Applications of peptide arrays prepared by the SPOT-technology. *Curr. Opin. Biotechnol.*, **12**, 59–64.
- Rhee, S. G. (2001). Regulation of phosphoinositide-specific phospholipase C. *Annu. Rev. Biochem.*, **70**, 281–312.
- Richter, J. D. & Sonenberg, N. (2005). Regulation of cap-dependent translation by eIF4E inhibitory proteins. *Nature*, **433**, 477–480.
- Rockman, H. A., Koch, W. J., & Lefkowitz, R. J. (2002). Seven-transmembrane-spanning receptors and heart function. *Nature*, **415**, 206–212.
- Rodrigues, G. A., Falasca, M., Zhang, Z., Ong, S. H., & Schlessinger, J. (2000). A novel positive feedback loop mediated by the docking protein Gab1 and phosphatidylinositol 3-kinase in epidermal growth factor receptor signaling. *Mol. Cell. Biol.*, **20**, 1448–1459.
- Rodriguez-Viciana, P., Warne, P. H., Dhand, R., Vanhaesebroeck, B., Gout, I., Fry, M. J., Waterfield, M. D., & Downward, J. (1994). Phosphatidylinositol-3-OH kinase as a direct target of Ras. *Nature*, **370**, 527–532.
- Rodriguez-Viciana, P., Warne, P. H., Vanhaesebroeck, B., Waterfield, M. D., & Downward, J. (1996). Activation of phosphoinositide 3-kinase by interaction with Ras and by point mutation. *EMBO J.*, **15**, 2442–2451.
- Rodriguez-Viciana, P., Sabatier, C., & McCormick, F. (2004). Signaling specificity by Ras family GTPases is determined by the full spectrum of effectors they regulate. *Mol. Cell. Biol.*, **24**, 4943–4954.
- Rodríguez-Borlado, L., Barber, D. F., Hernández, C., Rodríguez-Marcos, M. A., Sánchez, A., Hirsch, E., Wymann, M., Martínez-A., C., & Carrera, A. C. (2003). Phosphatidylinositol 3-kinase regulates the CD4/CD8 T cell differentiation ratio. *J. Immunol.*, **170**, 4475–4482.
- Rohrschneider, L. R., Fuller, J. F., Wolf, I., Liu, Y., & Lucas, D. M. (2000). Structure, function, and biology of SHIP proteins. *Genes Dev.*, **14**, 505–520.
- Rondinone, C. M., Carvalho, E., Rahn, T., Manganiello, V. C., Degerman, E., & Smith, U. P. (2000). Phosphorylation of PDE3B by phosphatidylinositol 3-kinase associated with the insulin receptor. *J. Biol. Chem.*, **275**, 10093–10098.
- Rossman, K. L., Der, C. J., & Sondek, J. (2005). GEF means go: Turning on Rho GTPases with guanine nucleotide-exchange factors. *Nat. Rev. Mol. Cell Biol.*, **6**, 167–180.
- Russell, R. B., Alber, F., Aloy, P., Davis, F. P., Korkin, D., Pichaud, M., Topf, M., & Sali, A. (2004). A structural perspective on protein-protein interactions. *Curr. Opin. Struct. Biol.*, **14**, 313–324.

REFERENCES

- Rückle, T., Schwarz, M. K., & Rommel, C. (2006). PI3K γ inhibition: towards an 'aspirin of the 21st century'?. *Nat. Rev. Drug Disc.*, **5**, 903–918.
- Sadhu, C., Masinovsky, B., Dick, K., Sowell, C. G., & Staunton, D. E. (2003). Essential role of phosphoinositide 3-kinase δ in neutrophil directional movement. *J. Immunol.*, **170**, 2647–2654.
- Salazar, N. C., Chen, J., & Rockman, H. A. (2007). Cardiac GPCRs: GPCR signaling in healthy and failing hearts. *Biochim. Biophys. Acta*, **1768**, 1006–1018.
- Sambrook, J. & Russel, D. W. (2001). *Molecular cloning: A laboratory manual* (3rd edition). Cold Spring Harbor Press, New York.
- Samuels, Y., Wang, Z., Bardelli, A., Silliman, N., Ptak, J., Szabo, S., Yan, H., Gazdar, A., Powell, S. M., Riggins, G. J., Willson, J. K. V., Markowitz, S., Kinzler, K. W., Vogelstein, B., & Velculescu, V. E. (2004). High frequency of mutations of the *PIK3CA* gene in human cancers. *Science*, **304**, 554.
- Santy, L. C., Ravichandran, K. S., & Casanova, J. E. (2005). The DOCK180/Elmo complex couples ARNO-mediated Arf6 activation to the downstream activation of Rac1. *Curr. Biol.*, **15**, 1749–1754.
- Sarbassov, D. D., Ali, S. M., & Sabatini, D. M. (2005a). Growing roles for the mTOR pathway. *Curr. Opin. Cell Biol.*, **17**, 596–603.
- Sarbassov, D. D., Guertin, D. A., Ali, S. M., & Sabatini, D. M. (2005b). Phosphorylation and regulation of Akt/PKB by the rictor-mTOR complex. *Science*, **307**, 1098–1101.
- Sasaki, A. T., Chun, C., Takeda, K., & Firtel, R. A. (2004). Localized Ras signaling at the leading edge regulates PI3K, cell polarity, and directional cell movement. *J. Cell Biol.*, **167**, 505–518.
- Sasaki, T., Irie-Sasaki, J., Jones, R. G., Oliveira-dos Santos, A. J., Stanford, W. L., Bolon, B., Wakeham, A., Itie, A., Bouchard, D., Kozieradzki, I., Joza, N., Mak, T. W., Ohashi, P. S., Suzuki, A., & Penninger, J. M. (2000). Function of PI3K γ in thymocyte development, T cell activation, and neutrophil migration. *Science*, **287**, 1040–1046.
- Schlessinger, J. (2000). Cell signaling by receptor tyrosine kinases. *Cell*, **103**, 211–225.
- Schultz, J., Milpetz, F., Bork, P., & Ponting, C. P. (1998). SMART, a simple modular architecture research tool: Identification of signaling domains. *Proc. Natl. Acad. Sci. U.S.A.*, **95**, 5857–5864.
- Scott, J. K., Huang, S. F., Gangadhar, B. P., Samoriski, G. M., Clapp, P., Gross, R. A., Tausig, R., & Smrcka, A. V. (2001). Evidence that a protein-protein interaction 'hot spot' on

- heterotrimeric G protein $\beta\gamma$ subunits is used for recognition of a subclass of effectors. *EMBO J.*, **20**, 767–776.
- Sen, G. L. & Blau, H. M. (2006). A brief history of RNAi: the silence of the genes. *FASEB J.*, **20**, 1293–1299.
- Sengeløv, H., Kjeldsen, L., Diamond, M. S., Springer, T. A., & Borregaard, N. (1993). Subcellular localization and dynamics of Mac-1 ($\alpha_m\beta_2$) in human neutrophils. *J. Clin. Invest.*, **92**, 1467–1476.
- Servant, G., Weiner, O. D., Herzmark, P., Balla, T., Sedat, J. W., & Bourne, H. R. (2000). Polarization of chemoattractant receptor signaling during neutrophil chemotaxis. *Science*, **287**, 1037–1040.
- Shaw, R. J. & Cantley, L. C. (2006). Ras, PI(3)K and mTOR signalling controls tumor cell growth. *Nature*, **441**, 424–430.
- Shekar, S. C., Wu, H., Fu, Z., Yip, S. C., Nagajyothi, Cahill, S. M., Girvin, M. E., & Backer, J. M. (2005). Mechanism of constitutive phosphoinositide 3-kinase activation by oncogenic mutants of the p85 regulatory subunit. *J. Biol. Chem.*, **280**, 27850–27855.
- Shepherd, M. C., Baillie, G. S., Stirling, D. I., & Houslay, M. D. (2004). Remodelling of the PDE4 cAMP phosphodiesterase isoform profile upon monocyte-macrophage differentiation of human U937 cells. *Br. J. Pharmacol.*, **142**, 339–351.
- Shinohara, M., Terada, Y., Iwamatsu, A., Shinohara, A., Mochizuki, N., Higuchi, M., Gotoh, Y., Ihara, S., Nagata, S., Itoh, H., Fukui, Y., & Jessberger, R. (2002). SWAP-70 is a guanine-nucleotide exchange factor that mediates signalling of membrane ruffling. *Nature*, **416**, 759–763.
- Sleeman, M. W., Wortley, K. E., Lai, K.-M. V., Gowen, L. C., Kintner, J., O’Kline, W., Garcia, K., Sitt, T. N., Yancopoulos, G. D., Wiegand, S. J., & Glass, D. J. (2005). Absence of the lipid phosphatase SHIP2 confers resistance to dietary obesity. *Nat. Med.*, **11**, 199–205.
- Soltoff, S. P. & Cantley, L. C. (1996). p120^{cb1} is a cytosolic adapter protein that associates with phosphoinositide 3-kinase in response to epidermal growth factor in PC12 and other cells. *J. Biol. Chem.*, **271**, 563–567.
- Songyang, Z., Shoelson, S. E., Chaudhuri, M., Gish, G., Pawson, T., Haser, W. G., King, F., Roberts, T., Ratnofsky, S., Lechleider, R. J., Neel, B. G., Birge, R. B., Fajardo, J. E., Chou, M. M., Hanafusa, H., Schaffhausen, B., & Cantley, L. C. (1993). SH2 domains recognize specific phosphopeptide sequences. *Cell*, **72**, 767–778.

REFERENCES

- Sprang, S. R. (1997). G protein mechanisms: Insights from structural analysis. *Annu. Rev. Biochem.*, **66**, 639–678.
- Stenmark, H., Aasland, R., & Driscoll, P. C. (2002). The phosphatidylinositol 3-phosphate-binding FYVE finger. *FEBS Lett.*, **513**, 77–84.
- Stephens, L., Eguinoa, A., Corey, S., Jackson, T., & Hawkins, P. T. (1993). Receptor stimulated accumulation of phosphatidylinositol (3,4,5)-trisphosphate by G-protein mediated pathways in human myeloid derived cells. *EMBO J.*, **12**, 2265–2273.
- Stephens, L. R., Eguinoa, A., Erdjument-Bromage, H., Lui, M., Cooke, F., Coadwell, J., Smrcka, A. S., Thelen, M., Cadwallader, K., Tempst, P., & Hawkins, P. T. (1997). The $G\beta\gamma$ sensitivity of a PI3K is dependent upon a tightly associated adaptor, p101. *Cell*, **89**, 105–114.
- Stoyanov, B., Volinia, S., Hanck, T., Rubio, I., Loubtchenkov, M., Malek, D., Stoyanova, S., Vanhaesebroeck, B., Dhand, R., Nürnberg, B., Gierschik, P., Seedorf, K., Hsuan, J. J., Waterfield, M. D., & Wetzker, R. (1995). Cloning and characterization of a G protein-activated human phosphoinositide 3-kinase. *Science*, **269**, 690–693.
- Stoyanova, S., Bulgarelli-Leva, G., Kirsch, C., Hanck, T., Klinger, R., Wetzker, R., & Wymann, M. P. (1997). Lipid kinase and protein kinase activities of G-protein-coupled phosphoinositide 3-kinase γ : structure–activity analysis and interactions with wortmannin. *Biochem. J.*, **324**, 489–495.
- Stryer, L. (1978). Fluorescence energy transfer as a spectroscopic ruler. *Annu. Rev. Biochem.*, **47**, 819–846.
- Suire, S., Hawkins, P. T., & Stephens, L. R. (2002). Activation of phosphoinositide 3-kinase γ by Ras. *Curr. Biol.*, **12**, 1068–1075.
- Suire, S., Coadwell, J., Ferguson, G. J., Davidson, K., Hawkins, P., & Stephens, L. (2005). p84, a new $G\beta\gamma$ -activated regulatory subunit of the type IB phosphoinositide 3-kinase p110 γ . *Curr. Biol.*, **15**, 566–570.
- Suire, S., Condliffe, A. M., Ferguson, G. J., Ellson, C. D., Guillou, H., Davidson, K., Welch, H., Coadwell, J., Turner, M., Chilvers, E. R., Hawkins, P. T., & Stephens, L. (2006). $G\beta\gamma$ s and the Ras binding domain of p110 γ are both important regulators of PI3K γ signalling in neutrophils. *Nat. Cell Biol.*, **8**, 1303–1309.
- Takesono, A., Finkelstein, L. D., & Schwartzberg, P. L. (2002). Beyond calcium: new signaling pathways for Tec family kinases. *J. Cell Sci.*, **115**, 3039–3048.

- Thomas, C. C., Deak, M., Alessi, D. R., & van Aalten, D. M. (2002). High-resolution structure of the pleckstrin homology domain of protein kinase B/Akt bound to phosphatidylinositol (3,4,5)-trisphosphate. *Curr. Biol.*, **12**, 1256–1262.
- Thompson, J. D., Higgins, D. G., & Gibson, T. J. (1994). CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res.*, **22**, 4673–4680.
- Thompson, W. J. & Appleman, M. M. (1971). Multiple cyclic nucleotide phosphodiesterase activities from rat brain. *Biochemistry*, **10**, 311–316.
- Touhara, K., Inglese, J., Pitcher, J. A., Shaw, G., & Lefkowitz, R. J. (1994). Binding of G protein $\beta\gamma$ -subunits to pleckstrin homology domains. *J. Biol. Chem.*, **269**, 10217–10220.
- Tovar, C., Rosinski, J., Filipovic, Z., Higgins, B., Kolinsky, K., Hilton, H., Zhao, X., Vu, B. T., Qing, W., Packman, K., Myklebost, O., Heimbrook, D. C., & Vassilev, L. T. (2006). Small-molecule MDM2 antagonists reveal aberrant p53 signaling in cancer: Implications for therapy. *Proc. Natl. Acad. Sci. U.S.A.*, **103**, 1888–1893.
- Traynor-Kaplan, A. E., Harris, A. L., Thompson, B. L., Taylor, P., & Sklar, L. A. (1988). An inositol tetrakisphosphate-containing phospholipid in activated neutrophils. *Nature*, **334**, 353–356.
- Tsien, R. Y. (1998). The green fluorescent protein. *Annu. Rev. Biochem.*, **67**, 509–544.
- Turner, S. J., Domin, J., Waterfield, M. D., Ward, S. G., & Westwick, J. (1998). The CC chemokine monocyte chemoattractant peptide-1 activates both the class I p85/p110 phosphatidylinositol 3-kinase and the class II PI3K-C2 α . *J. Biol. Chem.*, **273**, 25987–25995.
- Ueki, K., Fruman, D. A., Brachmann, S. M., Tseng, Y. H., Cantley, L. C., & Kahn, C. (2002). Molecular balance between the regulatory and catalytic subunits of phosphoinositide 3-kinase regulates cell signaling and survival. *Mol. Cell. Biol.*, **22**, 965–977.
- van Haastert, P. J. M. & Devreotes, P. N. (2004). Chemotaxis: Signalling the way forward. *Nat. Rev. Mol. Cell Biol.*, **5**, 626–634.
- Vanhaesebroeck, B., Leever, S. J., Ahmadi, K., Timms, J., Katso, R., Driscoll, P. C., Woscholski, R., Parker, P. J., & Waterfield, M. D. (2001). Synthesis and function of 3-phosphorylated inositol lipids. *Annu. Rev. Biochem.*, **70**, 535–602.
- Vanhaesebroeck, B., Welham, M. J., Kotani, K., Stein, R., Warne, P. H., Zvelebil, M. J., Higashi, K., Volinia, S., Downward, J., & Waterfield, M. D. (1997). p110 δ , a novel phosphoinositide 3-kinase in leukocytes. *Proc. Natl. Acad. Sci. U.S.A.*, **94**, 4330–4335.

REFERENCES

- Vanhaesebroeck, B., Higashi, K., Raven, C., Welham, M., Anderson, S., Brennan, P., G.Ward, S., & Waterfield, M. D. (1999). Autophosphorylation of p110 δ phosphoinositide 3-kinase: a new paradigm for the regulation of lipid kinases *in vitro* and *in vivo*. *EMBO J.*, **18**, 1292–1302.
- Vassilev, L. T., Vu, B. T., Graves, B., Carvajal, D., Podlaski, F., Filipovic, Z., Kong, N., Kammlott, U., Lukacs, C., Klein, C., Fotouhi, N., & Liu, E. A. (2004). *In vivo* activation of the p53 pathway by small-molecule antagonists of MDM2. *Science*, **303**, 844–848.
- Vaughn, J. L., Goodwin, R. H., Tompkins, G. J., & McCawley, P. (1977). The establishment of two cell lines from the insect *Spodoptera frugiperda* (Lepidoptera; Noctuidae). *In Vitro*, **13**, 213–217.
- Vecchione, C., Patrucco, E., Marino, G., Barberis, L., Poulet, R., Aretini, A., Maffei, A., Gentile, M. T., Storto, M., Azzolino, O., Brancaccio, M., Colussi, G. L., Bettarini, U., Altruda, F., Silengo, L., Tarone, G., Wymann, M. P., Hirsch, E., & Lembo, G. (2005). Protection from angiotensin II-mediated vasculotoxic and hypertensive response in mice lacking PI3K γ . *J. Exp. Med.*, **201**, 1217–1228.
- Viard, P., Exner, T., Maier, U., Mironneau, J., Nürnberg, B., & Macrez, N. (1999). G $\beta\gamma$ dimers stimulate vascular L-type Ca²⁺ channels via phosphoinositide 3-kinase. *FASEB J.*, **13**, 685–694.
- Viard, P., Butcher, A. J., Halet, G., Davies, A., Nürnberg, B., Hebllich, F., & Dolphin, A. C. (2004). PI3K promotes voltage-dependent calcium channel trafficking to the plasma membrane. *Nat. Neurosci.*, **7**, 939–946.
- Vlahos, C. J., Matter, W. F., Hui, K. Y., & Brown, R. F. (1994). A specific inhibitor of phosphatidylinositol 3-kinase, 2-(4-morpholinyl)-8-phenyl-4H-1-benzopyran-4-one (LY294002). *J. Biol. Chem.*, **269**, 5241–5248.
- Voigt, P. (2003). *Charakterisierung funktionell relevanter Domänen der regulatorischen Untereinheit der PI-3-Kinase γ* . Diplomarbeit, Institut für Biochemie, Fachbereich Biologie, Chemie, Pharmazie, Freie Universität Berlin.
- Volinia, S., Dhand, R., Vanhaesebroeck, B., MacDougall, L. K., Stein, R., Zvelebil, M. J., Domin, J., Panaretou, C., & Waterfield, M. D. (1995). A human phosphatidylinositol 3-kinase complex related to the yeast Vps34p-Vps15p protein sorting system. *EMBO J.*, **14**, 3339–3348.
- Walker, E. H., Pacold, M. E., Perisic, O., Stephens, L., Hawkins, P. T., Wymann, M. P., & Williams, R. L. (2000). Structural determinants of phosphoinositide 3-kinase inhibition by wortmannin, LY294002, querticin, myricetin, and staurosporine. *Mol. Cell*, **6**, 909–919.

- Walker, E. H., Perisic, O., Ried, C., Stephens, L., & Williams, R. L. (1999). Structural insights into phosphoinositide 3-kinase catalysis and signalling. *Nature*, **402**, 313–320.
- Wang, F., Herzmark, P., Weiner, O. D., Srinivasan, S., Servant, G., & Bourne, H. R. (2002). Lipid products of PI(3)Ks maintain persistent cell polarity and directed motility in neutrophils. *Nat. Cell Biol.*, **4**, 513–518.
- Wang, T., Pentylala, S., Rebecchi, M. J., & Scarlata, S. (1999). Differential association of the pleckstrin homology domains of phospholipases C- β_1 , C- β_2 , and C- δ_1 with lipid bilayers and the $\beta\gamma$ subunits of heterotrimeric G proteins. *Biochemistry*, **38**, 1517–1524.
- Ward, S. G. (2004). Do phosphoinositide 3-kinases direct lymphocyte navigation?. *Trends Immunol.*, **25**, 67–74.
- Weiner, O. D. (2002). Regulation of cell polarity during eukaryotic chemotaxis: the chemotactic compass. *Curr. Opin. Cell Biol.*, **14**, 196–202.
- Welch, H. C. E., Coadwell, W. J., Ellson, C. D., Ferguson, G. J., Andrews, S. R., Erdjument-Bromage, H., Tempst, P., Hawkins, P. T., & Stephens, L. R. (2002). P-Rex1, a PtdIns(3,4,5)P₃- and G $\beta\gamma$ -regulated guanine-nucleotide exchange factor for Rac. *Cell*, **108**, 809–821.
- Welch, H. C. E., Coadwell, W. J., Stephens, L. R., & Hawkins, P. T. (2003). Phosphoinositide 3-kinase-dependent activation of Rac. *FEBS Lett.*, **546**, 93–97.
- Wettschureck, N. & Offermanns, S. (2005). Mammalian G proteins and their cell type specific functions. *Physiol. Rev.*, **85**, 1159–1204.
- Wick, M. J., Ramos, F. J., Chen, H., Quon, M. J., Dong, L. Q., & Liu, F. (2003). Mouse 3-phosphoinositide-dependent protein kinase-1 undergoes dimerization and *trans*-phosphorylation in the activation loop. *J. Biol. Chem.*, **278**, 42913–42919.
- Williams, M. R., Arthur, J. S. C., Balendran, A., van der Kaay, J., Poli, V., Cohen, P., & Alessi, D. R. (2000). The role of 3-phosphoinositide-dependent protein kinase 1 in activating AGC kinases defined in embryonic stem cells. *Curr. Biol.*, **10**, 439–448.
- Wolfe, J. T., Wang, H., Howard, J., Garrison, J. C., & Barrett, P. Q. (2003). T-type calcium channel regulation by specific G-protein $\beta\gamma$ subunits. *Nature*, **424**, 209–213.
- Wong, W. & Scott, J. D. (2004). AKAP signalling complexes: Focal points in space and time. *Nat. Rev. Mol. Cell Biol.*, **5**, 959–970.
- Woodgett, J. R. (2001). Judging a protein by more than its name: GSK-3. *Science STKE*, http://www.stke.org/cgi/content/full/OC_sigtrans;2001/100/re12.

REFERENCES

- Woodgett, J. R. (2005). Recent advances in the protein kinase B signaling pathway. *Curr. Opin. Cell Biol.*, **17**, 150–157.
- Wullschleger, S., Loewith, R., & Hall, M. N. (2006). TOR signaling in growth and metabolism. *Cell*, **124**, 471–484.
- Wymann, M. P., Bulgarelli-Leva, G., Zvelebil, M. J., Pirola, L., Vanhaesebroeck, B., Waterfield, M. D., & Panayotou, G. (1996). Wortmannin inactivates phosphoinositide 3-kinase by covalent modification of Lys-802, a residue involved in the phosphate transfer reaction. *Mol. Cell. Biol.*, **16**, 1722–1733.
- Yang, J., Cron, P., Good, V. M., Thompson, V., Hemmings, B. A., & Barford, D. (2002a). Crystal structure of an activated Akt/protein kinase B ternary complex with GSK3-peptide and AMP-PNP. *Nat. Struct. Biol.*, **9**, 940–944.
- Yang, J., Cron, P., Thompson, V., Good, V. M., Hess, D., Hemmings, B. A., & Barford, D. (2002b). Molecular mechanism for the regulation of protein kinase B/Akt by hydrophobic motif phosphorylation. *Mol. Cell*, **9**, 1227–1240.
- Ye, K., Aghdasi, B., Luo, H. R., Moriarity, J. L., Wu, F. Y., Hong, J. J., Hurt, K. J., Bae, S. S., Suh, P.-G., & Snyder, S. H. (2002). Phospholipase C γ 1 is a physiological guanine nucleotide exchange factor for the nuclear GTPase PIKE. *Nature*, **415**, 541–544.
- Ye, K., Hurt, K. J., Wu, F. Y., Fang, M., Luo, H. R., Hong, J. J., Blackshaw, S., Ferris, C. D., & Snyder, S. H. (2000). PIKE: A nuclear GTPase that enhances PI3Kinase activity and is regulated by protein 4.1N. *Cell*, **103**, 919–930.
- Yu, J., Wjasow, C., & Backer, J. M. (1998a). Regulation of the p85 α /p110 α phosphatidylinositol 3'-kinase. Distinct roles for the N-terminal and C-terminal SH2 domains. *J. Biol. Chem.*, **273**, 30199–30203.
- Yu, J., Zhang, Y., McIlroy, J., Rordorf-Nikolic, T., Orr, G. A., & Backer, J. M. (1998b). Regulation of the p85/p110 phosphatidylinositol 3'-kinase: Stabilization and inhibition of the p110 α catalytic subunit by the p85 regulatory subunit. *Mol. Cell. Biol.*, **18**, 1379–1387.
- Zheng, Y., Bagrodia, S., & Cerione, R. A. (1994). Activation of phosphoinositide 3-kinase activity by Cdc42Hs binding to p85. *J. Biol. Chem.*, **269**, 18727–18730.
- Zimmermann, S. & Moelling, K. (1999). Phosphorylation and regulation of Raf by Akt (protein kinase B). *Science*, **286**, 1741–1744.