

8. Literaturverzeichnis

1. Wright J P, Potter P: Psyche and Soma: Physicians and Metaphysics on Mind-Body Problem from Antiquity to Enlightenment. Oxford, University Press, 2000
2. Bräutigam W, Christian P, von Rad M: Psychosomatische Medizin. Stuttgart, Georg Thieme, 1997
3. Kiecolt-Glaser J K, McGuire L, Robles T F, Glaser R. Psychoneuroimmunology and psychosomatic medicine: back to the future. *Psychosom Med* 2002; 64: 15-28.
4. Fleshner M, Laudenslager M L. Psychoneuroimmunology: then and now. *Behav Cogn Neurosci Rev* 2004; 3: 114-130.
5. Dantzer R. Somatization: A psychoneuroimmune perspective. *Psychoneuroendocrinology* 2005; 7: 7.
6. Koblenzer C S. Psychosomatic concepts in dermatology. A dermatologist-psychoanalyst's viewpoint. *Arch Dermatol* 1983; 119: 501-512.
7. Panconesi E, Hautmann G. Psychophysiology of stress in dermatology. The psychobiologic pattern of psychosomatics. *Dermatol Clin* 1996; 14: 399-421.
8. Black P H. Stress and the inflammatory response: A review of neurogenic inflammation. *Brain Behav Immun* 2002; 16: 622-653.
9. Carrasco G A, Van de Kar L D. Neuroendocrine pharmacology of stress. *Eur J Pharmacol* 2003; 463: 235-272.
10. Levin D M, Solomon G F. The discursive formation of the body in the history of medicine. *J Med Philos* 1990; 15: 515-537.
11. Elenkov I J, Wilder R L, Chrousos G P, Vizi E S. The sympathetic nerve--an integrative interface between two supersystems: the brain and the immune system. *Pharmacol Rev* 2000; 52: 595-638.
12. Frieri M. Neuroimmunology and inflammation: implications for therapy of allergic and autoimmune diseases. *Ann Allergy Asthma Immunol* 2003; 90: 34-40.
13. Lotti T, Hautmann G, Panconesi E. Neuropeptides in skin. *J Am Acad Dermatol* 1995; 33: 482-496.
14. Ansel J C, Kaynard A H, Armstrong C A, Olerud J, Bunnett N, Payan D. Skin-nervous system interactions. *J Invest Dermatol* 1996; 106: 198-204.
15. Levi-Montalcini R, Skaper S D, Dal Toso R, Petrelli L, Leon A. Nerve growth factor: from neurotrophin to neurokine. *Trends Neurosci* 1996; 19: 514-520.
16. Baluk P. Neurogenic inflammation in skin and airways. *J Investig Dermatol Symp Proc* 1997; 2: 76-81.
17. Paus R, Peters E M, Eichmüller S, Botchkarev V A. Neural mechanisms of hair growth control. *J Investig Dermatol Symp Proc* 1997; 2: 61-68.
18. Scholzen T, Armstrong C A, Bunnett N W, Luger T A, Olerud J E, Ansel J C. Neuropeptides in the skin: interactions between the neuroendocrine and the skin immune systems. *Exp Dermatol* 1998; 7: 81-96.
19. Bauer O, Razin E. Mast Cell-Nerve Interactions. *News Physiol Sci* 2000; 15: 213-218.
20. Levite M. Nerve-driven immunity. The direct effects of neurotransmitters on T-cell function. *Ann N Y Acad Sci* 2000; 917: 307-321.
21. Slominski A, Wortsman J. Neuroendocrinology of the skin. *Endocr Rev* 2000; 21: 457-487.
22. Ansel J C. Neuromediators and inflammation. *Exp Dermatol* 2001; 10: 352-353.
23. Steinhoff M, Stander S, Seeliger S, Ansel J C, Schmelz M, Luger T. Modern aspects of cutaneous neurogenic inflammation. *Arch Dermatol* 2003; 139: 1479-1488.

24. Alleva E, Santucci D. Psychosocial vs. "physical" stress situations in rodents and humans: role of neurotrophins. *Physiol Behav* 2001; 73: 313-320.
25. Blalock J E. The syntax of immune-neuroendocrine communication. *Immunol Today* 1994; 15: 504-511.
26. Misery L. Skin, immunity and the nervous system. *Br J Dermatol* 1997; 137: 843-850.
27. Arck P C. Stress and pregnancy loss: role of immune mediators, hormones and neurotransmitters. *Am J Reprod Immunol* 2001; 46: 117-123.
28. Peters E M J, Arck P C, Paus R. Review Article: Hair growth inhibition by psychoemotional stress: a model for neural mechanisms in hair growth control. *Exp Dermatol* 2005; (*in press*).
29. Henz B M, Kerl H, Sterry W: *Dermatologie Und Venerologie*. Berlin, Walter de Gruyter GmbH & Co.KG, 1997
30. Buhk H, Muthny F A. [Psychophysiologic and psychoneuroimmunologic studies in neurodermatitis. Overview and critical evaluation]. *Hautarzt* 1997; 48: 5-11.
31. Buske-Kirschbaum A, Hellhammer D H. Endocrine and immune responses to stress in chronic inflammatory skin disorders. *Ann N Y Acad Sci* 2003; 992: 231-240.
32. Tobin D J, Paus R. Graying: gerontobiology of the hair follicle pigmentary unit. *Exp Gerontol* 2001; 36: 29-54.
33. Bahmer F A. [The phenomena of sudden graying of the hair in the poem "Die Fusse im Feuer" by Conrad Ferdinand Meyer]. *Hautarzt* 2002; 53: 492-494.
34. Botchkarev V A, Eichmüller S, Johansson O, Paus R. Hair cycle-dependent plasticity of skin and hair follicle innervation in normal murine skin. *J Comp Neurol* 1997; 386: 379-395.
35. Müller-Röver S, Handjiski B, van der Veen C, Eichmuller S, Foitzik K, McKay I A, Stenn K S, Paus R. A comprehensive guide for the accurate classification of murine hair follicles in distinct hair cycle stages. *J Invest Dermatol* 2001; 117: 3-15.
36. Stenn K S, Paus R. Controls of hair follicle cycling. *Physiol Rev* 2001; 81: 449-494.
37. Peters E M J, Botchkarev V A, Moll I, Arck P, Paus R: Neurobiology of the Murine Pelage Hair Follicle. In: (eds.) K I Baumann, Z Halata , I Moll: *The Merkel cell - structure - development - function - and Merkel cell carcinoma. Proceedings of the International Merkel Cell Symposium Hamburg 2002.* , Springer-Verlag, Berlin, Heidelberg, New York, Barcelona, Hong Kong, London, Milan, Paris, Tokyo, 2003, pp 79-86
38. Takahashi K, Nakanishi S, Imamura S. Direct effects of cutaneous neuropeptides on adenylyl cyclase activity and proliferation in a keratinocyte cell line: stimulation of cyclic AMP formation by CGRP and VIP/PHM, and inhibition by NPY through G protein-coupled receptors. *J Invest Dermatol* 1993; 101: 646-651.
39. Paus R, Heinzelmann T, Robicsek S, Czarnetzki B M, Maurer M. Substance P stimulates murine epidermal keratinocyte proliferation and dermal mast cell degranulation *in situ*. *Arch Dermatol Res* 1995; 287: 500-502.
40. Schallreuter K U, Lemke K R, Pittelkow M R, Wood J H, Korner C, Malik R. Catecholamines in human keratinocyte differentiation. *J Invest Dermatol* 1995; 104: 953-957.
41. Bothwell M. Neurotrophin function in skin. *J Investig Dermatol Symp Proc* 1997; 2: 27-30.
42. Grando S A. Biological functions of keratinocyte cholinergic receptors. *J Invest Dermatol Symp Proc* 1997; 2: 41-48.
43. Huang I T, Lin W M, Shun C T, Hsieh S T. Influence of cutaneous nerves on keratinocyte proliferation and epidermal thickness in mice. *Neuroscience* 1999; 94: 965-973.
44. Pincelli C, Marconi A. Autocrine nerve growth factor in human keratinocytes. *J Dermatol Sci* 2000; 22: 71-79.
45. Kurzen H, Schallreuter K U. Novel aspects in cutaneous biology of acetylcholine synthesis and acetylcholine receptors. *Exp Dermatol* 2004; 4: 27-30.
46. Ibanez C F. Jekyll-Hyde neurotrophins: the story of proNGF. *Trends Neurosci* 2002; 25: 284-286.

47. Botchkarev V A, Botchkareva N V, Peters E M, Paus R. Epithelial growth control by neurotrophins: leads and lessons from the hair follicle. *Prog Brain Res* 2004; 146: 493-513.
48. Donaldson D J, Mahan J T. Influence of catecholamines on epidermal cell migration during wound closure in adult newts. *Comp Biochem Physiol C* 1984; 78: 267-270.
49. Parenti A, Amerini S, Ledda F, Maggi C A, Ziche M. The tachykinin NK1 receptor mediates the migration-promoting effect of substance P on human skin fibroblasts in culture. *Naunyn Schmiedebergs Arch Pharmacol* 1996; 353: 475-481.
50. Springer J, Geppetti P, Fischer A, Groneberg D A. Calcitonin gene-related peptide as inflammatory mediator. *Pulm Pharmacol Ther* 2003; 16: 121-130.
51. Chernyavsky A I, Arredondo J, Marubio L M, Grando S A. Differential regulation of keratinocyte chemokinesis and chemotaxis through distinct nicotinic receptor subtypes. *J Cell Sci* 2004; 117: 5665-5679.
52. Pascual D W, Bost K L, Xu-Amano J, Kiyono H, McGhee J R. The cytokine-like action of substance P upon B cell differentiation. *Reg Immunol* 1992; 4: 100-104.
53. Luger T A. Neuromediators--a crucial component of the skin immune system. *J Dermatol Sci* 2002; 30: 87-93.
54. Toyoda M, Nakamura M, Morohashi M. Neuropeptides and sebaceous glands. *Eur J Dermatol* 2002; 12: 422-427.
55. Grantham R, Fridkinb M, Rubinrautb S, Gozes I. VIP-derived sequences modified by N-terminal stearyl moiety induce cell death: the human keratinocyte as a model. *FEBS Lett* 2000; 475: 71-77.
56. Del Rey A, Kabiersch A, Petzoldt S, Besedovsky H O. Sympathetic abnormalities during autoimmune processes: potential relevance of noradrenaline-induced apoptosis. *Ann N Y Acad Sci* 2003; 992: 158-167.
57. Santoni G, Amantini C, Lucciarini R, Perfumi M, Pompei P, Piccoli M. Neonatal capsaicin treatment affects rat thymocyte proliferation and cell death by modulating substance P and neurokinin-1 receptor expression. *Neuroimmunomodulation* 2004; 11: 160-172.
58. Schallreuter K U. Epidermal adrenergic signal transduction as part of the neuronal network in the human epidermis. *J Investig Dermatol Symp Proc* 1997; 2: 37-40.
59. Bae S, Matsunaga Y, Tanaka Y, Katayama I. Autocrine induction of substance P mRNA and peptide in cultured normal human keratinocytes. *Biochem Biophys Res Commun* 1999; 263: 327-333.
60. Peters E M, Handjiski B, Kuhlmei A, Hagen E, Bielas H, Braun A, Klapp B F, Paus R, Arck P C. Neurogenic inflammation in stress-induced termination of murine hair growth is promoted by nerve growth factor. *Am J Pathol* 2004; 165: 259-271.
61. Peters E M J, Ericson M, Hosoi J, Seiffert K, Hordinsky M, Ansel J C, Paus R, Scholzen T E. Neuropeptidergic controls in cutaneous biology: Physiological and clinical significance. *J Invest Dermatol* 2005; (*invited review, in preparation*).
62. Ising H, Braun C. Acute and chronic endocrine effects of noise: Review of the research conducted at the Institute for Water, Soil and Air Hygiene. *Noise Health* 2000; 2: 7-24.
63. Kimata H. Enhancement of allergic skin wheal responses in patients with atopic eczema/dermatitis syndrome by playing video games or by a frequently ringing mobile phone. *Eur J Clin Invest* 2003; 33: 513-517.
64. Ising H, Kruppa B. Health effects caused by noise: evidence in the literature from the past 25 years. *Noise Health* 2004; 6: 5-13.
65. Hebert S, Beland R, Dionne-Fournelle O, Crete M, Lupien S J. Physiological stress response to video-game playing: the contribution of built-in music. *Life Sci* 2005; 76: 2371-2380.
66. Niemann H, Maschke C, Hecht K. [Noise induced annoyance and morbidity. Results from the pan European LARES-survey]. *Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz* 2005; 48: 315-328.
67. Evans G W. The built environment and mental health. *J Urban Health* 2003; 80: 536-555.

68. Melamed S, Fried Y, Froom P. The joint effect of noise exposure and job complexity on distress and injury risk among men and women: the cardiovascular occupational risk factors determination in Israel study. *J Occup Environ Med* 2004; 46: 1023-1032.
69. Schulz C, Babisch W, Becker K, Durkop J, Rosskamp E, Seiwert M, Steiner M, Szewzyk R, Ullrich D, Englert N, Seifert B, Eis D. [Environmental Survey for Children-- the environmental module of KiGGS. I. Design and research program]. *Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz* 2004; 47: 1066-1072.
70. Maslach C, Schaufeli W B, Leiter M P. Job burnout. *Annu Rev Psychol* 2001; 52: 397-422.
71. McVicar A. Workplace stress in nursing: a literature review. *J Adv Nurs* 2003; 44: 633-642.
72. Vezina M, Bourbonnais R, Brisson C, Trudel L. Workplace prevention and promotion strategies. *Healthc Pap* 2004; 5: 32-44.
73. Selye H: The Stress of Life. New York, McGraw-Hill, 1976
74. Mechiel Korte S, De Boer S F. A robust animal model of state anxiety: fear-potentiated behaviour in the elevated plus-maze. *Eur J Pharmacol* 2003; 463: 163-175.
75. Levine S. Developmental determinants of sensitivity and resistance to stress. *Psychoneuroendocrinology* 2005; 13: 13.
76. Bracha H S. Freeze, flight, fight, fright, faint: adaptationist perspectives on the acute stress response spectrum. *CNS Spectr* 2004; 9: 679-685.
77. Herman J P, Figueiredo H, Mueller N K, Ulrich-Lai Y, Ostrander M M, Choi D C, Cullinan W E. Central mechanisms of stress integration: hierarchical circuitry controlling hypothalamo-pituitary-adrenocortical responsiveness. *Front Neuroendocrinol* 2003; 24: 151-180.
78. De Kloet E R. Hormones and the stressed brain. *Ann N Y Acad Sci* 2004; 1-15.
79. Singh L K, Pang X, Alexacos N, Letourneau R, Theoharides T C. Acute immobilization stress triggers skin mast cell degranulation via corticotropin releasing hormone, neuropeptides, and substance P: A link to neurogenic skin disorders. *Brain Behav Immun* 1999; 13: 225-239.
80. Hosoi J, Tanida M, Tsuchiya T. Regulation of skin mast cells by stress and odorant. *J Invest Dermatol* 2002; to be published in Juli: (abstract).
81. Eskandari F, Sternberg E M. Neural-immune interactions in health and disease. *Ann N Y Acad Sci* 2002; 966: 20-27.
82. Dhabhar F S. Stress, leukocyte trafficking, and the augmentation of skin immune function. *Ann N Y Acad Sci* 2003; 992: 205-217.
83. Dhabhar F S, Satoskar A R, Bluethmann H, David J R, McEwen B S. Stress-induced enhancement of skin immune function: A role for gamma interferon. *Proc Natl Acad Sci U S A* 2000; 97: 2846-2851.
84. Li T, Harada M, Tamada K, Abe K, Nomoto K. Repeated restraint stress impairs the antitumor T cell response through its suppressive effect on Th1-type CD4+ T cells. *Anticancer Res* 1997; 17: 4259-4268.
85. Iwakabe K, Shimada M, Ohta A, Yahata T, Ohmi Y, Habu S, Nishimura T. The restraint stress drives a shift in Th1/Th2 balance toward Th2-dominant immunity in mice. *Immunol Lett* 1998; 62: 39-43.
86. Buske-Kirschbaum A, Geiben A, Hollig H, Morschhauser E, Hellhammer D. Altered responsiveness of the hypothalamus-pituitary-adrenal axis and the sympathetic adrenomedullary system to stress in patients with atopic dermatitis. *J Clin Endocrinol Metab* 2002; 87: 4245-4251.
87. Buske-Kirschbaum A, Jobst S, Hellhammer D H. Altered reactivity of the hypothalamus-pituitary-adrenal axis in patients with atopic dermatitis: pathologic factor or symptom? *Ann N Y Acad Sci* 1998; 840: 747-754.
88. Buske-Kirschbaum A, Fischbach S, Rauh W, Hanker J, Hellhammer D. Increased responsiveness of the hypothalamus-pituitary-adrenal (HPA) axis to stress in newborns with atopic disposition. *Psychoneuroendocrinology* 2004; 29: 705-711.

89. Mantyh P W. Neurobiology of substance P and the NK1 receptor. *J Clin Psychiatry* 2002; 11: 6-10.
90. Duffy R A. Potential therapeutic targets for neurokinin-1 receptor antagonists. *Expert Opin Emerg Drugs* 2004; 9: 9-21.
91. McLean S. Do substance P and the NK1 receptor have a role in depression and anxiety? *Curr Pharm Des* 2005; 11: 1529-1547.
92. Seidah N G, Benjannet S, Pareek S, Savaria D, Hamelin J, Goulet B, Laliberte J, Lazure C, Chretien M, Murphy R A. Cellular processing of the nerve growth factor precursor by the mammalian pro-protein convertases. *Biochem J* 1996; 314: 951-960.
93. Cheng L, Khan M, Mudge A W. Calcitonin gene-related peptide promotes Schwann cell proliferation. *J Cell Biol* 1995; 129: 789-796.
94. O'Connor T M, O'Connell J, O'Brien D I, Goode T, Bredin C P, Shanahan F. The role of substance P in inflammatory disease. *J Cell Physiol* 2004; 201: 167-180.
95. Steinhoff M, Buddenkotte J, Shpacovitch V, Rattenholl A, Moormann C, Vergnolle N, Luger T A, Hollenberg M D. Proteinase-activated receptors: transducers of proteinase-mediated signaling in inflammation and immune response. *Endocr Rev* 2005; 26: 1-43.
96. Ziche M, Parenti A, Amerini S, Zawieja D, Maggi C A, Ledda F. Effect of the non-peptide blocker (+/-) CP 96,345 on the cellular mechanism involved in the response to NK1 receptor stimulation in human skin fibroblasts. *Neuropeptides* 1996; 30: 345-354.
97. Lindsey K Q, Caughman S W, Olerud J E, Bunnett N W, Armstrong C A, Ansel J C. Neural regulation of endothelial cell-mediated inflammation. *J Investig Dermatol Symp Proc* 2000; 5: 74-78.
98. Quinlan K L, Naik S M, Cannon G, Armstrong C A, Bunnett N W, Ansel J C, Caughman S W. Substance P activates coincident NF-AT- and NF-kappa B-dependent adhesion molecule gene expression in microvascular endothelial cells through intracellular calcium mobilization. *J Immunol* 1999; 163: 5656-5665.
99. Fuller R W, Conradson T B, Dixon C M, Crossman D C, Barnes P J. Sensory neuropeptide effects in human skin. *Br J Pharmacol* 1987; 92: 781-788.
100. Ansel J C, Armstrong C A, Song I, Quinlan K L, Olerud J E, Caughman S W, Bunnett N W. Interactions of the skin and nervous system. *J Investig Dermatol Symp Proc* 1997; 2: 23-26.
101. Wallengren J. Vasoactive peptides in the skin. *J Investig Dermatol Symp Proc* 1997; 2: 49-55.
102. Carlton S M, Zhou S, Coggeshall R E. Localization and activation of substance P receptors in unmyelinated axons of rat glabrous skin. *Brain Res* 1996; 734: 103-108.
103. Andoh T, Nagasawa T, Satoh M, Kuraishi Y. Substance P induction of itch-associated response mediated by cutaneous NK1 tachykinin receptors in mice. *J Pharmacol Exp Ther* 1998; 286: 1140-1145.
104. Heyer G R, Hornstein O P. Recent studies of cutaneous nociception in atopic and non-atopic subjects. *J Dermatol* 1999; 26: 77-86.
105. Yamamoto T, Sakashita Y. The role of the spinal opioid receptor like1 receptor, the NK-1 receptor, and cyclooxygenase-2 in maintaining postoperative pain in the rat. *Anesth Analg* 1999; 89: 1203-1208.
106. Payan D G, Brewster D R, Missirian-Bastian A, Goetzl E J. Substance P recognition by a subset of human T lymphocytes. *J Clin Invest* 1984; 74: 1532-1539.
107. Tanaka T, Danno K, Ikai K, Imamura S. Effects of substance P and substance K on the growth of cultured keratinocytes. *J Investig Dermatol* 1988; 90: 399-401.
108. Pincelli C, Fantini F, Romualdi P, Sevignani C, Lesa G, Benassi L, Giannetti A. Substance P is diminished and vasoactive intestinal peptide is augmented in psoriatic lesions and these peptides exert disparate effects on the proliferation of cultured human keratinocytes. *J Investig Dermatol* 1992; 98: 421-427.
109. Morbidelli L, Maggi C A, Ziche M. Effect of selective tachykinin receptor antagonists on the growth of human skin fibroblasts. *Neuropeptides* 1993; 24: 335-341.

110. Kahler C M, Sitte B A, Reinisch N, Wiedermann C J. Stimulation of the chemotactic migration of human fibroblasts by substance P. *Eur J Pharmacol* 1993; 249: 281-286.
111. Feistritzer C, Clausen J, Sturn D H, Djanani A, Gunsilius E, Wiedermann C J, Kahler C M. Natural killer cell functions mediated by the neuropeptide substance P. *Regul Pept* 2003; 116: 119-126.
112. Walsh D A, Hu D E, Mapp P I, Polak J M, Blake D R, Fan T P. Innervation and neurokinin receptors during angiogenesis in the rat sponge granuloma. *Histochem J* 1996; 28: 759-769.
113. Seegers H C, Hood V C, Kidd B L, Cruwys S C, Walsh D A. Enhancement of angiogenesis by endogenous substance P release and neurokinin-1 receptors during neurogenic inflammation. *J Pharmacol Exp Ther* 2003; 306: 8-12.
114. Lowman M A, Benyon R C, Church M K. Characterization of neuropeptide-induced histamine release from human dispersed skin mast cells. *Br J Pharmacol* 1988; 95: 121-130.
115. Church M K, Lowman M A, Robinson C, Holgate S T, Benyon R C. Interaction of neuropeptides with human mast cells. *Int Arch Allergy Appl Immunol* 1989; 88: 70-78.
116. Calvo C F, Chavanel G, Senik A. Substance P enhances IL-2 expression in activated human T cells. *J Immunol* 1992; 148: 3498-3504.
117. Marriott I, Bost K L. IL-4 and IFN-gamma up-regulate substance P receptor expression in murine peritoneal macrophages. *J Immunol* 2000; 165: 182-191.
118. Marriott I, Bost K L. Substance P receptor mediated macrophage responses. *Adv Exp Med Biol* 2001; 493: 247-254.
119. Viac J, Gueniche A, Doutremepuich J D, Reichert U, Claudy A, Schmitt D. Substance P and keratinocyte activation markers: an in vitro approach. *Arch Dermatol Res* 1996; 288: 85-90.
120. Quinlan K L, Song I S, Naik S M, Letran E L, Olerud J E, Bunnett N W, Armstrong C A, Caughman S W, Ansel J C. VCAM-1 expression on human dermal microvascular endothelial cells is directly and specifically up-regulated by substance P. *J Immunol* 1999; 162: 1656-1661.
121. Kang B N, Kim H J, Jeong K S, Park S J, Kim S H, Kim S R, Kim T H, Ryu S Y. Regulation of leukocyte function-associated antigen 1-mediated adhesion by somatostatin and substance P in mouse spleen cells. *Neuroimmunomodulation* 2004; 11: 84-92.
122. Quinlan K L, Song I S, Bunnett N W, Letran E, Steinhoff M, Harten B, Olerud J E, Armstrong C A, Wright Caughman S, Ansel J C. Neuropeptide regulation of human dermal microvascular endothelial cell ICAM-1 expression and function. *Am J Physiol* 1998; 275: C1580-1590.
123. Levite M, Cahalon L, Herschkoviz R, Steinman L, Lider O. Neuropeptides, via specific receptors, regulate T cell adhesion to fibronectin. *J Immunol* 1998; 160: 993-1000.
124. Peters E M, Botchkarev V A, Botchkareva N V, Tobin D J, Paus R. Hair-cycle-associated remodeling of the peptidergic innervation of murine skin, and hair growth modulation by neuropeptides. *J Invest Dermatol* 2001; 116: 236-245.
125. Iumatov E A, Poppai M, Ratsak R. [Changes in the substance P content of the blood and hypothalamus during experimental emotional stress]. *Biull Eksp Biol Med* 1985; 99: 397-399.
126. Culman J, Unger T. Central tachykinins: mediators of defence reaction and stress reactions. *Can J Physiol Pharmacol* 1995; 73: 885-891.
127. Chancellor-Freeland C, Zhu G F, Kage R, Beller D I, Leeman S E, Black P H. Substance P and stress-induced changes in macrophages. *Ann N Y Acad Sci* 1995; 771: 472-484.
128. Arck P C, Handjiski B, Peters E M, Peter A S, Hagen E, Fischer A, Klapp B F, Paus R. Stress inhibits hair growth in mice by induction of premature catagen development and deleterious perifollicular inflammatory events via neuropeptide substance P-dependent pathways. *Am J Pathol* 2003; 162: 803-814.
129. Peters E M, Kuhlmei A, Tobin D J, Muller-Rover S, Klapp B F, Arck P C. Stress exposure modulates peptidergic innervation and degranulates mast cells in murine skin. *Brain Behav Immun* 2005; 19: 252-262.

130. Tobin D J, Nabarro G, Baart de la Faille H, van Vloten W A, van der Putte S C, Schuurman H J. Increased number of immunoreactive nerve fibers in atopic dermatitis. *J Allergy Clin Immunol* 1992; 90: 613-622.
131. Scholzen T E, Steinhoff M, Bonaccorsi P, Klein R, Amadesi S, Geppetti P, Lu B, Gerard N P, Olerud J E, Luger T A, Bunnett N W, Grady E F, Armstrong C A, Ansel J C. Neutral endopeptidase terminates substance P-induced inflammation in allergic contact dermatitis. *J Immunol* 2001; 166: 1285-1291.
132. Toyoda M, Nakamura M, Makino T, Hino T, Kagoura M, Morohashi M. Nerve growth factor and substance P are useful plasma markers of disease activity in atopic dermatitis. *Br J Dermatol* 2002; 147: 71-79.
133. Huang C H, Kuo I C, Xu H, Lee Y S, Chua K Y. Mite allergen induces allergic dermatitis with concomitant neurogenic inflammation in mouse. *J Invest Dermatol* 2003; 121: 289-293.
134. Jarvikallio A, Harvima I T, Naukkarinen A. Mast cells, nerves and neuropeptides in atopic dermatitis and nummular eczema. *Arch Dermatol Res* 2003; 295: 2-7.
135. Ohmura T, Tsunenari I, Hayashi T, Satoh Y, Konomi A, Nanri H, Kawachi M, Morikawa M, Kadota T, Satoh H. Role of Substance P in an NC/Nga Mouse Model of Atopic Dermatitis-Like Disease. *Int Arch Allergy Immunol* 2004; 133: 389-397.
136. Weigent D A, Carr D J, Blalock J E. Bidirectional communication between the neuroendocrine and immune systems. Common hormones and hormone receptors. *Ann N Y Acad Sci* 1990; 579: 17-27.
137. Gordon D J, Ostlere L S, Holden C A. Neuropeptide modulation of Th1 and Th2 cytokines in peripheral blood mononuclear leucocytes in atopic dermatitis and non-atopic controls. *Br J Dermatol* 1997; 137: 921-927.
138. Lambrecht B N, Germonpre P R, Everaert E G, Carro-Muino I, De Veerman M, de Felipe C, Hunt S P, Thielemans K, Joos G F, Pauwels R A. Endogenously produced substance P contributes to lymphocyte proliferation induced by dendritic cells and direct TCR ligation. *Eur J Immunol* 1999; 29: 3815-3825.
139. Lambrecht B N. Immunologists getting nervous: neuropeptides, dendritic cells and T cell activation. *Respir Res* 2001; 2: 133-138.
140. Arck P C, Handjiski B, Kuhlmei A, Peters E M, Knackstedt M, Peter A, Hunt S P, Klapp B F, Paus R. Mast cell deficient and neurokinin-1 receptor knockout mice are protected from stress-induced hair growth inhibition. *J Mol Med* 2005; 10: 10.
141. Paus R, Luftl M, Czarnetzki B M. Nerve growth factor modulates keratinocyte proliferation in murine skin organ culture. *Br J Dermatol* 1994; 130: 174-180.
142. Muller Y, Tangre K, Clos J. Autocrine regulation of apoptosis and bcl-2 expression by nerve growth factor in early differentiating cerebellar granule neurons involves low affinity neurotrophin receptor [In Process Citation]. *Neurochem Int* 1997; 31: 177-191.
143. Virdee K, Xue L, Hemmings B A, Goemans C, Heumann R, Tolokovsky A M. Nerve growth factor-induced PKB/Akt activity is sustained by phosphoinositide 3-kinase dependent and independent signals in sympathetic neurons. *Brain Res* 1999; 837: 127-142.
144. Edstrom A, Ekstrom P A. Role of phosphatidylinositol 3-kinase in neuronal survival and axonal outgrowth of adult mouse dorsal root ganglia explants. *J Neurosci Res* 2003; 74: 726-735.
145. Huang E J, Reichardt L F. Trk receptors: roles in neuronal signal transduction. *Annu Rev Biochem* 2003; 72: 609-642.
146. Segal R A. Selectivity in neurotrophin signaling: theme and variations. *Annu Rev Neurosci* 2003; 26: 299-330.
147. Teng K K, Hempstead B L. Neurotrophins and their receptors: signaling trios in complex biological systems. *Cell Mol Life Sci* 2004; 61: 35-48.
148. Roux P P, Barker P A. Neurotrophin signaling through the p75 neurotrophin receptor. *Prog Neurobiol* 2002; 67: 203-233.

149. Sachs M, Weidner K M, Brinkmann V, Walther I, Obermeier A, Ullrich A, Birchmeier W. Motogenic and morphogenic activity of epithelial receptor tyrosine kinases. *J Cell Biol* 1996; 133: 1095-1107.
150. Botchkarev V A, Yaar M, Botchkareva N V, Peters E M J, Marconi A, Raychaudhuri S P, Raychaudhuri S P, Paus R, Pincelli C. Neurotrophins in Skin Biology. *J Invest Dermatol* 2005: (*invited review, under review*).
151. Barker P A. p75NTR is positively promiscuous: novel partners and new insights. *Neuron* 2004; 42: 529-533.
152. Dechant G, Barde Y A. The neurotrophin receptor p75(NTR): novel functions and implications for diseases of the nervous system. *Nat Neurosci* 2002; 5: 1131-1136.
153. Peters E M J, Hansen M G, Liezman C, Overall R W, Nakamura M, Hagen E, Klapp B F, Arck P C, Paus P. p75 neurotrophin receptor-mediated signalling promotes human hair follicle regression (catagen). *Am J Pathol* 2005: (*under review*).
154. Botchkarev V A, Eichmüller S, Peters E M, Pietsch P, Johansson O, Maurer M, Paus R. A simple immunofluorescence technique for simultaneous visualization of mast cells and nerve fibers reveals selectivity and hair cycle--dependent changes in mast cell-nerve fiber contacts in murine skin. *Arch Dermatol Res* 1997; 289: 292-302.
155. Peters E M, Maurer M, Botchkarev V A, Gordon D S, Paus R. Hair growth-modulation by adrenergic drugs. *Exp Dermatol* 1999; 8: 274-281.
156. Peters E M J, Botchkarev V A, Paus R: Neurotrophins act as regulators of hair follicle morphogenesis and cycling. In: (eds.) D Van Neste: *Hair Science and Technology*, Skinterface, Tournai-Belgium, 2003, pp 353-366
157. Peters E M, Hansen M G, Overall R W, Nakamura M, Pertile P, Klapp B F, Arck P C, Paus R. Control of human hair growth by neurotrophins: brain-derived neurotrophic factor inhibits hair shaft elongation, induces catagen, and stimulates follicular transforming growth factor beta2 expression. *J Invest Dermatol* 2005; 124: 675-685.
158. Peters E M J, Hendrix S, Götz G, Klapp B F, Arck P C, Paus R. Nerve growth factor and its precursor regulate hair cycle progression in mice. *J Histochem Cytochem* 2005: (*accepted*).
159. Arck P C, Handjiski B, Peters E M, Hagen E, Klapp B F, Paus R. Topical minoxidil counteracts stress-induced hair growth inhibition in mice. *Exp Dermatol* 2003; 12: 580-590.
160. Peters E M J, Tobin D J, Hendrix S, Hagen E, Peter A S, Knackstedt M, Klapp B F, Arck P C. Neurogenic inflammation drives the worsening of peripheral inflammatory disease by stress. *J Clin Invest* 2005: (*submitted*).
161. Huntley A, White A R, Ernst E. Relaxation therapies for asthma: a systematic review. *Thorax* 2002; 57: 127-131.
162. Peters E M, Botchkarev V A, Muller-Rover S, Moll I, Rice F L, Paus R. Developmental timing of hair follicle and dorsal skin innervation in mice. *J Comp Neurol* 2002; 448: 28-52.
163. Ek L, Theodorsson E. Tachykinins and calcitonin gene-related peptide in oxazolone-induced allergic contact dermatitis in mice. *J Invest Dermatol* 1990; 94: 761-763.
164. Holzer P. Neurogenic vasodilatation and plasma leakage in the skin. *Gen Pharmacol* 1998; 30: 5-11.
165. Caulfield J P, el Lati S, Thomas G, Church M K. Dissociated human foreskin mast cells degranulate in response to anti-IgE and substance P. *Lab Invest* 1990; 63: 502-510.
166. Arck P C, Handjiski B, Hagen E, Joachim R, Klapp B F, Paus R. Indications for a 'brain-hair follicle axis (BHA)': inhibition of keratinocyte proliferation and up-regulation of keratinocyte apoptosis in telogen hair follicles by stress and substance P. *Faseb J* 2001; 15: 2536-2538.
167. Hordinsky M, Ericson M, Snow D, Boeck C, Lee W S. Peribulbar innervation and substance P expression following nonpermanent injury to the human scalp hair follicle. *J Investig Dermatol Symp Proc* 1999; 4: 316-319.
168. Schulze E, Witt M, Fink T, Hofer A, Funk R H. Immunohistochemical detection of human skin nerve fibers. *Acta Histochem* 1997; 99: 301-309.

169. Toyoda M, Makino T, Kagoura M, Morohashi M. Expression of neuropeptide-degrading enzymes in alopecia areata: an immunohistochemical study. *Br J Dermatol* 2001; 144: 46-54.
170. Staniek V, Doutremepuich J, Schmitt D, Claudy A, Misery L. Expression of substance P receptors in normal and psoriatic skin. *Pathobiology* 1999; 67: 51-54.
171. Kimata H. Laughter counteracts enhancement of plasma neurotrophin levels and allergic skin wheal responses by mobile phone-mediated stress. *Behav Med* 2004; 29: 149-152.
172. Hordinsky M K, Kennedy W, Wendelschafer Crabb G, Lewis S. Structure and function of cutaneous nerves in alopecia areata. *J Invest Dermatol* 1995; 104: 28S-29S.
173. Kimata H. Suckling reduces allergic skin responses and plasma levels of neuropeptide and neurotrophin in lactating women with atopic eczema/dermatitis syndrome. *Int Arch Allergy Immunol* 2003; 132: 380-383.
174. Huntley A, White A R, Ernst E. Relaxation therapies for asthma: a systematic review. *Thorax* 2002; 57: 127-131.
175. Aboussafy D, Campbell T S, Lavoie K, Aboud F E, Ditto B. Airflow and autonomic responses to stress and relaxation in asthma: The impact of stressor type. *Int J Psychophysiol* 2005; 20: 20.
176. Lonne-Rahm S, Berg M, Marin P, Nordlind K. Atopic dermatitis, stinging, and effects of chronic stress: a pathocausal study. *J Am Acad Dermatol* 2004; 51: 899-905.
177. El-Nour H, Lundeberg L, Boman A, Beck O, Harvima I T, Theodorsson E, Nordlind K. Study of innervation, sensory neuropeptides, and serotonin in murine contact allergic skin. *Immunopharmacol Immunotoxicol* 2005; 27: 67-76.
178. Bigliardi-Qi M, Lipp B, Sumanovski L T, Buechner S A, Bigliardi P L. Changes of epidermal mu-opiate receptor expression and nerve endings in chronic atopic dermatitis. *Dermatology* 2005; 210: 91-99.
179. Herbert M K, Holzer P. [Neurogenic inflammation. I. Basic mechanisms, physiology and pharmacology]. *Anesthesiol Intensivmed Notfallmed Schmerzther* 2002; 37: 314-325.
180. Holzer P. Capsaicin as a tool for studying sensory neuron functions. *Adv Exp Med Biol* 1991; 298: 3-16.
181. Heyer G. Abnormal cutaneous neurosensitivity in atopic skin. *Acta Derm Venereol Suppl* 1992; 176: 93-94.
182. Pincelli C, Fantini F, Massimi P, Girolomoni G, Seidenari S, Giannetti A. Neuropeptides in skin from patients with atopic dermatitis: an immunohistochemical study. *Br J Dermatol* 1990; 122: 745-750.
183. Ostlere L S, Cowen T, Rustin M H. Neuropeptides in the skin of patients with atopic dermatitis. *Clin Exp Dermatol* 1995; 20: 462-467.
184. Sugiura H, Omoto M, Hirota Y, Danno K, Uehara M. Density and fine structure of peripheral nerves in various skin lesions of atopic dermatitis. *Arch Dermatol Res* 1997; 289: 125-131.
185. Urashima R, Mihara M. Cutaneous nerves in atopic dermatitis. A histological, immunohistochemical and electron microscopic study. *Virchows Arch* 1998; 432: 363-370.
186. Toyoda M, Morohashi M. Morphological assessment of the effects of cyclosporin A on mast cell-nerve relationship in atopic dermatitis. *Acta Derm Venereol* 1998; 78: 321-325.
187. Botchkarev V A, Sharov A A, Syska S, Maurer M, Gilchrest B A. Involvement of p53 and Fas in Hair Follicle Apoptosis in Human Skin/SCID Model for Chemotherapy-Induced Hair Loss. *J Invest Dermatol* 2002; 119: 286.
188. Chatterjee A D, Raychaudhuri S P, Raychaudhuri S K, Farber E M. SCID mouse model of psoriasis: a unique tool to develop novel anti-inflammatory drugs. *J Invest Dermatol* 2002; to be published in Juli: (*abstract*).
189. Gilhar A, Landau M, Assy B, Shalaginov R, Serafimovich S, Kalish R S. Mediation of Alopecia Areata by Cooperation Between CD4+ and CD8+ T Lymphocytes: Transfer to Human Scalp Explants on Prkdcscid Mice. *Arch Dermatol* 2002; 138: 916-922.

190. Raychaudhuri S P, Dutt S, Raychaudhuri S K, Sanyal M, Farber E M. Severe combined immunodeficiency mouse-human skin chimeras: a unique animal model for the study of psoriasis and cutaneous inflammation. *Br J Dermatol* 2001; 144: 931-939.
191. Tsuboi H, Tanei R, Fujimura T, Ohta Y, Katsuoka K. Characterization of infiltrating T cells in human scalp explants from alopecia areata to SCID nude mice: possible role of the disappearance of CD8+ T lymphocytes in the process of hair regrowth. *J Dermatol* 1999; 26: 797-802.
192. Weinstock J V, Blum A, Metwali A, Elliott D, Arsenescu R. IL-18 and IL-12 signal through the NF-kappa B pathway to induce NK-1R expression on T cells. *J Immunol* 2003; 170: 5003-5007.
193. Jeanjean A P, Moussaoui S M, Maloteaux J M, Laduron P M. Interleukin-1 beta induces long-term increase of axonally transported opiate receptors and substance P. *Neuroscience* 1995; 68: 151-157.
194. Steinman L. Elaborate interactions between the immune and nervous systems. *Nat Immunol* 2004; 5: 575-581.
195. Ansel J C, Brown J R, Payan D G, Brown M A. Substance P selectively activates TNF-alpha gene expression in murine mast cells. *J Immunol* 1993; 150: 4478-4485.
196. Saban M R, Saban R, Bjoerling D, Haak-Freudscho M. Involvement of leukotrienes, TNF-alpha, and the LFA-1/ICAM-1 interaction in substance P-induced granulocyte infiltration. *J Leukoc Biol* 1997; 61: 445-451.
197. van der Kleij H P, Ma D, Redegeld F A, Kraneveld A D, Nijkamp F P, Bienenstock J. Functional expression of neurokinin 1 receptors on mast cells induced by IL-4 and stem cell factor. *J Immunol* 2003; 171: 2074-2079.
198. Foster A P, Cunningham F M. Substance P induces activation, adherence and migration of equine eosinophils. *J Vet Pharmacol Ther* 2003; 26: 131-138.
199. Gutwald J, Goebeler M, Sorg C. Neuropeptides enhance irritant and allergic contact dermatitis. *J Invest Dermatol* 1991; 96: 695-698.
200. Goebeler M, Henseleit U, Roth J, Sorg C. Substance P and calcitonin gene-related peptide modulate leukocyte infiltration to mouse skin during allergic contact dermatitis. *Arch Dermatol Res* 1994; 286: 341-346.
201. Schijns V E, Tangeras A. Vaccine adjuvant technology: from theoretical mechanisms to practical approaches. *Dev Biol* 2005; 121: 127-134.
202. Matzinger P. The danger model: a renewed sense of self. *Science* 2002; 296: 301-305.
203. Beresford L, Orange O, Bell E B, Miyan J A. Nerve fibres are required to evoke a contact sensitivity response in mice. *Immunology* 2004; 111: 118-125.
204. Weisshaar E, Heyer G, Forster C, Handwerker H O. Effect of topical capsaicin on the cutaneous reactions and itching to histamine in atopic eczema compared to healthy skin. *Arch Dermatol Res* 1998; 290: 306-311.
205. Olivry T, Mueller R S. Evidence-based veterinary dermatology: a systematic review of the pharmacotherapy of canine atopic dermatitis. *Vet Dermatol* 2003; 14: 121-146.
206. Hordinsky M, Ericson M. Autoimmunity: alopecia areata. *J Investig Dermatol Symp Proc* 2004; 9: 73-78.
207. Marsella R, Nicklin C F, Melloy C. The effects of capsaicin topical therapy in dogs with atopic dermatitis: a randomized, double-blinded, placebo-controlled, cross-over clinical trial. *Vet Dermatol* 2002; 13: 131-139.
208. Takano N, Arai I, Hashimoto Y, Kurachi M. Evaluation of antipruritic effects of several agents on scratching behavior by NC/Nga mice. *Eur J Pharmacol* 2004; 495: 159-165.
209. Hosoi J, Murphy G F, Egan C L, Lerner E A, Grabbe S, Asahina A, Granstein R D. Regulation of Langerhans cell function by nerves containing calcitonin gene-related peptide. *Nature* 1993; 363: 159-163.

210. Kawaguchi Y, Okada T, Konishi H, Fujino M, Asai J, Ito M. Reduction of the DTH response is related to morphological changes of Langerhans cells in mice exposed to acute immobilization stress. *Clin Exp Immunol* 1997; 109: 397-401.
211. Ahluwalia A, De Felipe C, O'Brien J, Hunt S P, Perretti M. Impaired IL-1 β -induced neutrophil accumulation in tachykinin NK1 receptor knockout mice. *Br J Pharmacol* 1998; 124: 1013-1015.
212. Schuiling M, Zuidhof A B, Zaagsma J, Meurs H. Role of tachykinin NK1 and NK2 receptors in allergen-induced early and late asthmatic reactions, airway hyperresponsiveness, and airway inflammation in conscious, unrestrained guinea pigs. *Clin Exp Allergy* 1999; 2: 48-52.
213. Castagliuolo I, Riegler M, Pasha A, Nikulasson S, Lu B, Gerard C, Gerard N P, Pothoulakis C. Neurokinin-1 (NK-1) receptor is required in *Clostridium difficile*- induced enteritis. *J Clin Invest* 1998; 101: 1547-1550.
214. Evangelista S, Meli A. Influence of capsaicin-sensitive fibres on experimentally-induced colitis in rats. *J Pharm Pharmacol* 1989; 41: 574-575.
215. Wallengren J, Ekman R, Moller H. Capsaicin enhances allergic contact dermatitis in the guinea pig. *Contact Dermatitis* 1991; 24: 30-34.
216. Reinshagen M, Flamig G, Ernst S, Geerling I, Wong H, Walsh J H, Eysselein V E, Adler G. Calcitonin gene-related peptide mediates the protective effect of sensory nerves in a model of colonic injury. *J Pharmacol Exp Ther* 1998; 286: 657-661.
217. Zachariae H, Henningsen S J. Histamine and plasma kinins in delayed allergic skin reactions. *Allerg Asthma* 1970; 16: 9-14.
218. McGillis J P, Organist M L, Payan D G. Substance P and immunoregulation. *Fed Proc* 1987; 46: 196-199.
219. Niizeki H, Kurimoto I, Streilein J W. A substance p agonist acts as an adjuvant to promote hapten-specific skin immunity. *J Invest Dermatol* 1999; 112: 437-442.
220. Babu R J, Kikwai L, Jaiani L T, Kanikkannan N, Armstrong C A, Ansel J C, Singh M. Percutaneous absorption and anti-inflammatory effect of a substance P receptor antagonist: spantide II. *Pharm Res* 2004; 21: 108-113.
221. Murozono M, Matsumoto S, Matsumoto E, Isshiki A, Watanabe Y. Neuroprotective and neurotoxic effects of cyclosporine A on transient focal ischemia in mdr1a knockout mice. *Eur J Pharmacol* 2004; 498: 115-118.
222. Tachibana T, Shiiya N, Kunihara T, Wakamatsu Y, Kudo A F, Ooka T, Watanabe S, Yasuda K. Immunophilin ligands FK506 and cyclosporine A improve neurologic and histopathologic outcome after transient spinal cord ischemia in rabbits. *J Thorac Cardiovasc Surg* 2005; 129: 123-128.
223. Paus R, Stenn K S, Link R E. The induction of anagen hair growth in telogen mouse skin by cyclosporine A administration. *Lab Invest* 1989; 60: 365-369.
224. Maurer M, Handjiski B, Paus R. Hair growth modulation by topical immunophilin ligands: induction of anagen, inhibition of massive catagen development, and relative protection from chemotherapy-induced alopecia. *Am J Pathol* 1997; 150: 1433-1441.
225. Hart P H, Townley S L, Grimaldeston M A, Khalil Z, Finlay-Jones J J. Mast cells, neuropeptides, histamine, and prostaglandins in UV-induced systemic immunosuppression. *Methods* 2002; 28: 79-89.
226. Seiffert K, Granstein R D. Neuropeptides and neuroendocrine hormones in ultraviolet radiation-induced immunosuppression. *Methods* 2002; 28: 97-103.
227. Zhang H T, Wang Q, Greene M I, Murali R. New perspectives on anti-HER2/neu therapeutics. *Drug News Perspect* 2000; 13: 325-329.
228. Kazlow Stern D, Tripp J M, Ho V C, Lebwohl M. The use of systemic immune moderators in dermatology: an update. *Dermatol Clin* 2005; 23: 259-300.
229. Mease P J. Psoriatic arthritis therapy advances. *Curr Opin Rheumatol* 2005; 17: 426-432.

230. Orsal AS, Blois S, Labuz D, Peters EMJ, Schaefer M, Arck PC: The Progesterone derivative dydrogesterone down-regulates Neurokinin 1 receptor expression on lymphocytes, induces a TH2 skew and exerts hypoalgesic effects in mice. *J Mol Med* (accepted). 2005