

List of Figures

Fig. 1	Sketch of tetraazaporphyrin or porphyrazine (TAP), phthalocyanine (Pc), naphthalocyanine (Nc) and anthracyanine (Ac) molecules. The sketches of porphyrin and chlorophyll molecules, as well as of the heme group in hemoglobin, are also added.	2
Fig. 1.2-1	Cu ₂ p _{3/2} spectra for Cu ₂ O (bottom) and CuO (top) normalized at peak height. The data are taken from Ref. [GTE88].....	13
Fig. 1.2-2	Schematic description of the processes which lead to the final states corresponding to the main line (ligand-to-metal charge transfer) and to the satellite (no charge transfer) at the Cu ₂ p photoionization of a compound with a copper 3d ⁹ configuration in the ground state.....	14
Fig. 1.2-3	Cu ₂ p photoemission spectra for copper dihalides. The data are taken from Ref. [LWH81]	15
Fig. 2.1-1	Schematic of the Max-Born-Institute undulator beamline at BESSY II.....	20
Fig. 2.1-2	Principle of the monochromator	21
Fig. 2.2-1	Schematic layout of the MBI UHV surface apparatus.....	22
Fig. 2.2-2	Sketch of the measurement geometry used in the present experiments. θ_i represents the incidence angle, which was 83°, and θ_p represents the angle between the synchrotron light electric vector and the sample normal (θ_p was 80° or 90°). The detection of the photoelectrons was normal to the sample surface.....	23
Fig. 2.2-3	View of the UHV apparatus at MBI User facility at BESSY II. In the photography the manipulator is set to serve the preparation chamber.....	25
Fig. 2.2-4	View of the UHV chamber at MBI User facility at BESSY II. The manipulator was rotated for bringing the sample into the analysis chamber.	26
Fig. 2.3-1	Optical layout of the U49/2 BTUC PGM beamline.....	27
Fig. 3.1-1	Molecular arrangements in copper phthalocyanine crystals of the α form and β form. The data are taken from Ref. [LeL89].	30
Fig. 3.2-1	The investigated set of porphyrazine molecules.....	32
Fig. 3.3-1	QMS spectrum for P0-Cu compound.	35
Fig. 3.3-2	Absorption spectra in solution for P0-Cu, P1-Cu, P2-Cu and P3-Cu compounds (also in [FrM86])......	36
Fig. 3.3-3	Results of absorption measurements for a P2-Cu film deposited on quartz, for the same film dissolved in toluene, and analogous data for P2-Cu prior to evaporation dissolved in toluene.	37
Fig. 3.3-4	Absorption spectrum of a P3-Cu film obtained by sublimation in UHV and deposited on quartz, of the same film dissolved in benzene, and of the P3-Cu compound in benzene prior to evaporation	38

Fig. 3.3-5	The photoemission spectrum of a P3-Cu film obtained by sublimation in UHV and of a P3-Cu film prepared by the wet chemical method. In both cases the excitation energy was 75.8 eV.	39
Fig. 3.3-6	Absorption spectrum of a P2-Zn film deposited on quartz by sublimation in UHV, of the same film dissolved in dioxane and of the P2-Zn compound in dioxane prior evaporation.	41
Fig. 3.3-7	Absorption spectra of a dissolved P2-H ₂ film from the Au(111) crystal surface and of the P2-H ₂ compound in benzene prior evaporation.	42
Fig. 4.1-1	Photoemission spectra from the unsubstituted H ₂ Pc film (top) and the P1-H ₂ film (bottom), both obtained with 90.6 eV photon energy. Electron binding energies are given with respect to vacuum level, and intensities are normalized in the 30-40 eV binding energy range. The labels for the top spectrum directly correspond to the nomenclature in [OrB92]; the respective photoemission feature assignment is presented in Table 4.1-1. In order to account for binding energy changes, the bottom spectrum is labeled by the corresponding capital letters. The dashed spectrum at the bottom is identical to the top spectrum.	44
Fig. 4.1-2	Molecular structure of H ₂ Pc with the same notations as used in [OrB92] and for the peaks assignment. N _p indicates a pyrrole nitrogen, and N _a and N _m denote pyrrole aza and meso-bridging aza nitrogens, respectively.	45
Fig. 4.1-3	The atomic orbital composition (marked by open and closed circles) of the two highest occupied molecular orbitals of H ₂ Pc, as taken from ref [OCP96].	47
Fig. 4.1-4	Photoemission spectra of condensed benzene multilayers. The data are taken from Ref. [GTM98]. The spectra were obtained with 50 eV photon energy at normal light incidence.	48
Fig. 4.1-5	Photoemission spectra of P0-H ₂ , P1-H ₂ , and P2-H ₂ compounds. Electron binding energies are given with respect to the vacuum level. The dashed spectrum overlapped on the P0-H ₂ spectrum and on the P2-H ₂ spectrum is the spectrum of P1-H ₂ compound (the middle spectrum in the figure), this being given for facilitating the comparison.	49
Fig. 4.1-6	Photoemission spectra of P0-H ₂ , P1-H ₂ and P2-H ₂ compounds at the edge of the valence region. The binding energy of feature A (HOMO) decreases with the extension of the π-electron system of the molecule. Labels mark the respective HOMO binding energies.	52
Fig. 4.1-7	Comparison between the one-electron energy levels obtained in ref. [OCP96] for metal-free tetraazaporphyrin (H ₂ TAP), metal-free phthalocyanine (H ₂ Pc), metal-free naphthalocyanine (H ₂ Nc) and HOMO and LUMO binding energies deduced in the present experiments for P0-H ₂ , P1-H ₂ , and P2-H ₂ . The theoretical energy scale was shifted by 0.52 eV in order to align the theoretical energy of HOMO for H ₂ Pc with the experimental binding energy for P1-H ₂ . Thick lines symbolize the experimental data while thinner lines symbolize the theoretical values.	54
Fig. 4.1-8	(a) Experimentally determined HOMO binding energies (solid squares) as a function of the number of benzo-units fused per pyrrole group for P0-H ₂ , P1-H ₂ and P2-H ₂ . The corresponding theoretical values obtained in Ref. [OCP96] for nonsubstituted compounds and shifted by 0.52 eV are added for comparison (open triangles). (b) Dependence of the 1 st absorption maximum measured in benzene as a function of the number of benzo-units fused per pyrrole group for P0-H ₂ , P1-H ₂ and P2-H ₂	55
Fig. 4.2-1	Changes in the photoemission spectra of the P0-H ₂ compound due to synchrotron radiation exposure. (a) spectra measured on a same spot and vertically translated for clarity. (b) two overlapped spectra measured after different beam exposure times, again on an identical spot but which is different from that in figure (a).	56

Fig. 4.2-2	Photoemission spectra of the P1-H ₂ compound for two different synchrotron radiation exposure times. The photon energy was 75.4 eV.	57
Fig. 5.1-1	Photoemission spectra from unsubstituted CuPc film (top) and P1-Cu film (bottom), both taken at 70.7 eV photon energy.	60
Fig. 5.1-2	Photoemission spectra from a P2-Cu film prepared by sublimation in UHV and from a P2-Cu film obtained by the wet chemical method described in Chapter 3.	61
Fig. 5.1-3	Comparison of photoemission spectra of unsubstituted CuPc and H ₂ Pc.	62
Fig. 5.1-4	Photoemission spectra from the Cu compounds in the series along with their metal-free counter parts. The data were taken for 70.7 eV excitation energy. The P3-H ₂ spectrum is missing due to its instability at film preparation. The P0-Cu, P1-Cu and P2-Cu films were produced by sublimation, while that of P3-Cu is prepared by the wet chemical method.	63
Fig. 5.1-5	Sketch of the HOMO for CuPc.	64
Fig. 5.1-6	Photoemission spectra of the P1-Cu compound for 69.7, 75.3 and 78.1 eV excitation energy. The intensity of band C (containing contributions from Cu3d direct emission) has an increase relative to the ligand features in the spectrum at the increase of the excitation energy, in the photon energy range applied in the present experiments. Feature F' shows a resonant behavior.	65
Fig. 5.1-7	Photoemission spectra of the copper porphyrazines in the series obtained for 70.7 eV excitation energy.	66
Fig. 5.1-8	Photoemission spectra of the valence region for P0-Cu, P1-Cu and P2-Cu compounds (expanded from Fig. 5.1-7). Energies for band A are labeled for P1-Cu and P2-Cu. A ± 0.07 eV uncertainty must be considered for the given values. The position of band A in the case of P0-Cu can not be identified unequivocally due to peak overlap.	68
Fig. 5.1-9	The result of fitting the 4.0-9.4 eV binding energy range for the P0-Cu spectrum. It was concluded that the feature found at about 6.8 eV binding energy includes both peaks A/HOMO-1 and A'/HOMO. The binding energy of feature A for P0-Cu was determined to be 6.50 ± 0.1 eV.	69
Fig. 5.2-1	Photoemission spectra of P0-Cu after various synchrotron radiation exposure times.	71
Fig. 5.2-2	Photoemission spectra of P1-Cu upon various synchrotron radiation exposure times.	72
Fig. 5.3-1	Important energy levels for Cu ²⁺ and Pc ²⁻ dianion in CuPc.	74
Fig. 5.3-2	Photoemission spectra for thin films of the Cu porphyrazine compounds at 70.7 and 75.8 eV photon energy. Arrows mark the Cu3d satellite peaks near 18 eV binding energy.	75
Fig. 5.3-3	Blow-up of the Cu satellite region for P0-Cu and P1-Cu. The spectra were obtained at several excitation energies around resonance. The position of the satellite peak at resonance is marked.	76
Fig. 5.3-4	Sketch of the copper ground state, of the final state corresponding to the satellite and of the final state reached after 3p-3d resonant excitation, in the case of CuPc and of copper porphyrazines with incomplete d shell. Considered here is the particular case when the satellite corresponds to the ionization of 3d _{xy} orbital and accompanying the resonant excitation, the Auger decay occurs from 3d _{xy} with the emission of an electron found initially in 3d _{x²-y²} . For simplification, the spin-orbit splitting of Cu 3p level after 3p hole creation is neglected.	78

Fig. 5.3-5	Sketch of the areas used for determining the overall dependence of the Cu satellite intensity on excitation energy.....	79
Fig. 5.3-6	The intensity change in the region of Cu satellite peaks as a function of the excitation energy for P0-Cu, P1-Cu and P2-Cu.....	80
Fig. 5.3-7	NEXAFS spectra for the P2-Cu and P3-Cu compounds. The spectra are vertically translated for clarity.....	82
Fig. 5.3-8	XPS spectra at the Cu2p core levels (top) and at the N1s level (bottom) for a P2-Cu film obtained by sublimation in UHV and for a P2-Cu film prepared by wet chemical method.....	83
Fig. 5.3-9	Cu2p XPS spectra for the investigated Cu porphyrazines. A linear background subtraction was performed.....	85
Fig. 5.3-10	XPS spectra of N1s core level for the investigated Cu porphyrazines after Shirley background subtraction.....	86
Fig. 5.4-1	The intensity ratio between the Cu2p _{1/2} satellite and main line for P0-Cu, P1-Cu and P2-Cu.....	87
Fig. 5.4-2	The XPS spectra for P0-Cu and P2-Cu in the region of Cu 2p core levels. The difference between the satellite intensities in the two compounds is visible.....	88
Fig. 5.4-3	The original spectrum, the calculated background, and the spectrum after background subtraction for P1-Cu for 75.3 eV photon energy.....	89
Fig. 5.4-4	The original spectrum, the calculated background, and the spectrum after background subtraction for P1-H ₂ for 75.3 eV photon energy.....	90
Fig. 5.4-5	The overlap of the P1-Cu and P1-H ₂ spectra, both after background subtraction, and their difference curve resulted by subtracting the P1-H ₂ spectrum from the corresponding copper one.....	90
Fig. 5.4-6	The overlap between the spectra of CuPc, P0-H ₂ and P2-H ₂ and those of the corresponding metal-free compounds after background subtraction.....	91
Fig. 5.4-7	Intensity ratio between the satellite at the position of band F' and main line for P1-Cu and CuPc at 70.7, 75.3, 75.8 and 80.9 eV excitation energy. The thicker lines with larger cap widths represent the error bars for CuPc while the thinner lines with shorter cap widths symbolize the error bars for P1-Cu.....	92
Fig. 5.4-8	The intensity ratio between the satellite and main line for P0-Cu, P1-Cu, and P2-Cu at 70.7, 75.3, 75.8 and 80.9 eV excitation energy.....	93
Fig. 5.4-9	The difference between the satellite to main line intensity ratios at resonant and off-resonant excitations (resonant enhancement). In these calculations were considered the satellites at the position of band F'. The values R(75.8)-R(70.7) mean the difference between the satellite to main line intensity ratio for 75.8 eV photon energy and the satellite to main line intensity ratio for 70.7 eV. R(75.3) and R(80.9) represent the satellite to main line intensity ratio for 75.3 and 80.9 eV photon energy, respectively.....	95
Fig. 6.1-1	Photoemission spectra from standard ZnPc and P1-Zn thin films at 98.7 eV excitation energy (see text for details). The dotted line represents the ZnPc spectrum, plotted on top of the P1-Zn one.....	100
Fig. 6.1-2	Photoemission spectra from unsubstituted ZnPc and H ₂ Pc. The main zinc contributions in the spectra are evidenced. They are the Zn3d direct emission line (at the position of band E) and Zn satellite peaks (as labeled).....	102

Fig. 6.1-3	Photoemission spectra of P0-Zn, P1-Zn and P2-Zn in comparison with the spectra of the respective metal-free compounds. Photon energy was 89.6 eV for all spectra.	103
Fig. 6.1-4	Photoemission spectra of P1-Zn for different excitation energies evidencing the intensity increase of band E (which includes Zn3d direct photoemission signal) relative to the ligand peaks for increasing photon energy.	105
Fig. 6.1-5	Photoemission spectra of P0-Zn, P1-Zn, and P2-Zn at 91.7 eV excitation energy. Notice the intensity increase of the benzene features relative to the Zn ones with the increase of the ligand size.	106
Fig. 6.1-6	Photoemission spectra for P0-Zn, P1-Zn, and P2-Zn at the edge of the valence region. All spectra were obtained for 70.4 eV photon energy. The binding energy of feature A decreases when the ligand gets larger. The labels mark the respective binding energies. All values are given with ± 0.07 eV uncertainty.	107
Fig. 6.2-1	Photoemission spectra of P0-Zn for different synchrotron radiation exposure times. The spectra were measured at 75.4 eV photon energy. They were normalized at the background and then vertically translated for clarity. Band A (derived from HOMO) is one of the features in the spectrum changing strongly as a result of photodegradation.	109
Fig. 6.2-2	Photoemission spectra of P1-Zn for different synchrotron radiation exposure times. The changes of the photoemission features induced by the synchrotron radiation exposure can be traced.	110
Fig. 6.3-1	N1s core level in the XPS spectra of P0-Zn, P1-Zn and P2-Zn	111
Fig. 6.3-2	C1s core level in the XPS spectra of P0-Zn, P1-Zn and P2-Zn	112
Fig. 6.3-3	Zn2p core level region of the XPS spectra of P0-Zn, P1-Zn and P2-Zn. No shift is detected for the two spin-orbit split components, Zn2p _{1/2} and Zn2p _{3/2} , at the enlargement of the ligand.	113
Fig. 6.4-1	Photoemission spectra of P0-Zn, P1-Zn, and P2-Zn for 91.7 eV (solid line) and 87.1 eV (dotted line) photon energy.	115
Fig. 6.4-2	Photoemission spectra (smoothed) in the region of Zn satellite peaks for ZnPc, P0-Zn, P1-Zn and P2-Zn for the photon energies used in the present experiments (from bottom to top: 86.6, 87.1, 87.6, 88.1, 88.6, 89.1, 89.6, 90.1, 90.6, 91.1, 91.7, 92.2, 92.7, 93.2, 93.7, 94.2, 94.7, 95.1, 95.7, 96.2, 96.8, 97.2, 97.8, 98.2, 98.7, 99.2, 99.8, 100.2, 100.7, 101.2, 101.7 eV). The graph for P0-Zn is missing the 100.7 eV photon energy curve, and the ZnPc graph is missing the spectra for 86.6 and 87.1 eV photon energy. One can note the evolution of the satellite peaks intensity at the variation of photon energy.	116
Fig. 6.4-3	Photoemission spectra of ZnPc and P1-Zn in the region of Zn satellite peaks obtained for 86.6, 91.7 and 100.2 eV photon energy. For ZnPc the spectrum at 85.6 eV is also shown. One can note that a weak satellite signal is visible in the ZnPc spectra, even for 85.6, 86.6, and 100.2 eV photon energy. The dashed line in the upper graph serves to guide the eye.	117
Fig. 6.4-4	Sketch of the ground state, of the final state reached after the 3d photoionization with shake-up, and of the final state reached after 3p-4s excitation for the Zn-porphyrine compounds in the series. For simplification, the spin-orbit splitting of Zn3p level after 3p hole creation is neglected.	119
Fig. 6.4-5	Photoemission spectra of P0-Zn at (from bottom to top) 87.1, 88.1, 89.1, 89.6, 90.6, 91.1, 91.7, 92.2, 92.7, 93.7, 94.7, 95.7, 96.8, 98.7 eV photon energy. The Auger signal in the photoemission spectra is indicated.	120

Fig. 6.4-6	The relative area of the largest Zn satellite peak as a function of the excitation energy for P0-Zn, P1-Zn and P2-Zn.	121
Fig. 6.4-7	The original spectrum, the calculated background, and the spectrum after background subtraction for P2-Zn at 94.2 eV excitation energy.	123
Fig. 6.4-8	The original spectrum, the calculated background, and the spectrum after background subtraction for P2-H ₂ at 94.2 eV excitation energy.	123
Fig. 6.4-9	The overlapped background-subtracted P2-Zn and P2-H ₂ spectra. The dashed curve represents the difference between the background-subtracted P2-Zn and P2-H ₂ spectra.	124
Fig. 6.4-10	Comparison of the ratios between the area of the Zn satellite peaks and the area of Zn3d main line as a function of the excitation energy for P0-Zn, P1-Zn and P2-Zn. The top panel displays the ratio between the area of the larger satellite peak, centered at ~37.8 eV binding energy, and the area of Zn3d main line. The lower panel shows the ratio between the sum area of both satellite peaks and the area of the Zn3d main line. For each graph the error bars are identical for all three compounds, in all points.	125
Fig. 6.5-1	Photoemission spectra of P1-Zn and P3-Zn films, prepared by wet chemical method, along with the corresponding spectra obtained for a UHV sublimed P1-Zn film.	127
Fig. 6.5-2	Photoemission spectra in the region of Zn satellite peaks from P1-Zn (top) and P3-Zn (bottom) at several photon energies. Both films have been deposited ex-situ, using a wet chemical method. Peak B represents most probably satellite signal but peak A doesn't originate from satellite signal.	128