

# List of Tables

Table 3.3-1	Temperatures used for the evaporation of CuPc, P0-Cu, P1-Cu and P2-Cu compounds.....	35
Table 3.3-2	Temperatures used for the evaporation of ZnPc, P0-Zn, P1-Zn, and P2-Zn compounds .....	40
Table 3.3-3	Temperatures used for the evaporation of H <sub>2</sub> Pc, P0-H <sub>2</sub> , P1-H <sub>2</sub> and P2-H <sub>2</sub> compounds. ....	42
Table 4.1-1	Measured electron binding energies relative to vacuum level (in eV) for H <sub>2</sub> Pc and P1-H <sub>2</sub> respectively, as obtained from Fig. 4.1-1 and similar data. The corresponding error bars are 0.07 eV. For comparison and peak assignment, data reported in [OrB92] are also included. Labels in small letters refer to unsubstituted H <sub>2</sub> Pc and capital letters refer to the P1-H <sub>2</sub> compound, respectively. ....	46
Table 4.1-2	Binding energies in eV and orbital symmetries as determined from calculations of the electronic structure for unsubstituted metal-free tetraazaporphyrin.....	51
Table 4.1-3	Measured binding energies for feature A (HOMO) together with the values corresponding to the 1 <sup>st</sup> absorption maximum of the compound dissolved in benzene. Assuming for simplicity that $E_{\text{HOMO}} - E_{\text{abs}} = E_{\text{LUMO}}$ , an approximate value for the LUMO binding energy can be determined. For comparison, the computed energy values for HOMO and LUMO, determined for the compounds without tert-butyl substitution in [OCP96], are also given.....	54
Table 5.1-1	Binding energy of feature A (ligand character) relative to the vacuum level for copper compounds in the series together with the values obtained for the metal-free complexes (written in italics). All the values are given within $\pm 0.07$ eV error range, except for P0-Cu where the uncertainty is $\pm 0.1$ eV. Note that the same orbitals contribute to feature A in a copper- and in its corresponding metal-free compound.....	70