

7. Literature

- (1). Nesterenko TH, Aly H. Fetal and neonatal programming: evidence and clinical implications. *Am J Perinatol*. 2009;26:191-8.
- (2). Barker DJ, Winter PD, Osmond C, Margetts B, Simmonds SJ. Weight in infancy and death from ischaemic heart disease. *Lancet* 1989;2:577–580
- (3). Kajantie E, Osmond C, Barker DJ, Forsen T, Phillips DI, Eriksson JG. Size at birth as a predictor of mortality in adulthood: a follow-up of 350 000 person-years. *Int J Epidemiol* 2005;34:655–663
- (4). Nathanielsz PW. *Life in Womb: The Origin of Health and Disease*. Ithaca, NY: Promethean Press; 1999
- (5). Cheung YF, Wong KY, Lam BC, Tsoi NS. Relation of arterial stiffness with gestational age and birth weight. *Arch Dis Child* 2004;89:217–221
- (6). Simmons R. Fetal origin of adult disease: concepts and controversies. *Neoreview*. 2004;5:511–515
- (7). Hovi P, Andersson S, Ericsson JG, et al. Glucose regulation in young adults with very low birth weight. *N Engl J Med* 2007;356:2053–206
- (8). Nijland MJ, Ford SP, Nathanielsz PW. Prenatal origins of adult disease. *Curr Opin Obstet Gynecol*. 2008;20:132-8.
- (9). Thone-Reineke C, Kalk P, Dorn M, et al. High-protein nutrition during pregnancy and lactation programs blood pressure, food efficiency, and body weight of the offspring in a sex-dependent manner.. *Am J Physiol Regul Integr Comp Physiol* 2006;291:1025–1030.
- (10). Singh RS, Cullen-McEwen LA, Kett MM, et al. Prenatal corticosterone exposure results in altered AT1/AT2, nephron deficit and hypertension in the rat offspring. *J Physiol* 2007;579:503–513.
- (11). Hocher B. Fetal programming of cardiovascular diseases in later life - mechanisms beyond maternal undernutrition. *J Physiol*. 2007;579:287-8.
- (12). Meyer K, Lubo Zhang. Fetal programming of cardiac function and disease. *Reprod Sci*. 2007 ;14:209-16
- (13). Fowden AL, Giussani DA, Forhead AJ. Intrauterine programming of physiological systems: causes and consequences. *Physiology (Bethesda)*. 2006;21:29-37

- (14). Waterland RA, Michels KB. Epigenetic epidemiology of the developmental origins hypothesis. *Annu Rev Nutr.* 2007;27:363-88.
- (15). Gardiner HM. Early environmental influences on vascular development. *Early Hum Dev.* 2007;83:819-23
- (16). Louey S, Thornburg KL. The prenatal environment and later cardiovascular disease. *Early Hum Dev.* 2005;81:745-51
- (17). Stefani L, Toncelli L, Gianassi M, et al. Two-dimensional tracking and TDI are consistent methods for evaluating myocardial longitudinal peak strain in left and right ventricle basal segments in athletes. *Cardiovasc Ultrasound.* 2007;7:5-7
- (18). Pirat B, Khoury DS, Hartley CJ, et al. A novel feature-tracking echocardiographic method for the quantitation of regional myocardial function: validation in an animal model of ischemia-reperfusion. *J Am Coll Cardiol.* 2008;51(6):651-9
- (19). Younoszai AK, Saudek DE, Emery SP, Thomas JD. Evaluation of myocardial mechanics in the fetus by velocity vector imaging. *J Am Soc Echocardiogr.* 2008;21:470-4
- (20). Barker PC, Houle H, Li JS, Miller S, Herlong JR, Camitta MG. Global longitudinal cardiac strain and strain rate for assessment of fetal cardiac function: novel experience with velocity vector imaging. *Echocardiography.* 2009;26:28-36
- (21). Murata Y, Takemura H, Kurachi K. Observation of fetal cardiac motion by M-mode ultrasonic cardiography. *Am J Obstet Gynecol.* 1971;111:287-94
- (22). DeVore GR. Assessing fetal cardiac ventricular function. *Semin Fetal Neonatal Med.* 2005;10:515-41
- (23). DeVore GR, Siassi B, Platt LD. The use of the abdominal circumference as a means of assessing M-mode ventricular dimensions during the second and third trimesters of pregnancy in the normal human fetus. *J Ultrasound Med* 1985;4:175-82.
- (24). DeVore GR, Siassi B, Platt LD. Use of femur length as a means of assessing M-mode ventricular dimensions during second and third trimesters of pregnancy in normal fetus. *J Clin Ultrasound* 1985;13:619-25.
- (25). DeVore GR, Siassi B, Platt LD. Fetal echocardiography. IV. M-mode assessment of ventricular size and contractility during the second and third trimesters of pregnancy in the normal fetus. *Am J Obstet Gynecol* 1984;150:981-8.

- (26). Sharland GK, Allan LD. Normal fetal cardiac measurements derived by cross-sectional echocardiography. *Ultrasound Obstet Gynecol* 1992;2:175-81.
- (27). Tan J, Silverman NH, Hoffman JI, Villegas M, Schmidt KG. Cardiac dimensions determined by cross-sectional echocardiography in the normal human fetus from 18 weeks to term. *Am J Cardiol* 1992;70:1459-67.
- (28). Shapiro I, Degani S, Leibovitz Z, Ohel G, Tal Y, Abinader EG. Fetal cardiac measurements derived by transvaginal and transabdominal cross-sectional echocardiography from 14 weeks of gestation to term. *Ultrasound Obstet Gynecol* 1998;12:404-18.
- (29). Firpo C, Hoffman JI, Silverman NH. Evaluation of fetal heart dimensions from 12 weeks to term. *Am J Cardiol* 2001;87:594-600.
- (30). Harada K, Rice MJ, Shiota T, Ishii M, McDonald RW, Reller MD, et al. Gestational age- and growth-related alterations in fetal right and left ventricular diastolic filling patterns. *Am J Cardiol* 1997;79:173-7.
- (31). Carceller-Blanchard AM, Fouron JC. Determinants of the Doppler flow velocity profile through the mitral valve of the human fetus. *Br Heart J* 1993;70:457-60
- (32). Romero R, Espinoza J, Goncalves LF, Gomez R, Medina L, Silva M, et al. Fetal cardiac dysfunction in preterm premature rupture of membranes. *J Matern Fetal Neonatal Med* 2004;16:146-57.
- (33). Weiner Z, Zloczower M, Lerner A, Zimmer E, Itskovitz- Eldor J. Cardiac compliance in fetuses of diabetic women. *Obstet Gynecol* 1999;93:948-51.
- (34). Tsyvian P, Malkin K, Artemieva O, Wladimiroff JW. Assessment of left ventricular filling in normally grown fetuses, growth-restricted fetuses and fetuses of diabetic mothers. *Ultrasound Obstet Gynecol* 1998;12:33-8.
- (35). Rizzo G, Arduini D. Fetal cardiac function in intrauterine growth retardation. *Am J Obstet Gynecol* 1991;165:876-82.
- (36). De Smedt MC, Visser GH, Meijboom EJ. Fetal cardiac output estimated by Doppler echocardiography during mid- and late gestation. *Am J Cardiol* 1987;60:338-42.
- (37). Mielke G, Benda N. Cardiac output and central distribution of blood flow in the human fetus. *Circulation* 2001;103:1662-8.
- (38). Severi FM, Rizzo G, Bocchi C, D'Antona D, Verzuri MS, Arduini D. Intrauterine growth retardation and fetal cardiac function. *Fetal Diagn Ther* 2000;15:8-19.

- (39). Di Renzo GC, Luzi G, Cucchia GC, Caserta G, Fusaro P, Perdikaris A, et al. The role of Doppler technology in the evaluation of fetal hypoxia. *Early Hum Dev* 1992;29:259-67.
- (40). Friedman D, Buyon J, Kim M, Glickstein JS. Fetal cardiac function assessed by Doppler myocardial performance index (Tei Index). *Ultrasound Obstet Gynecol* 2003; 21:33–36.
- (41). Hernandez-Andrade E, López-Tenorio J, Figueroa-Diesel H, et al. A modified myocardial performance (Tei) index based on the use of valve clicks improves reproducibility of fetal left cardiac function assessment. *Ultrasound Obstet Gynecol.* 2005;26:227-32
- (42). Tei C, Ling LH, Hodge DO, et al. New index of combined systolic and diastolic myocardial performance: a simple and reproducible measure of cardiac function--a study in normals and dilated cardiomyopathy. *J Cardiol.* 1995 Dec;26:357-66
- (43). Szwast A, Tian Z, McCann M, Donaghue D, Rychik J. Right ventricular performance in the fetus with hypoplastic left heart syndrome. *Ann Thorac Surg.* 2009;87:1214-9
- (44). Van Mieghem T, Klaritsch P, Doné E, et al. Assessment of fetal cardiac function before and after therapy for twin-to-twin transfusion syndrome. *Am J Obstet Gynecol.* 2009;200:1-7
- (45). Wong ML, Wong WH, Cheung YF. Fetal myocardial performance in pregnancies complicated by gestational impaired glucose tolerance. *Ultrasound Obstet Gynecol.* 2007;29:395-400
- (46). Tsutsumi T, Ishii M, Eto G, Hota M, Kato H. Serial evaluation for myocardial performance in fetuses and neonates using a new Doppler index. *Pediatr Int.* 1999;41:722-7
- (47). Eidem BW, Edwards JM, Cetta F. Quantitative assessment of fetal ventricular function: establishing normal values of the myocardial performance index in the fetus. *Echocardiography.* 2001;18:9-13
- (48). Van Mieghem T, Gucciardo L, Lewi P, et al. Validation of the fetal myocardial performance index in the second and third trimesters of gestation. *Ultrasound Obstet Gynecol.* 2009;33:58-63.

- (49). Crispi F, Hernandez-Andrade E, Pelsers MM, et al. Cardiac dysfunction and cell damage across clinical stages of severity in growth-restricted fetuses. *Am J Obstet Gynecol.* 2008;199:1-8
- (50). Yagel S, Cohen SM, Shapiro I, Valsky DV. 3D and 4D ultrasound in fetal cardiac scanning: a new look at the fetal heart. *Ultrasound Obstet Gynecol.* 2007;29:81-95
- (51). Esh-Broder E, Ushakov FB, Imbar T, Yagel S. Application of free-hand three-dimensional echocardiography in the evaluation of fetal cardiac ejection fraction: a preliminary study. *Ultrasound Obstet Gynecol* 2004;23:546–551
- (52). Meyer-Wittkopf M, Cole A, Cooper SG, Schmidt S, Sholler GF. Three dimensional quantitative echocardiographic assessment of ventricular volume in healthy human fetuses and in fetuses with congenital heart disease. *J Ultrasound Med* 2001;20:317–327
- (53). Messing B, Cohen SM, Valsky DV, et al. Fetal cardiac ventricle volumetry in the second half of gestation assessed by 4D ultrasound using STIC combined with inversion mode. *Ultrasound Obstet Gynecol.* 2007;30:142-51
- (54). Molina FS, Faro C, Sotiriadis A, Dagklis T, Nicolaides KH. Heart stroke volume and cardiac output by four-dimensional ultrasound in normal fetuses. *Ultrasound Obstet Gynecol.* 2008;32:181-7
- (55). Rizzo G, Capponi A, Cavicchioni O, Vendola M, Arduini D. Fetal cardiac stroke volume determination by four-dimensional ultrasound with spatio-temporal image correlation compared with two-dimensional and Doppler ultrasonography. *Prenat Diagn.* 2007;27:1147-50
- (56). Tutschek B, Sahn DJ. Semi-automatic segmentation of fetal cardiac cavities: progress towards an automated fetal echocardiogram. *Ultrasound Obstet Gynecol.* 2008;32:176-80.
- (57). Tutschek B, Zimmermann T, Buck T, Bender HG. Fetal tissue Doppler echocardiography: detection rates of cardiac structures and quantitative assessment of the fetal heart. *Ultrasound Obstet Gynecol.* 2003;21:26-32
- (58). Twining P. Myocardial motion imaging: a new application of power color flow and frequency-based color flow Doppler in fetal echocardiography. *Ultrasound Obstet Gynecol.* 1999;13:255-9

- (59). Harada K, Tsuda A, Orino T, Tanaka T, Takada G. Tissue Doppler imaging in the normal fetus. *Int J Cardiol.* 1999;71:227-34.
- (60). Paladini D, Lamberti A, Teodoro A, Arienzo M, Tartaglione A, Martinelli P. Tissue Doppler imaging of the fetal heart. *Ultrasound Obstet Gynecol.* 2000;16:530-5.
- (61). Aoki M, Harada K, Ogawa M, Tanaka T. Quantitative assessment of right ventricular function using doppler tissue imaging in fetuses with and without heart failure. *J Am Soc Echocardiogr.* 2004;17:28-35.
- (62). Chan LY, Fok WY, Wong JT, Yu CM, Leung TN, Lau TK. Reference charts of gestation-specific tissue Doppler imaging indices of systolic and diastolic functions in the normal fetal heart. *Am Heart J.* 2005;150:750-5.
- (63). Di Salvo G, Russo MG, Paladini D, et al. Quantification of regional left and right ventricular longitudinal function in 75 normal fetuses using ultrasound-based strain rate and strain imaging. *Ultrasound Med Biol.* 2005;31:1159-62
- (64). Larsen LU, Petersen OB, Norrild K, Sorensen K, Uldbjerg N, Sloth E. Strain rate derived from color Doppler myocardial imaging for assessment of fetal cardiac function. *Ultrasound Obstet Gynecol.* 2006;27:210-3
- (65). Gardiner HM, Pasquini L, Wolfenden J, et al. Myocardial tissue Doppler and long axis function in the fetal heart. *Int J Cardiol.* 2006;113:39-47
- (66). Perles Z, Nir A, Gavri S, Rein AJ. Assessment of fetal myocardial performance using myocardial deformation analysis. *Am J Cardiol.* 2007;99:993-6
- (67). Ta-Shma A, Perles Z, Gavri S, et al. Analysis of segmental and global function of the fetal heart using novel automatic functional imaging. *J Am Soc Echocardiogr.* 2008;21:146-50
- (68). Di Salvo G, Russo MG, Paladini D, et al. Two-dimensional strain to assess regional left and right ventricular longitudinal function in 100 normal foetuses. *Eur J Echocardiogr.* 2008;9:754-6
- (69). Bijmens BH, Cikes M, Claus P, Sutherland GR. Velocity and deformation imaging for the assessment of myocardial dysfunction. *Eur J Echocardiogr.* 2009;10:216-26.
- (70). D'hooge J, Heimdal A, Jamal F, et al. Regional strain and strain rate measurements by cardiac ultrasound: principles, implementation and limitations. *Eur J Echocardiogr.* 2000;1:154-70

- (71). Yip G, Abraham T, Belohlavek M, Khandheria BK. Clinical applications of strain rate imaging. *J Am Soc Echocardiogr.* 2003;16:1334-42.
- (72). Pedrizzetti G, Tonti G. Endocardial Border Kinematics by Automatic Tracking of Echocardiographic Imaging. Personal Communication
- (73). Heimdal A, Støylen A, Torp H, Skjaerpe T. Real-time strain rate imaging of the left ventricle by ultrasound. *J Am Soc Echocardiogr.* 1998;11:1013-9.
- (74). Peng QH, Zhou QC, Zeng S, et al. Evaluation of regional left ventricular longitudinal function in 151 normal fetuses using velocity vector imaging. *Prenat Diagn.* 2009
- (75). Langeland S, Wouters PF, Claus P, et al. Experimental assessment of a new research tool for the estimation of two-dimensional myocardial strain. *Ultrasound Med Biol.* 2006;32:1509-13
- (76). Yue Y, Clark Jr JW, Khoury DS. Speckle Tracking in Intracardiac Echocardiography for the Assessment of Myocardial Deformation *IEEE Trans. on Biomed. Engr.* 2009;56:416-425
- (77). Korinek J, Wang J, Sengupta PP, et al. Two-dimensional strain--a Doppler-independent ultrasound method for quantitation of regional deformation: validation in vitro and in vivo. *J Am Soc Echocardiogr.* 2005;18:1247-53
- (78). Kirkpatrick SJ, Wnag RK, Duncan DD. OCT-based elastography for large and small deformations. *Optics Express* 2006;14:11585-11597

8. Curriculum Vitae and Acknowledgement

8.1. Curriculum Vitae

Mein Lebenslauf wird aus datenschutzrechtlichen Gründen in der elektronischen Version meiner Arbeit nicht veröffentlicht.

My curriculum vitae is for privacy reasons in the electronic version of the work not published.

8.2. Acknowledgement

I would like to thank Prof. Dr. Joachim W. Dudenhausen for allowance to conduct the study in his clinic and for the financial aid to the congress participations in Germany. I do not know how to thank Prof. Dr. Wolfgang Henrich, who offered me this doctoral thesis and played a crucial role in my understanding not only of the fetal echocardiography but of all territories of ultrasound. He was for me there during working hours and after work, too. I am indebted to him for his endless support. My dear mother has always been a pushing power in almost every instance of my life. Her support during the last two difficult years are very valuable for me. I thank her from the deepest part of my heart. Senior physicians from our clinic Dr. Martin Brauer and Assoc. Prof. Dr. Ines Schönborn, residents of the Department of Obstetrics at Campus Virchow Klinikum, especially Dr. Jens Stupin, Dr. Elsa Hollatz, Dr. Nina Salomon, Dr. Ute Braig-Scherer and Dr. Nicole Gehrman helped me to recruit as many patients for our study as possible. I thank them a lot as they gave in their busy daily routine a thought to the hardships I may encounter if I will not have enough number of cases to evaluate. The doctor`s assistants and midwives in our clinic were always there with their kindness and helpfulness. When I was in a desperate condition they helped me out of it. I will never forget their help. My biggest thank goes to the mothers who allowed me to “look at the heart of their babies”. I experienced the most beautiful moments of my life with them scanning their babies and was delighted to see their curiosity as I explained them our research. It was a pleasure and privilege to have them.

Selbstständigkeitserklärung

„Ich, Onur Önügören, erkläre, dass ich die vorgelegte Dissertation mit dem Thema: Assessment of Long Axis Ventricular Function in the Fetal Heart based on 2D Grey Scale Image Processing with a Tissue Tracking Algorithm selbst verfasst und keine anderen als die angegebenen Quellen und Hilfsmittel benutzt, ohne die (unzulässige) Hilfe Dritter verfasst und auch in Teilen keine Kopien anderer Arbeiten dargestellt habe.“

15.03.2010, Ibbenbüren

Onur Önügören

Unterschrift