

8. Literaturverzeichnis

- Abrams J. M. (1999) An emerging blueprint for apoptosis in Drosophila. *Trends Cell Biol* **9**, 435-40.
- Acehan D., Jiang X., Morgan D.G., Heuser J.E., Wang X. and C.W. A. (2002) Three-dimensional structure of the apoptosome: implications for assembly, pro caspase-9 binding, and activation. *Mol Cell* **2**, 423-32.
- Adams J. M. (2003) Ways of dying: multiple pathways to apoptosis. *Genes Dev* **17**, 2481-95.
- Adams J. M. and Cory S. (1998) The Bcl-2 protein family: arbiters of cell survival. *Science* **281**, 1322-6.
- Adams J. M. and Cory S. (2001) Life-or-death decisions by the Bcl-2 protein family. *Trends Biochem Sci* **26**, 61-6.
- Adesanya M. R., Redman R. S., Baum B. J. and O'Connell B. C. (1996) Immediate inflammatory responses to adenovirus-mediated gene transfer in rat salivary glands. *Hum Gene Ther* **7**, 1085-93.
- Alnemri E. S., Livingston D. J., Nicholson D. W., Salvesen G., Thornberry N. A., Wong W. W. and Yuan J. (1996) Human ICE/CED-3 protease nomenclature. *Cell* **87**, 171.
- Annis M. G., Zamzami N., Zhu W., Penn L. Z., Kroemer G., Leber B. and Andrews D. W. (2001) Endoplasmic reticulum localized Bcl-2 prevents apoptosis when redistribution of cytochrome c is a late event. *Oncogene* **20**, 1939-52.
- Antonsson B., Montessuit S., Lauper S., Eskes R. and Martinou J. C. (2000) Bax oligomerization is required for channel-forming activity in liposomes and to trigger cytochrome c release from mitochondria. *Biochem J* **345 Pt 2**, 271-8.
- Arnt C. R., Chiorean M. V., Heldebrant M. P., Gores G. J. and Kaufmann S. H. (2002) Synthetic Smac/DIABLO peptides enhance the effects of chemotherapeutic agents by binding XIAP and cIAP1 in situ. *J Biol Chem* **277**, 44236-43.
- Ashkenazi A. and Dixit V. M. (1998) Death receptors: signaling and modulation. *Science* **281**, 1305-8.
- Barry M., Heibein J. A., Pinkoski M. J., Lee S. F., Moyer R. W., Green D. R. and Bleackley R. C. (2000) Granzyme B short-circuits the need for caspase 8 activity during granule-mediated cytotoxic T-lymphocyte killing by directly cleaving Bid. *Mol Cell Biol* **20**, 3781-94.
- Berridge M. J., Lipp P. and Bootman M. D. (2000) The versatility and universality of calcium signalling. *Nat Rev Mol Cell Biol* **1**, 11-21.
- Birnbaum M. J., Clem R. J. and Miller L. K. (1994) An apoptosis-inhibiting gene from a nuclear polyhedrosis virus encoding a polypeptide with Cys/His sequence motifs. *J Virol* **68**, 2521-8.
- Birnboim H. C. and Doly J. (1979) A rapid alkaline extraction procedure for screening recombinant plasmid DNA. *Nucleic Acids Res* **7**, 1513-23.
- Boatright K. M. and Salvesen G. S. (2003) Mechanisms of caspase activation. *Curr Opin Cell Biol* **15**, 725-31.
- Boatright K. M., Renatus M., Scott F. L., Sperandio S., Shin H., Pedersen I. M., Ricci J., Edris W. A., Sutherlin D. P., Green D. R. and Salvesen G. S. (2003) A unified model for caspase activation. *Mol Cell* **11**, 529-41.
- Boehning D., Patterson R. L., Sedaghat L., Glebova N. O., Kurosaki T. and Snyder S. H. (2003) Cytochrome c binds to inositol (1,4,5) trisphosphate receptors, amplifying calcium-dependent apoptosis. *Nat Cell Biol* **5**, 1051-61.
- Borner C. (2003) The Bcl-2 protein family: sensors and checkpoints for life-or-death decisions. *Mol Immunol* **39**, 615-47.

- Borner C., Martinou J., Mattmann C., Irmler M., Schaefer E., Martinou J. C. and Tschopp J. (1994a) The protein bcl-2 alpha does not require membrane attachment, but two conserved domains to suppress apoptosis. *J Cell Biol* **126**, 1059-68.
- Borner C., Olivier R., Martinou J., Mattmann C., Tschopp J. and Martinou J. C. (1994b) Dissection of functional domains in Bcl-2 alpha by site-directed mutagenesis. *Biochem Cell Biol* **72**, 463-9.
- Bossy-Wetzel, E., Newmeyer, D. D. and Green, D. R. (1998) Mitochondrial cytochrome c release in apoptosis occurs upstream of DEVD-specific caspase activation and independently of mitochondrial transmembrane depolarization. *EMBO J* **17**, 37-49.
- Brand K., Klocke R., Possling A., Paul D. and Strauss M. (1999) Induction of apoptosis and G2/M arrest by infection with replication-deficient adenovirus at high multiplicity of infection. *Gene Ther* **6**, 1054-63.
- Bratton S. B., Walker G., Srinivasula S. M., Sun X. M., Butterworth M., Alnemri E. S. and Cohen G. M. (2001) Recruitment, activation and retention of caspases-9 and -3 by Apaf-1 apoptosome and associated XIAP complexes. *Embo J* **20**, 998-1009.
- Breckenridge D. G., Stojanovic M., Marcellus R. C. and Shore G. C. (2003) Caspase cleavage product of BAP31 induces mitochondrial fission through endoplasmic reticulum calcium signals, enhancing cytochrome c release to the cytosol. *J Cell Biol* **160**, 1115-27.
- Brockstedt E., Rickers A., Kostka S., Laubersheimer A., Dorken B., Wittmann-Liebold B., Bommert K. and Otto A. (1998) Identification of apoptosis-associated proteins in a human Burkitt lymphoma cell line. Cleavage of heterogeneous nuclear ribonucleoprotein A1 by caspase 3. *J Biol Chem* **273**, 28057-64.
- Carothers A. M., Steigerwalt R. W., Urlaub G., Chasin L. A. and D. G. (1989) DNA base changes and RNA levels in N-acetoxy-2-acetylaminofluorene-induced dihydrofolate reductase mutants of Chinese hamster ovary cells. *J Mol Biol* **208**, 417-28.
- Cartron P. F., Juin P., Oliver L., Martin S., Meflah K. and Vallette F. M. (2003) Nonredundant role of Bax and Bak in Bid-mediated apoptosis. *Mol Cell Biol* **23**, 4701-12.
- Chai J., Du C., Wu J. W., Kyin S., Wang X. and Shi Y. (2000) Structural and biochemical basis of apoptotic activation by Smac/DIABLO. *Nature* **406**, 855-62.
- Chang D. W., Xing Z., Pan Y., Algeciras-Schimminich A., Barnhart B. C., Yaish-Ohad S., Peter M. E. and Yang X. (2002) c-FLIP(L) is a dual function regulator for caspase-8 activation and CD95-mediated apoptosis. *Embo J* **21**, 3704-14.
- Chartier C., Degryse E., Gantzer M., Dieterle A., Pavirani A. and Mehtali M. (1996) Efficient generation of recombinant adenovirus vectors by homologous recombination in *Escherichia coli*. *J Virol* **70**, 4805-10.
- Chen P., Nordstrom W., Gish B. and Abrams J. M. (1996) grim, a novel cell death gene in *Drosophila*. *Genes Dev* **10**, 1773-82.
- Cheng E. H., Sheiko T., Fisher J. K., Craigen W. J. and Korsmeyer S. J. (2003) VDAC2 inhibits Bak Activation and Mitochondrial Apoptosis. *Science* **301**, 513-17.
- Cheng E. H., Wei M. C., Weiler S., Flavell R. A., Mak T. W., Lindsten T. and Korsmeyer S. J. (2001) BCL-2, BCL-X(L) sequester BH3 domain-only molecules preventing BAX- and BAK-mediated mitochondrial apoptosis. *Mol Cell* **8**, 705-11.
- Chinnaiyan A., O'Rourke K., Lane BR and VM. D. (1997) Interaction of CED-4 with CED-3 and CED-9: a molecular framework for cell death. *Science* **275**, 1122-6.
- Chittenden T., Harrington E. A., O'Connor R., Flemington C., Lutz R. J., Evan G. I. and Guild B. C. (1995) Induction of apoptosis by the Bcl-2 homologue Bak. *Nature* **374**, 733-6.
- Cohen G. M. (1997) Caspases: the executioners of apoptosis. *Biochem J* **326** (Pt 1), 1-16.

- Cosset F. L., Takeuchi Y., Battini J. L., Weiss R. A. and Collins M. K. (1995) High-titer packaging cells producing recombinant retroviruses resistant to human serum. *J Virol* **69**, 7430-6.
- Crook N. E., Clem R. J. and Miller L. K. (1993) An apoptosis-inhibiting baculovirus gene with a zinc finger-like motif. *J Virol* **67**, 2168-74.
- Cryns V. L., Bergeron L., Zhu H., Li H. and Yuan J. (1996) Specific cleavage of alpha-fodrin during Fas- and tumor necrosis factor-induced apoptosis is mediated by an interleukin-1beta-converting enzyme/Ced-3 protease distinct from the poly(ADP-ribose) polymerase protease. *J Biol Chem* **271**, 31277-82.
- Custodio J. B., Cardoso J. M., Madeira V. M. and Almeida L. M. (2001) Mitochondrial permeability transition induced by the anticancer drug etoposide. *Toxicol In Vitro* **15**, 265-70.
- Dai Z., Zhu W. G., Morrison C. D., Brenna R. M., Smiraglia D. J., Raval A., Wu Y. Z., Rush L. J., Ross P., Molina J. R., Otterson G. A. and Plass C. (2003) A comprehensive search for DNA amplification in lung cancer identifies inhibitors of apoptosis cIAP1 and cIAP2 as candidate oncogenes. *Hum Mol Genet* **12**, 791-801.
- Daniel P. T., Schulze-Osthoff K., Belka C. and Guner D. (2003) Guardians of cell death: the Bcl-2 family proteins. *Essays Biochem* **39**, 73-88.
- De Giorgi F., Lartigue L., Bauer M. K., Schubert A., Grimm S., Hanson G. T., Remington S. J., Youle R. J. and Ichas F. (2002) The permeability transition pore signals apoptosis by directing Bax translocation and multimerization. *Faseb J* **16**, 607-9.
- Desagher S., Osen-Sand A., Nichols A., Eskes R., Montessuit S., Lauper S., Maundrell K., Antonsson B. and Martinou J. C. (1999) Bid-induced conformational change of Bax is responsible for mitochondrial cytochrome c release during apoptosis. *J Cell Biol* **144**, 891-901.
- Deveraux Q. L., Takahashi R., Salvesen G. S. and Reed J. C. (1997) X-linked IAP is a direct inhibitor of cell-death proteases. *Nature* **388**, 300-4.
- Distelhorst C. W. and McCormick T. S. (1996) Bcl-2 acts subsequent to and independent of Ca²⁺ fluxes to inhibit apoptosis in thapsigargin- and glucocorticoid-treated mouse lymphoma cells. *Cell Calcium* **19**, 473-83.
- Donepudi M., Mac Sweeney A., Briand C. and Grutter M. G. (2003) Insights into the regulatory mechanism for caspase-8 activation. *Mol Cell* **11**, 543-9.
- Du C., Fang M., Li Y., Li L. and Wang X. (2000) Smac, a mitochondrial protein that promotes cytochrome c-dependent caspase activation by eliminating IAP inhibition. *Cell* **102**, 33-42.
- Duckett C. S., Nava V. E., Gedrich R. W., Clem R. J., Van Dongen J. L., Gilfillan M. C., Shiels H., Hardwick J. M. and Thompson C. B. (1996) A conserved family of cellular genes related to the baculovirus iap gene and encoding apoptosis inhibitors. *Embo J* **15**, 2685-94.
- Earnshaw W. C., Martins L. M. and Kaufmann S. H. (1999) Mammalian caspases: structure, activation, substrates, and functions during apoptosis. *Annu Rev Biochem* **68**, 383-424.
- Eckhart L., Ban J., Fischer H. and Tschachler E. (2000a) Caspase-14: analysis of gene structure and mRNA expression during keratinocyte differentiation. *Biochem Biophys Res Commun* **277**, 655-9.
- Eckhart L., Declercq W., Ban J., Rendl M., Lengauer B., Mayer C., Lippens S., Vandenabeele P. and Tschachler E. (2000b) Terminal differentiation of human keratinocytes and stratum corneum formation is associated with caspase-14 activation. *J Invest Dermatol* **115**, 1148-51.
- Ellis R. E., Jacobson D. M. and Horvitz H. R. (1991) Genes required for the engulfment of cell corpses during programmed cell death in *Caenorhabditis elegans*. *Genetics* **129**, 79-94.

- Enari M., Sakahira H., Yokoyama H., Okawa K., Iwamatsu A. and Nagata S. (1998) A caspase-activated DNase that degrades DNA during apoptosis, and its inhibitor ICAD. *Nature* **391**, 43-50.
- Eskes R., Desagher S., Antonsson B. and Martinou J. C. (2000) Bid induces the oligomerization and insertion of Bax into the outer mitochondrial membrane. *Mol Cell Biol* **20**, 929-35.
- Faccio L., Fusco C., Chen A., Martinotti S., Bonventre J. V. and Zervos A. S. (2000) Characterization of a novel human serine protease that has extensive homology to bacterial heat shock endoprotease HtrA and is regulated by kidney ischemia. *J Biol Chem* **275**, 2581-8.
- Faccio L., Fusco C., Viel A. and Zervos A. S. (2000) Tissue-specific splicing of Omi stress-regulated endoprotease leads to an inactive protease with a modified PDZ motif. *Genomics* **68**, 343-7.
- Farrow S. N., White J. H., Martinou I., Raven T., Pun K. T., Grinham C. J., Martinou J. C. and Brown R. (1995) Cloning of a bcl-2 homologue by interaction with adenovirus E1B 19K. *Nature* **374**, 731-3.
- Fischer U., Janicke R. U. and Schulze-Osthoff K. (2003) Many cuts to ruin: a comprehensive update of caspase substrates. *Cell Death Differ* **10**, 76-100.
- Fischer H., Koenig U., Eckhart L. and Tschachler E. (2002) Human caspase 12 has acquired deleterious mutations. *Biochem Biophys Res Commun* **293**, 722-6.
- Fraser A. G., James C., Evan G. I. and Hengartner M. O. (1999) Caenorhabditis elegans inhibitor of apoptosis protein (IAP) homologue BIR-1 plays a conserved role in cytokinesis. *Curr Biol* **9**, 292-301.
- Freemont P. S., Hanson I. M. and Trowsdale J. (1991) A novel cysteine-rich sequence motif. *Cell* **64**, 483-4.
- Friedrich K., Wieder T., von Haefen C., Radetzki S., Schulze-Osthoff K., Jänicke R., Dörken B. and Daniel P. T. (2001) Overexpression of caspase-3 restores sensitivity for drug-induced apoptosis in breast cancer cells with acquired drug resistance. *Oncogene* **20**, 2749-2760.
- Fu J., Jin Y. and Arends L. J. (2003) Smac3, a Novel Smac/DIABLO Splicing Variant, Attenuates the Stability and Apoptosis-inhibiting Activity of X-linked Inhibitor of Apoptosis Protein. *JBC* **278**, 52660-52672.
- Fulda S., Wick W., Weller M. and Debatin K. M. (2002) Smac agonists sensitize for Apo2L/TRAIL- or anticancer drug-induced apoptosis and induce regression of malignant glioma *in vivo*. *Nat Med* **8**, 808-15.
- Gillissen B., Essmann F., Graupner V., Starck L., Radetzki S., Dorken B., Schulze-Osthoff K. and Daniel P. T. (2003) Induction of cell death by the BH3-only Bcl-2 homolog NbK/Bik is mediated by an entirely Bax-dependent mitochondrial pathway. *Embo J* **22**, 3580-90.
- Goldstein J. C. Waterhouse, N. J. Juin P., Evan, G. I and Green, D. R. (2000) The coordinate release of cytochrome c during apoptosis is rapid, complete and kinetically invariant *Nat Cell Biol*, **2**, 156-62.
- Goodwin E. C. and Rottman F. M. (1992) The 3'-flanking sequence of the bovine growth hormone gene contains novel elements required for efficient and accurate polyadenylation. *J Biol Chem* **267**, 16330-4.
- Graham F. L., Smiley J., Russell W. C. and Nairn R. (1977) Characteristics of a human cell line transformed by DNA from human adenovirus type 5. *J Gen Virol* **36**, 59-74.
- Graham, F.L., and A.J. van der Eb. 1973. Transformation of rat cells by DNA of human adenovirus 5. *Virology*. 54:536-9.

- Grether M. E., Abrams J. M., Agapite J., White K. and Steller H. (1995) The head involution defective gene of *Drosophila melanogaster* functions in programmed cell death. *Genes Dev* **9**, 1694.
- Gross A., Yin X. M., Wang K., Wei M. C., Jockel J., Milliman C., Erdjument-Bromage H., Tempst P. and Korsmeyer S. J. (1999) Caspase cleaved BID targets mitochondria and is required for cytochrome c release, while BCL-XL prevents this release but not tumor necrosis factor-R1/Fas death. *J Biol Chem* **274**, 1156-63.
- Guo F., Nimmanapalli R., Paranawithana S., Wittman S., Griffin D., Bali P., O'Bryan E., Fumero C., Wang H. G. and Bhalla K. (2002) Ectopic overexpression of second mitochondria-derived activator of caspases (Smac/DIABLO) or cotreatment with N-terminus of Smac/DIABLO peptide potentiates epothilone B derivative-(BMS 247550) and Apo-2L/TRAIL-induced apoptosis. *Blood* **99**, 3419-26.
- Halenbeck R., MacDonald H., Roulston A., Chen T. T., Conroy L. and Williams L. T. (1998) CPAN, a human nuclease regulated by the caspase-sensitive inhibitor DFF45. *Curr Biol* **8**, 537-40.
- Halestrap A. P., McStay G. P. and Clarke S. J. (2002) The permeability transition pore complex: Another view. *Biochimie* **84**, 153-166.
- Hanada M., Aime-Sempe C., Sato T. and Reed J. C. (1995) Structure-function analysis of Bcl-2 protein. Identification of conserved domains important for homodimerization with Bcl-2 and heterodimerization with Bax. *J Biol Chem* **270**, 11962-9.
- Harris B. Z. and Lim W. A. (2001) Mechanism and role of PDZ domains in signaling complex assembly. *J Cell Sci* **114**, 3219-31.
- Hay B. A., Wassarman D. A. and Rubin G. M. (1995) *Drosophila* homologs of baculovirus inhibitor of apoptosis proteins function to block cell death. *Cell* **83**, 1253-62.
- Hegde R., Srinivasula S. M., Zhang Z., Wassell R., Mukattash R., Cilenti L., DuBois G., Lazebnik Y., Zervos A. S., Fernandes-Alnemri T. and Alnemri E. S. (2002) Identification of Omi/HtrA2 as a mitochondrial apoptotic serine protease that disrupts inhibitor of apoptosis protein-caspase interaction. *J Biol Chem* **277**, 432-8.
- Hegde R., Srinivasula S. M., Datta P., Madesh M., Wassell R., Zhang Z., Cheong N., Nejmeh J., Fernandes-Alnemri T., Hoshino S. and Alnemri E. S. (2003) The polypeptide chain-releasing factor GSPT1/eRF3 is proteolytically processed into an IAP-binding protein. *J Biol Chem* **278**, 38699-706.
- Hemmati, P., Gillissen B., von Haefen C., Wendt J., Stärck L., Güner D., Dörken B., Daniel P.T. (2002) Adenovirus-mediated overexpression of p14ARF induces p53 and Bax-independent apoptosis. *Oncogene* **21**, 3149-61
- Hinds M. G., Norton R. S., Vaux D. L. and Day C. L. (1999) Solution structure of a baculoviral inhibitor of apoptosis (IAP) repeat. *Nat Struct Biol* **6**, 648-51.
- Hitomi J., Katayama T., Eguchi Y., Kudo T., Taniguchi M., Koyama Y., Manabe T., Yamagishi S., Bando Y., Imaizumi K., Tsujimoto Y. and Tohyama M. (2004) Involvement of caspase-4 in endoplasmic reticulum stress-induced apoptosis and A{beta}-induced cell death. *J Cell Biol* **165**, 347-56
- Holcik M. and Korneluk R. G. (2000) Functional characterization of the X-linked inhibitor of apoptosis (XIAP) internal ribosome entry site element: role of La autoantigen in XIAP translation. *Mol Cell Biol* **20**, 4648-57.
- Holcik M. and Korneluk R. G. (2001) XIAP, the guardian angel. *Nat Rev Mol Cell Biol* **2**, 550-6.
- Holley C. L., Olson M. R., Colon-Ramos D. A. and Kornbluth S. (2002) Reaper eliminates IAP proteins through stimulated IAP degradation and generalized translational inhibition. *Nat Cell Biol* **4**, 439-44.

- Hsu S. Y., Kaipia A., McGee E., Lomeli M. and Hsueh A. J. (1997) Bok is a pro-apoptotic Bcl-2 protein with restricted expression in reproductive tissues and heterodimerizes with selective anti-apoptotic Bcl-2 family members. *Proc Natl Acad Sci U S A* **94**, 12401-6.
- Hu S. and Yang X. (2003) Cellular inhibitor of apoptosis 1 and 2 are ubiquitin ligases for the apoptosis inducer Smac/DIABLO. *J Biol Chem* **278**, 10055-60.
- Huang H., Joazeiro C. A., Bonfoco E., Kamada S., Leverson J. D. and Hunter T. (2000) The inhibitor of apoptosis, cIAP2, functions as a ubiquitin-protein ligase and promotes in vitro monoubiquitination of caspases 3 and 7. *J Biol Chem* **275**, 26661-4.
- Humke E. W., Ni J. and Dixit V. M. (1998) ERICE, a novel FLICE-activatable caspase. *J Biol Chem* **273**, 15702-7.
- Hunter J. J. and Parslow T. G. (1996) A peptide sequence from Bax that converts Bcl-2 into an activator of apoptosis. *J Biol Chem* **271**, 8521-4.
- Hunter A. M., Kottachchi D., Lewis J., Duckett C. S., Korneluk R. G. and Liston P. (2003) A novel ubiquitin fusion system bypasses the mitochondria and generates biologically active Smac/DIABLO. *J Biol Chem* **278**, 7494-9.
- Inohara N., Ekhterae D., Garcia I., Carrio R., Merino J., Merry A., Chen S. and Nunez G. (1998) Mtd, a novel Bcl-2 family member activates apoptosis in the absence of heterodimerization with Bcl-2 and Bcl-XL. *J Biol Chem* **273**, 8705-10.
- Jackson P. K., Eldridge A. G., Freed E., Furstenthal L., Hsu J. Y., Kaiser B. K. and Reimann J. D. (2000) The lore of the RINGs: substrate recognition and catalysis by ubiquitin ligases. *Trends Cell Biol* **10**, 429-39.
- Jänicke R. U., Ng P., Sprengart M. L. and Porter A. G. (1998) Caspase-3 is required for alpha-fodrin cleavage but dispensable for cleavage of other death substrates in apoptosis. *J Biol Chem* **273**, 15540-5.
- Jänicke R. U., Sprengart M. L., Wati M. R. and Porter A. G. (1998) Caspase-3 is required for DNA fragmentation and morphological changes associated with apoptosis. *J Biol Chem* **273**, 9357-60.
- Joazeiro C. A. and Weissman A. M. (2000) RING finger proteins: mediators of ubiquitin ligase activity. *Cell* **102**, 549-52.
- Johnson D. E., Gastman B. R., Wieckowski E., Wang G. Q., Amoscato A., Delach S. M. and Rabinowich H. (2000) Inhibitor of apoptosis protein hILP undergoes caspase-mediated cleavage during T lymphocyte apoptosis. *Cancer Res* **60**, 1818-23.
- Jordan B. W., Dinev D., LeMellay V., Troppmair J., Gotz R., Wixler L., Sendtner M., Ludwig S. and Rapp U. R. (2001) Neurotrophin receptor-interacting mage homologue is an inducible inhibitor of apoptosis protein-interacting protein that augments cell death. *J Biol Chem* **276**, 39985-9.
- Joza N., Susin S. A., Daugas E., Stanford W. L., Cho S. K., Li C. Y., Sasaki T., Elia A. J., Cheng H. Y., Ravagnan L., Ferri K. F., Zamzami N., Wakeham A., Hakem R., Yoshida H., Kong Y. Y., Mak T. W., Zuniga-Pflucker J. C., Kroemer G. and Penninger J. M. (2001) Essential role of the mitochondrial apoptosis-inducing factor in programmed cell death. *Nature* **410**, 549-54.
- Jurgensmeier J. M., Xie Z., Deveraux Q., Ellerby L., Bredesen D. and Reed J. C. (1998) Bax directly induces release of cytochrome c from isolated mitochondria. *Proc Natl Acad Sci U S A* **95**, 4997-5002.
- Keij J. F., Bell-Prince C. and Steinkamp J. A. (2000) Staining of mitochondrial membranes with 10-nonyl acridine orange, MitoFluor Green, and MitoTracker Green is affected by mitochondrial membrane potential altering drugs. *Cytometry* **39**, 203-10.
- Kerr J. F., Winterford C. M. and Harmon B. V. (1994) Apoptosis. Its significance in cancer and cancer therapy. *Cancer* **73**, 2013-26.

- Koenig U., Eckhart L. and Tschachler E. (2001) Evidence that caspase-13 is not a human but a bovine gene. *Biochem Biophys Res Commun* **285**, 1150-4.
- Laemmli U. K. (1970) Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* **227**, 680-5.
- Lamkanfi M., Declercq W., Kalai M., Saelens X. and P. V. (2002) Alice in caspase land. A phylogenetic analysis of caspases from worm to man. *Cell Death Differ.* **4**, 358-61.
- Lazebnik Y. A., Takahashi A., Moir R. D., Goldman R. D., Poirier G. G., Kaufmann S. H. and Earnshaw W. C. (1995) Studies of the lamin proteinase reveal multiple parallel biochemical pathways during apoptotic execution. *Proc Natl Acad Sci U S A* **92**, 9042-6.
- Lee L. G., Connell C. R., Woo S. L., Cheng R. D., McArdle B. F., Fuller C. W., Halloran N. D. and Wilson R. K. (1992) DNA sequencing with dye-labeled terminators and T7 DNA polymerase: effect of dyes and dNTPs on incorporation of dye-terminators and probability analysis of termination fragments. *Nucleic Acids Res* **20**, 2471-83.
- Li H., Zhu H., Xu C. J. and Yuan J. (1998) Cleavage of BID by caspase 8 mediates the mitochondrial damage in the Fas pathway of apoptosis. *Cell* **94**, 491-501.
- Li L. Y., Luo X. and Wang X. (2001) Endonuclease G is an apoptotic DNase when released from mitochondria. *Nature* **412**, 95-9.
- Li W., Srinivasula S. M., Chai J., Li P., Wu J. W., Zhang Z., Alnemri E. S. and Shi Y. (2002) Structural insights into the pro-apoptotic function of mitochondrial serine protease HtrA2/Omi. *Nat Struct Biol* **9**, 436-41.
- Lindsten T., Ross A. J., King A., Zong W. X., Rathmell J. C., Shiels H. A., Ulrich E., Waymire K. G., Mahar P., Frauwirth K., Chen Y., Wei M., Eng V. M., Adelman D. M., Simon M. C., Ma A., Golden J. A., Evan G., Korsmeyer S. J., MacGregor G. R. and Thompson C. B. (2000) The combined functions of proapoptotic Bcl-2 family members bak and bax are essential for normal development of multiple tissues. *Mol Cell* **6**, 1389-99.
- Liston P., Fong W. G., Kelly N. L., Toji S., Miyazaki T., Conte D., Tamai K., Craig C. G., McBurney M. W. and Korneluk R. G. (2001) Identification of XAF1 as an antagonist of XIAP anti-Caspase activity. *Nat Cell Biol* **3**, 128-33.
- Liu X., Li P., Widlak P., Zou H., Luo X., Garrard W. T. and Wang X. (1998) The 40-kDa subunit of DNA fragmentation factor induces DNA fragmentation and chromatin condensation during apoptosis. *Proc Natl Acad Sci U S A* **95**, 8461-6.
- Liu X., Zou H., Slaughter C. and Wang X. (1997) DFF, a heterodimeric protein that functions downstream of caspase-3 to trigger DNA fragmentation during apoptosis. *Cell* **89**, 175-84.
- Loeffler M., Daugas E., Susin S. A., Zamzami N., Metivier D., Nieminen A. L., Brothers G., Penninger J. M. and Kroemer G. (2001) Dominant cell death induction by extramitochondrially targeted apoptosis-inducing factor. *Faseb J* **15**, 758-67.
- Louis N., Evelegh C. and Graham F. L. (1997) Cloning and sequencing of the cellular-viral junctions from the human adenovirus type 5 transformed 293 cell line. *Virology* **233**, 423-9.
- Luo X., Budihardjo I., Zou H., Slaughter C. and Wang X. (1998) Bid, a Bcl2 interacting protein, mediates cytochrome c release from mitochondria in response to activation of cell surface death receptors. *Cell* **94**, 481-90.
- Ly J.D., Grubb D.R. and Lawen A. (2003) The mitochondrial membrane potential ($\Delta\psi_m$) in apoptosis; an update. *Apoptosis* **8**, 115-28
- Mandic A., Viktorsson K., Strandberg L., Heiden T., Hansson J., Linder S. and Shoshan M. C. (2002) Calpain-Mediated Bid Cleavage and Calpain-Independent Bak Modulation: Two Separate Pathways in Cisplatin-Induced Apoptosis. *Mol Cell Biol* **22**, 3003-13

- Martinou J. C. and Green D. R. (2001) Breaking the mitochondrial barrier. *Nat Rev Mol Cell Biol* **2**, 63-7.
- Martins L. M., Iaccarino I., Tenev T., Gschmeissner S., Totty N. F., Lemoine N. R., Savopoulos J., Gray C. W., Creasy C. L., Dingwall C. and Downward J. (2002) The serine protease Omi/HtrA2 regulates apoptosis by binding XIAP through a reaper-like motif. *J Biol Chem* **277**, 439-44.
- Martins L. M., Turk B. E., Cowling V., Borg A., Jarrell E. T., Cantley L. C. and Downward J. (2003) Binding Specificity and Regulation of the Serine Protease and PDZ Domains of HtrA2/Omi. *JBC* **278**, 49417-49427.
- Marzo I., Brenner C., Zamzami N., Susin S. A., Beutner G., Brdiczka D., Remy R., Xie Z. H., Reed J. C. and Kroemer G. (1998) The permeability transition pore complex: a target for apoptosis regulation by caspases and bcl-2-related proteins. *J Exp Med* **187**, 1261-71.
- Mashima T., Naito M., Noguchi K., Miller D. K., Nicholson D. W. and Tsuruo T. (1997) Actin cleavage by CPP-32/apopain during the development of apoptosis. *Oncogene* **14**, 1007-12.
- McNeish I. A., Bell S., McKay T., Tenev T., Marani M. and Lemoine N. R. (2003) Expression of Smac/DIABLO in ovarian carcinoma cells induces apoptosis via caspase-9-mediated pathway. *Experimental Cell Research* **286**, 186-198.
- Micheau O., Thome M., Schneider P., Holler N., Tschopp J., Nicholson D. W., Briand C. and Grutter M. G. (2002) The long form of FLIP is an activator of caspase-8 at the Fas death-inducing signaling complex. *J Biol Chem* **277**, 45162-71.
- Miller L. K. (1999) An exegesis of IAPs: salvation and surprises from BIR motifs. *Trends Cell Biol* **9**, 323-8.
- Miramar M. D., Costantini P., Ravagnan L., Saraiva L. M., Haouzi D., Brothers G., Penninger J. M., Peleato M. L., Kroemer G. and Susin S. A. (2001) NADH oxidase activity of mitochondrial apoptosis-inducing factor. *J Biol Chem* **276**, 16391-8.
- Miura M., Zhu H., Rotello R., Hartwieg E. A. and Yuan J. (1993) Induction of apoptosis in fibroblasts by IL-1 beta-converting enzyme, a mammalian homolog of the *C. elegans* cell death gene ced-3. *Cell* **75**, 653-60.
- Mrozek A., Petrowsky H., Sturm I., Krauss J., Hermann S., Hauptmann S., Lorenz M. and Daniel P. (2003) Combined p53/Bax mutation results in extremely poor prognosis in gastric carcinoma with low microsatellite instability. *Cell Death Differ* **in press**.
- Muchmore S. W., Sattler M., Liang H., Meadows R. P., Harlan J. E., Yoon H. S., Nettesheim D., Chang B. S., Thompson C. B., Wong S. L., Ng S. L. and Fesik S. W. (1996) X-ray and NMR structure of human Bcl-xL, an inhibitor of programmed cell death. *Nature* **381**, 335-41.
- Mund T., Gewies A., Schoenfeld N., Bauer M. K. and Grimm S. (2003) Spike, a novel BH3-only protein, regulates apoptosis at the endoplasmic reticulum. *Faseb J* **17**, 696-8.
- Nakagawa T. and Yuan J. (2000) Cross-talk between two cysteine protease families. Activation of caspase-12 by calpain in apoptosis. *J Cell Biol* **150**, 887-94.
- Nakagawa T., Zhu H., Morishima N., Li E., Xu J., Yankner B. A. and Yuan J. (2000) Caspase-12 mediates endoplasmic-reticulum-specific apoptosis and cytotoxicity by amyloid-beta. *Nature* **403**, 98-103.
- Nakano K. and Vousden K. H. (2001) PUMA, a novel proapoptotic gene, is induced by p53. *Mol Cell* **7**, 683-94.
- Narita M., Shimizu S., Ito T., Chittenden T., Lutz R. J., Matsuda H. and Tsujimoto Y. (1998) Bax interacts with the permeability transition pore to induce permeability transition and cytochrome c release in isolated mitochondria. *Proc Natl Acad Sci U S A* **95**, 14681-6.

- Nath R., Raser K. J., Stafford D., Hajimohammadreza I., Posner A., Allen H., Talanian R. V., Yuen P., Gilbertsen R. B. and Wang K. K. (1996) Non-erythroid alpha-spectrin breakdown by calpain and interleukin 1 beta-converting-enzyme-like protease(s) in apoptotic cells: contributory roles of both protease families in neuronal apoptosis. *Biochem J* **319** (Pt 3), 683-90.
- Ng F. W. H., Nguyen M., Kwan T., Branton P. E., Nicholson D. W., Cromlish J. A. and Shore G. C. (1997) p28 Bap31, a Bcl-2/Bcl-X_L- and Procaspace-8-associated Protein in the Endoplasmic Reticulum. *J Cell Biol* **139**, 327-338.
- Nicholson D. W. (2001) Apoptosis. Baiting death inhibitors. *Nature* **410**, 33-4.
- Oda E., Ohki R., Murasawa H., Nemoto J., Shibue T., Yamashita T., Tokino T., Taniguchi T. and Tanaka N. (2000) Noxa, a BH3-only member of the Bcl-2 family and candidate mediator of p53-induced apoptosis. *Science* **288**, 1053-8.
- Okada H., Suh W. K., Jin J., Woo M., Du C., Elia A., Duncan G. S., Wakeham A., Itie A., Lowe S. W., Wang X. and Mak T. W. (2002) Generation and characterization of Smac/DIABLO-deficient mice. *Mol Cell Biol* **22**, 3509-17.
- Olson M. R., Holley C. L., Yoo S. J., Huh J. R., Hay B. A. and Kornbluth S. (2003) Reaper is regulated by IAP-mediated ubiquitination. *J Biol Chem* **278**, 4028-34.
- Oltvai Z. N., Milliman C. L. and Korsmeyer S. J. (1993) Bcl-2 heterodimerizes in vivo with a conserved homolog, Bax, that accelerates programmed cell death. *Cell* **74**, 609-19.
- Orth K., Chinnaiyan A. M., Garg M., Froelich C. J. and Dixit V. M. (1996) The CED-3/ICE-like protease Mch2 is activated during apoptosis and cleaves the death substrate lamin A. *J Biol Chem* **271**, 16443-6.
- Parrish J., Li L., Klotz K., Ledwich D., Wang X. and Xue D. (2001) Mitochondrial endonuclease G is important for apoptosis in *C. elegans*. *Nature* **412**, 90-4.
- Paszty K., Verma A. K., Padanyi R., Filoteo A. G., Penniston J. T. and Enyedi A. (2002) Plasma membrane Ca²⁺-ATPase isoform 4b is cleaved and activated by caspase-3 during the early phase of apoptosis. *J Biol Chem* **277**, 6822-9.
- Pörn-Ares M. I., Samali A. and Orrenius S. (1998) Cleavage of the calpain inhibitor, calpastatin, during apoptosis. *Cell Death Differ* **5**, 1028-33.
- Prober J. M., Trainor G.L., Dam R.J., Hobbs F.W., Robertson C.W., Zagursky R.J., Cocuzza A.J., Jensen M.A. and K. B. (1987) A system for rapid DNA sequencing with fluorescent chain-terminating dideoxynucleotides. *Science* **238**, 336-41.
- Puthalakath H., Huang D. C., O'Reilly L. A., King S. M. and Strasser A. (1999) The proapoptotic activity of the Bcl-2 family member Bim is regulated by interaction with the dynein motor complex. *Mol Cell* **3**, 287-96.
- Puthalakath H., Villunger A., O'Reilly L. A., Beaumont J. G., Coultras L., Cheney R. E., Huang D. C. and Strasser A. (2001) Bmf: a proapoptotic BH3-only protein regulated by interaction with the myosin V actin motor complex, activated by anoikis. *Science* **293**, 1829-32.
- Raisova M., Hossini A., Eberle J., Riebeling C., Wieder T., Sturm I., Daniel P. T., Orfanos C. E. and Geilen C. C. (2001) The Bax/Bcl-2 ratio determines the susceptibility of human melanoma cells to CD95/Fas-mediated apoptosis. *J Invest Dermatol* **177**, 333-340.
- Rao L., Perez D. and White E. (1996) Lamin proteolysis facilitates nuclear events during apoptosis. *J Cell Biol* **135**, 1441-55.
- Rasmussen U., Brogger Christensen S. and F. S. (1978) Thapsigargin and thapsigarginine, two new histamine liberators from *Thapsia gargarica* L. *Acta Pharm Suec* **15**, 133-40.
- Ravagnan L., Gurbuxani S., Susin S. A., Maisse C., Daugas E., Zamzami N., Mak T., Jaattela M., Penninger J. M., Garrido C. and Kroemer G. (2001) Heat-shock protein 70 antagonizes apoptosis-inducing factor. *Nat Cell Biol* **3**, 839-43.

- Raynal P. and Pollard H. B. (1994) Annexins: the problem of assessing the biological role for a gene family of multifunctional calcium- and phospholipid-binding proteins. *Biochim Biophys Acta* **1197**, 63-93.
- Renatus M., Stennicke H. R., Scott F. L., Liddington R. C. and Salvesen G. S. (2001) Dimer formation drives the activation of the cell death protease caspase 9. *Proc Natl Acad Sci U S A* **98**, 14250-14255.
- Roberts D. L., Merrison W., MacFarlane M. and Cohen G. M. (2001) The inhibitor of apoptosis protein-binding domain of Smac is not essential for its proapoptotic activity. *J Cell Biol* **153**, 221-8.
- Rodriguez J. and Lazebnik Y. (1999) Caspase-9 and APAF-1 form an active holoenzyme. *Genes Dev* **13**, 3179-84.
- Roy N., Mahadevan M. S., McLean M., Shutler G., Yaraghi Z., Farahani R., Baird S., Besner-Johnston A., Lefebvre C., Kang X. and et al. (1995) The gene for neuronal apoptosis inhibitory protein is partially deleted in individuals with spinal muscular atrophy. *Cell* **80**, 167-78.
- Rudner J., Lepple-Wienhues A., Budach W., Berschauer J., Friedrich B., Wesselborg S., Schulze-Osthoff K. and Belka C. (2001) Wild-type, mitochondrial and ER-restricted Bcl-2 inhibit DNA damage-induced apoptosis but do not affect death receptor-induced apoptosis. *J Cell Sci* **114**, 4161-72.
- Ryoo H. D., Bergmann A., Gonen H., Ciechanover A. and Steller H. (2002) Regulation of Drosophila IAP1 degradation and apoptosis by reaper and ubcD1. *Nat Cell Biol* **4**, 432-8.
- Saelens X., Festjens N., Vande Walle L., van Gurp M., van Loo G. and Vandenebeele P. (2004) Toxic proteins released from mitochondria in cell death. *Oncogene* **23**, 2861-2874.
- Saiki R. K., Chang C. A., Levenson C. H., Warren T. C., Boehm C. D., Kazazian H. H., Jr. and Erlich H. A. (1988) Diagnosis of sickle cell anemia and beta-thalassemia with enzymatically amplified DNA and nonradioactive allele-specific oligonucleotide probes. *N Engl J Med* **319**, 537-41.
- Saito M., Korsmeyer S. J. and Schlesinger P. H. (2000) BAX-dependent transport of cytochrome c reconstituted in pure liposomes. *Nat Cell Biol* **275**, 12321-12325.
- Sakahira H., Enari M. and Nagata S. (1998) Cleavage of CAD inhibitor in CAD activation and DNA degradation during apoptosis. *Nature* **391**, 96-9.
- Salvesen G. S. and Duckett C. S. (2002) IAP proteins: blocking the road to death's door. *Nat Rev Mol Cell Biol* **3**, 401-10.
- Sanger F., Nicklen S. and Coulson A. R. (1977) DNA sequencing with chain-terminating inhibitors. *Proc Natl Acad Sci U S A* **74**, 5463-7.
- Savopoulos J. W., Carter P. S., Turconi S., Pettman G. R., Karan E. H., Gray C. W., Ward R. V., Jenkins O. and Creasy C. L. (2000) Expression, purification, and functional analysis of the human serine protease HtrA2. *Protein Expr Purif* **19**, 227-34.
- Schulze-Osthoff K. (1994) The Fas/APO-1 receptor and its deadly ligand. *Trends Cell Biol* **4**, 421-6.
- Schwab B. L., Guerini D., Didzun C., Bano D., Ferrando-May E., Fava E., Tam J., Xu D., Xanthoudakis S., Nicholson D. W., Carafoli E. and Nicotera P. (2002) Cleavage of plasma membrane calcium pumps by caspases: a link between apoptosis and necrosis. *Cell Death Differ* **9**, 818-31.
- Scorrano L., Oakes S. A., Opferman J. T., Cheng E. H., Sorcinelli M. D., Pozzan T. and Korsmeyer S. J. (2003) BAX and BAK regulation of endoplasmic reticulum Ca²⁺: a control point for apoptosis. *Science* **300**, 135-9.
- Shimizu S., Narita M. and Tsujimoto Y. (1999) Bcl-2 family proteins regulate the release of apoptogenic cytochrome c by the mitochondrial channel VDAC. *Nature* **399**, 483-7.

- Soul H. D., Vazquez J., Long A., Albert S. and Brennan M. (1973) A human cell line from a pleural effusion derived from a breast carcinoma. *J Natl Cancer Inst* **51**, 1409-16
- Spector M. S., Desnoyers S., Hoeppner D. J. and Hengartner M. O. (1997) Interaction between the *C. elegans* cell-death regulators CED-9 and CED-4. *Nature* **385**, 653-6.
- Srinivasula S. M., Hegde R., Saleh A., Datta P., Shiozaki E., Chai J., Lee R. A., Robbins P. D., Fernandes-Alnemri T., Shi Y. and Alnemri E. S. (2001) A conserved XIAP-interaction motif in caspase-9 and Smac/DIABLO regulates caspase activity and apoptosis. *Nature* **410**, 112-6.
- Srinivasula S. M., Pinaki Datta, Masatomo Kobayashi, Jia-Wei Wu, Miki Fujioka, Ramesh Hegde, ZhiJia Zhang, Rula Mukattash, Teresa Fernandes-Alnemri, Yigong Shi, James B. Jaynes and Alnemri E. S. (2002) sickle, a Novel Drosophila Death Gene in the reaper/hid/grim Region, Encodes an IAP-Inhibitory Protein. *Current Biology* **12**, 125–130.
- Steller H., Abrams J. M., Grether M. E. and White K. (1994) Programmed cell death in Drosophila. *Philos Trans R Soc Lond B Biol Sci* **345**, 247-50.
- Stennicke H. R. and Salvesen G. S. (2000) Caspases - controlling intracellular signals by protease zymogen activation. *Biochim Biophys Acta* **1477**, 299-306.
- Stone K. R., Mickey D. D. and Wunderli H. (1978) Isolation of a human prostate carcinoma cell line (DU145). *Int J Cancer* **21**, 274-81
- Strasser A., O'Connor L. and Dixit V. M. (2000) Apoptosis signaling. *Annu Rev Biochem* **69**, 217-45.
- Sturm I., Petrowsky H., Volz R., Lorenz M., Radetzki S., Hillebrand T., Wolff G., Hauptmann S., Dörken B. and Daniel P. T. (2001) Analysis of p53/BAX/p16ink4a/CDKN2 in esophageal squamous cell carcinoma: High BAX and p16ink4a/CDKN2 identifies patients with good prognosis. *J Clin Oncol* **19**, 2272-2281.
- Susin S. A., Lorenzo H. K., Zamzami N., Marzo I., Snow B. E., Brothers G. M., Mangion J., Jacotot E., Costantini P., Loeffler M., Larochette N., Goodlett D. R., Aebersold R., Siderovski D. P., Penninger J. M. and Kroemer G. (1999) Molecular characterization of mitochondrial apoptosis-inducing factor. *Nature* **397**, 441-6.
- Susin S. A., Zamzami N., Castedo M., Daugas E., Wang H. G., Geley S., Fassy F., Reed J. C. and Kroemer G. (1997) The central executioner of apoptosis: multiple connections between protease activation and mitochondria in Fas/APO-1/CD95- and ceramide-induced apoptosis. *J Exp Med* **186**, 25-37.
- Suzuki Y., Nakabayashi Y. and Takahashi R. (2001) Ubiquitin-protein ligase activity of X-linked inhibitor of apoptosis protein promotes proteasomal degradation of caspase-3 and enhances its anti-apoptotic effect in Fas-induced cell death. *Proc Natl Acad Sci U S A* **98**, 8662-7.
- Suzuki Y., Imai Y., Nakayama H., Takahashi K., Takio K. and Takahashi R. (2001) A serine protease, HtrA2, is released from the mitochondria and interacts with XIAP, inducing cell death. *Mol Cell* **8**, 613-21.
- Suzuki Y., Takahashi-Niki K., Akagi T., Hashikawa T. and Takahashi R. (2004) Mitochondrial protease Omi/HtrA2 enhances caspase activation through multiple pathways. *Cell Death Differ* **11**, 208-16.
- Takahashi R., Devereaux Q., Tamm I., Welsh K., Assa-Munt N., Salvesen G. S. and Reed J. C. (1998) A single BIR domain of XIAP sufficient for inhibiting caspases. *J Biol Chem* **273**, 7787-90.
- Tamm I., Wang Y., Sausville E., Scudiero D. A., Vigna N., Oltersdorf T. and Reed J. C. (1998) IAP-family protein survivin inhibits caspase activity and apoptosis induced by Fas (CD95), Bax, caspases, and anticancer drugs. *Cancer Res* **58**, 5315-20.

- Tenev T., Zachariou A., Wilson R., Paul A. and Meier P. (2002) Jafrac2 is an IAP antagonist that promotes cell death by liberating Dronc from DIAP1. *Embo J* **21**, 5118-5129.
- Thome M. and Tschopp J. (2001) Regulation of lymphocyte proliferation and death by FLIP. *Nat Rev Immunol* **1**, 50-8.
- Thomenius M. J., Wang N. S., Reineks E. Z., Wang Z. and Distelhorst C. W. (2003) Bcl-2 on the endoplasmic reticulum regulates Bax activity by binding to BH3-only proteins. *J Biol Chem* **278**, 6243-50.
- Thornberry N. A. (1997) The caspase family of cysteine proteases. *Br Med Bull* **53**, 478-90.
- Toyota H., Yanase N., Yoshimoto T., Moriyama M., Sudo T. and Mizuguchi J. (2003) Calpain-induced Bax-cleavage product is a more potent inducer of apoptotic cell death than wild-type Bax. *Canc Lett* **189**, 221-30
- Tinel A. and Tschopp (2004) The PIDDosome, a protein complex implicated in activation of caspase-2 in response to genotoxic stress. *Science* **304**, 843-46
- Tsujimoto Y., Finger L. R., Yunis J., Nowell P. C. and Croce C. M. (1984) Cloning of the chromosome breakpoint of neoplastic B cells with the t(14;18) chromosome translocation. *Science* **226**, 1097-9.
- Twiddy D., Brown D. G., Adrain C., Jukes R., Martin S. J., Cohen G. M., MacFarlane M. and Cain K. (2004) Pro-apoptotic proteins released from the mitochondria regulate the composition and caspase-processing activity of the native Apaf-1/caspase-9 apoptosome complex. In *J Biol Chem*.
- Uren A. G., Beilharz T., O'Connell M. J., Bugg S. J., van Driel R., Vaux D. L. and Lithgow T. (1999) Role for yeast inhibitor of apoptosis (IAP)-like proteins in cell division. *Proc Natl Acad Sci U S A* **96**, 10170-5.
- Van de Craen M., Van Loo G., Pype S., Van Criekinge W., Van den brande I., Molemans F., Fiers W., Declercq W. and Vandenabeele P. (1998) Identification of a new caspase homologue: caspase-14. *Cell Death Differ* **5**, 838-46.
- van der Heiden M. G., Chadel N. S., Williamson E. K., Schumacker P. T. and Thompson C. B. (1997) Bcl-xL regulates the membrane potential and volume homeostasis of mitochondria. *Cell* **91**, 627-37.
- van Loo G., Saelens X., van Gurp M., MacFarlane M., Martin S. J. and Vandenabeele P. (2002) The role of mitochondrial factors in apoptosis: a Russian roulette with more than one bullet. *Cell Death Differ* **9**, 1031-42.
- Verhagen A. M., Ekert P. G., Pakusch M., Silke J., Connolly L. M., Reid G. E., Moritz R. L., Simpson R. J. and Vaux D. L. (2000) Identification of DIABLO, a mammalian protein that promotes apoptosis by binding to and antagonizing IAP proteins. *Cell* **102**, 43-53.
- Verhagen A. M., Silke J., Ekert P. G., Pakusch M., Kaufmann H., Connolly L. M., Day C. L., Tikoo A., Burke R., Wrobel C., Moritz R. L., Simpson R. J. and Vaux D. L. (2002) HtrA2 promotes cell death through its serine protease activity and its ability to antagonize inhibitor of apoptosis proteins. *J Biol Chem* **277**, 445-54.
- von Haefen C., Gillissen B., Hemmati P. G., Wendt J., Güner D., Mrozek A., Belka C., Dörken B. and Daniel P. T. (2004) The multidomain Bcl-2 homolog Bx but not Bak mediates synergistic induction of apoptosis by TRAIL and 5-FU through the mitochondrial apoptosis pathway. *Oncogene in press*
- von Haefen C., Wieder T., Gillissen B., Stärck L., Graupner V., Dörken B. and Daniel P. T. (2002) Ceramide induces mitochondrial activation and apoptosis via a Bax-dependent pathway in human carcinoma cells. *Oncogene* **21**, 4009-4019.
- Vucic D., Kaiser W. J. and Miller L. K. (1998) Inhibitor of apoptosis proteins physically interact with and block apoptosis induced by Drosophila proteins HID and GRIM. *Mol Cell Biol* **18**, 3300-9.

- Wang K. K., Posmantur R., Nadimpalli R., Nath R., Mohan P., Nixon R. A., Talanian R. V., Keegan M., Herzog L. and Allen H. (1998a) Caspase-mediated fragmentation of calpain inhibitor protein calpastatin during apoptosis. *Arch Biochem Biophys* **356**, 187-96.
- Wang K. K., Posmantur R., Nath R., McGinnis K., Whitton M., Talanian R. V., Glantz S. B. and Morrow J. S. (1998b) Simultaneous degradation of alphaII- and betaII-spectrin by caspase 3 (CPP32) in apoptotic cells. *J Biol Chem* **273**, 22490-7.
- Wei M. C., Lindsten T., Mootha V. K., Weiler S., Gross A., Ashiya M., Thompson C. B. and Korsmeyer S. J. (2000) tBID, a membrane-targeted death ligand, oligomerizes BAK to release cytochrome c. *Genes Dev* **14**, 2060-71.
- Wei M. C., Zong W. X., Cheng E. H., Lindsten T., Panoutsakopoulou V., Ross A. J., Roth K. A., MacGregor G. R., Thompson C. B. and Korsmeyer S. J. (2001) Proapoptotic BAX and BAK: a requisite gateway to mitochondrial dysfunction and death. *Science* **292**, 727-30.
- Weinmann P., Bommert K., Mapara M. Y., Dorken B. and Bargou R. C. (1997) Overexpression of the death-promoting gene bax-alpha sensitizes human BL-41 Burkitt lymphoma cells for surface IgM-mediated apoptosis. *Eur J Immunol* **27**, 2466-8.
- Wu D., Wallen HD and G. N. (1997) Interaction and regulation of subcellular localization of CED-4 by CED-9. *Science* **275**, 1126-9.
- Wyllie A. H. (1980) Glucocorticoid-induced thymocyte apoptosis is associated with endogenous endonuclease activation. *Nature* **284**, 555-6.
- Wyllie A. H., Kerr J. F. and Currie A. R. (1980) Cell death: the significance of apoptosis. *Int Rev Cytol* **68**, 251-306.
- Yamaguchi K., Nagai S., Ninomiya-Tsuji J., Nishita M., Tamai K., Irie K., Ueno N., Nishida E., Shibuya H. and Matsumoto K. (1999) XIAP, a cellular member of the inhibitor of apoptosis protein family, links the receptors to TAB1-TAK1 in the BMP signaling pathway. *Embo J* **18**, 179-87.
- Yang L., Mashima T., Sato S., Mochizuki M., Sakamoto H., Yamori T., Oh-Hara T. and Tsuruo T. (2003) Predominant suppression of apotosome by inhibitor of apoptosis protein in non-small cell lung cancer H460 cells: therapeutic effect of a novel polyarginine-conjugated Smac peptide. *Cancer Res* **63**, 831-7.
- Yang Q. H. and Du C. (2004) Smac/DIABLO selectively reduces the levels of c-IAP1 and c-IAP2 but not that of XIAP and livin in HeLa cells. *J Biol Chem* **279**, 16963-70.
- Yin X. M., Oltvai Z. N. and Korsmeyer S. J. (1994) BH1 and BH2 domains of Bcl-2 are required for inhibition of apoptosis and heterodimerization with Bax. *Nature* **369**, 321-3.
- Yoo S. J., Huh J. R., Muro I., Yu H., Wang L., Wang S. L., Feldman R. M., Clem R. J., Muller H. A. and Hay B. A. (2002) Hid, Rpr and Grim negatively regulate DIAP1 levels through distinct mechanisms. *Nat Cell Biol* **4**, 416-24.
- Yu J., Zhang L., Hwang P. M., Kinzler K. W. and Vogelstein B. (2001) PUMA induces the rapid apoptosis of colorectal cancer cells. *Mol Cell* **7**, 673-82.
- Zamzami N., El Hamel C., Maisse C., Brenner C., Munoz-Pinedo C., Belzacq A. S., Costantini P., Vieira H., Loeffler M., Molle G. and Kroemer G. (2000) Bid acts on the permeability transition pore complex to induce apoptosis. *Oncogene* **19**, 6342-50.
- Zha J., Harada H., Yang E., Jockel J. and Korsmeyer S. J. (1996) Serine phosphorylation of death agonist BAD in response to survival factor results in binding to 14-3-3 not BCL-X(L). *Cell* **87**, 619-28.
- Zhang L., Yu J., Park B. H., Kinzler K. W. and Vogelstein B. (2000) Role of BAX in the apoptotic response to anticancer agents. *Science* **290**, 989-92.