

6 Literaturverzeichnis

- Adams, M.D., S.E. Celniker, R.A. Holt, C.A. Evans, J.D. Gocayne, P.G. Amanatides, S.E. Scherer, P.W. Li, R.A. Hoskins, R.F. Galle, R.A. George, S.E. Lewis, S. Richards, M. Ashburner, S.N. Henderson, G.G. Sutton, J.R. Wortman, M.D. Yandell, Q. Zhang, L.X. Chen, R.C. Brandon, Y.H. Rogers, R.G. Blazej, M. Champe, B.D. Pfeiffer, K.H. Wan, C. Doyle, E.G. Baxter, G. Helt, C.R. Nelson, G.L. Gabor, J.F. Abril, A. Agbayani, H.J. An, C. Andrews-Pfannkoch, D. Baldwin, R.M. Ballew, A. Basu, J. Baxendale, L. Bayraktaroglu, E.M. Beasley, K.Y. Beeson, P.V. Benos, B.P. Berman, D. Bhandari, S. Bolshakov, D. Borkova, M.R. Botchan, J. Bouck, P. Brokstein, P. Brottier, K.C. Burtis, D.A. Busam, H. Butler, E. Cadieu, A. Center, I. Chandra, J.M. Cherry, S. Cawley, C. Dahlke, L.B. Davenport, P. Davies, B. de Pablos, A. Delcher, Z. Deng, A.D. Mays, I. Dew, S.M. Dietz, K. Dodson, L.E. Doup, M. Downes, S. Dugan-Rocha, B.C. Dunkov, P. Dunn, K.J. Durbin, C.C. Evangelista, C. Ferraz, S. Ferriera, W. Fleischmann, C. Fosler, A.E. Gabrielian, N.S. Garg, W.M. Gelbart, K. Glasser, A. Glodek, F. Gong, J.H. Gorrell, Z. Gu, P. Guan, M. Harris, N.L. Harris, D. Harvey, T.J. Heiman, J.R. Hernandez, J. Houck, D. Hostin, K.A. Houston, T.J. Howland, M.H. Wei, C. Ibegwam, et al. 2000. The genome sequence of *Drosophila melanogaster*. *Science*. 287:2185-95.
- Adamski, F.M., M.Y. Zhu, F. Bahiraei, and B.H. Shieh. 1998. Interaction of eye protein kinase C and INAD in *Drosophila*. Localization of binding domains and electrophysiological characterization of a loss of association in transgenic flies. *J Biol Chem.* 273:17713-9.
- Agazie, Y.M., and M.J. Hayman. 2003. Molecular mechanism for a role of SHP2 in epidermal growth factor receptor signaling. *Mol Cell Biol.* 23:7875-86.
- Airaksinen, M.S., A. Titievsky, and M. Saarma. 1999. GDNF family neurotrophic factor signaling: four masters, one servant? *Mol Cell Neurosci.* 13:313-25.
- Alberti, L., M.G. Borrello, S. Ghizzoni, F. Torriti, M.G. Rizzetti, and M.A. Pierotti. 1998. Grb2 binding to the different isoforms of Ret tyrosine kinase. *Oncogene*. 17:1079-87.

- Arighi, E., L. Alberti, F. Torriti, S. Ghizzoni, M.G. Rizzetti, G. Pelicci, B. Pasini, I. Bongarzone, C. Piutti, M.A. Pierotti, and M.G. Borrello. 1997. Identification of Shc docking site on Ret tyrosine kinase. *Oncogene*. 14:773-82.
- Asai, N., H. Murakami, T. Iwashita, and M. Takahashi. 1996. A mutation at tyrosine 1062 in MEN2A-Ret and MEN2B-Ret impairs their transforming activity and association with shc adaptor proteins. *J Biol Chem*. 271:17644-9.
- Ausubel, F.M., Brent,R., Kingston,R.E., Moore,D.D., Seidman,I.G., Smith,J.A., and Struhl,K. 1987. Current protocols in molecular biology. *Greene Publishing Associates and Wiley-Interscience, New York*.
- Baloh, R.H., M.G. Tansey, E.M. Johnson, Jr., and J. Milbrandt. 2000. Functional mapping of receptor specificity domains of glial cell line-derived neurotrophic factor (GDNF) family ligands and production of GFRalpha1 RET-specific agonists. *J Biol Chem*. 275:3412-20.
- Balter, M. 1995. Chernobyl's thyroid cancer toll. *Science*. 270:1758-9.
- Bardelli, A., P. Longati, D. Gramaglia, M.C. Stella, and P.M. Comoglio. 1997. Gab1 coupling to the HGF/Met receptor multifunctional docking site requires binding of Grb2 and correlates with the transforming potential. *Oncogene*. 15:3103-11.
- Barres, B.A., B.E. Silverstein, D.P. Corey, and L.L. Chun. 1988. Immunological, morphological, and electrophysiological variation among retinal ganglion cells purified by panning. *Neuron*. 1:791-803.
- Barros, E.J., O.F. Santos, K. Matsumoto, T. Nakamura, and S.K. Nigam. 1995. Differential tubulogenic and branching morphogenetic activities of growth factors: implications for epithelial tissue development. *Proc Natl Acad Sci U S A*. 92:4412-6.
- Baverstock, K., B. Egloff, A. Pinchera, C. Ruchti, and D. Williams. 1992. Thyroid cancer after Chernobyl. *Nature*. 359:21-2.
- Basset, V., R.P. Scott, and C.F. Ibanez. 2000. Signaling complexes and protein-protein interactions involved in the activation of the Ras and phosphatidylinositol 3-kinase pathways by the c-Ret receptor tyrosine kinase. *J Biol Chem*. 275:39159-66.

- Bezprozvanny, I., and A. Maximov. 2001. Classification of PDZ domains. *FEBS Lett.* 509:457-62.
- Bibel, M., E. Hoppe, and Y.A. Barde. 1999. Biochemical and functional interactions between the neurotrophin receptors trk and p75NTR. *Embo J.* 18:616-22.
- Bier, E. 2000. Drawing lines in the Drosophila wing: initiation of wing vein development. *Curr Opin Genet Dev.* 10:393-8.
- Birchmeier, C., W. Birchmeier, E. Gherardi, and G.F. Vande Woude. 2003. Met, metastasis, motility and more. *Nat Rev Mol Cell Biol.* 4:915-25.
- Birchmeier, C., and E. Gherardi. 1998. Developmental roles of HGF/SF and its receptor, the c-Met tyrosine kinase. *Trends Cell Biol.* 8:404-10.
- Bladt, F., D. Riethmacher, S. Isenmann, A. Aguzzi, and C. Birchmeier. 1995. Essential role for the c-met receptor in the migration of myogenic precursor cells into the limb bud. *Nature.* 376:768-71.
- Blume-Jensen, P., and T. Hunter. 2001. Oncogenic kinase signalling. *Nature.* 411:355-65.
- Boccaccio, C., M. Ando, and P.M. Comoglio. 2002. A differentiation switch for genetically modified hepatocytes. *Faseb J.* 16:120-2.
- Bockers, T.M., M.G. Mameza, M.R. Kreutz, J. Bockmann, C. Weise, F. Buck, D. Richter, E.D. Gundelfinger, and H.J. Kreienkamp. 2001. Synaptic scaffolding proteins in rat brain. Ankyrin repeats of the multidomain Shank protein family interact with the cytoskeletal protein alpha-fodrin. *J Biol Chem.* 276:40104-12.
- Bockmann, J., M.R. Kreutz, E.D. Gundelfinger, and T.M. Bockers. 2002. ProSAP/Shank postsynaptic density proteins interact with insulin receptor tyrosine kinase substrate IRSp53. *J Neurochem.* 83:1013-7.
- Boeckers, T.M., M.R. Kreutz, C. Winter, W. Zuschratter, K.H. Smalla, L. Sanmarti-Vila, H. Wex, K. Langnaese, J. Bockmann, C.C. Garner, and E.D. Gundelfinger. 1999a. Proline-rich synapse-associated protein-1/cortactin binding protein 1

- (ProSAP1/CortBP1) is a PDZ-domain protein highly enriched in the postsynaptic density. *J Neurosci.* 19:6506-18.
- Boeckers, T.M., C. Winter, K.H. Smalla, M.R. Kreutz, J. Bockmann, C. Seidenbecher, C.C. Garner, and E.D. Gundelfinger. 1999b. Proline-rich synapse-associated proteins ProSAP1 and ProSAP2 interact with synaptic proteins of the SAPAP/GKAP family. *Biochem Biophys Res Commun.* 264:247-52.
- Bongarzone, I., N. Monzini, M.G. Borrello, C. Carcano, G. Ferraresi, E. Arighi, P. Mondellini, G. Della Porta, and M.A. Pierotti. 1993. Molecular characterization of a thyroid tumor-specific transforming sequence formed by the fusion of ret tyrosine kinase and the regulatory subunit RI alpha of cyclic AMP-dependent protein kinase A. *Mol Cell Biol.* 13:358-66.
- Bonn, D. 1996. RET oncogene puzzle in Chernobyl thyroid tumours. *Lancet.* 347:1176.
- Borg, J.P., S. Marchetto, A. Le Bivic, V. Ollendorff, F. Jaulin-Bastard, H. Saito, E. Fournier, J. Adelaide, B. Margolis, and D. Birnbaum. 2000. ERBIN: a basolateral PDZ protein that interacts with the mammalian ERBB2/HER2 receptor. *Nat Cell Biol.* 2:407-14.
- Borrello, M.G., L. Alberti, E. Arighi, I. Bongarzone, C. Battistini, A. Bardelli, B. Pasini, C. Piutti, M.G. Rizzetti, P. Mondellini, M.T. Radice, and M.A. Pierotti. 1996. The full oncogenic activity of Ret/ptc2 depends on tyrosine 539, a docking site for phospholipase C γ . *Mol Cell Biol.* 16:2151-63.
- Brinkmann, V., H. Foroutan, M. Sachs, K.M. Weidner, and W. Birchmeier. 1995. Hepatocyte growth factor/scatter factor induces a variety of tissue-specific morphogenic programs in epithelial cells. *J Cell Biol.* 131:1573-86.
- Buchert, M., S. Schneider, V. Meskenaite, M.T. Adams, E. Canaani, T. Baechi, K. Moelling, and C.M. Hovens. 1999. The junction-associated protein AF-6 interacts and clusters with specific Eph receptor tyrosine kinases at specialized sites of cell-cell contact in the brain. *J Cell Biol.* 144:361-71.

- Buday, L., and J. Downward. 1993. Epidermal growth factor regulates p21ras through the formation of a complex of receptor, Grb2 adapter protein, and Sos nucleotide exchange factor. *Cell*. 73:611-20.
- Buechner, M., D.H. Hall, H. Bhatt, and E.M. Hedgecock. 1999. Cystic canal mutants in *Caenorhabditis elegans* are defective in the apical membrane domain of the renal (excretory) cell. *Dev Biol*. 214:227-41.
- Cao, T.T., H.W. Deacon, D. Reczek, A. Bretscher, and M. von Zastrow. 1999. A kinase-regulated PDZ-domain interaction controls endocytic sorting of the beta2-adrenergic receptor. *Nature*. 401:286-90.
- Carlomagno, F., G. De Vita, M.T. Berlingieri, V. de Franciscis, R.M. Melillo, V. Colantuoni, M.H. Kraus, P.P. Di Fiore, A. Fusco, and M. Santoro. 1996. Molecular heterogeneity of RET loss of function in Hirschsprung's disease. *Embo J*. 15:2717-25.
- Cho, K.O., C.A. Hunt, and M.B. Kennedy. 1992. The rat brain postsynaptic density fraction contains a homolog of the *Drosophila* discs-large tumor suppressor protein. *Neuron*. 9:929-42.
- Cik, M., S. Masure, A.S. Lesage, I. Van Der Linden, P. Van Gompel, M.N. Pangalos, R.D. Gordon, and J.E. Leysen. 2000. Binding of GDNF and neurturin to human GDNF family receptor alpha 1 and 2. Influence of cRET and cooperative interactions. *J Biol Chem*. 275:27505-12.
- Cohen, A.R., D.F. Woods, S.M. Marfatia, Z. Walther, A.H. Chishti, J.M. Anderson, and D.F. Wood. 1998. Human CASK/LIN-2 binds syndecan-2 and protein 4.1 and localizes to the basolateral membrane of epithelial cells. *J Cell Biol*. 142:129-38.
- Crowder, R.J., H. Enomoto, M. Yang, E.M. Johnson, Jr., and J. Milbrandt. 2004. Dok-6, a novel p62 Dok family member, promotes Ret-mediated neurite outgrowth. *J Biol Chem*.
- Cybulsky, A.V., P.R. Goodyer, and A.J. McTavish. 1994. Epidermal growth factor receptor activation in developing rat kidney. *Am J Physiol*. 267:F428-36.

- Datta, S.R., A. Brunet, and M.E. Greenberg. 1999. Cellular survival: a play in three Akts. *Genes Dev.* 13:2905-27.
- de Graaff, E., S. Srinivas, C. Kilkenny, V. D'Agati, B.S. Mankoo, F. Costantini, and V. Pachnis. 2001. Differential activities of the RET tyrosine kinase receptor isoforms during mammalian embryogenesis. *Genes Dev.* 15:2433-44.
- Doyle, D.A., A. Lee, J. Lewis, E. Kim, M. Sheng, and R. MacKinnon. 1996. Crystal structures of a complexed and peptide-free membrane protein-binding domain: molecular basis of peptide recognition by PDZ. *Cell.* 85:1067-76.
- Du, Y., S.A. Weed, W.C. Xiong, T.D. Marshall, and J.T. Parsons. 1998. Identification of a novel cortactin SH3 domain-binding protein and its localization to growth cones of cultured neurons. *Mol Cell Biol.* 18:5838-51.
- Durbec, P., C.V. Marcos-Gutierrez, C. Kilkenny, M. Grigoriou, K. Wartiowaara, P. Suvanto, D. Smith, B. Ponder, F. Costantini, M. Saarma, and et al. 1996. GDNF signalling through the Ret receptor tyrosine kinase. *Nature.* 381:789-93.
- Durick, K., R.Y. Wu, G.N. Gill, and S.S. Taylor. 1996. Mitogenic signaling by Ret/ptc2 requires association with enigma via a LIM domain. *J Biol Chem.* 271:12691-4.
- Edlund, H. 2002. Pancreatic organogenesis--developmental mechanisms and implications for therapy. *Nat Rev Genet.* 3:524-32.
- Elowe, S., S.J. Holland, S. Kulkarni, and T. Pawson. 2001. Downregulation of the Ras-mitogen-activated protein kinase pathway by the EphB2 receptor tyrosine kinase is required for ephrin-induced neurite retraction. *Mol Cell Biol.* 21:7429-41.
- Enomoto, H., T. Araki, A. Jackman, R.O. Heuckeroth, W.D. Snider, E.M. Johnson, Jr., and J. Milbrandt. 1998. GFR alpha1-deficient mice have deficits in the enteric nervous system and kidneys. *Neuron.* 21:317-24.
- Fanning, A.S., and J.M. Anderson. 1999. Protein modules as organizers of membrane structure. *Curr Opin Cell Biol.* 11:432-9.

- Frisen, J., J. Holmberg, and M. Barbacid. 1999. Ephrins and their Eph receptors: multitalented directors of embryonic development. *Embo J.* 18:5159-65.
- Fugazzola, L., M.A. Pierotti, E. Vigano, F. Pacini, T.V. Vorontsova, and I. Bongarzone. 1996. Molecular and biochemical analysis of RET/PTC4, a novel oncogenic rearrangement between RET and ELE1 genes, in a post-Chernobyl papillary thyroid cancer. *Oncogene*. 13:1093-7.
- Fukuda, N., M. Ichihara, T. Morinaga, K. Kawai, H. Hayashi, Y. Murakumo, S. Matsuo, and M. Takahashi. 2003. Identification of a novel GDNF-inducible gene required for renal branching morphogenesis. *J Biol Chem*.
- Fukuda, T., K. Kiuchi, and M. Takahashi. 2002. Novel mechanism of regulation of Rac activity and lamellipodia formation by RET tyrosine kinase. *J Biol Chem*. 277:19114-21.
- Garratt, A.N., S. Britsch, and C. Birchmeier. 2000. Neuregulin, a factor with many functions in the life of a schwann cell. *Bioessays*. 22:987-96.
- Glynnne, P.A., and T.J. Evans. 2002. Role of the PDZ scaffolding protein in tubule cells in maintenance of polarised function. *Exp Nephrol*. 10:307-12.
- Grieco, M., M. Santoro, M.T. Berlingieri, R.M. Melillo, R. Donghi, I. Bongarzone, M.A. Pierotti, G. Della Porta, A. Fusco, and G. Vecchio. 1990. PTC is a novel rearranged form of the ret proto-oncogene and is frequently detected in vivo in human thyroid papillary carcinomas. *Cell*. 60:557-63.
- Grimm, J. 2001. Funktionelle Interaktion von neuen Mitgliedern der dok-Familie mit der Rezeptor-Tyrosinkinase c-Ret. *Dissertation, HU Berlin*.
- Grimm, J., M. Sachs, S. Britsch, S. Di Cesare, T. Schwarz-Romond, K. Alitalo, and W. Birchmeier. 2001. Novel p62dok family members, dok-4 and dok-5, are substrates of the c-Ret receptor tyrosine kinase and mediate neuronal differentiation. *J Cell Biol*. 154:345-54.
- Guy, G.R., P. Yusoff, D. Bangarusamy, C.W. Fong, and E.S. Wong. 2002. Dockers at the crossroads. *Cell Signal*. 14:11-20.

- Hanahan, D., and R.A. Weinberg. 2000. The hallmarks of cancer. *Cell.* 100:57-70.
- Hayashi, H., M. Ichihara, T. Iwashita, H. Murakami, Y. Shimono, K. Kawai, K. Kurokawa, Y. Murakumo, T. Imai, H. Funahashi, A. Nakao, and M. Takahashi. 2000. Characterization of intracellular signals via tyrosine 1062 in RET activated by glial cell line-derived neurotrophic factor. *Oncogene.* 19:4469-75.
- Hayashi, Y., T. Iwashita, H. Murakamai, Y. Kato, K. Kawai, K. Kurokawa, I. Tohnai, M. Ueda, and M. Takahashi. 2001. Activation of BMK1 via tyrosine 1062 in RET by GDNF and MEN2A mutation. *Biochem Biophys Res Commun.* 281:682-9.
- Hiles, I.D., M. Otsu, S. Volinia, M.J. Fry, I. Gout, R. Dhand, G. Panayotou, F. Ruiz-Larrea, A. Thompson, N.F. Totty, and et al. 1992. Phosphatidylinositol 3-kinase: structure and expression of the 110 kd catalytic subunit. *Cell.* 70:419-29.
- Hilpela, P., M.K. Vartiainen, and P. Lappalainen. 2004. Regulation of the actin cytoskeleton by PI(4,5)P₂ and PI(3,4,5)P₃. *Curr Top Microbiol Immunol.* 282:117-63.
- Hogan, B.L., and P.A. Kolodziej. 2002. Organogenesis: molecular mechanisms of tubulogenesis. *Nat Rev Genet.* 3:513-23.
- Holt, K.H., L. Olson, W.S. Moye-Rowley, and J.E. Pessin. 1994. Phosphatidylinositol 3-kinase activation is mediated by high-affinity interactions between distinct domains within the p110 and p85 subunits. *Mol Cell Biol.* 14:42-9.
- Homma, S., R.W. Oppenheim, H. Yaginuma, and S. Kimura. 2000. Expression pattern of GDNF, c-ret, and GFRalphas suggests novel roles for GDNF ligands during early organogenesis in the chick embryo. *Dev Biol.* 217:121-37.
- Huang, C., Y. Ni, T. Wang, Y. Gao, C.C. Haudenschild, and X. Zhan. 1997. Down-regulation of the filamentous actin cross-linking activity of cortactin by Src-mediated tyrosine phosphorylation. *J Biol Chem.* 272:13911-5.
- Huang, Y.Z., M. Zang, W.C. Xiong, Z. Luo, and L. Mei. 2003. Erbin suppresses the MAP kinase pathway. *J Biol Chem.* 278:1108-14.

- Hunter, T. 2000. Signaling--2000 and beyond. *Cell*. 100:113-27.
- Ishizaka, Y., F. Itoh, T. Tahira, I. Ikeda, T. Sugimura, J. Tucker, A. Fertitta, A.V. Carrano, and M. Nagao. 1989. Human ret proto-oncogene mapped to chromosome 10q11.2. *Oncogene*. 4:1519-21.
- Ivanchuk, S.M., C. Eng, W.K. Cavenee, and L.M. Mulligan. 1997. The expression of RET and its multiple splice forms in developing human kidney. *Oncogene*. 14:1811-8.
- Ivanchuk, S.M., S.M. Myers, and L.M. Mulligan. 1998. Expression of RET 3' splicing variants during human kidney development. *Oncogene*. 16:991-6.
- Jhiang, S.M. 2000. The RET proto-oncogene in human cancers. *Oncogene*. 19:5590-7.
- Jones, N., and D.J. Dumont. 1999. Recruitment of Dok-R to the EGF receptor through its PTB domain is required for attenuation of Erk MAP kinase activation. *Curr Biol*. 9:1057-60.
- Jung Kang, Y., E. Su Jeon, H. Jin Lee, Y.S. Oh, P.G. Suh, J. Sup Jung, M. Donowitz, and J. Ho Kim. 2004. NHERF2 increases platelet-derived growth factor-induced proliferation through PI-3-kinase/Akt-, ERK-, and Src family kinase-dependent pathway. *Cell Signal*. 16:791-800.
- Kaech, S.M., C.W. Whitfield, and S.K. Kim. 1998. The LIN-2/LIN-7/LIN-10 complex mediates basolateral membrane localization of the *C. elegans* EGF receptor LET-23 in vulval epithelial cells. *Cell*. 94:761-71.
- Kawaida, K., K. Matsumoto, H. Shimazu, and T. Nakamura. 1994. Hepatocyte growth factor prevents acute renal failure and accelerates renal regeneration in mice. *Proc Natl Acad Sci U S A*. 91:4357-61.
- Khwaja, A., K. Lehmann, B.M. Marte, and J. Downward. 1998. Phosphoinositide 3-kinase induces scattering and tubulogenesis in epithelial cells through a novel pathway. *J Biol Chem*. 273:18793-801.

- Kim, E., M. Niethammer, A. Rothschild, Y.N. Jan, and M. Sheng. 1995. Clustering of Shaker-type K⁺ channels by interaction with a family of membrane-associated guanylate kinases. *Nature*. 378:85-8.
- Kim, S.K. 1997. Polarized signaling: basolateral receptor localization in epithelial cells by PDZ-containing proteins. *Curr Opin Cell Biol*. 9:853-9.
- Kinashi, T., J.A. Escobedo, L.T. Williams, K. Takatsu, and T.A. Springer. 1995. Receptor tyrosine kinase stimulates cell-matrix adhesion by phosphatidylinositol 3 kinase and phospholipase C-gamma 1 pathways. *Blood*. 86:2086-90.
- Klugbauer, S., E.P. Demidchik, E. Lengfelder, and H.M. Rabes. 1998. Detection of a novel type of RET rearrangement (PTC5) in thyroid carcinomas after Chernobyl and analysis of the involved RET-fused gene RFG5. *Cancer Res*. 58:198-203.
- Kolch, W. 2003. Erbin: sorting out ErbB2 receptors or giving Ras a break? *Sci STKE*. 2003:pe37.
- Kroll, J., and J. Waltenberger. 1997. The vascular endothelial growth factor receptor KDR activates multiple signal transduction pathways in porcine aortic endothelial cells. *J Biol Chem*. 272:32521-7.
- Krugmann, S., I. Jordens, K. Gevaert, M. Driessens, J. Vandekerckhove, and A. Hall. 2001. Cdc42 induces filopodia by promoting the formation of an IRSp53:Mena complex. *Curr Biol*. 11:1645-55.
- Kullander, K., and R. Klein. 2002. Mechanisms and functions of Eph and ephrin signalling. *Nat Rev Mol Cell Biol*. 3:475-86.
- Kurokawa, K., T. Iwashita, H. Murakami, H. Hayashi, K. Kawai, and M. Takahashi. 2001. Identification of SNT/FRS2 docking site on RET receptor tyrosine kinase and its role for signal transduction. *Oncogene*. 20:1929-38.
- Kurokawa, K., K. Kawai, M. Hashimoto, Y. Ito, and M. Takahashi. 2003. Cell signalling and gene expression mediated by RET tyrosine kinase. *J Intern Med*. 253:627-33.

- Laemmli, U.K. 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature*. 227:680-5.
- Lanzi, C., M.G. Borrello, I. Bongarzone, A. Migliazza, A. Fusco, M. Grieco, M. Santoro, R.A. Gambetta, F. Zunino, G. Della Porta, and et al. 1992. Identification of the product of two oncogenic rearranged forms of the RET proto-oncogene in papillary thyroid carcinomas. *Oncogene*. 7:2189-94.
- Leevers, S.J., B. Vanhaesebroeck, and M.D. Waterfield. 1999. Signalling through phosphoinositide 3-kinases: the lipids take centre stage. *Curr Opin Cell Biol.* 11:219-25.
- Lim, S., S. Naisbitt, J. Yoon, J.I. Hwang, P.G. Suh, M. Sheng, and E. Kim. 1999. Characterization of the Shank family of synaptic proteins. Multiple genes, alternative splicing, and differential expression in brain and development. *J Biol Chem.* 274:29510-8.
- Lore, F., G. Di Cairano, and F. Talidis. 2000. Unilateral renal agenesis in a family with medullary thyroid carcinoma. *N Engl J Med.* 342:1218-9.
- Lore, F., F. Talidis, G. Di Cairano, and A. Renieri. 2001. Multiple endocrine neoplasia type 2 syndromes may be associated with renal malformations. *J Intern Med.* 250:37-42.
- Lorenzo, M.J., C. Eng, L.M. Mulligan, T.J. Stonehouse, C.S. Healey, B.A. Ponder, and D.P. Smith. 1995. Multiple mRNA isoforms of the human RET proto-oncogene generated by alternate splicing. *Oncogene*. 10:1377-83.
- Lorenzo, M.J., G.D. Gish, C. Houghton, T.J. Stonehouse, T. Pawson, B.A. Ponder, and D.P. Smith. 1997. RET alternate splicing influences the interaction of activated RET with the SH2 and PTB domains of Shc, and the SH2 domain of Grb2. *Oncogene*. 14:763-71.
- Ma, L., Y.Z. Huang, G.M. Pitcher, J.G. Valtschanoff, Y.H. Ma, L.Y. Feng, B. Lu, W.C. Xiong, M.W. Salter, R.J. Weinberg, and L. Mei. 2003. Ligand-dependent recruitment of the ErbB4 signaling complex into neuronal lipid rafts. *J Neurosci.* 23:3164-75.

- Malumbres, M., and A. Pellicer. 1998. RAS pathways to cell cycle control and cell transformation. *Front Biosci.* 3:d887-912.
- Manser, E., T.H. Loo, C.G. Koh, Z.S. Zhao, X.Q. Chen, L. Tan, I. Tan, T. Leung, and L. Lim. 1998. PAK kinases are directly coupled to the PIX family of nucleotide exchange factors. *Mol Cell.* 1:183-92.
- Maroun, C.R., M.A. Naujokas, M. Holgado-Madruga, A.J. Wong, and M. Park. 2000. The tyrosine phosphatase SHP-2 is required for sustained activation of extracellular signal-regulated kinase and epithelial morphogenesis downstream from the met receptor tyrosine kinase. *Mol Cell Biol.* 20:8513-25.
- Marra, F., M. Pinzani, R. DeFranco, G. Laffi, and P. Gentilini. 1995. Involvement of phosphatidylinositol 3-kinase in the activation of extracellular signal-regulated kinase by PDGF in hepatic stellate cells. *FEBS Lett.* 376:141-5.
- Marshall, C.J. 1994. MAP kinase kinase kinase, MAP kinase kinase and MAP kinase. *Curr Opin Genet Dev.* 4:82-9.
- Marshall, C.J. 1995. Specificity of receptor tyrosine kinase signaling: transient versus sustained extracellular signal-regulated kinase activation. *Cell.* 80:179-85.
- Maudsley, S., A.M. Zamah, N. Rahman, J.T. Blitzer, L.M. Luttrell, R.J. Lefkowitz, and R.A. Hall. 2000. Platelet-derived growth factor receptor association with Na(+)/H(+) exchanger regulatory factor potentiates receptor activity. *Mol Cell Biol.* 20:8352-63.
- McIntyre, E., P. Bond, F. Douglas, T. Lennard, R. Peaston, and P. Perros. 2003. Multiple endocrine neoplasia type 2A: an unusual clinical presentation and association with renal dysplasia. *Cancer Genet Cytogenet.* 141:157-9.
- Melillo, R.M., M. Santoro, S.H. Ong, M. Billaud, A. Fusco, Y.R. Hadari, J. Schlessinger, and I. Lax. 2001. Docking protein FRS2 links the protein tyrosine kinase RET and its oncogenic forms with the mitogen-activated protein kinase signaling cascade. *Mol Cell Biol.* 21:4177-87.

- Meyer-Franke, A., M.R. Kaplan, F.W. Pfrieger, and B.A. Barres. 1995. Characterization of the signaling interactions that promote the survival and growth of developing retinal ganglion cells in culture. *Neuron*. 15:805-19.
- Miki, H., H. Yamaguchi, S. Suetsugu, and T. Takenawa. 2000. IRSp53 is an essential intermediate between Rac and WAVE in the regulation of membrane ruffling. *Nature*. 408:732-5.
- Mikoshiba, K., and M. Hattori. 2000. IP3 receptor-operated calcium entry. *Sci STKE*. 2000:PE1.
- Miyamoto, N., M. Yoshida, S. Kuratani, I. Matsuo, and S. Aizawa. 1997. Defects of urogenital development in mice lacking Emx2. *Development*. 124:1653-64.
- Mohammadi, M., J. Schlessinger, and S.R. Hubbard. 1996. Structure of the FGF receptor tyrosine kinase domain reveals a novel autoinhibitory mechanism. *Cell*. 86:577-87.
- Montesano, R., G. Schaller, and L. Orci. 1991. Induction of epithelial tubular morphogenesis in vitro by fibroblast-derived soluble factors. *Cell*. 66:697-711.
- Morais Cabral, J.H., C. Petosa, M.J. Sutcliffe, S. Raza, O. Byron, F. Poy, S.M. Marfatia, A.H. Chishti, and R.C. Liddington. 1996. Crystal structure of a PDZ domain. *Nature*. 382:649-52.
- Morgenstern, J.P., and H. Land. 1990. Advanced mammalian gene transfer: high titre retroviral vectors with multiple drug selection markers and a complementary helper-free packaging cell line. *Nucleic Acids Res*. 18:3587-96.
- Muller, B.M., U. Kistner, S. Kindler, W.J. Chung, S. Kuhlendahl, S.D. Fenster, L.F. Lau, R.W. Veh, R.L. Huganir, E.D. Gundelfinger, and C.C. Garner. 1996. SAP102, a novel postsynaptic protein that interacts with NMDA receptor complexes in vivo. *Neuron*. 17:255-65.
- Muller, M., A. Morotti, and C. Ponzetto. 2002. Activation of NF-kappaB is essential for hepatocyte growth factor-mediated proliferation and tubulogenesis. *Mol Cell Biol*. 22:1060-72.

- Mulligan, L.M., J.B. Kwok, C.S. Healey, M.J. Elsdon, C. Eng, E. Gardner, D.R. Love, S.E. Mole, J.K. Moore, L. Papi, and et al. 1993. Germ-line mutations of the RET proto-oncogene in multiple endocrine neoplasia type 2A. *Nature*. 363:458-60.
- Myers, S.M., C. Eng, B.A. Ponder, and L.M. Mulligan. 1995. Characterization of RET proto-oncogene 3' splicing variants and polyadenylation sites: a novel C-terminus for RET. *Oncogene*. 11:2039-45.
- Nagler, K., D.H. Mauch, and F.W. Pfrieger. 2001. Glia-derived signals induce synapse formation in neurones of the rat central nervous system. *J Physiol*. 533:665-79.
- Naisbitt, S., E. Kim, J.C. Tu, B. Xiao, C. Sala, J. Valtschanoff, R.J. Weinberg, P.F. Worley, and M. Sheng. 1999. Shank, a novel family of postsynaptic density proteins that binds to the NMDA receptor/PSD-95/GKAP complex and cortactin. *Neuron*. 23:569-82.
- Niemann, C., V. Brinkmann, E. Spitzer, G. Hartmann, M. Sachs, H. Naundorf, and W. Birchmeier. 1998. Reconstitution of mammary gland development in vitro: requirement of c-met and c-erbB2 signaling for branching and alveolar morphogenesis. *J Cell Biol*. 143:533-45.
- Niethammer, M., E. Kim, and M. Sheng. 1996. Interaction between the C terminus of NMDA receptor subunits and multiple members of the PSD-95 family of membrane-associated guanylate kinases. *J Neurosci*. 16:2157-63.
- Nose, A., A. Nagafuchi, and M. Takeichi. 1988. Expressed recombinant cadherins mediate cell sorting in model systems. *Cell*. 54:993-1001.
- O'Brien, L.E., K. Tang, E.S. Kats, A. Schutz-Geschwender, J.H. Lipschutz, and K.E. Mostov. 2004. ERK and MMPs sequentially regulate distinct stages of epithelial tubule development. *Dev Cell*. 7:21-32.
- Ohiwa, M., H. Murakami, T. Iwashita, N. Asai, Y. Iwata, T. Imai, H. Funahashi, H. Takagi, and M. Takahashi. 1997. Characterization of Ret-Shc-Grb2 complex induced by GDNF, MEN 2A, and MEN 2B mutations. *Biochem Biophys Res Commun*. 237:747-51.

- Ohmichi, H., U. Koshimizu, K. Matsumoto, and T. Nakamura. 1998. Hepatocyte growth factor (HGF) acts as a mesenchyme-derived morphogenic factor during fetal lung development. *Development*. 125:1315-24.
- Okkenhaug, K., and B. Vanhaesbroeck. 2003. PI3K in lymphocyte development, differentiation and activation. *Nat Rev Immunol*. 3:317-30.
- Ong, S.H., G.R. Guy, Y.R. Hadari, S. Laks, N. Gotoh, J. Schlessinger, and I. Lax. 2000. FRS2 proteins recruit intracellular signaling pathways by binding to diverse targets on fibroblast growth factor and nerve growth factor receptors. *Mol Cell Biol*. 20:979-89.
- Pachnis, V., B. Mankoo, and F. Costantini. 1993. Expression of the c-ret proto-oncogene during mouse embryogenesis. *Development*. 119:1005-17.
- Pandey, A., H. Duan, P.P. Di Fiore, and V.M. Dixit. 1995. The Ret receptor protein tyrosine kinase associates with the SH2-containing adapter protein Grb10. *J Biol Chem*. 270:21461-3.
- Pandey, A., X. Liu, J.E. Dixon, P.P. Di Fiore, and V.M. Dixit. 1996. Direct association between the Ret receptor tyrosine kinase and the Src homology 2-containing adapter protein Grb7. *J Biol Chem*. 271:10607-10.
- Paratcha, G., F. Ledda, L. Baars, M. Couplier, V. Basset, J. Anders, R. Scott, and C.F. Ibanez. 2001. Released GFRalpha1 potentiates downstream signaling, neuronal survival, and differentiation via a novel mechanism of recruitment of c-Ret to lipid rafts. *Neuron*. 29:171-84.
- Park, E., M. Na, J. Choi, S. Kim, J.R. Lee, J. Yoon, D. Park, M. Sheng, and E. Kim. 2003. The Shank family of postsynaptic density proteins interacts with and promotes synaptic accumulation of the beta PIX guanine nucleotide exchange factor for Rac1 and Cdc42. *J Biol Chem*.
- Pasini, B., M.G. Borrello, A. Greco, I. Bongarzone, Y. Luo, P. Mondellini, L. Alberti, C. Miranda, E. Arighi, R. Bocciardi, and et al. 1995. Loss of function effect of RET mutations causing Hirschsprung disease. *Nat Genet*. 10:35-40.

- Paulson, R.F., and A. Bernstein. 1995. Receptor tyrosine kinases and the regulation of hematopoiesis. *Semin Immunol.* 7:267-77.
- Pawson, T., and J.D. Scott. 1997. Signaling through scaffold, anchoring, and adaptor proteins. *Science.* 278:2075-80.
- Pelicci, G., F. Troglio, A. Bodini, R.M. Melillo, V. Pettirossi, L. Coda, A. De Giuseppe, M. Santoro, and P.G. Pelicci. 2002. The neuron-specific Rai (ShcC) adaptor protein inhibits apoptosis by coupling Ret to the phosphatidylinositol 3-kinase/Akt signaling pathway. *Mol Cell Biol.* 22:7351-63.
- Perantoni, A.O., L.F. Dove, and I. Karavanova. 1995. Basic fibroblast growth factor can mediate the early inductive events in renal development. *Proc Natl Acad Sci U S A.* 92:4696-700.
- Pollack, A.L., R.B. Runyan, and K.E. Mostov. 1998. Morphogenetic mechanisms of epithelial tubulogenesis: MDCK cell polarity is transiently rearranged without loss of cell-cell contact during scatter factor/hepatocyte growth factor-induced tubulogenesis. *Dev Biol.* 204:64-79.
- Ponting, C.P., C. Phillips, K.E. Davies, and D.J. Blake. 1997. PDZ domains: targeting signalling molecules to sub-membranous sites. *Bioessays.* 19:469-79.
- Porter, A.C., and R.R. Vaillancourt. 1998. Tyrosine kinase receptor-activated signal transduction pathways which lead to oncogenesis. *Oncogene.* 17:1343-52.
- Potempa, S., and A.J. Ridley. 1998. Activation of both MAP kinase and phosphatidylinositide 3-kinase by Ras is required for hepatocyte growth factor/scatter factor-induced adherens junction disassembly. *Mol Biol Cell.* 9:2185-200.
- Poteryaev, D., A. Titievsky, Y.F. Sun, J. Thomas-Crusells, M. Lindahl, M. Billaud, U. Arumae, and M. Saarma. 1999. GDNF triggers a novel ret-independent Src kinase family-coupled signaling via a GPI-linked GDNF receptor alpha1. *FEBS Lett.* 463:63-6.

- Qiao, J., R. Uzzo, T. Obara-Ishihara, L. Degenstein, E. Fuchs, and D. Herzlinger. 1999. FGF-7 modulates ureteric bud growth and nephron number in the developing kidney. *Development*. 126:547-54.
- Qualmann, B., T.M. Boeckers, M. Jeromin, E.D. Gundelfinger, and M.M. Kessels. 2004. Linkage of the actin cytoskeleton to the postsynaptic density via direct interactions of Abp1 with the ProSAP/Shank family. *J Neurosci*. 24:2481-95.
- Qui, M.S., and S.H. Green. 1992. PC12 cell neuronal differentiation is associated with prolonged p21ras activity and consequent prolonged ERK activity. *Neuron*. 9:705-17.
- Redecker, P., E.D. Gundelfinger, and T.M. Boeckers. 2001. The cortactin-binding postsynaptic density protein proSAP1 in non-neuronal cells. *J Histochem Cytochem*. 49:639-48.
- Robinson, M.J., and M.H. Cobb. 1997. Mitogen-activated protein kinase pathways. *Curr Opin Cell Biol*. 9:180-6.
- Rogers, S.A., G. Ryan, and M.R. Hammerman. 1992. Metanephric transforming growth factor-alpha is required for renal organogenesis in vitro. *Am J Physiol*. 262:F533-9.
- Romeo, G., P. Ronchetto, Y. Luo, V. Barone, M. Seri, I. Ceccherini, B. Pasini, R. Bocciardi, M. Lerone, H. Kaariainen, and et al. 1994. Point mutations affecting the tyrosine kinase domain of the RET proto-oncogene in Hirschsprung's disease. *Nature*. 367:377-8.
- Rongo, C., C.W. Whitfield, A. Rodal, S.K. Kim, and J.M. Kaplan. 1998. LIN-10 is a shared component of the polarized protein localization pathways in neurons and epithelia. *Cell*. 94:751-9.
- Rosario, M., and W. Birchmeier. 2003. How to make tubes: signaling by the Met receptor tyrosine kinase. *Trends Cell Biol*. 13:328-35.
- Sachs, M., H. Brohmann, D. Zechner, T. Muller, J. Hulsken, I. Walther, U. Schaeper, C. Birchmeier, and W. Birchmeier. 2000. Essential role of Gab1 for signaling by the c-Met receptor in vivo. *J Cell Biol*. 150:1375-84.

- Sachs, M., K.M. Weidner, V. Brinkmann, I. Walther, A. Obermeier, A. Ullrich, and W. Birchmeier. 1996. Motogenic and morphogenic activity of epithelial receptor tyrosine kinases. *J Cell Biol.* 133:1095-1107.
- Sakurai, H., and S.K. Nigam. 1997. Transforming growth factor-beta selectively inhibits branching morphogenesis but not tubulogenesis. *Am J Physiol.* 272:F139-46.
- Sala, C., V. Piech, N.R. Wilson, M. Passafaro, G. Liu, and M. Sheng. 2001. Regulation of dendritic spine morphology and synaptic function by Shank and Homer. *Neuron.* 31:115-30.
- Sambrook, E.F., Fritsch,E.F., and Maniatis,T. 1989. Molecular cloning: a laborarory handbook. *Cold Spring Harbor Laboratory Press, New York.*
- Sanchez, M.P., I. Silos-Santiago, J. Frisen, B. He, S.A. Lira, and M. Barbacid. 1996. Renal agenesis and the absence of enteric neurons in mice lacking GDNF. *Nature.* 382:70-3.
- Santoro, M., N.A. Dathan, M.T. Berlingieri, I. Bongarzone, C. Paulin, M. Grieco, M.A. Pierotti, G. Vecchio, and A. Fusco. 1994. Molecular characterization of RET/PTC3; a novel rearranged version of the RETproto-oncogene in a human thyroid papillary carcinoma. *Oncogene.* 9:509-16.
- Santoro, M., R. Rosati, M. Grieco, M.T. Berlingieri, G.L. D'Amato, V. de Franciscis, and A. Fusco. 1990. The ret proto-oncogene is consistently expressed in human pheochromocytomas and thyroid medullary carcinomas. *Oncogene.* 5:1595-8.
- Santos, O.F., E.J. Barros, X.M. Yang, K. Matsumoto, T. Nakamura, M. Park, and S.K. Nigam. 1994. Involvement of hepatocyte growth factor in kidney development. *Dev Biol.* 163:525-9.
- Santos, O.F., and S.K. Nigam. 1993. HGF-induced tubulogenesis and branching of epithelial cells is modulated by extracellular matrix and TGF-beta. *Dev Biol.* 160:293-302.
- Sariola, H. 2002. Nephron induction. *Nephrol Dial Transplant.* 17 Suppl 9:88-90.

- Sariola, H., and K. Sainio. 1997. The tip-top branching ureter. *Curr Opin Cell Biol.* 9:877-84.
- Saxen, L., and H. Sariola. 1987. Early organogenesis of the kidney. *Pediatr Nephrol.* 1:385-92.
- Schaeper, U., N.H. Gehring, K.P. Fuchs, M. Sachs, B. Kempkes, and W. Birchmeier. 2000. Coupling of Gab1 to c-Met, Grb2, and Shp2 mediates biological responses. *J Cell Biol.* 149:1419-32.
- Schlessinger, J. 2000. Cell signaling by receptor tyrosine kinases. *Cell.* 103:211-25.
- Schmidt, C., F. Bladt, S. Goedecke, V. Brinkmann, W. Zschiesche, M. Sharpe, E. Gherardi, and C. Birchmeier. 1995. Scatter factor/hepatocyte growth factor is essential for liver development. *Nature.* 373:699-702.
- Schober, A., U. Arumae, M. Saarma, and K. Unsicker. 2000. Expression of GFR alpha-1, GFR alpha-2, and c-Ret mRNAs in rat adrenal gland. *J Neurocytol.* 29:209-13.
- Schuchardt, A., V. D'Agati, L. Larsson-Blomberg, F. Costantini, and V. Pachnis. 1994. Defects in the kidney and enteric nervous system of mice lacking the tyrosine kinase receptor Ret. *Nature.* 367:380-3.
- Schuchardt, A., V. D'Agati, L. Larsson-Blomberg, F. Costantini, and V. Pachnis. 1995. RET-deficient mice: an animal model for Hirschsprung's disease and renal agenesis. *J Intern Med.* 238:327-32.
- Schuchardt, A., V. D'Agati, V. Pachnis, and F. Costantini. 1996. Renal agenesis and hypodysplasia in ret-k- mutant mice result from defects in ureteric bud development. *Development.* 122:1919-29.
- Seger, R., and E.G. Krebs. 1995. The MAPK signaling cascade. *Faseb J.* 9:726-35.
- Segouffin-Cariou, C., and M. Billaud. 2000. Transforming ability of MEN2A-RET requires activation of the phosphatidylinositol 3-kinase/AKT signaling pathway. *J Biol Chem.* 275:3568-76.

- Shelly, M., Y. Mosesson, A. Citri, S. Lavi, Y. Zwang, N. Melamed-Book, B. Aroeti, and Y. Yarden. 2003. Polar expression of ErbB-2/HER2 in epithelia. Bimodal regulation by Lin-7. *Dev Cell.* 5:475-86.
- Sheng, M., and E. Kim. 2000. The Shank family of scaffold proteins. *J Cell Sci.* 113 (Pt 11):1851-6.
- Shilo, B.Z. 1992. Roles of receptor tyrosine kinases in Drosophila development. *Faseb J.* 6:2915-22.
- Skolnik, E.Y., A. Batzer, N. Li, C.H. Lee, E. Lowenstein, M. Mohammadi, B. Margolis, and J. Schlessinger. 1993. The function of GRB2 in linking the insulin receptor to Ras signaling pathways. *Science.* 260:1953-5.
- Smith, D.P., C. Eng, and B.A. Ponder. 1994. Mutations of the RET proto-oncogene in the multiple endocrine neoplasia type 2 syndromes and Hirschsprung disease. *J Cell Sci Suppl.* 18:43-9.
- Soltau, M., K. Berhorster, S. Kindler, F. Buck, D. Richter, and H.J. Kreienkamp. 2004. Insulin receptor substrate of 53 kDa links postsynaptic shank to PSD-95. *J Neurochem.* 90:659-65.
- Soltau, M., D. Richter, and H.J. Kreienkamp. 2002. The Insulin Receptor Substrate IRSp53 Links Postsynaptic shank1 to the Small G-Protein cdc42. *Mol Cell Neurosci.* 21:575-83.
- Songyang, Z., A.S. Fanning, C. Fu, J. Xu, S.M. Marfatia, A.H. Chishti, A. Crompton, A.C. Chan, J.M. Anderson, and L.C. Cantley. 1997. Recognition of unique carboxyl-terminal motifs by distinct PDZ domains. *Science.* 275:73-7.
- Songyang, Z., S.E. Shoelson, J. McGlade, P. Olivier, T. Pawson, X.R. Bustelo, M. Barbacid, H. Sabe, H. Hanafusa, T. Yi, and et al. 1994. Specific motifs recognized by the SH2 domains of Csk, 3BP2, fps/fes, GRB-2, HCP, SHC, Syk, and Vav. *Mol Cell Biol.* 14:2777-85.
- Sparks, A.B., J.E. Rider, N.G. Hoffman, D.M. Fowlkes, L.A. Quillam, and B.K. Kay. 1996. Distinct ligand preferences of Src homology 3 domains from Src, Yes, Abl,

- Cortactin, p53bp2, PLCgamma, Crk, and Grb2. *Proc Natl Acad Sci U S A.* 93:1540-4.
- Srinivas, S., Z. Wu, C.M. Chen, V. D'Agati, and F. Costantini. 1999. Dominant effects of RET receptor misexpression and ligand-independent RET signaling on ureteric bud development. *Development.* 126:1375-86.
- Stork, P.J. 2003. Does Rap1 deserve a bad Rap? *Trends Biochem Sci.* 28:267-75.
- Stover, T., T.L. Gong, Y. Cho, R.A. Altschuler, and M.I. Lomax. 2000. Expression of the GDNF family members and their receptors in the mature rat cochlea. *Brain Res Mol Brain Res.* 76:25-35.
- Sundaram, M., and M. Han. 1996. Control and integration of cell signaling pathways during *C. elegans* vulval development. *Bioessays.* 18:473-80.
- Suzu, S., M. Tanaka-Douzono, K. Nomaguchi, M. Yamada, H. Hayasawa, F. Kimura, and K. Motoyoshi. 2000. p56(dok-2) as a cytokine-inducible inhibitor of cell proliferation and signal transduction. *Embo J.* 19:5114-22.
- Takahashi, M., J. Ritz, and G.M. Cooper. 1985. Activation of a novel human transforming gene, ret, by DNA rearrangement. *Cell.* 42:581-8.
- Takemura, T., S. Hino, M. Okada, Y. Murata, H. Yanagida, M. Ikeda, K. Yoshioka, and R.C. Harris. 2002. Role of membrane-bound heparin-binding epidermal growth factor-like growth factor (HB-EGF) in renal epithelial cell branching. *Kidney Int.* 61:1968-79.
- Tan, P.B., and S.K. Kim. 1999. Signaling specificity: the RTK/RAS/MAP kinase pathway in metazoans. *Trends Genet.* 15:145-9.
- The *C. elegans* Sequencing Consortium. 1998. Genome sequence of the nematode *C. elegans*: a platform for investigating biology. *Science.* 282:2012-8.
- Torres, R., B.L. Firestein, H. Dong, J. Staudinger, E.N. Olson, R.L. Huganir, D.S. Bredt, N.W. Gale, and G.D. Yancopoulos. 1998. PDZ proteins bind, cluster, and

- synaptically colocalize with Eph receptors and their ephrin ligands. *Neuron*. 21:1453-63.
- Traverse, S., N. Gomez, H. Paterson, C. Marshall, and P. Cohen. 1992. Sustained activation of the mitogen-activated protein (MAP) kinase cascade may be required for differentiation of PC12 cells. Comparison of the effects of nerve growth factor and epidermal growth factor. *Biochem J*. 288 (Pt 2):351-5.
- Trupp, M., E. Arenas, M. Fainzilber, A.S. Nilsson, B.A. Sieber, M. Grigoriou, C. Kilkenny, E. Salazar-Grueso, V. Pachnis, and U. Arumae. 1996. Functional receptor for GDNF encoded by the c-ret proto-oncogene. *Nature*. 381:785-9.
- Tu, J.C., B. Xiao, S. Naisbitt, J.P. Yuan, R.S. Petralia, P. Brakeman, A. Doan, V.K. Aakalu, A.A. Lanahan, M. Sheng, and P.F. Worley. 1999. Coupling of mGluR/Homer and PSD-95 complexes by the Shank family of postsynaptic density proteins. *Neuron*. 23:583-92.
- Utsumi, H., H. Chiba, Y. Kamimura, M. Osanai, Y. Igarashi, H. Tobioka, M. Mori, and N. Sawada. 2000. Expression of GFRalpha-1, receptor for GDNF, in rat brain capillary during postnatal development of the BBB. *Am J Physiol Cell Physiol*. 279:C361-8.
- Vaccaro, P., and L. Dente. 2002. PDZ domains: troubles in classification. *FEBS Lett*. 512:345-9.
- Vainio, S., and Y. Lin. 2002. Coordinating early kidney development: lessons from gene targeting. *Nat Rev Genet*. 3:533-43.
- van Heyningen, V. 1994. Genetics. One gene--four syndromes. *Nature*. 367:319-20.
- van Puijenbroek, A.A., D.H. van Weering, C.E. van den Brink, J.L. Bos, P.T. van der Saag, S.W. de Laat, and J. den Hertog. 1997. Cell scattering of SK-N-MC neuroepithelioma cells in response to Ret and FGF receptor tyrosine kinase activation is correlated with sustained ERK2 activation. *Oncogene*. 14:1147-57.
- van Weering, D.H., and J.L. Bos. 1998. Signal transduction by the receptor tyrosine kinase Ret. *Recent Results Cancer Res*. 154:271-81.

- Vanhaesebroeck, B., S.J. Leevers, K. Ahmadi, J. Timms, R. Katso, P.C. Driscoll, R. Woscholski, P.J. Parker, and M.D. Waterfield. 2001. Synthesis and function of 3-phosphorylated inositol lipids. *Annu Rev Biochem.* 70:535-602.
- Venter, J.C., M.D. Adams, E.W. Myers, P.W. Li, R.J. Mural, G.G. Sutton, H.O. Smith, M. Yandell, C.A. Evans, R.A. Holt, J.D. Gocayne, P. Amanatides, R.M. Ballew, D.H. Huson, J.R. Wortman, Q. Zhang, C.D. Kodira, X.H. Zheng, L. Chen, M. Skupski, G. Subramanian, P.D. Thomas, J. Zhang, G.L. Gabor Miklos, C. Nelson, S. Broder, A.G. Clark, J. Nadeau, V.A. McKusick, N. Zinder, A.J. Levine, R.J. Roberts, M. Simon, C. Slayman, M. Hunkapiller, R. Bolanos, A. Delcher, I. Dew, D. Fasulo, M. Flanigan, L. Florea, A. Halpern, S. Hannenhalli, S. Kravitz, S. Levy, C. Mobarry, K. Reinert, K. Remington, J. Abu-Threideh, E. Beasley, K. Biddick, V. Bonazzi, R. Brandon, M. Cargill, I. Chandramouliwaran, R. Charlaba, K. Chaturvedi, Z. Deng, V. Di Francesco, P. Dunn, K. Eilbeck, C. Evangelista, A.E. Gabrielian, W. Gan, W. Ge, F. Gong, Z. Gu, P. Guan, T.J. Heiman, M.E. Higgins, R.R. Ji, Z. Ke, K.A. Ketchum, Z. Lai, Y. Lei, Z. Li, J. Li, Y. Liang, X. Lin, F. Lu, G.V. Merkulov, N. Milshina, H.M. Moore, A.K. Naik, V.A. Narayan, B. Neelam, D. Nusskern, D.B. Rusch, S. Salzberg, W. Shao, B. Shue, J. Sun, Z. Wang, A. Wang, X. Wang, J. Wang, M. Wei, R. Wides, C. Xiao, C. Yan, et al. 2001. The sequence of the human genome. *Science.* 291:1304-51.
- Vincent, S., R. Wilson, C. Coelho, M. Affolter, and M. Leptin. 1998. The Drosophila protein Dof is specifically required for FGF signaling. *Mol Cell.* 2:515-25.
- Wehner, R., and W. Gehring. 1995. Zoologie. Thieme, Stuttgart - New York. 861 pp.
- Weidner, K.M., S. Di Cesare, M. Sachs, V. Brinkmann, J. Behrens, and W. Birchmeier. 1996. Interaction between Gab1 and the c-Met receptor tyrosine kinase is responsible for epithelial morphogenesis. *Nature.* 384:173-6.
- Weidner, K.M., M. Sachs, and W. Birchmeier. 1993. The Met receptor tyrosine kinase transduces motility, proliferation, and morphogenic signals of scatter factor/hepatocyte growth factor in epithelial cells. *J Cell Biol.* 121:145-54.

- West, G.B., J.H. Brown, and B.J. Enquist. 1999. The fourth dimension of life: fractal geometry and allometric scaling of organisms. *Science*. 284:1677-9.
- Whitfield, C.W., C. Benard, T. Barnes, S. Hekimi, and S.K. Kim. 1999. Basolateral localization of the *Caenorhabditis elegans* epidermal growth factor receptor in epithelial cells by the PDZ protein LIN-10. *Mol Biol Cell*. 10:2087-100.
- Widmann, C., S. Gibson, M.B. Jarpe, and G.L. Johnson. 1999. Mitogen-activated protein kinase: conservation of a three-kinase module from yeast to human. *Physiol Rev*. 79:143-80.
- Williams, D. 1994. Epidemiology. Chernobyl, eight years on. *Nature*. 371:556.
- Williams, R., J. Sanghera, F. Wu, D. Carbonaro-Hall, D.L. Campbell, D. Warburton, S. Pelech, and F. Hall. 1993. Identification of a human epidermal growth factor receptor-associated protein kinase as a new member of the mitogen-activated protein kinase/extracellular signal-regulated protein kinase family. *J Biol Chem*. 268:18213-7.
- Willott, E., M.S. Balda, A.S. Fanning, B. Jameson, C. Van Itallie, and J.M. Anderson. 1993. The tight junction protein ZO-1 is homologous to the *Drosophila* discs-large tumor suppressor protein of septate junctions. *Proc Natl Acad Sci U S A*. 90:7834-8.
- Wu, C., C.F. Lai, and W.C. Mobley. 2001. Nerve growth factor activates persistent Rap1 signaling in endosomes. *J Neurosci*. 21:5406-16.
- Yao, I., Y. Hata, K. Hirao, M. Deguchi, N. Ide, M. Takeuchi, and Y. Takai. 1999. Synamon, a novel neuronal protein interacting with synapse-associated protein 90/postsynaptic density-95-associated protein. *J Biol Chem*. 274:27463-6.
- Yenush, L., R. Fernandez, M.G. Myers, Jr., T.C. Grammer, X.J. Sun, J. Blenis, J.H. Pierce, J. Schlessinger, and M.F. White. 1996. The *Drosophila* insulin receptor activates multiple signaling pathways but requires insulin receptor substrate proteins for DNA synthesis. *Mol Cell Biol*. 16:2509-17.

- Yu, T., S. Scully, Y. Yu, G.M. Fox, S. Jing, and R. Zhou. 1998. Expression of GDNF family receptor components during development: implications in the mechanisms of interaction. *J Neurosci.* 18:4684-96.
- Zaccaro, M.C., L. Ivanisevic, P. Perez, S.O. Meakin, and H.U. Saragovi. 2001. p75 Co-receptors regulate ligand-dependent and ligand-independent Trk receptor activation, in part by altering Trk docking subdomains. *J Biol Chem.* 276:31023-9.
- Zhao, R., and Z.J. Zhao. 2000. Dissecting the interaction of SHP-2 with PZR, an immunoglobulin family protein containing immunoreceptor tyrosine-based inhibitory motifs. *J Biol Chem.* 275:5453-9.
- Zitzer, H., H.H. Honck, D. Bachner, D. Richter, and H.J. Kreienkamp. 1999. Somatostatin receptor interacting protein defines a novel family of multidomain proteins present in human and rodent brain. *J Biol Chem.* 274:32997-3001.