

VII. Anhang

C. Literatur

- Adesnik M, Salditt M, Thomas W, Darnell JE. (1972) Evidence that all messenger RNA molecules (except histone messenger RNA) contain Poly (A) sequences and that the Poly(A) has a nuclear function. *J Mol Biol.* 1972 Oct 28;71(1):21-30. PMID: 4539023
- Advani R, Peethambaram P, Lum BL, Fisher GA, Hartmann L, Long HJ, Halsey J, Holmlund JT, Dorr A, Sikic BI. A Phase II trial of aprinocarsen, an antisense oligonucleotide inhibitor of protein kinase C alpha, administered as a 21-day infusion to patients with advanced ovarian carcinoma. *Cancer.* 2004 Jan 15;100(2):321-6.
- Agrawal S, Temsamani J, Tang JY. Pharmacokinetics, biodistribution, and stability of oligodeoxynucleotide phosphorothioates in mice. *Proc Natl Acad Sci U S A.* 1991;88:7595–7599.[
- Ahlquist, 1948: A study of the adrenotropic receptors. A study of the adrenotropic receptors. *Am. J. Physiol.* 153:586-600.
- Akhtar S, Rossi JJ. Anti-HIV therapy with antisense oligonucleotides and ribozymes: realistic approaches or expensive myths? *J Antimicrob Chemother.* 1996;38:159–165.
- Altuvia S, Wagner EG. Switching on and off with RNA. *Proc Natl Acad Sci U S A.* 2000 Aug 29;97(18):9824-6.
- Ambros V. Development. Dicing up RNAs. *Science* 2001 Aug 3;293(5531):811-3
- Askew DS, Xu F. New insights into the function of noncoding RNA and its potential role in disease pathogenesis. *Histol Histopathol.* 1999 Jan;14(1):235-41.
- Bartsch H, Voigtsberger S, Baumann G, Morano I, Luther HP. Detection of a novel sense-antisense RNA-hybrid structure by RACE experiments on endogenous troponin I antisense RNA. *RNA.* 2004 Aug;10(8):1215-24.
- Bass BL, Weintraub H. A developmentally regulated activity that unwinds RNA duplexes. *Cell.* 1987 Feb 27;48(4):607-13.
- Bass BL. Double-stranded RNA as a template for gene silencing. *Cell.* 2000 Apr 28;101(3):235-8.
- Baumann G, Riess G, Erhardt WD, Felix SB, Ludwig L, Blumel G, Blomer H. Impaired beta-adrenergic stimulation in the uninvolved ventricle post-acute myocardial infarction: reversible defect due to excessive circulating catecholamine-induced decline in number and affinity of beta-receptors. *Am Heart J.* 1981 May;101(5):569-81.
- Beau SL, Tolley TK, Saffitz JE. Heterogeneous transmural distribution of beta-adrenergic receptor subtypes in failing human hearts. *Circulation.* 1993 Dec;88(6):2501-9. Related Articles, Links
- Bechler K. Influence of capping and polyadenylation on mRNA expression and on antisense RNA mediated inhibition of gene expression. *Biochem Biophys Res Commun* 1997 Dec 8;241(1):193-9
- Bernstein E, Caudy AA, Hammond SM, Hannon GJ. Role for a bidentate ribonuclease in the initiation step of RNA interference. *Nature* 2001-a Jan 18;409(6818):363-6

- Bernstein E, Denli AM, Hannon GJ. The rest is silence. *RNA* 2001-b Nov;7(11):1509-21
- Billy E, Brondani V, Zhang H, Muller U, Filipowicz W. Specific interference with gene expression induced by long, double-stranded RNA in mouse embryonal teratocarcinoma cell lines. *Proc Natl Acad Sci U S A*. 2001 Dec 4;98(25):14428-33. Epub 2001 Nov 27.
- Blaszczyk J, Tropea JE, Bubunencko M, Routzahn KM, Waugh DS, Court DL, Ji X. Crystallographic and modeling studies of RNase III suggest a mechanism for double-stranded RNA cleavage. *Structure (Camb)*. 2001 Dec;9(12):1225-36.
- Bohm M, Lohse MJ. Quantification of beta-adrenoceptors and beta-adrenoceptor kinase on protein and mRNA levels in heart failure. *Eur Heart J*. 1994 Dec;15 Suppl D:30-4.
- Book WM, Hott BJ. Beta-adrenergic Receptor Blockers in Heart Failure. *Curr Treat Options Cardiovasc Med*. 2003 Dec;5(6):475-485.
- Bosher, J.M. and Labouesse, M. 2000. RNA interference: Genetic wand and genetic watchdog. *Nat. Cell Biol*. 2: E31–E36
- Bristow MR, Ginsburg R, Minobe W, Cubicciotti RS, Sageman WS, Lurie K, Billingham ME, Harrison DC, Stinson EB. Decreased catecholamine sensitivity and beta-adrenergic-receptor density in failing human hearts. *N Engl J Med*. 1982 Jul 22;307(4):205-11.
- Bristow MR, Minobe WA, Reynolds MV, Port JD, Rasmussen R, Ray PE, Feldman AM. Reduced beta 1 receptor messenger RNA abundance in the failing human heart. *J Clin Invest*. 1993 Dec;92(6):2737-45.
- Brodde OE, Michel MC. Adrenergic and muscarinic receptors in the human heart. *Pharmacol Rev*. 1999 Dec;51(4):651-90.
- Brodde OE. Adrenoceptors and their signal transduction mechanisms. *J Auton Pharmacol*. 1994 Feb;14(1):3-4.
- Brodde OE. Beta 1- and beta 2-adrenoceptors in the human heart: properties, function, and alterations in chronic heart failure. *Pharmacol Rev*. 1991 Jun;43(2):203-42.
- Brodde OE. Beta-adrenoceptors in cardiac disease. *Pharmacol Ther*. 1993 Dec;60(3):405-30.
- Brown L, Deighton NM, Bals S, Sohlmann W, Zerkowski HR, Michel MC, Brodde OE. Spare receptors for beta-adrenoceptor-mediated positive inotropic effects of catecholamines in the human heart. *J Cardiovasc Pharmacol*. 1992;19:222–232.
- Brownie J, Shawcross S, Theaker J, Whitcombe D, Ferrie R, Newton C, Little S. The elimination of primer-dimer accumulation in PCR. *Nucleic Acids Res*. 1997 Aug 15;25(16):3235-41.
- Bustin SA. Absolute quantification of mRNA using real-time reverse transcription polymerase chain reaction assays. *J Mol Endocrinol*. 2000 Oct;25(2):169-93.
- Bylund DB, Eikenberg DC, Hieble JP, Langer SZ, Lefkowitz RJ, Minneman KP, Molinoff PB, Ruffolo RR Jr, Trendelenburg U. International Union of Pharmacology nomenclature of adrenoceptors. *Pharmacol Rev*. 1994 Jun;46(2):121-36.
- Caplen NJ, Parrish S, Imani F, Fire A, Morgan RA. Specific inhibition of gene expression by small double-stranded RNAs in invertebrate and vertebrate systems. *Proc Natl Acad Sci U S A*. 2001 Aug 14;98(17):9742-7. Epub 2001 Jul 31.
- Caplen, N.J., Fleenor, J., Fire, A., and Morgan, R.A. 2000. dsRNA-mediated gene silencing in cultured *Drosophila* cells: A tissue culture model for the analysis of RNA interference. *Gene* 252: 95–105.

- Castro C, Arnold JJ, Cameron CE. Incorporation fidelity of the viral RNA-dependent RNA polymerase: a kinetic, thermodynamic and structural perspective. *Virus Res.* 2005 Feb;107(2):141-9.
- Chi KN, Gleave ME, Klasa R, Murray N, Bryce C, Lopes de Menezes DE, D'Aloisio S, Tolcher AW. A phase I dose-finding study of combined treatment with an antisense Bcl-2 oligonucleotide (Genasense) and mitoxantrone in patients with metastatic hormone-refractory prostate cancer. *Clin Cancer Res.* 2001 Dec;7(12):3920-7.
- Chien A, Edgar DB, Trela JM. Deoxyribonucleic acid polymerase from the extreme thermophile *Thermus aquaticus*. *J Bacteriol.* 1976 Sep;127(3):1550-7.
- Chomczynski P, Sacchi N. Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. *Anal Biochem.* 1987 Apr;162(1):156-9.
- Collins S, Ostrowski J, Lefkowitz RJ. Cloning and sequence analysis of the human beta 1-adrenergic receptor 5'-flanking promoter region. *Biochim Biophys Acta.* 1993 Feb 20;1172(1-2):171-4.
- Cotter FE. Antisense therapy of hematologic malignancies. *Semin Hematol.* 1999 Oct;36(4 Suppl 6):9-14. Review.
- Crooke ST, Bennett CF. Progress in antisense oligonucleotide therapeutics. *Annu Rev Pharmacol Toxicol.* 1996;36:107-29.
- Crooke ST. An overview of progress in antisense therapeutics. *Antisense Nucleic Acid Drug Dev.* 1998 Apr;8(2):115-22.
- Crooke ST. Therapeutic applications of oligonucleotides. *Biotechnology (NY).* 1992 Aug;10(8):882-6.
- Dachs GU, Dougherty GJ, Stratford IJ, Chaplin DJ. Targeting gene therapy to cancer: a *Oncol Res.* 1997;9:313-325
- de la Taille A, Olsson CA, Katz AE. Molecular staging of prostate cancer: dream or reality? *Oncology (Huntingt).* 1999 Feb;13(2):187-94; discussion 194-8, 204-5 pas.
- DiPaola NR, Sweet WE, Stull LB, Francis GS, Schomisch Moravec C. Beta-adrenergic receptors and calcium cycling proteins in non-failing, hypertrophied and failing human hearts: transition from hypertrophy to failure. *J Mol Cell Cardiol.* 2001 Jun;33(6):1283-95.
- Dohlman HG, Bouvier M, Benovic JL, Caron MG, Lefkowitz RJ. The multiple membrane spanning topography of the beta 2-adrenergic receptor. Localization of the sites of binding, glycosylation, and regulatory phosphorylation by limited proteolysis. *J Biol Chem.* 1987 Oct 15;262(29):14282-8.
- Dreyfus M, Regnier P. The poly(A) tail of mRNAs: bodyguard in eukaryotes, scavenger in bacteria. *Cell* 2002 Nov 27;111(5):611-3
- Edmonds M. A history of poly A sequences: from formation to factors to function. *Prog Nucleic Acid Res Mol Biol.* 2002;71:285-389.
- Elbashir SM, Harborth J, Lendeckel W, Yalcin A, Weber K, Tuschl T. Duplexes of 21-nucleotide RNAs mediate RNA interference in cultured mammalian cells. *Nature.* 2001 May 24;411(6836):494-8.
- Elbashir SM, Lendeckel W, Tuschl T. RNA interference is mediated by 21- and 22-nucleotide RNAs. *Genes Dev.* 2001 Jan 15;15(2):188-200.
- Elbashir, S. M., Martinez, J., Patkaniowska, A., Lendeckel, W. & Tuschl, T. Functional anatomy

- of siRNAs for mediating efficient RNAi in *Drosophila melanogaster* embryo lysate. *EMBO J.* 20, 6877–6888 (2001-a).
- Emorine LJ, Fève B, Pairault J, Briend-Sutren MM, Marullo S, Delavier-Klutchko C, Strosberg DA. Structural basis for functional diversity of beta 1-, beta 2- and beta 3-adrenergic receptors. *Biochem Pharmacol.* 1991 Mar 15-Apr 1;41(6-7):853-9.
- Engelhardt S, Böhm M, Erdmann E, Lohse MJ. Analysis of beta-adrenergic receptor mRNA levels in human ventricular biopsy specimens by quantitative polymerase chain reactions: progressive reduction of beta 1-adrenergic receptor mRNA in heart failure. *J Am Coll Cardiol.* 1996 Jan;27(1):146-54.
- Erdmann VA, Barciszewska MZ, Szymanski M, Hochberg A, de Groot N, Barciszewski J. The non-coding RNAs as riboregulators. *Nucleic Acids Res.* 2001 Jan 1;29(1):189-93.
- Fedoroff N, Wellauer PK, Wall R. Intermolecular duplexes in heterogeneous nuclear RNA from HeLa cells. *Cell.* 1977 Apr;10(4):597-610.
- Feldman AM. Modulation of adrenergic receptors and G-transduction proteins in failing human ventricular myocardium. *Circulation.* 1993 May;87(5 Suppl):IV27-34.
- Fire A, Xu S, Montgomery MK, Kostas SA, Driver SE, Mello CC. Potent and specific genetic interference by double-stranded RNA in *Nature* 1998 Feb 19;391(6669):806-11
- Fire A. RNA-triggered gene silencing. *Trends Genet.* 1999 Sep;15(9):358-63.
- Frederich RC Jr, Waldo GL, Harden TK, Perkins JP. Characterization of agonist-induced beta-adrenergic receptor-specific desensitization in C62B glioma cells. *J Cyclic Nucleotide Protein Phosphor Res.* 1983;9(2):103-18.
- Freeman WM, Walker SJ, Vrana KE. Quantitative RT-PCR: pitfalls and potential. *Biotechniques.* 1999 Jan;26(1):112-22, 124-5.
- Frielle T, Collins S, Daniel KW, Caron MG, Lefkowitz RJ, Kobilka BK. Cloning of the cDNA for the human beta 1-adrenergic receptor. *Proc Natl Acad Sci U S A.* 1987 Nov;84(22):7920-4.
- Gala JL, Heusterspreute M, Loric S, Hanon F, Tombal B, Van Cangh P, De Nayer P, Philippe M. Expression of prostate-specific antigen and prostate-specific membrane antigen transcripts in blood cells: implications for the detection of hematogenous prostate cells and standardization. *Clin Chem.* 1998 Mar;44(3):472-81.
- Gerard CJ, Olsson K, Ramanathan R, Reading C, Hanania EG. Improved quantitation of minimal residual disease in multiple myeloma using real-time polymerase chain reaction and plasmid-DNA complementarity determining region III standards. *Cancer Res.* 1998 Sep 1;58(17):3957-64.
- Grishok A, Pasquinelli AE, Conte D, Li N, Parrish S, Ha I, Baillie DL, Fire A, Ruvkun G, Mello CC. Genes and mechanisms related to RNA interference regulate expression of the small temporal RNAs that control *C. elegans* developmental timing. *Cell.* 2001 Jul 13;106(1):23-34.
- Guest SJ, Hadcock JR, Watkins DC, Malbon CC. Beta 1- and beta 2-adrenergic receptor expression in differentiating 3T3-L1 cells. Independent regulation at the level of mRNA. *J Biol Chem.* 1990 Apr 5;265(10):5370-5.
- Gyurko R, Tran D, Phillips MI. Time course of inhibition of hypertension by antisense oligonucleotides targeted to AT1 angiotensin receptor mRNA in spontaneously hypertensive rats. *Am J Hypertens.* 1997;10:56S–62S.

- Ha I, Wightman B, Ruvkun G. A bulged lin-4/lin-14 RNA duplex is sufficient for *Caenorhabditis elegans* lin-14 temporal gradient formation. *Genes Dev.* 1996 Dec 1;10(23):3041-50.
- Hadcock JR, Malbon CC. Down-regulation of beta-adrenergic receptors: agonist-induced reduction in receptor mRNA levels. *Proc Natl Acad Sci U S A.* 1988 Jul;85(14):5021-5.
- Hadcock JR, Malbon CC. Regulation of receptor expression by agonists: transcriptional and post-transcriptional controls. *Trends Neurosci.* 1991 Jun;14(6):242-7.
- Hadcock JR, Wang HY, Malbon CC. Agonist-induced destabilization of beta-adrenergic receptor mRNA. Attenuation of glucocorticoid-induced up-regulation of beta-adrenergic receptors. *J Biol Chem.* 1989 Nov 25;264(33):19928-33.
- Hammond SM, Bernstein E, Beach D, Hannon GJ. An RNA-directed nuclease mediates post-transcriptional gene silencing in *Drosophila* cells. *Nature.* 2000 Mar 16;404(6775):293-6.
- Hammond SM, Caudy AA, Hannon GJ. Post-transcriptional gene silencing by double-stranded RNA. *Nat Rev Genet.* 2001 Feb;2(2):110-9. Review.
- Hannon GJ. RNA interference. *Nature.* 2002 Jul 11;418(6894):244-51. Review.
- Harden TK. Agonist-induced desensitization of the beta-adrenergic receptor-linked adenylate cyclase. *Pharmacol Rev.* 1983 Mar;35(1):5-32.
- Hata JA, Williams ML, Koch WJ. Genetic manipulation of myocardial beta-adrenergic receptor activation and desensitization. *J Mol Cell Cardiol.* 2004 Jul;37(1):11-21.
- Hausdorff WP, Caron MG, Lefkowitz RJ. Turning off the signal: desensitization of beta-adrenergic receptor function. *FASEB J.* 1990 Aug;4(11):2881-9. Review. Erratum in: *FASEB J* 1990 Sep;4(12):3049.
- Heid CA, Stevens J, Livak KJ, Williams PM. Real time quantitative PCR. *Genome Res.* 1996 Oct;6(10):986-94.
- Hein L, Kobilka BK. Adrenergic receptor signal transduction and regulation. *Neuropharmacology.* 1995 Apr;34(4):357-66. Review.
- Herbert A, Rich A. RNA processing in evolution. The logic of soft-wired genomes. *Ann N Y Acad Sci.* 1999 May 18;870:119-32.
- Herbert A. The four Rs of RNA-directed evolution. *Nat Genet.* 2004 Jan;36(1):19-25.
- Higuchi R, Dollinger G, Walsh PS, Griffith R. Simultaneous amplification and detection of specific DNA sequences. *Biotechnology (N Y).* 1992 Apr;10(4):413-7.
- Higuchi R, Fockler C, Dollinger G, Watson R. Kinetic PCR analysis: real-time monitoring of DNA amplification reactions. *Biotechnology (N Y).* 1993 Sep;11(9):1026-30.
- Hilleren P, McCarthy T, Rosbash M, Parker R, Jensen TH. Quality control of mRNA 3'-end processing is linked to the nuclear exosome. *Nature* 2001 Oct 4;413(6855):538-42
- Hod Y. A simplified ribonuclease protection assay. *Biotechniques.* 1992 Dec;13(6):852-4.
- Homburger V, Lucas M, Cantau B, Barabe J, Penit J, Bockaert J. Further evidence that desensitization of beta-adrenergic-sensitive adenylate cyclase proceeds in two steps. Modification of the coupling and loss of beta-adrenergic receptors. *J Biol Chem.* 1980 Nov 10;255(21):10436-44.
- Hutvagner G, Zamore PD. RNAi: nature abhors a double-strand. *Curr Opin Genet Dev* 2002

- Apr;12(2):225-32
- Ihl-Vahl R, Eschenhagen T, Kubler W, Marquetant R, Nose M, Schmitz W, Scholz H, Strasser RH. Differential regulation of mRNA specific for beta 1- and beta 2-adrenergic receptors in human failing hearts. Evaluation of the absolute cardiac mRNA levels by two independent methods. *J Mol Cell Cardiol.* 1996 Jan;28(1):1-10.
- Ishikawa Y, Homcy CJ. The adenylyl cyclases as integrators of transmembrane signal transduction. *Circ Res.* 1997 Mar;80(3):297-304.
- Iversen P. In vivo studies with phosphorothioate oligonucleotides: pharmacokinetics prologue. *Anticancer Drug Des.* 1991 Dec;6(6):531-8.
- Johnson GL, Wolfe BB, Harden TK, Molinoff PB, Perkins JP. Role of beta-adrenergic receptors in catecholamine-induced desensitization of adenylyl cyclase in human astrocytoma cells. *J Biol Chem.* 1978 Mar 10;253(5):1472-80.
- Kallal L, Gagnon AW, Penn RB, Benovic JL. Visualization of agonist-induced sequestration and down-regulation of a green fluorescent protein-tagged beta2-adrenergic receptor. *J Biol Chem.* 1998 Jan 2;273(1):322-8.
- Kaumann AJ, Molenaar P. Modulation of human cardiac function through 4 beta-adrenoceptor populations. *Naunyn Schmiedebergs Arch Pharmacol.* 1997 Jun;355(6):667-81.
- Keilholz U, Willhauck M, Rimoldi D, Brasseur F, Dummer W, Rass K, de Vries T, Blaheta J, Voit C, Lethe B, Burchill S. Reliability of reverse transcription-polymerase chain reaction (RT-PCR)-based assays for the detection of circulating tumour cells: a quality-assurance initiative of the EORTC Melanoma Cooperative Group. *Eur J Cancer.* 1998 Apr;34(5):750-3.
- Kennerdell JR, Carthew RW. Use of dsRNA-mediated genetic interference to demonstrate that frizzled and frizzled 2 act in the wingless pathway. *Cell.* 1998 Dec 23;95(7):1017-26.
- Kerr IM, Brown RE. pppA2'p5'A2'p5'A: an inhibitor of protein synthesis synthesized with an enzyme fraction from interferon-treated cells. *Proc Natl Acad Sci U S A.* 1978 Jan;75(1):256-60.
- Ketting RF, Fischer SE, Bernstein E, Sijen T, Hannon GJ, Plasterk RH. Dicer functions in RNA interference and in synthesis of small RNA involved in developmental timing in *C. elegans*. *Genes Dev.* 2001 Oct 15;15(20):2654-9.
- Ketting RF, Plasterk RH. A genetic link between co-suppression and RNA interference in *C. elegans*. *Nature.* 2000 Mar 16;404(6775):296-8.
- Kimelman D, Kirschner MW. An antisense mRNA directs the covalent modification of the transcript encoding fibroblast growth factor in *Xenopus* oocytes. *Cell.* 1989 Nov 17;59(4):687-96.
- Knee R, Murphy PR. Regulation of gene expression by natural antisense RNA transcripts. *Neurochem Int.* 1997 Sep;31(3):379-92.
- Korneev SA, Park JH, O'Shea M. Neuronal expression of neural nitric oxide synthase (nNOS) protein is suppressed by an antisense RNA transcribed from an NOS pseudogene. *J Neurosci.* 1999 Sep 15;19(18):7711-20.
- Krystal GW, Armstrong BC, Battey JF. N-myc mRNA forms an RNA-RNA duplex with endogenous antisense transcripts. *Mol Cell Biol.* 1990 Aug;10(8):4180-91.
- Kukkonen SK, Vaheri A, Plyusnin A. L protein, the RNA-dependent RNA polymerase of han-

- taviruses. *Arch Virol.* 2004 Oct 20;
- Kumar M, Carmichael GG. Antisense RNA: function and fate of duplex RNA in cells of higher eukaryotes. *Microbiol Mol Biol Rev.* 1998 Dec;62(4):1415-34.
- Lagos-Quintana, M., Rauhut, R., Lendeckel, W. & Tuschl, T. Identification of novel genes coding for small expressed RNAs. *Science* 294, 853–858 (2001).
- Lai, E. C. Micro RNAs are complementary to 3' UTR sequence motifs that mediate negative posttranscriptional regulation. *Nature Genet.* 30, 363–364 (2002).
- Lau, N. C., Lim, L. P., Weinstein, E. G. & Bartel, D. P. An abundant class of tiny RNAs with probable regulatory roles in *Caenorhabditis elegans*. *Science* 294, 858–862 (2001).
- Laurila MR, Salgado PS, Makeyev EV, Nettelship J, Stuart DI, Grimes JM, Bamford DH. Gene silencing pathway RNA-dependent RNA polymerase of *Neurospora crassa*: yeast expression and crystallization of selenomethionated QDE-1 protein. *J Struct Biol.* 2005 Jan;149(1):111-5.
- Lavorgna G, Dahary D, Lehner B, Sorek R, Sanderson CM, Casari G. In search of antisense. *Trends Biochem Sci.* 2004 Feb;29(2):88-94.
- Lazar-Wesley E, Hadcock JR, Malbon CC, Kunos G, Ishac EJ. Tissue-specific regulation of alpha 1B, beta 1, and beta 2-adrenergic receptor mRNAs by thyroid state in the rat. *Endocrinology.* 1991 Aug;129(2):1116-8.
- Lefkowitz RJ, Caron MG. Adrenergic receptors: molecular mechanisms of clinically relevant regulation. *Clin Res.* 1985 Sep;33(3):395-406.
- Lefkowitz RJ, Caron MG. Adrenergic receptors. *Harvey Lect.* 1990;86:33-45.
- Lefkowitz RJ. G protein-coupled receptors. III. New roles for receptor kinases and beta-arrestins in receptor signaling and desensitization. *J Biol Chem.* 1998 Jul 24;273(30):18677-80.
- Lemoine H, Kaumann AJ. Regional differences of beta 1- and beta 2-adrenoceptor-mediated functions in feline heart. A beta 2-adrenoceptor-mediated positive inotropic effect possibly unrelated to cyclic AMP. *Naunyn Schmiedeberg's Arch Pharmacol.* 1991 Jul;344(1):56-69.
- Lewin B. *Molekularbiologie der Gene.* Spektrum akademischer Verlag Heidelberg – Berlin. 1998
- Lewin, Benjamin: *Molekularbiologie der Gene, Studienausgabe* Spektrum Akademischer Verlag (August 2002)
- Li AW, Murphy PR. Expression of alternatively spliced FGF-2 antisense RNA transcripts in the central nervous system: regulation of FGF-2 mRNA translation. *Mol Cell Endocrinol.* 2000 Dec 22;170(1-2):233-42.
- Linz U, Degenhardt H. Polymerase chain reaction: an overview. *Naturwissenschaften.* 1990 Nov;77(11):515-30.
- Lohse MJ, Benovic JL, Caron MG, Lefkowitz RJ. Multiple pathways of rapid beta 2-adrenergic receptor desensitization. Delineation with specific inhibitors. *J Biol Chem.* 1990 Feb 25;265(6):3202-11.
- Lohse MJ. Molecular mechanisms of membrane receptor desensitization. *Biochim Biophys Acta.* 1993 Nov 7;1179(2):171-88.
- Lowry OH, Rosebrough, Farr AL, Randall RJ. Protein measurement with the Folin phenol re-

- agent. *J Biol Chem*. 1951 Nov;193(1):265-75.
- Luther HP, Haase H, Hohaus A, Beckmann G, Reich J, Morano I. Characterization of naturally occurring myosin heavy chain antisense mRNA in rat heart. *J Cell Biochem*. 1998 Jul 1;70(1):110-20.
- Luther HP, Morwinski R, Wallukat G, Haase H, Morano I. Expression of sense and naturally occurring antisense mRNA of myosin heavy chain in rat heart tissue and cultivated cardiomyocytes. *J Mol Cell Cardiol*. 1997 Jan;29(1):27-35.
- Luther HP, Podlowski S, Hetzer R, Baumann G. Analysis of sense and naturally occurring antisense transcripts of myosin heavy chain in the human myocardium. *J Cell Biochem*. 2001;80(4):596-605.
- Lyons J. The polymerase chain reaction and cancer diagnostics. *Cancer*. 1992 Mar 15;69(6 Suppl):1527-31.
- Machida CA, Bunzow JR, Searles RP, Van Tol H, Tester B, Neve KA, Teal P, Nipper V, Civelli O. Molecular cloning and expression of the rat beta 1-adrenergic receptor gene. *J Biol Chem*. 1990 Aug 5;265(22):12960-5.
- Maniatis T, Fritsch EF, Sambrook J "Molecular Cloning - A laboratory manual. " New York Cold Spring Harbor Laboratory B. (1989)
- Marcucci G, Byrd JC, Dai G, Klisovic MI, Kourlas PJ, Young DC, Cataland SR, Fisher DB, Lucas D, Chan KK, Porcu P, Lin ZP, Farag SF, Frankel SR, Zwiebel JA, Kraut EH, Balcerzak SP, Bloomfield CD, Grever MR, Caligiuri MA. Phase 1 and pharmacodynamic studies of G3139, a Bcl-2 antisense oligonucleotide, in combination with chemotherapy in refractory or relapsed acute leukemia. *Blood*. 2003 Jan 15;101(2):425-32. Epub 2002 Aug 22.
- Marshall J, Chen H, Yang D, Figueira M, Bouker KB, Ling Y, Lippman M, Frankel SR, Hayes DF. A phase I trial of a Bcl-2 antisense (G3139) and weekly docetaxel in patients with advanced breast cancer and other solid tumors. *Ann Oncol*. 2004 Aug;15(8):1274-83.
- Mattick JS, Gagen MJ. The evolution of controlled multitasked gene networks: the role of introns and other noncoding RNAs in the development of complex organisms. *Mol Biol Evol*. 2001 Sep;18(9):1611-30.
- Matzke MA, Matzke AJ, Pruss GJ, Vance VB. RNA-based silencing strategies in plants. *Curr Opin Genet Dev*. 2001 Apr;11(2):221-7.
- McDevitt DG. In vivo studies on the function of cardiac beta-adrenoceptors in man. *Eur Heart J*. 1989 Jun;10 Suppl B:22-8.
- Melton DA, Krieg PA, Rebagliati MR, Maniatis T, Zinn K, Green MR. Efficient in vitro synthesis of biologically active RNA and RNA hybridization probes from plasmids containing a bacteriophage SP6 promoter. *Nucleic Acids Res*. 1984 Sep 25;12(18):7035-56.
- Michaelis M, Green M, Rieger R. *Glossary of Genetics: Classical and Molecular*. (1991) 5th ed Springer-Verlag
- Mills SE. Implication of feedback regulation of beta-1 adrenergic signalling. (2002) *J. Anim. Sci*. 80(E. Suppl. 1):E30–E35
- Mitchusson KD, Blaxall BC, Pende A, Port JD. Agonist-mediated destabilization of human beta1-adrenergic receptor mRNA: role of the 3' untranslated translated region. *Biochem Biophys Res Commun*. 1998 Nov 18;252(2):357-62.

- Mittmann C, Munstermann U, Weil J, Bohm M, Herzig S, Nienaber C, Eschenhagen T. Analysis of gene expression patterns in small amounts of human ventricular myocardium by a multiplex RNase protection assay. *Mol Med*. 1998 Feb;76(2):133-40.
- Moradpour D, Brass V, Bieck E, Friebe P, Gosert R, Blum HE, Bartenschlager R, Penin F, Lohmann V. Membrane association of the RNA-dependent RNA polymerase is essential for hepatitis C virus RNA replication. *J Virol*. 2004 Dec;78(23):13278-84.
- Morris MJ, Tong WP, Cordon-Cardo C, Drobnjak M, Kelly WK, Slovin SF, Terry KL, Siedlecki K, Swanson P, Rafi M, DiPaola RS, Rosen N, Scher HI. Phase I trial of BCL-2 antisense oligonucleotide (G3139) administered by continuous intravenous infusion in patients with advanced cancer. *Clin Cancer Res*. 2002 Mar;8(3):679-83.
- Morrison TB, Weis JJ, Wittwer CT. Quantification of low-copy transcripts by continuous SYBR Green I monitoring during amplification. *Biotechniques*. 1998 Jun;24(6):954-8, 960, 962.
- Mülhardt C. *Der Experimentator: Molekularbiologie* Stuttgart, Jena, Lübeck, Ulm. (1999) Gustav Fischer Verlag, Berlin 1. Aufl.
- Murphree SS, Saffitz JE. Distribution of beta-adrenergic receptors in failing human myocardium. Implications for mechanisms of down-regulation. *Circulation*. 1989 Jun;79(6):1214-25.
- Murphy PR, Knee RS. Identification and characterization of an antisense RNA transcript (gfg) from the human basic fibroblast growth factor gene. *Mol Endocrinol*. 1994 Jul;8(7):852-9.
- Newton CR, Graham A. *PCR*. Spektrum Akademischer Verlag (Januar 1997)
- Ngo, H., Tschudi, C., Gull, K., and Ullu, E. 1998. Double stranded RNA induces mRNA degradation in *Trypanosoma brucei*. *Proc. Natl. Acad. Sci*. 95: 14687–14692.
- Nykanen, A., Haley, B. & Zamore, P. D. ATP requirements and small interfering RNA structure in the RNA interference pathway. *Cell* 107, 309–321 (2001).
- Oelgeschlager, M., Larrain, J. Geissert, D., De Robertis, E.M. 2000. The evolutionarily conserved BMP-binding protein Twisted gastrulation promotes BMP signalling. *Nature* 405: 757–763.
- Ogino T, Kobayashi M, Iwama M, Mizumoto K. Sendai virus RNA-dependent RNA polymerase L protein catalyzes cap methylation of virus-specific mRNA. *J Biol Chem*. 2004 Nov 30;
- Oliveira JG, Prados RZ, Guedes AC, Ferreira PC, Kroon EG. The housekeeping gene glyceraldehyde-3-phosphate dehydrogenase is inappropriate as internal control in comparative studies between skin tissue and cultured skin fibroblasts using Northern blot analysis. *Arch Dermatol Res*. 1999 Dec;291(12):659-61.
- Orlando C, Pinzani P, Pazzagli M. Developments in quantitative PCR. *Clin Chem Lab Med*. 1998 May;36(5):255-69.
- Paddison PJ, Caudy AA, Bernstein E, Hannon GJ, Conklin DS.: Short hairpin RNAs (shRNAs) induce sequence-specific silencing in mammalian cells. *Genes Dev* 2002 Apr 15;16(8):948-58
- Parker RM, Barnes NM. mRNA: detection by in Situ and northern hybridization. *Methods Mol Biol*. 1999;106:247-83.

- Peccoud J, Jacob C. Theoretical uncertainty of measurements using quantitative polymerase chain reaction. *Biophys J*. 1996 Jul;71(1):101-8.
- Pende A, Tremmel KD, DeMaria CT, Blaxall BC, Minobe WA, Sherman JA, Bisognano JD, Bristow MR, Brewer G, Port J. Regulation of the mRNA-binding protein AUF1 by activation of the beta-adrenergic receptor signal transduction pathway. *J Biol Chem*. 1996 Apr 5;271(14):8493-501.
- Pietri-Rouxel F, Lenzen G, Kapoor A, Drumare MF, Archimbault P, Strosberg AD, Manning BS. Molecular cloning and pharmacological characterization of the bovine beta 3-adrenergic receptor. *Eur J Biochem*. 1995 May 15;230(1):350-8.
- Ping P, Hammond HK. Diverse G protein and beta-adrenergic receptor mRNA expression in normal and failing porcine hearts. *Am J Physiol*. 1994 Nov;267(5 Pt 2):H2079-85
- Podlowski S. Expression und Funktion kardialer β -Adrenozeptoren unter Berücksichtigung von β 1-Adrenozeptor-Autoantikörpern bei der dilatativen Kardiomyopathie. Dissertation. Mathematisch-Naturwissenschaftliche Fakultät I der Humboldt-Universität zu Berlin 1997
- Podlowski S, Bramlage P, Baumann G, Morano I, Luther HP. Cardiac troponin I sense-antisense RNA duplexes in the myocardium. *J Cell Biochem*. 2002;85(1):198-207.
- Podlowski S, Wenzel K, Luther HP, Muller J, Bramlage P, Baumann G, Felix SB, Speer A, Hetzer R, Kopke K, Hoehe MR, Wallukat G. Beta1-adrenoceptor gene variations: a role in idiopathic dilated cardiomyopathy? *J Mol Med*. 2000;78(2):87-93.
- Podlowski S, Luther HP, Morwinski R, Muller J, Wallukat G. Agonistic anti-beta1-adrenergic receptor autoantibodies from cardiomyopathy patients reduce the beta1-adrenergic receptor expression in neonatal rat cardiomyocytes. *Circulation*. 1998 Dec 1;98(22):2470-6.
- Port JD, Huang LY, Malbon CC. Beta-adrenergic agonists that down-regulate receptor mRNA up-regulate a M(r) 35,000 protein(s) that selectively binds to beta-adrenergic receptor mRNAs. *J Biol Chem*. 1992 Nov 25;267(33):24103-8.
- Proud CG. PKR: a new name and new roles. *Trends Biochem Sci*. 1995 Jun;20(6):241-6.
- Proudfoot NJ, Furger A, Dye MJ. Integrating mRNA processing with transcription. *Cell*. 2002 Feb 22;108(4):501-12.
- Rao S, Watkins D, Cunningham D, Dunlop D, Johnson P, Selby P, Hancock BW, Fegan C, Culligan D, Schey S, Morris TC, Lissitchkov T, Oliver JW, Holmlund JT. Phase II study of ISIS 3521, an antisense oligodeoxynucleotide to protein kinase C alpha, in patients with previously treated low-grade non-Hodgkin's lymphoma. *Ann Oncol*. 2004 Sep;15(9):1413-8.
- Rapacciuolo A, Suvarna S, Barki-Harrington L, Luttrell LM, Cong M, Lefkowitz RJ, Rockman HA. Protein kinase A and G protein-coupled receptor kinase phosphorylation mediates beta-1 adrenergic receptor endocytosis through different pathways. *J Biol Chem*. 2003 Sep 12;278(37):35403-11. Epub 2003 Jun 23.
- Rappolee DA. Optimizing the sensitivity of RT-PCR. *Amplifications A Forum for PCR Users*, (1990) 7:1-4.
- Rathz DA, Brown KM, Kramer LA, Liggett SB. Amino acid 49 polymorphisms of the human beta1-adrenergic receptor affect agonist-promoted trafficking. *J Cardiovasc Pharmacol*. 2002 Feb;39(2):155-60.

- Ririe KM, Rasmussen RP, Wittwer CT. Product differentiation by analysis of DNA melting curves during the polymerase chain reaction. *Anal Biochem.* 1997 Feb 15;245(2):154-60.
- Rosok O, Sioud M. Systematic identification of sense-antisense transcripts in mammalian cells. *Nat Biotechnol.* 2004 Jan;22(1):104-8. Epub 2003 Dec 14.
- Rudin CM, Kozloff M, Hoffman PC, Edelman MJ, Karnauskas R, Tomek R, Szeto L, Vokes EE. Phase I study of G3139, a bcl-2 antisense oligonucleotide, combined with carboplatin and etoposide in patients with small-cell lung cancer. *J Clin Oncol.* 2004 Mar 15;22(6):1110-7.
- Ruiz-Gomez A, Mayor F Jr. Beta-adrenergic receptor kinase (GRK2) colocalizes with beta-adrenergic receptors during agonist-induced receptor internalization. *J Biol Chem.* 1997 Apr 11;272(15):9601-4.
- Sacomanno CF, Bordonaro M, Chen JS, Nordstrom JL. A faster ribonuclease protection assay. *Biotechniques.* 1992 Dec;13(6):846-50.
- Sato Y, Adachi-Akahane S, Prados P, Imai K, Nagao T. Desensitization and selective down-regulation of rat cardiac beta 1-adrenoceptors by prolonged in vivo infusion of T-0509, a beta 1-adrenoceptor full agonist. *Jpn J Pharmacol.* 1995 Dec;69(4):343-50.
- Sharp PA. RNA interference--2001. *Genes Dev.* 2001 Mar 1;15(5):485-90.
- Sharp PA. RNAi and double-strand RNA. *Genes Dev.* 1999 Jan 15;13(2):139-41.
- Shear M, Insel PA, Melmon KL, Coffino P. Agonist-specific refractoriness induced by isoproterenol. Studies with mutant cells. *J Biol Chem.* 1976 Dec 10;251(23):7572-6.
- Shi, H. et al. Genetic interference in *Trypanosoma brucei* by heritable and inducible double-stranded RNA. *RNA* 6, 1069–1076 (2000).
- Sibley DR, Benovic JL, Caron MG, Lefkowitz RJ. Regulation of transmembrane signaling by receptor phosphorylation. *Cell.* 1987 Mar 27;48(6):913-22.
- Sibley DR, Lefkowitz RJ. Molecular mechanisms of receptor desensitization using the beta-adrenergic receptor-coupled adenylate cyclase system as a model. *Nature.* 1985 Sep 12-18;317(6033):124-9.
- Sijen T, Kooter JM. Post-transcriptional gene-silencing: RNAs on the attack or on the defense? *Bioessays.* 2000 Jun;22(6):520-31.
- Sokoloff MH, Tso CL, Kaboo R, Nelson S, Ko J, Dorey F, Figlin RA, Pang S, deKernion J, Belldegrun A. Quantitative polymerase chain reaction does not improve preoperative prostate cancer staging: a clinicopathological molecular analysis of 121 patients. *J Urol.* 1996 Nov;156(5):1560-6.
- Sperelakis N. Regulation of calcium slow channels of heart by cyclic nucleotides and effects of ischemia. *Adv Pharmacol.* 1994;31:1-24.
- Stein CA, Cheng Y-C. Antisense oligonucleotides: is the bullet really magical? *Science.* 1993; 261 : 1004 –1012.
- Strosberg AD, Pietri-Rouxel F. Function and regulation of the beta 3-adrenoceptor. *Trends Pharmacol Sci.* 1996 Oct;17(10):373-81. Review.
- Strosberg AD. Structure, function, and regulation of adrenergic receptors. *Protein Sci.* 1993 Aug;2(8):1198-209. Review.

- Su YF, Harden TK, Perkins JP. Catecholamine-specific desensitization of adenylate cyclase. Evidence for a multistep process. *J Biol Chem*. 1980 Aug 10;255(15):7410-9. No abstract available.
- Su YF, Harden TK, Perkins JP. Isoproterenol-induced desensitization of adenylate cyclase in human astrocytoma cells. Relation of loss of hormonal responsiveness and decrement in beta-adrenergic receptors. *J Biol Chem*. 1979 Jan 10;254(1):38-41.
- Svoboda, P., Stein, P., Hayashi, H. & Schultz, R. M. Selective reduction of dormant maternal mRNAs in mouse oocytes by RNA interference. *Development* 127, 4147–4156 (2000).
- Swedberg K, Eneroth P, Kjekshus J, Wilhelmsen L. Hormones regulating cardiovascular function in patients with severe congestive heart failure and their relation to mortality. CONSENSUS Trial Study Group. *Circulation*. 1990 Nov;82(5):1730-6.
- Sylvén C, Mischa G, Jansson E, Sotonyi P, Fu LX, Waagstein F, Bronnegard M. Beta 1 and beta 2 adrenoceptor ligand and mRNA expression in dilated cardiomyopathy. *Biol Pharm Bull*. 1995 Oct;18(10):1430-4.
- Tada M, Katz AM. Phosphorylation of the sarcoplasmic reticulum and sarcolemma. *Annu Rev Physiol*. 1982;44:401-23.
- Thellin O, Zorzi W, Lakaye B, De Borman B, Coumans B, Hennen G, Grisar T, Igout A, Heinen E. Housekeeping genes as internal standards: use and limits. *J Biotechnol*. 1999 Oct 8;75(2-3):291-5.
- Timmons L, Fire A. Specific interference by ingested dsRNA. *Nature*. 1998 Oct 29;395(6705):854.
- Tolcher AW, Kuhn J, Schwartz G, Patnaik A, Hammond LA, Thompson I, Fingert H, Bushnell D, Malik S, Kreisberg J, Izbicka E, Smetzer L, Rowinsky EK. A Phase I pharmacokinetic and biological correlative study of oblimersen sodium (genasense, g3139), an antisense oligonucleotide to the bcl-2 mRNA, and of docetaxel in patients with hormone-refractory prostate cancer. *Clin Cancer Res*. 2004 Aug 1;10(15):5048-57.
- Tucker M, Valencia-Sanchez MA, Staples RR, Chen J, Denis CL, Parker R. The transcription factor associated Ccr4 and Caf1 proteins are components of the major cytoplasmic mRNA deadenylase in *Saccharomyces cerevisiae*. *Cell*. 2001 Feb 9;104(3):377-86.
- Tuschl T, Zamore PD, Lehmann R, Bartel DP, Sharp PA. Targeted mRNA degradation by double-stranded RNA in vitro. *Genes Dev*. 1999 Dec 15;13(24):3191-7.
- Ungerer M, Böhm M, Elce JS, Erdmann E, Lohse MJ. Altered expression of beta-adrenergic receptor kinase and beta 1-adrenergic receptors in the failing human heart. *Circulation*. 1993 Feb;87(2):454-63.
- Vanhee-Brossollet C, Vaquero C. Do natural antisense transcripts make sense in eukaryotes? *Gene*. 1998 Apr 28;211(1):1-9. Review.
- Villalona-Calero MA, Ritch P, Figueroa JA, Otterson GA, Belt R, Dow E, George S, Leonardo J, McCachren S, Miller GL, Modiano M, Valdivieso M, Geary R, Oliver JW, Holmlund J. A phase I/II study of LY900003, an antisense inhibitor of protein kinase C-alpha, in combination with cisplatin and gemcitabine in patients with advanced non-small cell lung cancer. *Clin Cancer Res*. 2004 Sep 15;10(18 Pt 1):6086-93.
- Wagner EG, Simons RW. Antisense RNA control in bacteria, phages, and plasmids. *Annu Rev Microbiol*. 1994;48:713-42.

- Wagner RW. Gene inhibition using antisense oligodeoxynucleotides. *Nature*. 1994 Nov 24;372(6504):333-5.
- Wang T, Brown MJ. mRNA quantification by real time TaqMan polymerase chain reaction: validation and comparison with RNase protection. *Anal Biochem*. 1999 Apr 10;269(1):198-201.
- Wang Y, Liu XF, Cornish KG, Zucker IH, Patel KP. Effects of nNOS antisense in the paraventricular nucleus on blood pressure and heart rate in rats with heart failure. *Am J Physiol Heart Circ Physiol*. 2005 Jan;288(1):H205-13. Epub 2004 Aug 26.
- Waters JS, Webb A, Cunningham D, Clarke PA, Raynaud F, di Stefano F, Cotter FE. Phase I clinical and pharmacokinetic study of bcl-2 antisense oligonucleotide therapy in patients with non-Hodgkin's lymphoma. *J Clin Oncol*. 2000 May;18(9):1812-23.
- Weis JH, Tan SS, Martin BK, Wittwer CT. Detection of rare mRNAs via quantitative RT-PCR. *Trends Genet*. 1992 Aug;8(8):263-4.
- Wianny F, Zernicka-Goetz M. Specific interference with gene function by double-stranded RNA in early mouse development. *Nat Cell Biol*. 2000 Feb;2(2):70-5.
- Wilusz CJ, Wormington M, Peltz SW. The cap-to-tail guide to mRNA turnover. *Nat Rev Mol Cell Biol* 2001 Apr;2(4):237-46
- Wink MU, Wehrle H. (1994): 3.10.8. Praktische Hinweise. In: PCR im medizinischen und biologischen Labor -Handbuch für den Praktiker, Darmstadt, GIT Verlag, 227-228.
- Winston WM, Molodowitch C, Hunter CP. Systemic RNAi in *C. elegans* requires the putative transmembrane protein SID-1. *Science*. 2002 Mar 29;295(5564):2456-9. Epub 2002 Feb 07.
- Wittwer CT, Herrmann MG, Moss AA, Rasmussen RP. Continuous fluorescence monitoring of rapid cycle DNA amplification. *Biotechniques*. 1997 Jan;22(1):130-1, 134-8.
- Wu DY, Ugozzoli L, Pal BK, Qian J, Wallace RB. The effect of temperature and oligonucleotide primer length on the specificity and efficiency of amplification by the polymerase chain reaction. *DNA Cell Biol*. 1991 Apr;10(3):233-8.
- Yelin R, Dahary D, Sorek R, Levanon EY, Goldstein O, Shoshan A, Diber A, Biton S, Tamir Y, Khosravi R, Nemzer S, Pinner E, Walach S, Bernstein J, Savitsky K, Rotman G. Widespread occurrence of antisense transcription in the human genome. *Nat Biotechnol*. 2003 Apr;21(4):379-86. Epub 2003 Mar 17.
- Yla Herttuala S. Vascular gene transfer. *Curr Opin Lipidol*. 1997 Apr;8(2):72-6.
- Zamecnik PC and Stephenson, M.L. (1978) Inhibition of Rous sarcoma virus replication and cell transformation by a specific oligodeoxynucleotide. *Proc. Natl Acad. Sci. USA*, 75, 289-294.
- Zamorano PL, Mahesh VB, Brann DW. Quantitative RT-PCR for neuroendocrine studies. A minireview. *Neuroendocrinology*. 1996 May;63(5):397-407.
- Zamore PD, Tuschl T, Sharp PA, Bartel DP. RNAi: double-stranded RNA directs the ATP-dependent cleavage of mRNA at 21 to 23 nucleotide intervals. *Cell*. 2000 Mar 31;101(1):25-33.
- Zhang J, Desai M, Ozanne SE, Doherty C, Hales CN, Byrne CD. Two variants of quantitative reverse transcriptase PCR used to show differential expression of alpha-, beta- and gamma-fibrinogen genes in rat liver lobes. *Biochem J*. 1997 Feb 1;321 (Pt 3):769-75.

- Zhang R, Diasio RB, Lu Z, Liu T, Jiang Z, Galbraith WM, Agrawal S. Pharmacokinetics and tissue distribution in rats of an oligodeoxynucleotide phosphorothioate (GEM 91) developed as a therapeutic agent for human immunodeficiency virus type-1. *Biochem Pharmacol.* 1995;49:929–939.
- Zhang Y, Kimura B, Shen L, Phillips MI. New beta-blocker: prolonged reduction in high blood pressure with beta(1) antisense oligodeoxynucleotides. *Hypertension.* 2000 Jan;35(1 Pt 2):219-24.
- Zuniga Mejia Borja A, Meijers C, Zeller R. Expression of alternatively spliced bFGF first coding exons and antisense mRNAs during chicken embryogenesis. *Dev Biol.* 1993 May;157(1):110-8.