

6 Literaturverzeichnis

Albrecht, S. und J. Schuttler (1999). "Postoperative pain management after intraoperative remifentanil." *Anesth Analg* 89: 40-45.

Ali, N. (1986). "Hyperalgesic response in a patient receiving high concentrations of spinal morphine." *Anesthesiology* 65: 449-450.

Angst, M. und W. Koppert (2003). "Short-term infusion of the μ -opioid agonist remifentanil in humans causes hyperalgesia during withdrawal." *Pain* 106: 49-57.

Arendt-Nielsen, L. (1997). "Induction and assessment of experimental pain from human skin, muscle and viscera." *Proceedings of the 8th World Congress on Pain. Edited by Jensen TS, Turner JA, Wiesenfeld-Hallin Z. Seattle, IASP Press: 393-425.*

Ashton, H. und I. Ebenezer (1984). "Effects of acupuncture and transcutaneous electrical nerve stimulation on cold-induced pain in normal subjects." *J Psychosom Res* 28: 301-308.

Bayer, T. und J. Coverdale (1998). "The role of prior pain experience and expectancy in psychologically and physically induced pain." *Pain* 74: 327-331.

Birbaumer, N. und R. Schmidt (1999). "Biologische Psychologie." *Springer Verlag* 4. Auflage.

Black, M. und J. Hill (1999). "Behavioral and physiological effects of remifentanil and alfentanil in healthy volunteers." *Anesthesiology* 90(3): 718-726.

Brennum, J. (1989). "Measurements of human pressure - pain thresholds on fingers and toes." *Pain* 38: 211-217.

Bullinger, M. und D. Naber (1984). "Endocrine effects of the cold pressor test: relationships to subjective pain appraisal and coping." *Psychiatry Res* 12: 227-233.

Campero, M. und J. Serra (1996). "C-polymodal nociceptors activated by noxious low temperature in human skin." *J Physiol Lond* 497: 565-572.

Celerier, E. und J. Laulin (2001). "Progressive enhancement of delayed hyperalgesia induced by repeated heroin administration: a sensitization process." *J Neurosci* 21: 4074-4080.

Celerier, E. und C. Rivat (2000). "Long-lasting hyperalgesia induced by fentanyl in rats: preventive effect of ketamine." *Anesthesiology* 92(2): 465-472.

Chen, A. und S. Dworkin (1989). "Human pain responsivity in a tonic pain model: psychological determinants." *Pain* 1989(37): 143-160.

Chia, Y. und K. Liu (1999). "Intraoperative high dose fentanyl induces postoperative fentanyl tolerance." *Can J Anaesth* 46: 872-877.

- Coderre, T. und J. Katz (1993).** "Contribution of central neuroplasticity to pathological pain: review of clinical and experimental evidence." *Pain* 52(3): 259-285.
- Compton, P. und V. Charuvastra (2000).** "Pain responses in methadone-maintained opioid abusers." *J Pain Symptom Manage* 20: 237-245.
- Compton, P. und V. Charuvastra (2001).** "Pain intolerance in opioid-maintained former opiate addicts: effect of long-acting maintenance agent." *Drug and Alcohol Dependence* 63: 139-146.
- Cortinez, L. und V. Brandes (2001).** "No clinical evidence of acute opioid tolerance after remifentanyl-based anaesthesia." *Br J Anaesth* 87(6): 866-869.
- Curatolo, M. und F. Petersen (1997).** "Epidural epinephrine and clonidine: segmental analgesia and effects on different pain modalities." *Anesthesiology* 87: 785-794.
- Dershwitz, M. und G. Randel (1995).** "Initial clinical experience with remifentanyl, a new opioid metabolized by esterases." *Anesth Analg* 81: 619-623.
- Dickenson, A. (1995).** "Spinal cord pharmacology of pain." *Br J Anaesth* 75: 193-200.
- Doverly, M. und J. White (2001).** "Hyperalgesic responses in methadone maintenance patients." *Pain* 90: 91-96.
- Dymond, D. und J. Caplin (1984).** "Temporal evolution of changes in left ventricular function induced by cold pressor stimulation: an assessment with radionuclide angiography and gold 195m." *Br Heart J* 51: 557-564.
- Eisenach, J. (2000).** "Preemptive hyperalgesia, not analgesia? (editorial)." *Anesthesiology* 92: 308-309.
- Fillingim, R. und W. Maixner (1995).** "Gender differences in response to noxious stimuli." *Pain Forum* 4: 209-221.
- Fletcher, D. und M. Pinaud (2000).** "The efficacy of intravenous 0.15 versus 0.25 mg/kg intraoperative morphine for immediate postoperative analgesia after remifentanyl-based anesthesia for major surgery." *Anesth Analg* 90: 666-671.
- Freye, E. und L. Latasch (2003).** "Toleranzentwicklung unter Opioidgabe - Molekulare Mechanismen und klinische Bedeutung." *Anesthesiol Intensivmed Notfallmed Schmerzther* 38: 14-26.
- Gardmark, M. und M. Ekblom (1993).** "Quantification of effect delay and acute tolerance development to morphine in the rat." *J Pharmacol Exp Ther* 267: 1061-1067.
- Glass, P. und T. Gan (1999).** "A review of the pharmacokinetics and pharmacodynamics of remifentanyl." *Anesth Analg* 89: 7-14.
- Goldstein, A. und S. Tachibana (1979).** "Dynorphin-(1-13), an extraordinarily potent opioid peptide." *Proc Natl Acad Sci USA* 76: 6666-6670.

Gormsen, L. und A. Riisgard Ribe (2004). "Pain thresholds during and after treatment of severe depression with electroconvulsive therapy." *Eur J Pain* 8(5): 487-493.

Guignard, B. und A. Bossard (2000). "Acute opioid tolerance: intraoperative remifentanil increases postoperative pain and morphine requirement." *Anesthesiology* 93: 409-417.

Guignard, B. und C. Coste (2002). "Supplementing desflurane-remifentanil anesthesia with small-dose ketamine reduces perioperative opioid analgesic requirements." *Anesth Analg* 95: 103-108.

Gustorff, B. und G. Nahlik (2002). "The absence of acute tolerance during remifentanil infusion in volunteers." *Anesth Analg* 94: 1223-1228.

Gutstein, H. (1996). "The effects of pain on opioid tolerance: how do we resolve the controversy?" *Pharmacol Rev* 48: 403-407, 409-411.

Hines, E. J. und G. Brown (1933). "A standard test for measuring the variability of blood pressure: its significance as an index of the prehypertensive state." *Ann Intern Med* 7: 209-217.

Ho, S. und J. Wang (1999). "Surgical pain attenuates acute morphine tolerance in rats." *Br J Anaesth* 82: 112-116.

Ho, S. und J. Wang (2002). "The magnitude of acute tolerance to morphine analgesia: concentration-dependent or time-dependent?" *Anesth Analg* 95: 948-951.

Hood, D. und R. Curry (2003). "Intravenous remifentanil produces withdrawal hyperalgesia in volunteers with capsaicin-induced hyperalgesia." *Anesth Analg* 97: 810-815.

Jones, S. und H. McQuay (1988). "Morphine and ibuprofen compared using the cold pressor test." *Pain* 34: 117-122.

Kaplan, H. und H. Fields (1991). "Hyperalgesia during acute opioid abstinence: evidence for a nociceptive facilitating function of the rostral ventromedial medulla." *J Neurosci* 11: 1433-1439.

Kest, B. und E. Sarton (2000). "Gender differences in opioid-mediated analgesia: animal and human studies." *Anesthesiology* 93: 539-547.

Kissin, I. und C. Bright (2000). "The effect of ketamine on opioid-induced acute tolerance: can it explain reduction of opioid consumption with ketamine-opioid analgesic combinations?" *Anesth Analg* 91: 1483-1488.

Kissin, I. und C. Bright (2001). "Can inflammatory pain prevent the development of acute tolerance to alfentanil?" *Anesth Analg* 91: 110-116.

Kissin, I. und S. Lee (1996). "Time course characteristics of acute tolerance development to continuously infused alfentanil in rats." *Anesth Analg* 83: 600-605.

Kochs, E. und D. Côté (2000). "Postoperative pain management and recovery after remifentanil-based anaesthesia with isoflurane or propofol for major abdominal surgery." *Br J Anaesth* 84(2): 169-173.

Koppert, W. (2004). "Opioid-induzierte Hyperalgesie." *Anaesthesist* 53: 455-466.

Koppert, W. und R. Sittl (2003). "Differential modulation of remifentanil-induced analgesia and postinfusion hyperalgesia by s-ketamine and clonidine in humans." *Anesthesiology* 99(1): 152-159.

Larcher, A. und J. Laulin (1998). "Acute tolerance associated with a single opiate administration: involvement of N-methyl-D-aspartate-dependent pain facilitatory systems." *Neuroscience* 84: 583-589.

Larsen, R. (1999). "Anästhesie." *Urban und Schwarzenberg Verlag* 6. Auflage: 101-106, 405.

Laulin, J. und A. Larcher (1998). "Long-lasting increased pain sensitivity in rat following exposure to heroin for the first time." *Eur J Neurosci* 10: 782-785.

Li, X. und M. Angst (2001). "Opioid-induced hyperalgesia and incisional pain." *Anesth Analg* 93: 204-209.

Lötsch, J. und M. Angst (2003). "The μ -opioid agonist remifentanil attenuates hyperalgesia evoked by blunt and punctuated stimuli with different potency: a pharmacological evaluation of the freeze lesion in humans." *Pain* 102: 151-161.

Lovallo, W. (1975). "The cold pressor test and autonomic function: a review and integration." *Psychophysiology* 12: 268-282.

Luginbühl, M. und A. Gerber (2003). "Modulation of remifentanil-induced analgesia, hyperalgesia, and tolerance by small-dose ketamine in humans." *Anesth Analg* 96: 726-732.

Luginbühl, M. und T. Schnider (2001). "Comparison of five experimental pain tests to measure analgesic effects of alfentanil." *Anesthesiology* 95(1): 22-29.

Lyness, W. und F. Smith (1989). "Morphine self-administration in the rat during adjuvant-induced arthritis." *Life Sci* 45: 2217-2224.

Mao, J. und D. Price (1994). "Thermal hyperalgesia in association with the development of morphine tolerance in rats: roles of excitatory amino acid receptors and protein kinase C." *J Neurosci* 14: 2301-2312.

Mao, J. und D. Price (1995). "Mechanisms of hyperalgesia and morphine tolerance: a current view of their possible interactions." *Pain* 62: 259-274.

Mc Mahon, S. und M. Koltzenburg (1990). "Novel classes of nociceptors beyond Sherrington." *Trends in Neurosciences* 13.

- Michelsen, L. und C. Hug, Jr (1996).** "The pharmacokinetics of remifentanyl." *J Clin Anesth* 8(8): 679-82.
- Mitchell, L. und A. Raymond (2004).** "Temperature and the cold pressor test." *J Pain* 5(4): 233-238.
- Posner, J. und A. Telekes (1985).** "Effects of an opiate on cold-induced pain and the CNS in healthy volunteers." *Pain* 23: 73-82.
- Przewlocki, R. und H. Machelska (1993).** "Inhibition of nitric oxide synthase enhances morphine antinociception in the rat spinal cord." *Life Sci* 53: 1-5.
- Rivat, C. und J. Laulin (2002).** "Fentanyl enhancement of carrageenan-induced long-lasting hyperalgesia in rats: prevention by the N-methyl-D-aspartate receptor antagonist ketamine." *Anesthesiology* 96: 381-391.
- Schmelz, M. und R. Schmidt (1996).** "Limitation of sensitization to injured parts of receptive fields in human skin C-nociceptors." *Exp Brain Res* 109: 141-147.
- Schmidt, R. und M. Schmelz (1995).** "Novel classes of responsive and unresponsive C-nociceptors in human skin." *J Neurosci* 15: 333-341.
- Simonnet, G. und C. Rivat (2003).** "Opioid-induced hyperalgesia: abnormal or normal pain." *Neuroreport* 14: 1-7.
- Solomon, R. (1980).** "The opponent-process theory of acquired motivation: the costs of pleasure and the benefits of pain." *Am Psychol.* 35(8): 691-712.
- Solomon, R. und J. Corbit (1974).** "An opponent-process theory of motivation. I. Temporal dynamics of affect." *Psychol Rev* 81(2): 119-145.
- Staats, P. und H. Hekmat (1998).** "Suggestion/placebo effects on pain: negative as well as positive." *J Pain Symptom Manage* 15: 235-243.
- Sun, H. (1998).** "Naloxone-precipitated acute opioid withdrawal syndrome after epidural morphine." *Anesth Analg* 86: 544-545.
- Sweitzer, S. und C. Allen (2004).** "Mechanical allodynia and thermal hyperalgesia upon acute opioid withdrawal in the neonatal rat." *Pain* 110: 269-280.
- Treede, R. und W. Magerl (2000).** "Multiple mechanisms of secondary hyperalgesia." *Prog Brain Res* 129: 331-341.
- Trujillo, K. und H. Akil (1991).** "Inhibition of morphine tolerance and dependence by the NMDA receptor antagonist MK-801." *Science* 251: 85-87.
- Vanderah, T. und M. Ossipov (2001).** "Mechanisms of opioid-induced pain and antinociceptive tolerance: descending facilitation and spinal dynorphin." *Pain* 92: 5-9.

- Vanderah, T. und N. Suenaga (2001).** "Tonic descending facilitation from the rostral ventromedial medulla mediates opioid- induced abnormal pain and antinociceptive tolerance." *J Neurosci* 21: 279-286.
- Vinik, H. und I. Kissin (1998).** "Rapid development of tolerance to analgesia during remifentanil infusion in humans." *Anesth Analg* 86(6): 1307-1311.
- Wolf, B. (1984).** "Methods of testing pain mechanisms in normal man." *In: Wall PD, Melzack R, editors. Textbook of pain. New York: Churchill Livingstone: 186-194.*
- Woolf, C. und S. Thompson (1991).** "The induction and maintenance of central sensitization is dependent on N-methyl-D-aspartic acid receptor activation; implications for the treatment of post-injury pain hypersensitivity states." *Pain* 44: 293-299.
- Yarmush, J. und R. D`Angelo (1997).** "A comparison of remifentanil and morphine sulfate for acute postoperative analgesia after total intravenous anesthesia with remifentanil and propofol." *Anesthesiology* 87: 235-243.
- Yuan, C. und T. Karrison (1998).** "Dose-related effects of oral acetaminophen on cold-induced pain: A double-blind, randomized, placebo-controlled trial." *Clin Pharmacol Ther* 63(3): 379-383.
- Zarate, E. und M. Sa Rego (1999).** "Comparison of adenosine and remifentanil infusions as adjuvants to desflurane anesthesia." *Anesthesiology* 90: 956-963.
- Ziegler, E. und W. Magerl (1999).** "Secondary hyperalgesia to punctate mechanical stimuli. Central sensitization to A-fibre nociceptor input." *Brain* 122: 2245-2257.