

Acknowledgements

"Don't cry because it's over. Smile because it happened."

Theodor Seuss Geisel

This is going to be hard. Because there are so many people to whom I need to thank, and I fear I will not remember all of them. But I'll try.

First of all, I would like to thank Prof. Dr. Eckart Rühl, for giving me the chance to work in his group. I am indebted to him for teaching me many things, and, most of all, for having the patience of explaining things twice, and sometimes even thrice.

Many thanks are addressed to Dr. Roman Flesch, for the nice beam times we spent at BESSY and for the many times he spared some of his time just to give an useful advice. Of course, sometimes advice is not enough, so, when this was the case, he spent time explaining and showing how things should be done.

Also, many thanks go to Kai Godehusen and Olaf Schwarzkopf and to the BESSY staff, in Berlin who made sure the beam lines are usable and in use.

I want to express my gratitude towards Dr. Thomas Schlathöler and Dr. Fresia Alvarado and all the people from the "Morgenstern working group" at the KVI Groningen, which were so hard working and entertaining at the same time during my beam time at their facility. It was a one-of-a-kind experience, proving that despite all the problems that can occur, a beam time can still be successful.

To the "Frankfurt gang", and especially to Till, Markus, Lutz, Sven and Katharina I am indebted for their support during the interesting beam time at BESSY we shared. Actually, I guess the appropriate description of that beam time is "roller-coaster", considering all the ups and downs of those two weeks. And I guess I'll always remember what should be the motto of every beam time, especially when things seem hopeless: "we can handle it"¹

¹"It's gonna be hard, but..we can handle it"

I am indebted to the workshop personnel from Institute of Physical Chemistry of the University of Würzburg for their help in building up my experimental setup. Special thanks go to Thomas Schreckling, Wolfgang Liebler, Ralf Kohrmann, Peter Lang, Gerhard Bömmel, Rainer Eck, Stefan Schwarz and Jürgen Zimmermann.

Of course, as they were the ones to put up with me for the past three years, all the people in the "Rühl working group" deserve to be mentioned in particular.

Harald has gained the right to be the first one mentioned at this point for the countless coffees², beers and pizzas we consumed chatting. To him I owe a better understanding of the German culture and economy and a valuable friendship. And, of course, he gets to be blamed for the time I spent solving the problems using L^AT_EX brought up.

Renè gets the blame for my nickname. And gets to be thanked for the many good times³ we had when, while having a beer or a coffee, he tried convincing me that kayaking is a nice sport. I guess in the end I understood that :-). And, of course, together with Harald, he's responsible for my improved German. Thanks for being my friend, Renè.

I'd like to thank Jürgen for spending some of his time to advise me as I started doing the SimIon simulations and the GSCF3 calculations. I need to thank him also for the interesting discussions we had throughout the last three years, especially in what regards the differences between a MAC and a PC.

To Christina, Andreas Hoffmann and Dr. Sofia Dembski I am very grateful for the many interesting discussions and laughters we shared having coffees and cakes on the 'red couch'.

Mathias, Eddie and Andreas Wirsing are thanked for the numerous talks "over a coffee" and for the loads of good time we shared together.

I'd like to thank Dr. B. Wassermann for the patience he proved in explaining me how a simulation program needs to be handled, and for the enlightening scientific discussions we shared.

Of course I need to thank Andrée for the help she always gave me in sorting out the paperwork which needed to be done in Würzburg. And she deserves special thanks for the excellent cakes she baked and shared with us, especially the rhubarb one :-). For this as well as for the countless times she managed to put a smile on my face.

To Sabine I am grateful for the many mature discussions we shared. She is also responsible for my better understanding of Würzburg and of the people living in the area. And for always listening to me and giving me good advice she deserves special thanks.

To Prof. Dr. John Neville I am grateful for the many enlightening discussions and for the beam time at BESSY we shared together

²"Show me the coffee!!!"

³"Try dragging it!!!"

I can not but thank also my friends who offered me support and comfort through the hard moments I've had so far: Nicu, Horia, M@rio, Radu, Bernd, Ste , Ste en and Ioana.

Special thanks are addressed to my parents, my sister and my favorite uncle who always supported my actions (the good ones) and criticized the bad ones. Without their support I would not be where and who I am today.

And, last, but not least, I would like to thank my wife, Niculina, for the patience and dedication with which she stands by me every day.

Erklärung

Hiermit erkläre ich an Eides statt, dass ich die Dissertation 'Photoionization and investigation of chalcogene clusters' selbständig angefertigt und keine anderen als die von mir angegebenen Quellen und Hilfsmittel benutzt habe.

Ich erkläre außerdem, dass diese Dissertation weder in gleicher oder anderer Form bereits in einem anderen Prüfungsverfahren vorgelegen hat.

Ich habe früher außer den mit dem Zulassungsgesuch urkundlich vorgelegten Graden keine weiteren akademischen Grade erworben oder zu erwerben versucht.

Berlin, den

Unterschrift
Tiberiu Arion

Publications and Conference Contributions

Parts of this thesis have been already presented in scientific journals and/or at conferences.

Publications

- T. Arion, R. Flesch, T. Schlathölter, F. Alvarado, R. Hoekstra, R. Morgenstern and E. Rühl
"Collision induced fragmentation of free sulfur clusters"
submitted (2008)
- T. Arion, R. Flesch, J. Neville, J. Plenge and E. Rühl
"Investigation of the core-level excitation and fragmentation of free selenium aggregates"
In preparation (2008)
- T. Arion, T. Jahnke, M. Schöer, S. Schöbler, K. Kreidi, L. Foucar, R. Dörner and E. Rühl
"COLTRIMS investigation of the fragmentation of multiply charged sulfur clusters"
In preparation (2008)
- T. Arion, T. Jahnke, M. Schöer, S. Schöbler, K. Kreidi, L. Foucar, R. Dörner and E. Rühl
"Fragmentation of multiply charged sulfur clusters investigated by COLTRIMS"
BESSY annual report (2007)
- T. Arion, R. Flesch, J. Neville and E. Rühl
"Inner-shell excitation and photo-fragmentation of free selenium clusters"
BESSY annual report (2006)
- R. Flesch, I. L. Bradeanu, J. Plenge, T. Arion and E. Rühl
"High-resolution spectroscopy of core-excited organic van der Waals-clusters"
BESSY annual report (2005)

Conference Contributions

- T. Arion, T. Jahnke, M. Schöber, S. Schöblier, K. Kreidi, L. Foucar, R. Dörner and E. Rühl
"*Fragmentation of multiply charged sulfur clusters investigated by COLTRIMS*"
poster, 26th BESSY User Meeting, Berlin, Germany (2007)
- T. Arion, R. Flesch, J. J. Neville and E. Rühl
"*Inner-shell excitation and fragmentation of free selenium clusters*"
talk, 15th International Conference on Vacuum Ultraviolet Radiation Physics, Berlin, Germany (2007)
- T. Arion, R. Flesch, J. Plenge, J. J. Neville and E. Rühl
"*Inner-shell excitation and fragmentation of free selenium clusters*"
poster, 52nd IUVSTA Workshop on "Structure and dynamics of free and supported nanoparticles using short wavelength radiation", Erice-Sicily, Italy (2007)
- T. Arion, R. Flesch, T. Schlathölter, F. Alvarado, R. Hoekstra, R. Morgenstern and E. Rühl
"*Highly charged ion induced fragmentation of free sulfur clusters*"
poster, Bunsentagung 2007: die 106. Hauptversammlung der Deutschen Bunsen-Gesellschaft für Physikalische Chemie, Graz, Austria (2007)
- T. Arion, R. Flesch, T. Schlathölter, F. Alvarado, R. Hoekstra, R. Morgenstern and E. Rühl
"*Collision-induced fragmentation of free sulfur clusters*"
talk, Deutsche Physikalische Gesellschaft Tagung, Düsseldorf, Germany (2007)
- I. L. Bradeanu, R. Flesch, J. Plenge, T. Arion, A. A. Pavlychev, N. Kosugi and E. Rühl
"*Inner-shell excitation of molecular van der Waals clusters*"
talk, The 13th International Conference on X-Ray Absorption Fine Structure, Stanford, USA (2006)
- R. Flesch, J. Plenge, T. Arion, I. Bradeanu, M. Meyer, A. A. Pavlychev and E. Rühl
"*Near-edge spectroscopy of atomic and molecular van der Waals clusters*"
talk, 354. Wilhelm und Else Heraeus-Seminar on "Structure and Dynamics of Free Clusters and Nanoparticles using Short Wavelength Radiation", Bad Honnef, Germany (2005)

List of Figures

2.1	Composition of sulfur vapor as a function of the temperature [19]	13
2.2	Composition of selenium vapor as a function of the temperature [32]	15
2.3	Schematics of creation and filling of a core hole subsequent to the absorption of a X-ray photon	16
2.4	The mean free path of the electrons in solid (the dashed curve is a theoretical result)[36]	17
2.5	K-edge NEXAFS spectrum of a diatomic molecule	18
2.6	Core-level excitation of a diatomic molecule, from the 1s level into unoccupied molecular orbitals	19
2.7	Schematics of a two-body dissociation, together with an ideal PEPIPICO signal shape	24
2.8	Schematics of a deferred charge separation mechanism, together with an ideal PEPIPICO signal shape	26
2.9	Schematics of a secondary decay mechanism, together with an ideal PEPIPICO signal shape	28
2.10	Schematics of a secondary decay in competition mechanism, together with an ideal PEPIPICO signal shape	30
3.1	Schematics of the two vacuum systems used	34
3.2	Schematics of two staged oven cluster source	35
3.3	Schematics of He (I) discharge lamp	38
3.4	Schematics of the ECR ion source at the KVI Groningen, The Netherlands	39
3.5	Schematics of an electron storage ring	40
3.6	Optical layout of the BESSY U49/2-PGM 1 beam line [82]	42
3.7	Schematics of a micro-channel plate detector	43

3.8	Schematics of a Wiley-McLaren type time of flight mass spectrometer	44
3.9	Schematics of the working principle of a COLTRIMS experimental setup	46
3.10	Trajectories of electrons and ions in the COLTRIMS spectrometer	47
3.11	Schematics of a quadratic (a), hexagonal (b) and side view (c) of a delay line anode [97]	49
3.12	Schematics of the delay line principle	50
4.1	Ion-ion-coincidence spectrum of sulfur clusters recorded after ionization with a Xe^{5+} ion beam	57
4.2	Ion-ion-coincidence spectrum of sulfur clusters recorded after ionization with Xe^{20+} ion beam	58
4.3	Time of flight mass spectra of sulfur vapor after ionization with $h\nu = 21.2\text{ eV}$ photons	59
4.4	Time of flight mass spectra of sulfur clusters recorded after ionization with various Xe^{n+} ion beam charges	61
4.5	Time of flight mass spectra of sulfur clusters recorded after ionization with various Xe^{n+} ion beam charges	63
4.6	Ion-ion-coincidence correlation plots of sulfur aggregates ionized via collisions with Xe^{5+} ions	65
4.7	The puckered ring structure of the S_8 cluster	67
4.8	Ion-ion-coincidence correlation plots of sulfur aggregates ionized via collisions with Xe^{10+} ions	71
4.9	Shapes of ion-ion-coincidence signals from sulfur ionic fragments resulting through ionization via collisions with Xe^{15+} ions	74
4.10	Ion-ion-coincidence correlation plots of sulfur aggregates ionized via collisions with Xe^{20+} ions	79
4.11	Photoion-photoion-coincidence spectra of sulfur clusters recorded at $T = 400\text{ K}$ (see text for details)	85
4.12	Radius of the detector vs. the time-of-flight of the ion (a) and the calculated energy resolution of the ion detector (b) [126] (see text for details)	86
4.13	Ion-ion coincidence spectrum after ionization with $h\nu = 419\text{ eV}$ (see text for details)	87
4.14	Momentum distribution and KER for the N^+ / N^+ channel after ionization with $h\nu = 419\text{ eV}$ (see text for details)	87

4.15	KER distribution of the N^+ / N^+ channel after ionization with $h\nu = 419$ eV as a function of the angle (see text for details)	88
4.16	Photoion-photoion-coincidence spectra of sulfur clusters as recorded after ionization with $h\nu = 158$ eV ((a) and (b)), $h\nu = 174$ eV ((c) and (d)) and $h\nu = 235$ eV ((e) and (f)).	89
4.17	KER distribution and KER distribution as a function of the angle for the S^+ / S^{2+} channel after ionization with $h\nu = 158$ eV ((a) and (b)), $h\nu = 174$ eV ((c) and (d)) and $h\nu = 235$ eV ((e) and (f)) (see text for details)	90
4.18	KER distribution and KER distribution as a function of the angle for the S^+ / S^+ channel after ionization with $h\nu = 174$ eV ((a) and (b)) and $h\nu = 235$ eV ((c) and (d)) (see text for details)	91
4.19	KER distribution and KER distribution as a function of the angle for the S^{2+} / S^{2+} channel after ionization with $h\nu = 235$ eV (see text for details)	93
4.20	Time of flight mass spectra of selenium clusters recorded after ionization with $h\nu = 21.22$ eV photons	95
4.21	Time of flight mass spectra of selenium clusters recorded at $T = 656$ K after ionization with $h\nu = 150$ eV and $h\nu = 1450$ eV	97
4.22	PEPIPICO spectrum of selenium clusters recorded after ionization with $h\nu = 150$ eV	99
4.23	Shapes of PEPIPICO signals from selenium ionic fragments resulting through ionization with $h\nu = 150$ eV photons	100
4.24	Total yield spectra of selenium clusters recorded at the 2p-ionization edge	103
4.25	The structure of the Se_5 (a) and Se_6 (b) clusters used for the simulations performed using the GSCF3 code	104
4.26	Total electron yield of the near-edge structure of selenium clusters recorded at the 2p-ionization edge	105

List of Tables

4.1	Ionization potentials of sulfur clusters represented as a function of the size of the cluster, as reported in [109] and [110]	56
4.2	Cation pairs identified after ionization with ions beams of Xe^{5+} and Xe^{20+} , respectively	57
4.3	Mass peaks' intensities normalized to the intensity of the S_8^+ mass peak, as recorded after ionization with $h\nu = 21.2$ eV photons at different temperatures of selenium vapor, as shown in Fig. 4.3	60
4.4	Mass peaks' intensities normalized to the intensity of the S_8^+ mass peak, as recorded after ionization with ionization with various Xe^{n+} ion beam charges, as shown in Fig. 4.4	62
4.5	Peak shape analysis of the ion-ion-coincidence signals recorded after ionization via collisions with Xe^{5+} ions. m and n represent the slopes of the long and of the short edge of the correlation plots, respectively	69
4.6	Peak shape analysis of the ion-ion-coincidence signals recorded after ionization via collisions with Xe^{10+} ions. m and n represent the slopes of the long and of the short edge of the correlation plots, respectively	72
4.7	Peak shape analysis of the ion-ion-coincidence signals recorded after ionization via collisions with Xe^{15+} ions. m and n represent the slopes of the long and of the short edge of the correlation plots, respectively	77
4.8	Peak shape analysis of the ion-ion-coincidence signals recorded after ionization via collisions with Xe^{20+} ions. m and n represent the slopes of the long and of the short edge of the correlation plots, respectively	82
4.9	Ionization potentials of selenium clusters represented as a function of the size of the cluster, as reported by Berkowitz [33]	94
4.10	Mass peaks intensities normalized to the intensity of the Se_2^+ mass peak, as recorded after ionization with $h\nu = 21.2$ eV photons at different temperatures of selenium vapor, as shown in Fig. 4.20	96

4.11	Mass peaks' intensities normalized to the intensity of the Se^+ mass peak, as recorded at $T = 656$ K after ionization with various energy photons, as shown in Fig. 4.21	98
4.12	Peak shape analysis of the ion-ion-coincidence signals shown in Fig.4.6. m represents the slope of the long edge of the correlation plots	102
4.13	Geometry optimization results as obtained from using the Gaussian98 [40] program, with different basis sets	107
4.14	Geometry optimization results as obtained from using the Gaussian98 [40] program, with different basis sets	108
4.15	Theoretical results obtained with the GSCF3 code for the Se_5 and Se_6 aggregates	109