

## 7. Literatur

- Ahnert-Hilger, G.; Holtje, M.; Pahner, I.; Winter, S.; Brunk, I. (2003) Regulation of vesicular neurotransmitter transporters. *Rev.Physiol Biochem.Pharmacol.* 150: 140-160
- Ahnert-Hilger, G.; Nurnberg, B.; Exner, T.; Schafer, T.; Jahn, R. (1998) The heterotrimeric G protein Go2 regulates catecholamine uptake by secretory vesicles. *Embo J* 17(2): 406-13
- Ahnert-Hilger, G.; Pahner, I.; Höltje, M. (2000) Pore-Forming Toxins as Cell-Biological and Pharmacological Tools. *Handb.Exp.Pharm.* 145: 557-573
- Ahnert-Hilger, G.; Schafer, T.; Spicher, K.; Grund, C.; Schultz, G.; Wiedenmann, B. (1994) Detection of G-protein heterotrimers on large dense core and small synaptic vesicles of neuroendocrine and neuronal cells. *Eur.J.Cell Biol.* 65(1): 26-38
- Alfonso, A.; Grundahl, K.; Duerr, J.S.; Han, H.P.; Rand, J.B. (1993) The Caenorhabditis elegans unc-17 gene: a putative vesicular acetylcholine transporter. *Science* 261(5121): 617-9
- Aronin, N.; DiFiglia, M. (1992) The subcellular localization of the G-protein Gi alpha in the basal ganglia reveals its potential role in both signal transduction and vesicle trafficking. *J Neurosci* 12(9): 3435-44
- Bae, Y.S.; Lee, T.G.; Park, J.C.; Hur, J.H.; Kim, Y.; Heo, K.; Kwak, J.Y.; Suh, P.G.; Ryu, S.H. (2003) Identification of a compound that directly stimulates phospholipase C activity. *Molecular Pharmacology* 63(5): 1043-1050
- Bai, L.; Xu, H.; Collins, J.F.; Ghishan, F.K. (2001) Molecular and functional analysis of a novel neuronal vesicular glutamate transporter. *J Biol Chem* 276(39): 36764-9
- Bedet, C.; Isambert, M.F.; Henry, J.P.; Gasnier, B. (2000) Constitutive phosphorylation of the vesicular inhibitory amino acid transporter in rat central nervous system. *J Neurochem* 75(4): 1654-63
- Bellocchio, E.E.; Reimer, R.J.; Fremeau, R.T., Jr.; Edwards, R.H. (2000) Uptake of glutamate into synaptic vesicles by an inorganic phosphate transporter. *Science* 289(5481): 957-60
- Bourne, H.R.; Sanders, D.A.; McCormick, F. (1990) The GTPase superfamily: a conserved switch for diverse cell functions. *Nature* 348(6297): 125-32
- Bruns, D.; Riedel, D.; Klingauf, J.; Jahn, R. (2000) Quantal release of serotonin. *Neuron* 28(1): 205-20
- Chaudhry, F.A.; Reimer, R.J.; Bellocchio, E.E.; Danbolt, N.C.; Osen, K.K.; Edwards, R.H.; Storm-Mathisen, J. (1998) The vesicular GABA transporter, VGAT, localizes to synaptic vesicles in sets of glycinergic as well as GABAergic neurons. *J Neurosci* 18(23): 9733-50
- Cowan, C.W.; Fariss, R.N.; Sokal, I.; Palczewski, K.; Wensel, T.G. (1998) High expression levels in cones of RGS9, the predominant GTPase accelerating protein of rods. *Proceedings of the National Academy of Sciences of the United States of America* 95(9): 5351-5356

- Cumbay, M.G.; Watts, V.J. (2001) Heterologous sensitization of recombinant adenylyl cyclases by activation of D(2) dopamine receptors. *J Pharmacol Exp Ther* 297(3): 1201-9
- Danbolt, N.C. (2001) Glutamate uptake. *Prog Neurobiol* 65(1): 1-105
- Daniel, H.; Levenes, C.; Crepel, F. (1998) Cellular mechanisms of cerebellar LTD. *Trends Neurosci* 21(9): 401-7
- De Vries, L.; Zheng, B.; Fischer, T.; Elenko, E.; Farquhar, M.G. (2000) The regulator of G protein signaling family. *Annual Review of Pharmacology and Toxicology* 40: 235-271
- Dell, E.J.; Blackmer, T.; Skiba, N.P.; Daaka, Y.; Shekter, L.R.; Rosal, R.; Reuveny, E.; Hamm, H.E. (2002) Defining G protein beta gamma specificity for effector recognition. *Methods Enzymol* 344: 421-34
- Dhingra, A.; Jiang, M.; Wang, T.L.; Lyubarsky, A.; Savchenko, A.; Bar-Yehuda, T.; Sterling, P.; Birnbaumer, L.; Vardi, N. (2002) Light response of retinal ON bipolar cells requires a specific splice variant of Galphao(o). *J Neurosci* 22(12): 4878-84
- Diverse-Pierluissi, M.A.; Fischer, T.; Jordan, J.D.; Schiff, M.; Ortiz, D.F.; Farquhar, M.G.; De Vries, L. (1999) Regulators of G protein signaling proteins as determinants of the rate of desensitization of presynaptic calcium channels. *Journal of Biological Chemistry* 274(20): 14490-14494
- Dolphin, A.C. (1998) Mechanisms of modulation of voltage-dependent calcium channels by G proteins. *J Physiol* 506 ( Pt 1): 3-11
- Druey, K.M.; Blumer, K.J.; Kang, V.H.; Kehrl, J.H. (1996) Inhibition of G protein-mediated MAP kinase activation by members of a novel mammalian gene family. *Journal of Investigative Medicine* 44(3): A271-A271
- Dumoulin, A.; Rostaing, P.; Bedet, C.; Levi, S.; Isambert, M.F.; Henry, J.P.; Triller, A.; Gasnier, B. (1999) Presence of the vesicular inhibitory amino acid transporter in GABAergic and glycinergic synaptic terminal boutons. *J Cell Sci* 112 ( Pt 6): 811-23
- Erickson, J.D.; Schafer, M.K.; Bonner, T.I.; Eiden, L.E.; Weihe, E. (1996) Distinct pharmacological properties and distribution in neurons and endocrine cells of two isoforms of the human vesicular monoamine transporter. *Proc Natl Acad Sci U S A* 93(10): 5166-71
- Erickson, J.D.; Varoqui, H.; Schafer, M.K.H.; Modi, W.; Diebler, M.F.; Weihe, E.; Rand, J.; Eiden, L.E.; Bonner, T.I.; Usdin, T.B. (1994) Functional Identification of A Vesicular Acetylcholine Transporter and Its Expression from A Cholinergic Gene Locus. *Journal of Biological Chemistry* 269(35): 21929-21932
- Exner, T.; Jensen, O.N.; Mann, M.; Kleuss, C.; Nurnberg, B. (1999) Posttranslational modification of Galphao1 generates Galphao3, an abundant G protein in brain. *Proc Natl Acad Sci U S A* 96(4): 1327-32
- Exton, J.H. (1996) Regulation of phosphoinositide phospholipases by hormones, neurotransmitters, and other agonists linked to G proteins. *Annu Rev Pharmacol Toxicol* 36: 481-509

- Fawcett, L.; Baxendale, R.; Stacey, P.; McGrouther, C.; Harrow, I.; Soderling, S.; Hetman, J.; Beavo, J.A.; Phillips, S.C. (2000) Molecular cloning and characterization of a distinct human phosphodiesterase gene family: PDE11A. *Proc.Natl.Acad.Sci.U.S.A* 97(7): 3702-3707
- Fisher, D.A.; Smith, J.F.; Pillar, J.S.; St Denis, S.H.; Cheng, J.B. (1998a) Isolation and characterization of PDE8A, a novel human cAMP-specific phosphodiesterase. *Biochem.Biophys.Res.Commun.* 246(3): 570-577
- Fisher, D.A.; Smith, J.F.; Pillar, J.S.; St Denis, S.H.; Cheng, J.B. (1998b) Isolation and characterization of PDE9A, a novel human cGMP-specific phosphodiesterase. *J.Biol.Chem.* 273(25): 15559-15564
- Floor, E.; Leventhal, P.S.; Wang, Y.L.; Meng, L.H.; Chen, W.Q. (1995) Dynamic Storage of Dopamine in Rat-Brain Synaptic Vesicles In-Vitro. *Journal of Neurochemistry* 64(2): 689-699
- Fon, E.A.; Pothos, E.N.; Sun, B.C.; Killeen, N.; Sulzer, D.; Edwards, R.H. (1997) Vesicular transport regulates monoamine storage and release but is not essential for amphetamine action. *Neuron* 19(6): 1271-1283
- Fremeau, R.T., Jr.; Burman, J.; Qureshi, T.; Tran, C.H.; Proctor, J.; Johnson, J.; Zhang, H.; Sulzer, D.; Copenhagen, D.R.; Storm-Mathisen, J.; Reimer, R.J.; Chaudhry, F.A.; Edwards, R.H. (2002) The identification of vesicular glutamate transporter 3 suggests novel modes of signaling by glutamate. *Proc Natl Acad Sci U S A* 99(22): 14488-93
- Fremeau, R.T., Jr.; Troyer, M.D.; Pahner, I.; Nygaard, G.O.; Tran, C.H.; Reimer, R.J.; Bellocchio, E.E.; Fortin, D.; Storm-Mathisen, J.; Edwards, R.H. (2001) The expression of vesicular glutamate transporters defines two classes of excitatory synapse. *Neuron* 31(2): 247-60
- Fujiyama, F.; Furuta, T.; Kaneko, T. (2001) Immunocytochemical localization of candidates for vesicular glutamate transporters in the rat cerebral cortex. *J Comp Neurol* 435(3): 379-87
- Garzon, M.; Pickel, V.M. (2000) Dendritic and axonal targeting of the vesicular acetylcholine transporter to membranous cytoplasmic organelles in laterodorsal and pedunculopontine tegmental nuclei. *Journal of Comparative Neurology* 419(1): 32-48
- Gold, S.J.; Ni, Y.G.; Dohlman, H.G.; Nestler, E.J. (1997) Regulators of G-protein signaling (RGS) proteins: Region-specific expression of nine subtypes in rat brain. *Journal of Neuroscience* 17(20): 8024-8037
- Gras, C.; Herzog, E.; Bellachi, G.C.; Bernard, V.; Ravassard, P.; Pohl, M.; Gasnier, B.; Giros, B.; El Mestikawy, S. (2002) A third vesicular glutamate transporter expressed by cholinergic and serotonergic neurons. *J Neurosci* 22(13): 5442-51
- Hansson, S.R.; Mezey, E.; Hoffman, B.J. (1998) Ontogeny of vesicular monoamine transporter mRNAs VMAT1 and VMAT2. II. Expression in neural crest derivatives and their target sites in the rat. *Brain Res Dev Brain Res* 110(1): 159-74
- Hayashi, M.; Haga, M.; Yatsushiro, S.; Yamamoto, A.; Moriyama, Y. (1999) Vesicular monoamine transporter 1 is responsible for storage of 5-hydroxytryptamine in rat pinealocytes. *J Neurochem* 73(6): 2538-45

- Hayashi, M.; Otsuka, M.; Morimoto, R.; Hirota, S.; Yatsushiro, S.; Takeda, J.; Yamamoto, A.; Moriyama, Y. (2001) Differentiation-associated Na<sup>+</sup>-dependent inorganic phosphate cotransporter (DNPI) is a vesicular glutamate transporter in endocrine glutamatergic systems. *J Biol Chem* 276(46): 43400-6
- He, W.; Cowan, C.W.; Wensel, T.G. (1998) RGS9, a GTPase accelerator for phototransduction. *Neuron* 20(1): 95-102
- Hell, J.W.; Jahn, R. (1998) Preparation of synaptic vesicles from mammalian brain. *Cell Biology: laboratory handbook*, 2nd edition: 102-110
- Hell, J.W.; Maycox, P.R.; Jahn, R. (1990) Energy dependence and functional reconstitution of the gamma-aminobutyric acid carrier from synaptic vesicles. *J Biol Chem* 265(4): 2111-7
- Hell, J.W.; Maycox, P.R.; Stadler, H.; Jahn, R. (1988) Uptake of GABA by rat brain synaptic vesicles isolated by a new procedure. *Embo J* 7(10): 3023-9
- Helms, J.B. (1995) Role of heterotrimeric GTP binding proteins in vesicular protein transport: indications for both classical and alternative G protein cycles. *FEBS Lett* 369(1): 84-8
- Hepler, J.R. (2003) RGS protein and G protein interactions: A little help from their friends. *Molecular Pharmacology* 64(3): 547-549
- Hepler, J.R.; Berman, D.M.; Gilman, A.G.; Kozasa, T. (1997) RGS4 and GAIP are GTPase-activating proteins for G(q alpha) and block activation of phospholipase C beta by gamma-thio-GTP-G(q alpha). *Proceedings of the National Academy of Sciences of the United States of America* 94(2): 428-432
- Hisano, S.; Hoshi, K.; Ikeda, Y.; Maruyama, D.; Kanemoto, M.; Ichijo, H.; Kojima, I.; Takeda, J.; Nogami, H. (2000) Regional expression of a gene encoding a neuron-specific Na<sup>(+)</sup>-dependent inorganic phosphate cotransporter (DNPI) in the rat forebrain. *Brain Res Mol Brain Res* 83(1-2): 34-43
- Hollinger, S.; Hepler, J.R. (2002) Cellular regulation of RGS proteins: Modulators and integrators of G protein signaling. *Pharmacological Reviews* 54(3): 527-559
- Höltje, M.; Pahner, I.; Lautenschlager, M.; Hortnagl, H.; Nurnberg, B.; Jahn, R.; Ahnert-Hilger, G. (2000) The neuronal monoamine transporter VMAT2 is regulated by the trimeric GTPase Go(2). *J.Neurosci.* 20(6): 2131-2141
- Höltje, M.; Winter, S.; Walther, D.; Pahner, I.; Hortnagl, H.; Ottersen, O.P.; Bader, M.; Ahnert-Hilger, G. (2003) The vesicular monoamine content regulates VMAT2 activity through Galphaq in mouse platelets. Evidence for autoregulation of vesicular transmitter uptake. *J.Biol.Chem.* 278(18): 15850-15858
- Jiang, M.; Gold, M.S.; Boulay, G.; Spicher, K.; Peyton, M.; Brabet, P.; Srinivasan, Y.; Rudolph, U.; Ellison, G.; Birnbaumer, L. (1998) Multiple neurological abnormalities in mice deficient in the G protein Go. *Proc Natl Acad Sci U S A* 95(6): 3269-74
- Jiang, M.; Spicher, K.; Boulay, G.; Wang, Y.; Birnbaumer, L. (2001) Most central nervous system D2 dopamine receptors are coupled to their effectors by Go. *Proc Natl Acad Sci U S A* 98(6): 3577-82
- Johnson, R.G., Jr. (1988) Accumulation of biological amines into chromaffin granules: a model for hormone and neurotransmitter transport. *Physiol Rev* 68(1): 232-307

- Kaneko, T.; Fujiyama, F. (2002) Complementary distribution of vesicular glutamate transporters in the central nervous system. *Neuroscience Research* 42(4): 243-250
- Kleuss, C.; Hescheler, J.; Ewel, C.; Rosenthal, W.; Schultz, G.; Wittig, B. (1991) Assignment of G-protein subtypes to specific receptors inducing inhibition of calcium currents. *Nature* 353(6339): 43-8
- Laemmli, U.K. (1970) Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 227(259): 680-5
- Lesch, K.P.; Gross, J.; Wolozin, B.L.; Murphy, D.L.; Riederer, P. (1993) Extensive sequence divergence between the human and rat brain vesicular monoamine transporter: possible molecular basis for species differences in the susceptibility to MPP+. *J Neural Transm Gen Sect* 93(1): 75-82
- Liu, G.; Choi, S.; Tsien, R.W. (1999) Variability of neurotransmitter concentration and nonsaturation of postsynaptic AMPA receptors at synapses in hippocampal cultures and slices. *Neuron* 22(2): 395-409
- Liu, Y.; Peter, D.; Roghani, A.; Schuldiner, S.; Prive, G.G.; Eisenberg, D.; Brecha, N.; Edwards, R.H. (1992) A cDNA that suppresses MPP+ toxicity encodes a vesicular amine transporter. *Cell* 70(4): 539-51
- Liu, Y.; Schweitzer, E.S.; Nirenberg, M.J.; Pickel, V.M.; Evans, C.J.; Edwards, R.H. (1994) Preferential localization of a vesicular monoamine transporter to dense core vesicles in PC12 cells. *J Cell Biol* 127(5): 1419-33
- Man-Son-Hing, H.J.; Codina, J.; Abramowitz, J.; Haydon, P.G. (1992) Microinjection of the alpha-subunit of the G protein Go2, but not Go1, reduces a voltage-sensitive calcium current. *Cell Signal* 4(4): 429-41
- Maycox, P.R.; Hell, J.W.; Jahn, R. (1990) Amino-Acid Neurotransmission - Spotlight on Synaptic Vesicles. *Trends in Neurosciences* 13(3): 83-87
- McIntire, S.L.; Reimer, R.J.; Schuske, K.; Edwards, R.H.; Jorgensen, E.M. (1997) Identification and characterization of the vesicular GABA transporter. *Nature* 389(6653): 870-6
- Montmayeur, J.P.; Borrelli, E. (1994) Targeting of G alpha i2 to the Golgi by alternative spliced carboxyl-terminal region. *Science* 263(5143): 95-8
- Morgenstern, E. (1995) The formation of compound granules from different types of secretory organelles in human platelets (dense granules and alpha-granules). A cryofixation/-substitution study using serial sections. *Eur J Cell Biol* 68(2): 183-90
- Mosharov, E.V.; Gong, L.W.; Khanna, B.; Sulzer, D.; Lindau, M. (2003) Intracellular patch electrochemistry: Regulation of cytosolic catecholamines in chromaffin cells. *Journal of Neuroscience* 23(13): 5835-5845
- Muller, S.; Lohse, M.J. (1995) The role of G-protein beta gamma subunits in signal transduction. *Biochem Soc Trans* 23(1): 141-8
- Nakanishi, N.; Onozawa, S.; Matsumoto, R.; Hasegawa, H.; Yamada, S. (1995a) Cyclic AMP-dependent modulation of vesicular monoamine transport in pheochromocytoma cells. *J Neurochem* 64(2): 600-7

- Nakanishi, N.; Onozawa, S.; Matsumoto, R.; Kurihara, K.; Ueha, T.; Hasegawa, H.; Minami, N. (1995b) Effects of protein kinase inhibitors and protein phosphatase inhibitors on cyclic AMP-dependent down-regulation of vesicular monoamine transport in pheochromocytoma PC12 cells. *FEBS Lett.* 368(3): 411-4
- Ni, B.; Rosteck, P.R., Jr.; Nadi, N.S.; Paul, S.M. (1994) Cloning and expression of a cDNA encoding a brain-specific Na(+)-dependent inorganic phosphate cotransporter. *Proc Natl Acad Sci U S A* 91(12): 5607-11
- Nicholls, D.G. (1994) Proteins, Transmitter and Synapses. Blackwell Science Ltd, Oxford, UK
- Nurnberg, B.; Ahnert-Hilger, G. (1996) Potential roles of heterotrimeric G proteins of the endomembrane system. *FEBS Lett.* 389(1): 61-65
- Offermanns, S.; Hashimoto, K.; Watanabe, M.; Sun, W.; Kurihara, H.; Thompson, R.F.; Inoue, Y.; Kano, M.; Simon, M.I. (1997a) Impaired motor coordination and persistent multiple climbing fiber innervation of cerebellar Purkinje cells in mice lacking Galphaq. *Proc Natl Acad Sci U S A* 94(25): 14089-94
- Offermanns, S.; Toombs, C.F.; Hu, Y.H.; Simon, M.I. (1997b) Defective platelet activation in G alpha(q)-deficient mice. *Nature* 389(6647): 183-6
- Pahner, I.; Holtje, M.; Winter, S.; Nurnberg, B.; Ottersen, O.P.; Ahnert-Hilger, G. (2002) Subunit composition and functional properties of G-protein heterotrimers on rat chromaffin granules. *Eur J Cell Biol.* 81(8): 449-456
- Palfreyman, M.N.; Souness, J.E. (1996) Phosphodiesterase type IV inhibitors. *Prog Med Chem.* 33: 1-52
- Peter, D.; Jimenez, J.; Liu, Y.; Kim, J.; Edwards, R.H. (1994) The chromaffin granule and synaptic vesicle amine transporters differ in substrate recognition and sensitivity to inhibitors. *J Biol Chem* 269(10): 7231-7
- Peter, D.; Liu, Y.; Sternini, C.; de Giorgio, R.; Brecha, N.; Edwards, R.H. (1995) Differential expression of two vesicular monoamine transporters. *J Neurosci* 15(9): 6179-88
- Reimer, R.J.; Fon, E.A.; Edwards, R.H. (1998) Vesicular neurotransmitter transport and the presynaptic regulation of quantal size. *Curr Opin Neurobiol* 8(3): 405-12
- Reimer, R.J.; Fremeau, R.T., Jr.; Bellocchio, E.E.; Edwards, R.H. (2001) The essence of excitation. *Curr Opin Cell Biol* 13(4): 417-21
- Ross, E.M.; Wilkie, T.M. (2000) GTPase-activating proteins for heterotrimeric G proteins: Regulators of G protein signaling (RGS) and RGS-like proteins. *Annual Review of Biochemistry* 69: 795-827
- Rudolph, U.; Brabet, P.; Hasty, P.; Bradley, A.; Birnbaumer, L. (1993) Disruption of the G(i2) alpha locus in embryonic stem cells and mice: a modified hit and run strategy with detection by a PCR dependent on gap repair. *Transgenic Res.* 2(6): 345-355
- Saeed, S.A.; Rasheed, H.; Fecto, F.A.W.; Achakzai, M.I.; Ali, R.; Connor, J.D.; Gilani, A.U.H. (2004) Signaling mechanisms mediated by G-protein coupled receptors in human platelets. *Acta Pharmacologica Sinica* 25(7): 887-892

- Sagne, C.; El Mestikawy, S.; Isambert, M.F.; Hamon, M.; Henry, J.P.; Giros, B.; Gasnier, B. (1997) Cloning of a functional vesicular GABA and glycine transporter by screening of genome databases. *FEBS Lett* 417(2): 177-83
- Sakata-Haga, H.; Kanemoto, M.; Maruyama, D.; Hoshi, K.; Mogi, K.; Narita, M.; Okado, N.; Ikeda, Y.; Nogami, H.; Fukui, Y.; Kojima, I.; Takeda, J.; Hisano, S. (2001) Differential localization and colocalization of two neuron-types of sodium-dependent inorganic phosphate cotransporters in rat forebrain. *Brain Res* 902(2): 143-55
- Schmidt, M.; Evellin, S.; Weernink, P.A.; von Dorp, F.; Rehmann, H.; Lomasney, J.W.; Jakobs, K.H. (2001) A new phospholipase-C-calcium signalling pathway mediated by cyclic AMP and a Rap GTPase. *Nat Cell Biol* 3(11): 1020-4
- Schuldiner, S.; Shirvan, A.; Linial, M. (1995) Vesicular neurotransmitter transporters: from bacteria to humans. *Physiol Rev* 75(2): 369-92
- Shen, X.M.; Xia, B.; Wrona, M.Z.; Dryhurst, G. (1996) Synthesis, redox properties, in vivo formation, and neurobehavioral effects of N-acetylcysteinyl conjugates of dopamine: Possible metabolites of relevance to Parkinson's disease. *Chemical Research in Toxicology* 9(7): 1117-1126
- Smith, P.K.; Krohn, R.I.; Hermanson, G.T.; Mallia, A.K.; Gartner, F.H.; Provenzano, M.D.; Fujimoto, E.K.; Goeke, N.M.; Olson, B.J.; Klenk, D.C. (1985) Measurement of protein using bicinchoninic acid. *Anal Biochem* 150(1): 76-85
- Soderling, S.H.; Bayuga, S.J.; Beavo, J.A. (1998) Identification and characterization of a novel family of cyclic nucleotide phosphodiesterases. *J.Biol.Chem.* 273(25): 15553-15558
- Soderling, S.H.; Bayuga, S.J.; Beavo, J.A. (1999) Isolation and characterization of a dual-substrate phosphodiesterase gene family: PDE10A. *Proc.Natl.Acad.Sci.U.S.A* 96(12): 7071-7076
- Sondek, J.; Siderovski, D.P. (2001) Ggamma-like (GGL) domains: new frontiers in G-protein signaling and beta-propeller scaffolding. *Biochem Pharmacol* 61(11): 1329-37
- Speidel, D.; Bruederle, C.E.; Enk, C.; Voets, T.; Varoqueaux, F.; Reim, K.; Becherer, U.; Fornal, F.; Ruggieri, S.; Holighaus, Y.; Weihe, E.; Bruns, D.; Brose, N.; Rettig, J. (2005) CAPS1 regulates catecholamine loading of large dense-core vesicles. *Neuron* 46(1): 75-88
- Spencer, J.P.E.; Whiteman, M.; Jenner, P.; Halliwell, B. (2002) 5-S-cysteinyl-conjugates of catecholamines induce cell damage, extensive DNA base modification and increases in caspase-3 activity in neurons. *Journal of Neurochemistry* 81(1): 122-129
- Su, C.Y.; Luo, D.G.; Terakita, A.; Shichida, Y.; Liao, H.W.; Kazmi, M.A.; Sakmar, T.P.; Yau, K.W. (2006) Parietal-eye phototransduction components and their potential evolutionary implications. *Science* 311(5767): 1617-1621
- Sulzer, D.; Edwards, R. (2000) Vesicles: equal in neurotransmitter concentration but not in volume. *Neuron* 28(1): 5-7
- Takahashi, N.; Miner, L.L.; Sora, I.; Ujike, H.; Revay, R.S.; Kostic, V.; JacksonLewis, V.; Przedborski, S.; Uhl, G.R. (1997) VMAT2 knockout mice: Heterozygotes display reduced amphetamine-conditioned reward, enhanced amphetamine locomotion, and enhanced MPTP toxicity. *Proceedings of the National Academy of Sciences of the United States of America* 94(18): 9938-9943

- Takamori, S.; Malherbe, P.; Broger, C.; Jahn, R. (2002) Molecular cloning and functional characterization of human vesicular glutamate transporter 3. *EMBO Rep* 3(8): 798-803
- Takamori, S.; Rhee, J.S.; Rosenmund, C.; Jahn, R. (2000a) Identification of a vesicular glutamate transporter that defines a glutamatergic phenotype in neurons. *Nature* 407(6801): 189-194
- Takamori, S.; Rhee, J.S.; Rosenmund, C.; Jahn, R. (2001) Identification of differentiation-associated brain-specific phosphate transporter as a second vesicular glutamate transporter (VGLUT2). *J Neurosci* 21(22): RC182-
- Takamori, S.; Riedel, D.; Jahn, R. (2000b) Immunoisolation of GABA-specific synaptic vesicles defines a functionally distinct subset of synaptic vesicles. *J Neurosci* 20(13): 4904-11
- Tang, M.; Wang, G.; Lu, P.; Karas, R.H.; Aronovitz, M.; Heximer, S.P.; Kaltenbronn, K.M.; Blumer, K.J.; Siderovski, D.P.; Zhu, Y.; Mendelsohn, M.E. (2003) Regulator of G-protein signaling-2 mediates vascular smooth muscle relaxation and blood pressure. *Nature Medicine* 9(12): 1506-1512
- Thomas, E.A.; Danielson, P.E.; Sutcliffe, J.G. (1998) RGS9: A regulator of G-protein signalling with specific expression in rat and mouse striatum. *Journal of Neuroscience Research* 52(1): 118-124
- Traver, S.; Bidot, C.; Spassky, N.; Baltauss, T.; De Tand, M.F.; Thomas, J.L.; Zalc, B.; Janoueix-Lerosey, I.; Gunzburg, J.D. (2000) RGS14 is a novel Rap effector that preferentially regulates the GTPase activity of galphao. *Biochem J* 350 Pt 1: 19-29
- Travis, E.R.; Wang, Y.M.; Michael, D.J.; Caron, M.G.; Wightman, R.M. (2000) Differential quantal release of histamine and 5-hydroxytryptamine from mast cells of vesicular monoamine transporter 2 knockout mice. *Proc Natl Acad Sci U S A* 97(1): 162-7
- Usdin, T.B.; Eiden, L.E.; Bonner, T.I.; Erickson, J.D. (1995) Molecular-Biology of the Vesicular Ach Transporter. *Trends in Neurosciences* 18(5): 218-224
- Van der Kloot, W. (1991) The regulation of quantal size. *Prog Neurobiol* 36(2): 93-130
- Varoqui, H.; Diebler, M.F.; Meunier, F.M.; Rand, J.B.; Usdin, T.B.; Bonner, T.I.; Eiden, L.E.; Erickson, J.D. (1994) Cloning and Expression of the Vesamicol Binding-Protein from the Marine Ray Torpedo - Homology with the Putative Vesicular Acetylcholine Transporter Unc-17 from *Caenorhabditis-Elegans*. *Febs Letters* 342(1): 97-102
- Varoqui, H.; Erickson, J.D. (1996) Active transport of acetylcholine by the human vesicular acetylcholine transporter. *J Biol Chem* 271(44): 27229-32
- Wang, Y.M.; Gainetdinov, R.R.; Fumagalli, F.; Xu, F.; Jones, S.R.; Bock, C.B.; Miller, G.W.; Wightman, R.M.; Caron, M.G. (1997) Knockout of the vesicular monoamine transporter 2 gene results in neonatal death and supersensitivity to cocaine and amphetamine. *Neuron* 19(6): 1285-1296
- Watts, V.J.; Wiens, B.L.; Cumbay, M.G.; Vu, M.N.; Neve, R.L.; Neve, K.A. (1998) Selective activation of Galphao by D2L dopamine receptors in NS20Y neuroblastoma cells. *J Neurosci* 18(21): 8692-9

- Weihe, E.; Tao-Cheng, J.H.; Schafer, M.K.; Erickson, J.D.; Eiden, L.E. (1996) Visualization of the vesicular acetylcholine transporter in cholinergic nerve terminals and its targeting to a specific population of small synaptic vesicles. Proc Natl Acad Sci U S A 93(8): 3547-52
- Weller, U.; Muller, L.; Messner, M.; Palmer, M.; Valeva, A.; Tranum-Jensen, J.; Agrawal, P.; Biermann, C.; Dobereiner, A.; Kehoe, M.A.; Bhakdi, S. (1996) Expression of active streptolysin O in Escherichia coli as a maltose-binding-protein--streptolysin-O fusion protein. The N-terminal 70 amino acids are not required for hemolytic activity. Eur J Biochem 236(1): 34-9
- Wettschureck, N.; Offermanns, S. (2005) Mammalian G proteins and their cell type specific functions. Physiol Rev. 85(4): 1159-1204
- Williams, J. (1997) How does a vesicle know it is full? Neuron 18(5): 683-6
- Winter, S.; Brunk, I.; Walther, D.J.; Höltje, M.; Jiang, M.; Peter, J.U.; Takamori, S.; Jahn, R.; Birnbaumer L.; Ahnert-Hilger G. (2005) Galphao<sub>2</sub> regulates vesicular glutamate transporter activity by changing its chloride dependence. J Neurosci 25(18): 4672-80
- Wolosker, H.; de Souza, D.O.; de Meis, L. (1996) Regulation of glutamate transport into synaptic vesicles by chloride and proton gradient. J Biol Chem 271(20): 11726-31
- Wurzinger, L.J. (1990) Histophysiology of the circulating platelet. Adv Anat Embryol Cell Biol 120: 1-96
- Yan, Y.B.; Chi, P.P.; Bourne, H.R. (1997) RGS4 inhibits G(q)-mediated activation of mitogen-activated protein kinase and phosphoinositide synthesis. Journal of Biological Chemistry 272(18): 11924-11927