

## 7. Literaturverzeichnis

### 7.1. Veröffentlichungen

- Allen, B.W. (1998): Mycobacteria. General Culture Methodology and Safety Considerations. *Methods Mol. Biol.*, 101, 15-30
- Alroy, I., Soussan, L., Seger, R. and Yarden, Y. (1999): Neu Differentiation Factor Stimulates Phosphorylation and Activation of the Sp1 Transcription Factor. *Mol. Cell. Biol.*, 19 (3), 1961-72
- Andersson, L.C., Jokinen, M. and Gahmberg, C.G. (1979): Induction of erythroid differentiation in the human leukaemia cell line K562. *Nature*, 278 (5702), 364-5
- Aplan, P.D., Jones, C.A., Chervinsky, D.S., Zhao, X., Ellsworth, M., Wu, C., McGuire, E.A. and Gross, K.W. (1997): An *scl* gene product lacking the transactivation domain induces bony abnormalities and co-operates with LMO1 to generate T-cell malignancies in transgenic mice. *EMBO J.*, 16 (9), 2408-19
- Aplan, P.D., Nakahara, K., Orkin, S.H. and Kirsch, I.R. (1992): The *SCL* gene product: a positive regulator of erythroid differentiation. *EMBO J.*, 11 (11), 4073-81
- Archer, V.E.V., Breton, J., Sanchez-Garcia, I., Osada, H., Forster, A., Thomson, A.J. and Rabbitts, T.H. (1994): Cystein-rich LIM domains of LIM-homeodomain and LIM-only proteins contain zinc but not iron. *Proc. Natl. Acad. Sci. USA*, 91, 316-20
- Ayoubi, T.A.Y. and Van de Ven, W.J.M. (1996): Regulation of gene expression by alternative promoters. *FASEB J.*, 10, 453-60
- Barr, F.G. (1998): Translocations, cancer and the puzzle of specificity. *Nature Genet.*, 19, 121-4
- Begley, C.G. and Green, A.R. (1999): The *SCL* gene: from case report to critical hematopoietic regulator. *Blood*, 93 (9), 2760-70
- Bielinska, A., Shivdasani, R.A., Zhang, L. and Nabel, G.J. (1990): Regulation of gene expression with double-stranded phosphorothioate oligonucleotides. *Science*, 250, 997-1000

- Bijsterbosch, M.K., Manoharan, M., Rump, E.T., De Vruet, R.L.C., van Veghel, R., Tivel, K.L., Biessen, E.A.L., Frank Bennet, C., Dan Cook, P. and van Berkel, T.J.C. (1997): In vivo fate of phosphorothioate antisense oligodeoxynucleotides: predominant uptake by scavenger receptors on endothelial liver cells. *Nucleic Acids Res.* 25 (16), 3290-6
- Birnboim, H.C. and Doly, J. (1979): A rapid alkaline lysis procedure for screening recombinant plasmid DNA. *Nucleic Acids Res.*, 7, 1513-22
- Bishop, J.M. (1991): Molecular Themes in Oncogenesis. *Cell*, 64, 235-48
- Boehm, T., Foroni, L., Kaneko, Y., Perutz, M.F. and Rabbitts, T.H. (1991): The rhombotin family of cystein-rich LIM-domain oncogenes: Distinct members are involved in T-cell translocations to human chromosome 11p15 and 11p13. *Proc. Natl. Acad. Sci. USA*, 88, 4367-71
- Boehm, T., Mengle-Gaw, L., Kees, U.R., Spurr, N., Lavenir, I., Forster, A. and Rabbitts, T.H. (1989): Alternating purine - pyrimidine tracts may promote chromosomal translocations seen in a variety of human lymphoid tumours. *EMBO J.*, 8 (9), 2621-31
- Boisclair, Y.R., Brown, A.L., Casola, S. and Bechler, M.M. (1993): Three Clustered Sp1 Sites Are Required for Efficient Transcription of the TATA-less Promoter of the Gene for Insulin-like Growth Factor-binding Protein 2 from the Rat. *J. Biol. Chem.*, 268 (33), 24892-901
- Bonifer, C., Jäggle, U. and Huber, M.C. (1997): The chicken lysozyme locus as a paradigm for the complex developmental regulation of eukaryotic gene loci. *J. Biol. Chem.*, 272, 26075-8
- Boyes, J., Byfield, P., Nakatani, Y. and Ogryzko, V. (1998b): Regulation of activity of the transcription factor GATA-1 by acetylation. *Nature*, 396, 594-8
- Boyes, J., Omichinski, J., Clark, D., Pikaart, M. and Felsenfeld, G. (1998a): Perturbation of Nucleosome Structure by the Erythroid Transcription Factor GATA-1. *J. Mol. Biol.*, 279, 529-44
- Briggs, M.R., Kadonaga, J.T., Bell, S.P. and Tjian, R. (1986): Purification and Biochemical Characterization Of the Promotor-Specific Transcription Factor, Sp1. *Science*, 234, 47-52
- Brown, P.C. and Silverman, J.A. (1996): Characterization of the rat mdr2 promoter and its regulation by the transcription factor Sp1. *Nucleic Acids Res.*, 24 (16), 3235-41

- Cahill, D.P., Kinzler, K.W., Vogelstein, B. and Lengauer, C. (1999): Genetic instability and darwinian selection in tumours. *Trends Gen.*, 15 (12), M57-M60 (Millenium issue)
- Calligaris, R., Bottardi, S., Cogoi, S., Apezteguia, I. and Santoro, C. (1995): Alternative translation initiation site usage results in two functionally distinct forms of the GATA-1 transcription factor. *Proc. Natl. Acad. Sci. USA*, 92, 11598-602
- Caput, D., Beutler, B., Hartog, K., Thayer, R., Brown-Shimer, S. and Cerami, A. (1986): Identification of a common nucleotide sequence in the 3' untranslated region of mRNA molecules specifying inflammatory mediators. *Proc. Natl. Acad. Sci. USA*, 83, 1670-4
- Carninci, P. et al. (1998): Thermostabilization and thermoactivation of thermolabile enzymes by trehalose and its application for the synthesis of full length cDNA. *Proc. Natl. Acad. Sci USA*, 95, 520-25
- Chen, C.Y.A. and Shyu, A.B. (1995): AU-rich elements: characterization and importance in mRNA degradation. *Trends Biochem.*, 20, 465-70
- Chen, X., Wright, K.L., Berkowitz, E.A., Azizkhan, J.C., Ting, J.P.Y. and Lee, D.C. (1994): Protein interactions at Sp1-like sites in the TGF $\alpha$  promoter as visualized by *in vivo* genomic footprinting. *Oncogene*, 9, 3179-87
- Chirgwin, J.M., Przybyla, A.E., MacDonald, R.J. and Rutter, W.J. (1979): Isolation of Biologically Active Ribonucleic Acid from Sources Enriched in Ribonuclease. *Biochemistry*, 18 (24), 5294-9
- Chu, B.C.F. and Orgel, L.E. (1992): The stability of different forms of double-stranded decoy DNA in serum and nuclear extracts. *Nucleic Acids Res.*, 20 (21), 5857-8
- Cohen, B., McGuffin, E., Pfeifle, C., Segal, D. and Cohen, S.M. (1992): *apterous*, a gene required for imaginal disc development in *Drosophila* encodes a member of the LIM family of developmental regulatory proteins. *Genes Dev.*, 6, 715-29
- Dawid, I.B. (1998): LIM protein interactions: *Drosophila* enters the stage. *Trends Genet.*, 14 (12), 480-2
- Dawid, I.B., Breen, J.J. and Toyama, R. (1998): LIM domains: multiple roles as adapters and functional modifiers in protein interaction. *Trends Genet.*, 14 (4), 156-62

- De Maria, R., Zeuner, A., Eramo, A., Domenichelli, C., Bonci, D., Grignani, F., Srinivasula, S.M., Alnemri, E.S., Testa, U. and Peschle, C. (1999): Negative regulation of erythropoiesis by caspase-mediated cleavage of GATA-1. *Nature*, 401, 489-93
- Dignam, J.D., Lebovitz, R.M. and Roeder, R.G. (1983): Accurate transcription initiation by RNA polymerase II in a soluble extract from isolated mammalian nuclei. *Nucleic Acids Res.*, 11 (5), 1475-89
- Dong, W.F., Billia, H.L., Atkins, H.L., Iscove, N.N. and Minden, M.D. (1996): Expression of rhombotin 2 in normal and leukaemic haematopoietic cells. *Br. J. Haematol.*, 93, 280-6
- Drechsler, M., Schumacher, V., Friedrich, S., Wildhardt, G, Giesler, S., Schroth, A., Bodem, J. and Royer-Pokora, B. (1999): Genomic structure, alternative transcripts and chromosome location of the human LIM domain binding protein 1 gene LDB1. *Cytogenet. Cell Genet.*, 87, 119-24
- Dusing, M.R. and Wiginton, D.A. (1994): Sp1 is essential for both enhancer-mediated and basal activation of the TATA-less human adenosine deaminase promoter. *Nucleic Acids Res.*, 22 (4), 669-77
- Dynan, W.S. and Tjian, R. (1983): Isolation of Transcription Factors That Discriminate between Different Promoters Recognized by RNA Polymerase II. *Cell*, 32, 669-80
- Dynan, W.S. and Tjian, R. (1985): Control of eukaryotic messenger RNA synthesis by sequence-specific DNA-binding proteins. *Nature*, 316, 774-8.
- Eckstein, F. (1979): Phosphorothioate analogues of nucleotides. *Accounts of Chem. Res.*, 12, 204-10
- Engel, I. and Murre, C. (1999): Transcription factors in hematopoiesis. *Curr. Opin. Gen. Dev.*, 9, 575-9
- Enver, T., Brewer, A.C. and Patient, R.K. (1985): Simian virus 40-mediated *cis* induction of the *Xenopus*  $\beta$ -globin DNaseI hypersensitive site. *Nature*, 318, 680-3
- Feinberg, A.P. and Vogelstein, B. (1983): A Technique for Radiolabelling DNA Restriction Endonuclease Fragments to High Specific Activity. *Anal. Biochem.*, 132, 6-13.

- Fisch, P., Boehm, T., Lavenir, I., Larson, T., Arno, J., Forster, A. and Rabbitts, T.H. (1992): T-cell acute lymphoblastic lymphoma induced in transgenic mice by the *RBTN1* and *RBTN2* LIM-domain genes. *Oncogene*, 7, 2389-97
- Foroni, L., Boehm, T., White, L., Forster, A., Sherrington, P., Liao, X.B., Brannan, C.I., Jenkins, N.A., Copeland, N.G. and Rabbitts, T.H. (1992): The Rhombotin Gene Family Encode Related LIM-domain Proteins whose Differing Expression Suggests Multiple Roles in Mouse Development. *J. Mol. Biol.*, 226, 747-61
- Fox, A.H., Liew, C., Holmes, M., Kowalski, K., Mackay, J. and Crossley, M. (1999): Transcriptional cofactors of the FOG family interact with GATA proteins by means of multiple zinc fingers. *EMBO J.*, 18 (10), 2812-22
- Fried, M. and Crothers, D.M. (1981): Equilibria and kinetics of lac repressor-operator interaction by polyacrylamide gel electrophoresis. *Nucleic Acids Res.*, 9 (23), 6505-25
- Gilar, M., Belenky, A., Smisek, D.L., Bourque, A. and Cohen, A.S. (1997): Kinetics of phosphorothioate oligonucleotide metabolism in biological fluids. *Nucleic Acids Res.*, 25 (18), 3615-20
- Graham, F.L. and Van der Eb, A.J. (1973): A New Technique for the Assay of Infectivity of Human Adenovirus 5 DNA. *Virology*, 52, 456-67
- Gregory, R.C., Taxman, D.J., Shesasayee, D., Kensinger, M.H., Biecker, J.J. and Wojchowski, D.M. (1996): Functional interaction of GATA1 with erythroid Kruppel-like factor and Sp1 at defined erythroid promoters. *Blood*, 87 (5), 1793-801
- Grosveld, F., Blom van Assendelft, G., Greaves, D.R. and Kollias, G. (1987): Position-Independent, High-Level Expression of the Human  $\beta$ -Globin Gene in Transgenic Mice. *Cell*, 51, 975-85
- Grutz, G., Forster, A. and Rabbitts, T.H. (1998): Identification of the *LMO4* gene encoding an interacting partner of the LIM-binding protein LDB1/NLI1: a candidate for displacement by LMO proteins in T cell acute leukaemia. *Oncogene*, 17, 2799-803
- Grütz, G.G., Bucher, K., Lavenir, I., Larson, T., Larson, R. and Rabbitts, T.H. (1998): The oncogenic T cell LIM-protein Lmo2 forms part of a DNA-binding complex specifically in immature T cells. *EMBO J.*, 17 (16), 4594-605

- Guvakova, M.A., Yakubov, L.A., Vlodavsky, I., Tonkinson, J.L. and Stein, C.A. (1995): Phosphorothioate oligodeoxynucleotides bind to basic fibroblast growth factor, inhibit its binding to cell surface receptors, and remove it from low affinity binding sites on extracellular matrix. *J. Biol. Chem.*, 270 (6), 2620-7
- Harber, D. and Harlow, E. (1997): Tumour-suppressor genes: evolving definitions in the genomic age. *Nature Genet.*, 16, 320-2
- Heerema, N.A. et al. (1998): Frequency and Clinical Significance of Cytogenetic Abnormalities in Pediatric T-Lineage Acute Lymphoblastic Leukemia: A Report From the Children's Cancer Group. *J. Clin. Oncol.*, 16 (4), 1270-78
- Heim, S. and Mitelman, F. (1995): Nonrandom Chromosome Abnormalities in Cancer -An Overview. In *Cancer Cytogenetics*, 19-32, Wiley-Liss.
- Henninghausen, L. and Lubon, H. (1987): Interaction of Protein with DNA in vitro. *Methods Enzymol.*, 152, 721-35
- Hentze, M.W. (1995): Translational regulation: versatile mechanisms for metabolic and developmental control. *Curr. Opin. Cell Biol.*, 7, 393-8
- Hinks, G.L., Shah, B., French, S.L., Campos, L.S., Staley, K., Hughes, J. and Sofroniew, M.V. (1997): Expression of LIM Protein Genes *Lmo1*, *Lmo2*, and *Lmo3* in Adult Mouse Hippocampus and Other Forebrain Regions: Differential Regulation by Seizure Activity. *J. Neuroscience*, 17 (14), 5549-59
- Hiom, K., Melek, M. and Gellert, M. (1998): DNA Transposition by the RAG1 and RAG2 Proteins: A Possible Source of Oncogenic Translocations. *Cell*, 94, 463-70
- Hofmann, W., Royer, H.D., Drechsler, M., Schneider, S. and Royer-Pokora, B. (1993): Characterization of the transcriptional regulatory region of the human *WT1* gene. *Oncogene*, 8, 3123-32
- Hsu, H.L., Cheng, J.T., Chen, Q. and Baer, R. (1991): Enhancer-binding activity of the tal-1 oncoprotein in association with the E47/E12 helix-loop-helix proteins. *Mol. Cell. Biol.*, 11 (6), 3037-42
- Irvine, K.D. and Vogt, T.F. (1997): Dorsal-ventral signalling in limb development. *Curr. Opin. Cell Biol.*, 9, 867-76

- Jiménez, G., Griffith, S.D., Ford, A.M., Greaves, M.F. and Enver, T. (1992): Activation of the  $\beta$ -globin locus control region precedes commitment to the erythroid lineage. *Proc. Natl. Acad. Sci. USA*, 89, 10618-22
- Jurata, L.W., Kenny, D.A. and Gill, G.N. (1996): Nuclear LIM interactor, a rhombotin and LIM homeodomain interacting protein, is expressed early in neuronal development. *Proc. Natl. Acad. Sci. USA*, 93, 11693-8
- Kao, W.Y., Briggs, J.A., Kinney, M.C., Jensen, R.A. and Briggs, R.C. (1997): Structure and Function Analysis of the Human Myeloid Cell Nuclear Differentiation Antigen Promotor: Evidence for the Role of Sp1 and Not of c-Myb or PU.1 in Myelomonocytic Lineage-Specific Expression. *J. Cell. Biochem.*, 65, 231-44
- Kenny, D.A., Jurata, L.W., Saga, Y. and Gill, G.N. (1998): Identification and characterization of *LMO4*, an LMO gene with a novel pattern of expression during embryogenesis. *Proc. Natl. Acad. Sci. USA*, 95, 11257-62
- Kollmar, R., Sukow, K.A., Sponagle, S.K. and Farnham, P.J. (1994): Start Site Selection at the TATA-less Carbamoyl-phosphate Synthase (Glutamine-hydrolyzing)/Aspartate Carbamoyltransferase/Dihydroorotase Promoter. *J. Biol. Chem.*, 269 (3), 2252-7
- Kornberg, R.D. (1999): Eukaryotic transcriptional control. *Trends Gen.*, 15 (12), M46-M49 (Millenium issue)
- Körner, K., Wolfrain, L.A., Lucibello, F.C. and Müller, R. (1997): Characterization of the TATA-less core promoter of the cell cycle-regulated *cdc25C* gene. *Nucleic Acids Res.*, 25 (24), 4933-9
- Korsmeyer, S.J. (1992): Chromosomal Translocations in Lymphoid Malignancies Reveal Novel Proto-oncogenes. *Annu. Rev. Immunol.*, 10, 785-807
- Labbaye, C., Valtieri, M., Barberi, T., Meccia, E., Masella, B., Pelosi, E, Gondorelli, G.L., Testa, U. and Peschle, C. (1995): Differential expression and functional role of GATA-2, NF-E2, and GATA-1 in normal adult hematopoiesis. *J. Clin. Invest.*, 95, 2346-58
- Lane, D., Prentki, P. and Chandler, M. (1992): Use of Gel Retardation To Analyze Protein-Nucleic Acid Interaction. *Microbiol. Rev.*, 56 (4), 509-28
- Larson, A.E. and Papayannopoulou, T. (1985): HEL and K562 Cells: Analysis of Proteins by Two-dimensional Gel Electrophoresis and Comparison to Erythroid and Nonerythroid Cells. *Exp. Hematol.*, 13, 129-35

- Larson, R.C., Fisch, P., Larson, T.A., Lavenir, I., Langford, T., King, G. and Rabbitts, T.H. (1994): T cell tumours of disparate phenotype in mice transgenic for *Rbtn-2*. *Oncogene*, 9, 3675-81
- Larson, R.C., Lavenir, I., Larson, T.A., Baer, R., Warren, A.J., Wadman, I., Nottage, K. and Rabbitts, T.H. (1996): Protein dimerization between Lmo2 (*Rbtn2*) and Tal1 alters thymocyte development and potentiates T cell tumorigenesis in transgenic mice. *EMBO J.*, 15 (5), 1021-7
- Larson, R.C., Osada, H., Larson, T.A., Lavenir, I. and Rabbitts, T.H. (1995): The oncogenic LIM protein *Rbtn2* causes thymic developmental aberrations that precede malignancy in transgenic mice. *Oncogene*, 11, 853-62
- Lasky, L.A. (1996): Hematopoiesis: Wandering progenitor cells. *Curr. Biol.*, 6(10), 1238-40
- Lengauer, C., Kinzler, K.W. and Vogelstein, B. (1998): Genetic instabilities in human cancers. *Nature*, 396, 643-9
- Li, Y. and Jaiswal, A.K. (1994): Human antioxidant-response-element-mediated regulation of type 1 NAD(P)H:quinone oxidoreductase gene expression. Effect of sulfhydryl modifying agents. *Eur. J. Biochem.*, 226, 31-9
- Look, A.T. (1997): Oncogenic transcription factors in the human acute leukemias. *Science*, 278, 1059-64
- Lozzio, C.B. and Lozzio, B.B. (1975): Human Chronic Myelogenous Leukemia Cell-Line With Positive Philadelphia Chromosome. *Blood*, 45 (3), 321-34
- Maeshima, Y., Kashihara, N., Yasuda, T., Sugiyama, H., Sekikawa, T., Okamoto, K., Kanao, K., Watanabe, Y., Kanwar, Y.S. and Makino, H. (1998): Inhibition of mesangial cell proliferation by E2F decoy oligodeoxynucleotide in vitro and in vivo. *J. Clin. Invest.*, 101 (11), 2589-97
- Malter, J.S. (1989): Identification of an AUUUA-Specific Messenger RNA Binding Protein. *Science*, 246, 664-6
- Mantovani, R. (1998): A survey of 178 NF-Y binding CCAAT boxes. *Nucleic Acids Res.*, 26 (5), 1135-43
- Mao, S., Neale, G.A.M. and Goorha, R.M. (1997a): T-cell Proto-oncogen Rhombotin-2 Is a Complex Transcription Regulator Containing Multiple Activation and Repression Domains. *J. Biol. Chem.*, 272 (9), 5594-9



- Mao, S., Neale, G.A.M. and Goorha, R.M. (1997b): T-cell oncogene rhombotin-2 interacts with retinoblastoma-binding protein 2. *Oncogene*, 14, 1531-9
- Marshall, C.J. (1991): Tumor Suppressor Genes. *Cell*, 64, 313-26
- Martin, D.I.K., Tsai, S.F. and Orkin, S.H. (1989): Increased  $\gamma$ -globin expression in a nondeletion HPFH mediated by an erythroid-specific DNA-binding factor. *Nature*, 338, 435-8
- Martin, P. and Papayannopoulou, T. (1982): HEL cells: A New Human Erythroleukemia Cell Line with Spontaneous and Induced Globin Expression. *Science*, 216, 1233-5
- McGuire, E.A., Rintoul, C.E., Sclar, G.M. and Korsmeyer, S.J. (1992): Thymic Overexpression of *Ttg-1* in Transgenic Mice Results in T-Cell Acute Lymphoblastic Leukemia/Lymphoma. *Mol. Cell. Biol.*, 12 (9), 4186-96
- Merika, M. and Orkin, S.H. (1995): Functional Synergy and Physical Interactions of the Erythroid Factor GATA-1 with the Krüppel Family Proteins Sp1 and EKLF. *Mol. Cell. Biol.*, 15 (5), 2437-47
- Michelsen, J.W., Schmeichel, K.L., Beckerle, M.C. and Winge, D.R. (1993): The LIM motif defines a specific zinc-binding protein domain. *Proc. Natl. Acad. Sci. USA*, 90, 4404-8
- Milán, M. and Cohen, S.M. (1999): Regulation of LIM Homeodomain Activity In Vivo: A Tetramer of dLDB and Apterous Confers Activity and Capacity for Regulation by dLMO. *Mol. Cell*, 4, 267-73
- Minegishi, N., Ohta, J., Suwabe, N., Nakauchi, H., Ishihara, H., Hayashi, N. and Yamamoto, M. (1998): Alternative Promoters Regulate Transcription of the Mouse GATA-2 Gene. *J. Biol. Chem.*, 273 (6), 3625-34
- Mitani, K. (1996): Chromosomal abnormalities and oncogenes. *Int. J. Hematol.*, 63, 81-93
- Mitelman, F., Mertens, F. and Johansson, B. (1997): A breakpoint map of recurrent chromosomal rearrangements in human neoplasia. *Nature Genet.*, 15, 417-74
- Morcillo, P., Rosen, C., Baylies, M.K. and Dorsett, D. (1997): Chip, a widely expressed chromosomal protein required for segmentation and activity of a remote wing margin enhancer in *Drosophila*. *Genes Dev.*, 11, 2729-40

- Morishita, R. et al. (1997): In vivo transfection of cis element „decoy“ against nuclear factor- $\kappa$ B binding site prevents myocardial infarction. *Nature Medicine*, 3 (8), 894-9
- Morishita, R., Gibbons, G.H., Horiuchi, M., Ellison, K.E., Nakajima, M., Zhang, L., Kaneda, Y., Ogihara, T. and Dzau, V.J. (1995): A gene therapy strategy using a transcription factor decoy of the E2F binding site inhibits smooth muscle proliferation in vivo. *Proc. Natl. Acad. Sci. USA*, 92, 5855-9
- Mullis, K.B. and Faloona, F.A. (1987): Specific Synthesis of DNA *in Vitro* via a Polymerase-Catalyzed Chain Reaction. *Methods in Enzymology*, 155, 335-50
- Mutoh, H., Fukuda, T., Kitamaoto, T., Masushige, S., Sasaki, H., Siimizu, T. and Kato, S. (1996): Tissue-specific response of the human platelet-activating factor receptor gene to retinoic acid and thyroid hormone by alternative promoter usage. *Proc. Natl. Acad. Sci. USA*, 93, 774-9
- Neale, G.A.M., Mao, S., Parham, D.M., Gopal Murti, K. and Goorha, R.M. (1995a): Expression of the Proto-oncogene Rhombotin-2 Is Identical to the Acute Phase Response Protein Metallothionein, Suggesting Multiple Functions. *Cell Growth Diff.*, 6, 587-96
- Neale, G.A.M., Rehg, J.E. and Goorha, R.M. (1995b): Ectopic Expression of Rhombotin-2 Causes Selective Expansion of CD4<sup>+</sup>CD8<sup>-</sup> Lymphocytes in the Thymus and T-Cell Tumors in Transgenic Mice. *Blood*, 86 (8), 3060-71
- Neale, G.A.M., Rehg, J.E. and Goorha, R.M. (1997): Disruption of T-cell differentiation precedes T-cell tumor formation in LMO-2 (Rhombotin-2) transgenic mice. *Leukemia, Suppl.* 3, 289-90
- Nishiyama, C., Yokota, Y., Okumura, K. and Ra, C. (1999): The Transcription Factors Elf-1 and GATA-1 Bind to Cell-Specific Enhancer Elements of Human High-Affinity IgE Receptor  $\alpha$ -Chain Gene. *J. Immunol.*, 163 (2), 623-30
- Niswander, L. (1997): Limb mutants: what can they tell us about normal limb development ?. *Curr. Opin. Gen. Dev.*, 7, 530-6
- O'Keefe, D.D., Thor, S. and Thomas, J.B. (1998): Function and specificity of LIM domains in *Drosophila* nervous system and wing development. *Development*, 125, 3915-23

- Orkin, S.H. (1995a): Hematopoiesis: how does it happen ?  
Curr. Opin. Cell Biol., 7, 870-77
- Orkin, S.H. (1995b): Transcription factors and hematopoietic development.  
J. Biol. Chem., 270 (10), 4955-8
- Orkin, S.H. (1996) : Development of the hematopoietic system.  
Curr. Opin. Gen. Dev., 6, 597-602
- Orkin, S.H., Porcher, C., Fujiwara, Y., Visvader, J. and Wang, L.C. (1999):  
Intersection between Blood Cell Development and Leukemia Genes.  
Cancer Res., (Suppl.), 59, 1784s-8s
- Osada, H., Grutz, G., Axelson, H, Forster, A. and Rabbitts, T.H. (1995):  
Association of erythroid transcription factors: Complexes involving the  
LIM protein RBTN2 and the zinc-finger protein GATA1.  
Proc. Natl. Acad. Sci. USA, 92, 9585-9
- Osada, H., Grutz, G.G., Axelson, H., Forster, A. and Rabbitts, T.H. (1997): LIM-  
only protein Lmo2 forms a protein complex with erythroid transcription  
factor GATA-1. Leukemia, 11, Suppl. 3, 307-12
- Partington, G.A. and Patient, R.K. (1999): Phosphorylation of GATA-1 increases  
its DNA-binding affinity and is correlated with induction of human K562  
erythroleukaemia cells. Nucleic Acids Res., 27 (4), 1168-75
- Philipsen, S. and Suske, G. (1999): A tale of three fingers: the family of  
mammalian Sp/XKLF transcription factors. Nucleic Acids Res., 27 (15),  
2991-3000
- Piechaczyk, M., Yang, J.Q., Blanchard, J.M., Jeanteur, P. and Marcu, K.B.  
(1985): Posttranscriptional Mechanisms Are Responsible for  
Accumulation of Truncated c-myc RNAs in Murine Plasma Cell Tumors.  
Cell, 42, 589-97
- Pruess, M. M., Drechsler, M. and Royer-Pokora, B. (2000): Promoter 1 of *LMO2*  
is active in erythroid cells and regulated by GATA-1.  
Manuskript in Vorbereitung
- Quandt, K., Frech, K., Karas, H., Wingender, E. and Werner, T. (1995): MatInd  
and MatInspector: new fast and versatile tools for detection of consensus  
matches in nucleotide sequence data.  
Nucleic Acids Res., 23 (23), 4878-84
- Rabbitts, T.H. (1991): Translocations, Master Genes, and Differences between the  
Origins of Acute and Chronic Leukemias. Cell, 67, 641-4

- Rabbitts, T.H. (1994): Chromosomal translocations in human cancer. *Nature*, 372, 143-9
- Rabbitts, T.H. (1997): Chromosomal breakpoints hit the spot. *Nature Med.*, 3 (5), 496-7
- Rabbitts, T.H. (1998): LMO T-cell translocation oncogenes typify genes activated by chromosomal translocations that alter transcription and developmental processes. *Genes Dev.*, 12, 2651-7
- Rabbitts, T.H., Bucher, K., Chung, G., Grutz, G., Warren, A. and Yamada, Y. (1999): The Effect of Chromosomal Translocations in Acute Leukemias: The LMO2 Paradigm in Transcription and Development. *Cancer Res. (Suppl.)*, 59, 1794s-1798s
- Raskind, W.H., Steinmann, L. and Najfeld, V. (1998): Clonal development of myeloproliferative disorders: clues to hematopoietic differentiation and multistep pathogenesis of cancer. *Leukemia*, 12, 108-16
- Reddy, P.M.S. and Shen, C.K.J. (1991): Protein-DNA interactions *in vivo* of an erythroid-specific, human  $\beta$ -globin locus enhancer. *Proc. Natl. Acad. Sci. USA*, 88, 8676-80
- Richardson, C., Moynaham, M.E. and Jasin, M. (1998): Double-strand break repair by interchromosomal recombination: suppression of chromosomal translocations. *Genes Dev.*, 12, 3831-42
- Robb, L. and Begley, C.C. (1997): The SCL/TAL1 gene: roles in normal and malignant haematopoiesis. *BioEssay*, 19 (7), 607-13
- Ross, J. (1996): Control of messenger RNA stability in higher eukaryotes. *Trends Genet.*, 12 (5), 171-5
- Roth, D.B. and Craig, N.L. (1998): VDJ Recombination: A Transposase Goes to Work. *Cell*, 94, 411-4
- Royer-Pokora, B., Fleischer, B., Ragg, S., Loos, U. and Williams, D. (1989): Molecular cloning of the translocation breakpoint in T-ALL 11;14 (p13;q11): genomic map of TCR alpha and delta region on chromosome 14q11 and long-range map of region 11p13. *Hum. Genet.*, 82, 264-70
- Royer-Pokora, B., Loos, U. and Ludwig, W.D. (1991): TTG-2, a new gene encoding a cystein-rich protein with the LIM motif, is overexpressed in acute T-cell leukaemia with the t(11;13)(p13;q11). *Oncogene*, 6, 1887-93

- Royer-Pokora, B., Rogers, M., Zhu, T.H., Schneider, S., Loos, U. and Bölit, U. (1995): The TTG-2/RBTN2 T cell oncogene encodes two alternative transcripts from two promoters: the distal promoter is removed by most 11p13 translocations in acute T cell leukaemia's (T-ALL). *Oncogene*, 10, 1353-60
- Sachs, L. (1996): The control of hematopoiesis and leukemia: From basic biology to the clinic. *Proc. Natl. Acad. Sci. USA*, 93, 4742-9
- Saiki, R., Scharf, S., Faloona, F., Mullis, K., Horn, G., Erlich, H. and Arnheim, N. (1985): Enzymatic amplification of  $\beta$ -globin genomic sequences and restriction site analysis for diagnosis of sickle cell anemia. *Science*, 230, 1350-4
- Saiki, R.K., Gelfand, D.H., Stoffel, S., Scharf, S.J., Higuchi, R., Horn, G.T., Mullis, K.B. and Ehrlich, H.A. (1988): Primer-Directed Enzymatic Amplification of DNA with a Thermostable DNA Polymerase. *Science*, 239, 487-91
- Sambrook, J., Fritsch, E.F. and Maniatis, T. (1989): *Molecular cloning: a laboratory manual* (2<sup>nd</sup> edition). Cold Spring Harbour Laboratory Press
- Sánchez-García, I. (1997): Consequences of chromosomal abnormalities in tumor development. *Annu. Rev. Genet.*, 31, 429-53
- Sánchez-García, I. and Rabbitts, T.H. (1994): The LIM domain: a new structural motif found in zinc-finger-like proteins. *Trends Genet.*, 10 (9), 315-20
- Sánchez-García, I., Axelson, H. and Rabbitts, T.H. (1995): Functional diversity of LIM proteins: Amino-terminal activation domains in the oncogenic proteins RBTN1 and RBTN2. *Oncogene*, 10, 1301-6
- Sanger, F., Nicklen, D. and Coulson, A.R. (1977): DNA sequencing with chain terminating inhibitors. *Proc. Natl. Acad. Sci. USA*, 74, 5463-7
- Sawyers, L.C., Denny, C.T. and Witte, O.N. (1991): Leukemia and the Disruption of Normal Hematopoiesis. *Cell*, 64, 337-50
- Schibler, U., Hagenbüchle, O., Wellauer, P.K. and Pittet, A.C. (1983): Two Promoters of Different Strengths Control the Transcription of the Mouse Alpha-Amylase Gene *Amy-1<sup>a</sup>* in the Parotid Gland and the Liver. *Cell*, 33, 501-8
- Schmeichel, K.L. and Beckerle, M.C. (1994): The LIM Domain Is a Modular Protein-Binding Interface. *Cell*, 79, 211-9

- Sharma, H.W., Perez, J.R., Higgins-Sochaski, K., Hsiao, R. and Narayanan, R. (1996): Transcription factor decoy approach to decipher the role of NF- $\kappa$ B in oncogenesis. *Anticancer Res.*, 16, 61-70
- Shaw, G. and Kamen, R. (1986): A Conserved AU Sequence from the 3' Untranslated Region of GM-CSF mRNA Mediates Selective mRNA Degradation. *Cell*, 46, 659-67
- Shivdasani, R.A., Mayer, E.L. and Orkin, S.H. (1995): Absence of blood formation in mice lacking the T-cell leukaemia oncprotein tal-1/SCL. *Nature*, 373, 432-4
- Shoresh, M., Orgad, S., Shmueli, O., Werczberger, R., Gelbaum, D., Abiri, S. and Segal, D. (1998): Overexpression Beadex mutations and loss-of-function heldup a mutations in *Drosophila* affect the 3' regulatory and coding components, respectively, of the *Dlmo* gene. *Genetics*, 150 (1), 283-99
- Shubin, N., Tabin, C. and Carroll, S. (1997): Fossils, genes and the evolution of animal limbs. *Nature*, 388, 639-48
- Sieweke, M.H. and Graf, T. (1998): A transcription factor party during blood cell differentiation. *Curr. Opin. Gen. Dev.*, 8, 545-51
- Silver, L. and Palis, J. (1997): Initiation of Murine Embryonic Erythropoiesis: A Spatial Analysis. *Blood*, 89(4), 1154-64
- Smith, C.L. and Hager, G.L. (1997): Transcriptional regulation of mammalian genes in vivo. A tale of two templates. *J. Biol. Chem.*, 272, 27493-6
- Smith, L.M., Sanders, J.Z., Kaiser, R.J., Hughes, P., Dodd, C., Connell, C.R., Heiner, C., Kent, S.B.H. and Hood, L.E. (1986): Fluorescence detection in automated DNA sequence analysis. *Nature*, 321, 674-9
- Smith, R.F., Wiese, B.A., Wojzynski, M.K., Davison, D.B. and Worley, K.B. (1996): BCM Search Launcher - An Integrated Interface to Molecular Biology Data Base Search and Analysis Services Available on the World Wide Web. *Genome Res.*, 6, 454-62
- Socolovsky, M., Lodish, H.F. and Daley, G.Q. (1998): Control of hematopoietic differentiation: Lack of specificity in signalling by cytokine receptors. *Proc. Natl. Acad. Sci. USA*, 95, 6573-5
- Sonenberg, N. (1994): mRNA translation: influence of the 5' and 3' untranslated regions. *Curr. Opin. Gen. Dev.*, 4, 310-5

- Sun, F.L. and Elgin, S.C.R. (1999): Putting Boundaries on Silence. *Cell*, 99, 459-62
- Tanaka, H., Vickart, P., Bertrand, J.R., Rayner, B., Morvan, F., Imbach, J.L., Paulin, D. and Malvy, C. (1994): Sequence-specific interaction of  $\alpha^*\beta$ -anomeric double-stranded DNA with the p50 subunit of NF $\kappa$ B: application to the decoy approach. *Nucleic Acids Res.*, 22 (15), 3069-74
- Thandla, S. and Aplan, P.D. (1997): Molecular biology of acute lymphocytic leukemia. *Semin. Oncol.*, 24 (1), 45-56
- Tjian, R. and Maniatis, T. (1994): Transcriptional Activation: A Complex Puzzle with Few Easy Pieces. *Cell*, 77, 5-8
- Travers, A. (1999): Chromatin modification by DNA tracking. *Proc. Natl. Acad. Sci., USA*, 96, 13634-7
- Tsai, S.H., Strauss, E. and Orkin, S.H. (1991): Functional analysis and in vivo footprinting implicate the erythroid transcription factor GATA-1 as a positive regulator of its own promoter. *Genes Dev.*, 5, 919-31
- Tsang, A.P., Visvader, J.E., Turner, C.A., Fujiwara, Y., Yu, C., Weiss, M.J., Crossley, M. and Orkin, S.H. (1997): FOG, a multiple zinc finger protein, acts as a cofactor for transcription factor GATA-1 in erythroid and megakaryocytic differentiation. *Cell*, 90, 109-19
- Tse, E. et al. (1999): Characterization of the *Lmo4* gene encoding a LIM-only protein: genomic organization and comparative chromosomal mapping. *Mamm. Genome*, 10, 1089-94
- Tyler, J.K. and Kadonaga, J.T. (1999): The „Dark Side“ of Chromatin Remodelling: Repressive Effects on Transcription. *Cell*, 99, 443-6
- Uckun, F.M., Sensel, M.G., Sun, L., Steinherz, P.G., Trigg, M.E., Heerema, N.A., Sather, H.N., Reaman, G.H. and Gaynon, P.S. (1998): Biology and Treatment of Childhood T-Lineage Acute Lymphoblastic Leukemia. *Blood*, 91 (3), 735-46
- Valge-Archer, V., Forster, A. and Rabbitts, T.H. (1998): The LMO1 and LDB1 proteins interact in human T cell acute leukaemia with the chromosomal translocation t(11;14)(p15;q11). *Oncogene*, 17, 3199-202
- Valge-Archer, V.E., Osada, H., Warren, A.J., Forster, A., Li, J., Baer, R. and Rabbitts, T.H. (1994): The LIM protein RBTN2 and the basic helix-loop-helix protein TAL1 are present in a complex in erythroid cells. *Proc. Natl. Acad. Sci. USA*, 91, 8617-21

- Varshavsky, A. (1987): Electrophoretic Assay for DNA-Binding Proteins. *Methods Enzymol.*, 151, 551-65
- Visvader, J.E., Mao, X., Fujiwara, Y., Hahm, K. and Orkin, S.H. (1997): The LIM-domain binding protein Ldb1 and its partner LMO2 act as negative regulators of erythroid differentiation. *Proc. Natl. Acad. Sci. USA*, 94, 13707-12
- Vyas, P., McDevitt, M.A., Cantor, A.B., Katz, S.G., Fujiwara, Y. and Orkin, S.H. (1999): Different sequence requirements for expression in erythroid and megakaryocytic cells within a regulatory element upstream of the *GATA-1* gene. *Development*, 126, 2799-811
- Wadman, I., Li, X., Bash, R.O., Forster, A., Osada, H., Rabbitts, T.H. and Baer, R. (1994): Specific *in vivo* association between the bHLH and LIM proteins implicated in human T cell leukemia. *EMBO J.*, 13 (29), 4831-9
- Wadman, I.A., Osada, H., Grütz, G.G., Agulnick, A.D., Westphal, H., Forster, A. and Rabbitts, T.H. (1997): The LIM-only protein Lmo2 is a bridging molecule assembling an erythroid, DNA-binding complex which includes the TAL1, E47, GATA-1 and Ldb1/NLI proteins. *EMBO J.*, 16 (11), 3145-57
- Wagner, R.W. (1994): Gene inhibition using antisense oligodeoxynucleotides. *Nature*, 372, 333-5
- Warren, A.J., Colledge, W.H., Carlton, M.B.L., Evans, M.J., Smith, A.J.H. and Rabbitts, T.H. (1994): The Oncogenic Cystein-Rich LIM Domain Protein Rbtn2 Is Essential for Erythroid Development. *Cell*, 78, 45-57
- Wilkinson, D.A., Neale, G.A.M., Mao, S., Naeve, C.W. and Goorha, R.M. (1997): Elf-2, a rhombotin-2 binding *ets* transcription factor: discovery and potential role in T cell leukemia. *Leukemia*, 11, 86-96
- Wood, W.G. (1996): The complexity of  $\beta$  globin gene regulation. *Trends Gen.*, 12 (6), 204-6
- Workman, J.L. and Kingston, R.E. (1998): Alteration of nucleosome structure as a mechanism of transcriptional regulation. *Annu. Rev. Biochem.*, 67, 545-79
- Wu, C. (1997): Chromatin remodelling and the control of gene expression. *J. Biol. Chem.*, 272, 28171-4



- Xu, X., Kang, S.H., Heidenreich, O., Brown, D.A. and Nerenberg, M.I. (1996): Sequence Requirements of ATF2 and CREB Binding to the Human T-Cell Leukemia Virus Type 1 LTR R Region. *Virology*, 218, 362-71
- Yamada, Y., Pannell, R., Forster, A. and Rabbitts, T.H. (2000): The oncogenic LIM-only transcription factor Lmo2 regulates angiogenesis but not vasculogenesis in mice. *Proc. Natl. Acad. Sci. USA*, 97, 320-4
- Yamada, Y., Warren, A.J., Dobson, C., Forster, A., Pannell, R. and Rabbitts, T.H. (1998): The T cell leukemia LIM protein Lmo2 is necessary for adult mouse hematopoiesis. *Proc. Natl. Acad. Sci. USA*, 95, 3890-5
- Zeng, C., Justice, N.J., Abdelilah, S., Chan, Y.M., Jan, L.Y. and Jan, Y.N. (1998): The *Drosophila* Lim-only gene, *dLMO*, is mutated in *Beadex* alleles and might represent an evolutionarily conserved function in appendage development. *Proc. Natl. Acad. Sci. USA*, 95, 10637-42
- Zhang, P., Behre, G, Pan, J., Iwama, A., Wara-Aswapati, N. Radomska, H.S., Auron, P.E., Tenen, D.G. and Sun, Z. (1999): Negative cross-talk between hematopoietic regulators: GATA proteins repress PU.1. *Proc. Natl. Acad. Sci. USA*, 96, 8705-10
- Zhu, T.H., Bodem, J., Keppel, E., Paro, R. and Royer-Pokora, B. (1995): A single ancestral gene of the human LIM domain oncogene family *LMO* in *Drosophila*: characterization of the *Drosophila Dlmo* gene. *Oncogene*, 11, 1283-90
- Zucman-Rossi, J., Legoix, P., Victor, J.M., Lopez, B. and Thomas, G. (1998): Chromosome translocation based on illegitimate recombination in human tumors. *Proc. Natl. Acad. Sci. USA*, 95, 11786-91

## 7.2. Eigene Veröffentlichungen

### Poster-Präsentationen:

Roest Crolius, H., Kleinkauf, N., Leser, U., Rak, K., **Pruess, M.** and Lehrach, H. (1997): Towards a clone and transcript map of the human X-chromosome. Congress of Molecular Medicine, Berlin, Germany 1997. J. Mol. Med. 75(5), B49, 1997

**Pruess, Maik M.**, Drechsler, M. and Royer-Pokora, B. (1998): GATA1 binds to the *LMO2* promotor: Implications for the transcriptional control of *LMO2*. 10. Jahrestagung der Deutschen Gesellschaft für Humangenetik, Jena, 25. - 28. März 1998. Medizinische Genetik, **10** (1) 136, 1998

**Pruess, Maik M.**, Drechsler, M. and Royer-Pokora, B. (2000): Lineage specific DNaseI hypersensitive sites located within the 5' flanking region of the human *LMO2* gene. 12. Jahrestagung der Deutschen Gesellschaft für Humangenetik, Lübeck, 22. - 25. März 2000. Medizinische Genetik, im Druck.

### Vortrag:

**Pruess, Maik M.:** „Analyse der transkriptionellen Regulation des putativen T-Zell Onkogens *LMO-2*“. 7. Oktober 1999  
Max-Planck-Institut für Molekulare Genetik in Berlin.

### Veröffentlichung in wissenschaftlicher Fachzeitschrift

**Pruess, Maik M.**, Drechsler, M. and Royer-Pokora, B. (2000): Promoter 1 of *LMO2* is active in erythroid cells and regulated by GATA-1.  
Manuskript in Vorbereitung