

6 REFERENCES

- Adriaensen, H., J. Gybels, H.O. Handwerker, and J. Van Hees. 1983. Response properties of thin myelinated (A-delta) fibers in human skin nerves. *J Neurophysiol.* 49:111-22.
- Airaksinen, M.S., M. Koltzenburg, G.R. Lewin, Y. Masu, C. Helbig, E. Wolf, G. Brem, K.V. Toyka, H. Thoenen, and M. Meyer. 1996. Specific subtypes of cutaneous mechanoreceptors require neurotrophin-3 following peripheral target innervation. *Neuron.* 16:287-95.
- Alberts, B.; Bray, D.; Lewis, J.; Raff, M.; Roberts, K. and Watson, J.D. 1994 The Cytoskeleton. In *Molecular Biology of the Cell*, Garland Publishing New York 3rd edn, pp787-861,
- Alvarez de la Rosa, D., P. Zhang, D. Shao, F. White, and C.M. Canessa. 2002. Functional implications of the localization and activity of acid-sensitive channels in rat peripheral nervous system. *Proc Natl Acad Sci U S A.* 99:2326-31.
- Anzai, N., E. Deval, L. Schaefer, V. Friend, M. Lazdunski, and E. Lingueglia. 2002. The multivalent PDZ domain-containing protein CIPP is a partner of acid-sensing ion channel 3 in sensory neurons. *J Biol Chem.* 277:16655-61.
- Bassilana, F., G. Champigny, R. Waldmann, J.R. de Weille, C. Heurteaux, and M. Lazdunski. 1997. The acid-sensitive ionic channel subunit ASIC and the mammalian degenerin MDEG form a heteromultimeric H⁺-gated Na⁺ channel with novel properties. *J Biol Chem.* 272:28819-22.
- Bendtsen, J.D., H. Nielsen, G. von Heijne, and S. Brunak. 2004. Improved prediction of signal peptides: SignalP 3.0. *J Mol Biol.* 340:783-95.
- Benson, C.J., J. Xie, J.A. Wemmie, M.P. Price, J.M. Henss, M.J. Welsh, and P.M. Snyder. 2002. Heteromultimers of DEG/ENaC subunits form H⁺-gated channels in mouse sensory neurons. *Proc Natl Acad Sci U S A.* 99:2338-43.
- Bickel, P.E., P.E. Scherer, J.E. Schnitzer, P. Oh, M.P. Lisanti, and H.F. Lodish. 1997. Flotillin and epidermal surface antigen define a new family of caveolae-associated integral membrane proteins. *J Biol Chem.* 272:13793-802.
- Birder, L.A., and E.R. Perl. 1994. Cutaneous sensory receptors. *J Clin Neurophysiol.* 11:534-52.

- Blatter, M.C., R.W. James, S. Messmer, F. Barja, and D. Pometta. 1993. Identification of a distinct human high-density lipoprotein subspecies defined by a lipoprotein-associated protein, K-45. Identity of K-45 with paraoxonase. *Eur J Biochem.* 211:871-9.
- Brown, A.G.a.I., A. 1967. A quantitative study of cutaneous receptors and afferent fibres in the cat and rabbit. *J.Physiol. Lond.:*707-733.
- Burgess, P.P., D; Warren, RM. 1968. Receptor types in cat hairy skin supplied by myelinated fibers. *J Neurophysiol.* 31:833-848.
- Burgess, P.R., and E.R. Perl. 1967. Myelinated afferent fibres responding specifically to noxious stimulation of the skin. *J Physiol.* 190:541-62.
- Burnstock, G. 1999. Release of vasoactive substances from endothelial cells by shear stress and purinergic mechanosensory transduction. *J Anat.* 194 (Pt 3):335-42.
- Cain, D.M., S.G. Khasabov, and D.A. Simone. 2001. Response properties of mechanoreceptors and nociceptors in mouse glabrous skin: an in vivo study. *J Neurophysiol.* 85:1561-74.
- Campbell, J.N., and R.A. Meyer. 2006. Mechanisms of neuropathic pain. *Neuron.* 52:77-92.
- Caterina, M.J., A. Leffler, A.B. Malmberg, W.J. Martin, J. Trafton, K.R. Petersen-Zeitz, M. Koltzenburg, A.I. Basbaum, and D. Julius. 2000. Impaired nociception and pain sensation in mice lacking the capsaicin receptor. *Science.* 288:306-13.
- Catton, W.T. 1970. Mechanoreceptor function. *Physiol Rev.* 50:297-318.
- Chalfie, M., and M. Au. 1989. Genetic control of differentiation of the *Caenorhabditis elegans* touch receptor neurons. *Science.* 243:1027-33.
- Chalfie, M., and J. Sulston. 1981. Developmental genetics of the mechanosensory neurons of *Caenorhabditis elegans*. *Dev Biol.* 82:358-70.
- Chalfie, M., J.E. Sulston, J.G. White, E. Southgate, J.N. Thomson, and S. Brenner. 1985. The neural circuit for touch sensitivity in *Caenorhabditis elegans*. *J Neurosci.* 5:956-64.
- Chambers, M.R., K.H. Andres, M. von Duering, and A. Iggo. 1972. The structure and function of the slowly adapting type II mechanoreceptor in hairy skin. *Q J Exp Physiol Cogn Med Sci.* 57:417-45.
- Chambers, M.R., and A. Iggo. 1967. Slowly-adapting cutaneous mechanoreceptors. *J Physiol.* 192:26P-27P.

- Chelur, D.S., G.G. Ernstrom, M.B. Goodman, C.A. Yao, L. Chen, O.H. R, and M. Chalfie. 2002. The mechanosensory protein MEC-6 is a subunit of the *C. elegans* touch-cell degenerin channel. *Nature*. 420:669-73.
- Chen, C.C., A.N. Akopian, L. Sivilotti, D. Colquhoun, G. Burnstock, and J.N. Wood. 1995. A P2X purinoceptor expressed by a subset of sensory neurons. *Nature*. 377:428-31.
- Corey, D.P., and A.J. Hudspeth. 1979. Response latency of vertebrate hair cells. *Biophys J*. 26:499-506.
- Dell'Orco, R.T., J.K. McClung, E.R. Jupe, and X.T. Liu. 1996. Prohibitin and the senescent phenotype. *Exp Gerontol*. 31:245-52.
- Deval, E., V. Friend, C. Thirant, M. Salinas, M. Jodar, M. Lazdunski, and E. Lingueglia. 2006. Regulation of sensory neuron-specific acid-sensing ion channel 3 by the adaptor protein Na⁺/H⁺ exchanger regulatory factor-1. *J Biol Chem*. 281:1796-807.
- Donier, E., F. Rugiero, K. Okuse, and J.N. Wood. 2005. Annexin II light chain p11 promotes functional expression of acid-sensing ion channel ASIC1a. *J Biol Chem*. 280:38666-72.
- Drummond, H.A., F.M. Abboud, and M.J. Welsh. 2000. Localization of beta and gamma subunits of ENaC in sensory nerve endings in the rat foot pad. *Brain Res*. 884:1-12.
- Drummond, H.A., M.J. Welsh, and F.M. Abboud. 2001. ENaC subunits are molecular components of the arterial baroreceptor complex. *Ann N Y Acad Sci*. 940:42-7.
- Dubreuil, A.S., H. Boukhaddaoui, G. Desmadryl, C. Martinez-Salgado, R. Moshourab, G.R. Lewin, P. Carroll, J. Valmier, and F. Scamps. 2004. Role of T-type calcium current in identified D-hair mechanoreceptor neurons studied in vitro. *J Neurosci*. 24:8480-4.
- Eisenhaber, B., P. Bork, and F. Eisenhaber. 1999. Prediction of potential GPI-modification sites in proprotein sequences. *J Mol Biol*. 292:741-58.
- Eisenhaber B, B.P., Eisenhaber F. 1998. Sequence properties of GPI-anchored proteins near the omega-site: constraints for the polypeptide binding site of the putative transamidase. *Protein Eng*. 11:1155-61.
- Ernstrom, G.G., and M. Chalfie. 2002. Genetics of sensory mechanotransduction. *Annu Rev Genet*. 36:411-53.
- Fain, G.L. Sensory Transduction, Sinauer Associates Inc. 2003
Cutaneous mechanoreceptors and nociceptors. *Handbook of Sensory Physiology*. Vol II, ed Iggo, A. Springer, Berlin, Heidelberg, New York

REFERENCES

- Fleischer, E., H.O. Handwerker, and S. Joukhadar. 1983. Unmyelinated nociceptive units in two skin areas of the rat. *Brain Res.* 267:81-92.
- Foster, L.J., C.L. De Hoog, and M. Mann. 2003. Unbiased quantitative proteomics of lipid rafts reveals high specificity for signaling factors. *Proc Natl Acad Sci U S A.* 100:5813-8.
- French, A.S. 1992. Mechanotransduction. *Annu Rev Physiol.* 54:135-52.
- Fricke, B., R. Lints, G. Stewart, H. Drummond, G. Dodt, M. Driscoll, and M. von During. 2000. Epithelial Na⁺ channels and stomatin are expressed in rat trigeminal mechanosensory neurons. *Cell Tissue Res.* 299:327-34.
- Fricke, B., G.W. Stewart, K.J. Treharne, A. Mehta, G. Knopfle, N. Friedrichs, K.M. Muller, and M. von During. 2003. Stomatin immunoreactivity in ciliated cells of the human airway epithelium. *Anat Embryol (Berl).* 207:1-7.
- Friede, R.L. 1986. Relation between myelin sheath thickness, internode geometry, and sheath resistance. *Exp Neurol.* 92:234-47.
- Friede, R.L., and R. Bischhausen. 1982. How are sheath dimensions affected by axon caliber and internode length? *Brain Res.* 235:335-50.
- Gallagher, P.G., and B.G. Forget. 1995. Structure, organization, and expression of the human band 7.2b gene, a candidate gene for hereditary hydrocytosis. *J Biol Chem.* 270:26358-63.
- Garcia-Anoveros, J., and D.P. Corey. 1997. The molecules of mechanosensation. *Annu Rev Neurosci.* 20:567-94.
- Garcia-Anoveros, J., C. Ma, and M. Chalfie. 1995. Regulation of *Caenorhabditis elegans* degenerin proteins by a putative extracellular domain. *Curr Biol.* 5:441-8.
- Garcia-Anoveros, J., T.A. Samad, L. Zuvella-Jelaska, C.J. Woolf, and D.P. Corey. 2001. Transport and localization of the DEG/ENaC ion channel BNaC1alpha to peripheral mechanosensory terminals of dorsal root ganglia neurons. *J Neurosci.* 21:2678-86.
- Gillespie, P.G., and R.G. Walker. 2001. Molecular basis of mechanosensory transduction. *Nature.* 413:194-202.
- Goldstein, B.J., H.M. Kulaga, and R.R. Reed. 2003. Cloning and characterization of SLP3: a novel member of the stomatin family expressed by olfactory receptor neurons. *J Assoc Res Otolaryngol.* 4:74-82.
- Goodman, M.B., G.G. Ernstrom, D.S. Chelur, R. O'Hagan, C.A. Yao, and M. Chalfie. 2002. MEC-2 regulates *C. elegans* DEG/ENaC channels needed for mechanosensation. *Nature.* 415:1039-42.

-
- Goodman, M.B., and E.M. Schwarz. 2003. Transducing touch in *Caenorhabditis elegans*. *Annu Rev Physiol.* 65:429-52.
- Guic-Robles, E., C. Valdivieso, and G. Guajardo. 1989. Rats can learn a roughness discrimination using only their vibrissal system. *Behav Brain Res.* 31:285-9.
- Gunthorpe, M.J., G.D. Smith, J.B. Davis, and A.D. Randall. 2001. Characterisation of a human acid-sensing ion channel (hASIC1a) endogenously expressed in HEK293 cells. *Pflugers Arch.* 442:668-74.
- Habler, H.J., W. Janig, and M. Koltzenburg. 1990. Activation of unmyelinated afferent fibres by mechanical stimuli and inflammation of the urinary bladder in the cat. *J Physiol.* 425:545-62.
- Halata, Z. 1993. Sensory innervation of the hairy skin (light- and electronmicroscopic study). *J Invest Dermatol.* 101:75S-81S.
- Hamill, O.P., and B. Martinac. 2001. Molecular basis of mechanotransduction in living cells. *Physiol Rev.* 81:685-740.
- Handwerker, H.O., S. Kilo, and P.W. Reeh. 1991. Unresponsive afferent nerve fibres in the sural nerve of the rat. *J Physiol.* 435:229-42.
- Harel, M., A. Aharoni, L. Gaidukov, B. Brumshtein, O. Khersonsky, R. Megeed, H. Dvir, R.B. Ravelli, A. McCarthy, L. Toker, I. Silman, J.L. Sussman, and D.S. Tawfik. 2004. Structure and evolution of the serum paraoxonase family of detoxifying and anti-atherosclerotic enzymes. *Nat Struct Mol Biol.* 11:412-9.
- Hiebl-Dirschmied, C.M., G.R. Adolf, and R. Prohaska. 1991a. Isolation and partial characterization of the human erythrocyte band 7 integral membrane protein. *Biochim Biophys Acta.* 1065:195-202.
- Hiebl-Dirschmied, C.M., B. Entler, C. Glotzmann, I. Maurer-Fogy, C. Stratowa, and R. Prohaska. 1991b. Cloning and nucleotide sequence of cDNA encoding human erythrocyte band 7 integral membrane protein. *Biochim Biophys Acta.* 1090:123-4.
- Hruska-Hageman, A.M., C.J. Benson, A.S. Leonard, M.P. Price, and M.J. Welsh. 2004. PSD-95 and Lin-7b interact with acid-sensing ion channel-3 and have opposite effects on H⁺-gated current. *J Biol Chem.* 279:46962-8.
- Hruska-Hageman, A.M., J.A. Wemmie, M.P. Price, and M.J. Welsh. 2002. Interaction of the synaptic protein PICK1 (protein interacting with C kinase 1) with the non-voltage gated sodium channels BNC1 (brain Na⁺ channel 1) and ASIC (acid-sensing ion channel). *Biochem J.* 361:443-50.
- Hu, J., and G.R. Lewin. 2006. Mechanosensitive currents in the neurites of cultured mouse sensory neurones. *J Physiol.*

REFERENCES

- Hu, J., N. Milenkovic, and G.R. Lewin. 2006. The high threshold mechanotransducer: a status report. *Pain*. 120:3-7.
- Hu, J.W., and B.J. Sessle. 1988. Properties of functionally identified nociceptive and nonnociceptive facial primary afferents and presynaptic excitability changes induced in their brain stem endings by raphe and orofacial stimuli in cats. *Exp Neurol*. 101:385-99.
- Huang, M., G. Gu, E.L. Ferguson, and M. Chalfie. 1995. A stomatin-like protein necessary for mechanosensation in *C. elegans*. *Nature*. 378:292-5.
- Hutson, K.A., and R.B. Masterton. 1986. The sensory contribution of a single vibrissa's cortical barrel. *J Neurophysiol*. 56:1196-223.
- Iggo, A. 1977. Cutaneous and subcutaneous sense organs. *Br Med Bull*. 33:97-102.
- Iggo, A., and K.H. Andres. 1982. Morphology of cutaneous receptors. *Annu Rev Neurosci*. 5:1-31.
- Iggo, A., and A.R. Muir. 1969. The structure and function of a slowly adapting touch corpuscle in hairy skin. *J Physiol*. 200:763-96.
- Innes, D.S., J.H. Sinard, D.M. Gilligan, L.M. Snyder, P.G. Gallagher, and J.S. Morrow. 1999. Exclusion of the stomatin, alpha-adducin and beta-adducin loci in a large kindred with dehydrated hereditary stomatocytosis. *Am J Hematol*. 60:72-4.
- Kamynina, E., and O. Staub. 2002. Concerted action of ENaC, Nedd4-2, and Sgk1 in transepithelial Na(+) transport. *Am J Physiol Renal Physiol*. 283:F377-87.
- Kellenberger, S., and L. Schild. 2002. Epithelial sodium channel/degenerin family of ion channels: a variety of functions for a shared structure. *Physiol Rev*. 82:735-67.
- Kelso, G.J., W.D. Stuart, R.J. Richter, C.E. Furlong, T.C. Jordan-Starck, and J.A. Harmony. 1994. Apolipoprotein J is associated with paraoxonase in human plasma. *Biochemistry*. 33:832-9.
- Kernan, M., D. Cowan, and C. Zuker. 1994. Genetic dissection of mechanosensory transduction: mechanoreception-defective mutations of *Drosophila*. *Neuron*. 12:1195-206.
- Kernan, M., and C. Zuker. 1995. Genetic approaches to mechanosensory transduction. *Curr Opin Neurobiol*. 5:443-8.
- Kobayakawa, K., R. Hayashi, K. Morita, K. Miyamichi, Y. Oka, A. Tsuboi, and H. Sakano. 2002. Stomatin-related olfactory protein, SRO, specifically expressed in the murine olfactory sensory neurons. *J Neurosci*. 22:5931-7.

-
- Koerber, H.M., LM 6. 1992. Functional Heterogeneity of Dorsal Root Ganglion Cells. pp77-96, pp.
- Koltzenburg, M., C.L. Stucky, and G.R. Lewin. 1997. Receptive properties of mouse sensory neurons innervating hairy skin. *J Neurophysiol.* 78:1841-50.
- Kress, M., M. Koltzenburg, P.W. Reeh, and H.O. Handwerker. 1992. Responsiveness and functional attributes of electrically localized terminals of cutaneous C-fibers in vivo and in vitro. *J Neurophysiology.* 68:581-595.
- Krishtal, O. 2003. The ASICs: signaling molecules? Modulators? *Trends Neurosci.* 26:477-83.
- Kronegg, J.B., Didier. 1999. Detection/prediction of GPI cleavage site (GPI-anchor) in a protein (DGPI. Retrieved [retrieve date] from <http://129.194.185.165/dgpi/>.
- Kruger, L., E.R. Perl, and M.J. Sedivec. 1981. Fine structure of myelinated mechanical nociceptor endings in cat hairy skin. *J Comp Neurol.* 198:137-54.
- Lai, C.C., K. Hong, M. Kinnell, M. Chalfie, and M. Driscoll. 1996. Sequence and transmembrane topology of MEC-4, an ion channel subunit required for mechanotransduction in *Caenorhabditis elegans*. *J Cell Biol.* 133:1071-81.
- Lawson, S. 1992. Morphological and Biochemical Cell Types of Sensory Neurons. *Sensory neurons: Diversity, Development, and Plasticity*, edited by S.A. Scott, New York: Oxford University Press:27-59.
- Lewin, G.R., S.J. Lisney, and L.M. Mendell. 1992a. Neonatal Anti-NGF Treatment Reduces the Delta- and C-Fibre Evoked Vasodilator Responses in Rat Skin: Evidence That Nociceptor Afferents Mediate Antidromic Vasodilatation. *Eur J Neurosci.* 4:1213-1218.
- Lewin, G.R., and S.B. McMahon. 1991. Physiological properties of primary sensory neurons appropriately and inappropriately innervating skin in the adult rat. *J Neurophysiol.* 66:1205-17.
- Lewin, G.R., and L.M. Mendell. 1994. Regulation of cutaneous C-fiber heat nociceptors by nerve growth factor in the developing rat. *J Neurophysiol.* 71:941-9.
- Lewin, G.R., and R. Moshourab. 2004. Mechanosensation and pain. *J Neurobiol.* 61:30-44.
- Lewin, G.R., A.M. Ritter, and L.M. Mendell. 1992b. On the role of nerve growth factor in the development of myelinated nociceptors. *J Neurosci.* 12:1896-905.

REFERENCES

- Loewenstein, W.R. 1959. The generation of electric activity in a nerve ending. *Ann N Y Acad Sci.* 81:367-87.
- Low, M.G., and A.R. Saltiel. 1988. Structural and functional roles of glycosyl-phosphatidylinositol in membranes. *Science.* 239:268-75.
- Lynn, B. 1991. 'Silent' nociceptors in the skin. *Trends Neurosci.* 14:95.
- Lynn, B., and S.E. Carpenter. 1982. Primary afferent units from the hairy skin of the rat hind limb. *Brain Res.* 238:29-43.
- Maingret, F., M. Fosset, F. Lesage, M. Lazdunski, and E. Honore. 1999. TRAAK is a mammalian neuronal mechano-gated K⁺ channel. *J Biol Chem.* 274:1381-7.
- Maklad, A., B. Fritsch, and L.A. Hansen. 2004. Innervation of the maxillary vibrissae in mice as revealed by anterograde and retrograde tract tracing. *Cell Tissue Res.* 315:167-80.
- Mannsfeldt, A.G., P. Carroll, C.L. Stucky, and G.R. Lewin. 1999. Stomatin, a MEC-2 like protein, is expressed by mammalian sensory neurons. *Mol Cell Neurosci.* 13:391-404.
- Meyer, R.A., K.D. Davis, R.H. Cohen, R.D. Treede, and J.N. Campbell. 1991. Mechanically insensitive afferents (MIAs) in cutaneous nerves of monkey. *Brain Res.* 561:252-61.
- Mochizuki, H., S.W. Scherer, T. Xi, D.C. Nickle, M. Majer, J.J. Huizenga, L.C. Tsui, and M. Prochazka. 1998. Human PON2 gene at 7q21.3: cloning, multiple mRNA forms, and missense polymorphisms in the coding sequence. *Gene.* 213:149-57.
- Moller, S., M.D. Croning, and R. Apweiler. 2001. Evaluation of methods for the prediction of membrane spanning regions. *Bioinformatics.* 17:646-53.
- Nakamura, F., and S.M. Strittmatter. 1996. P2Y1 purinergic receptors in sensory neurons: contribution to touch-induced impulse generation. *Proc Natl Acad Sci U S A.* 93:10465-70.
- Ng, C.J., D.J. Wadleigh, A. Gangopadhyay, S. Hama, V.R. Grijalva, M. Navab, A.M. Fogelman, and S.T. Reddy. 2001. Paraoxonase-2 is a ubiquitously expressed protein with antioxidant properties and is capable of preventing cell-mediated oxidative modification of low density lipoprotein. *J Biol Chem.* 276:44444-9.
- Nielsen, H.E., Jacob; Brunak, Søren and von Heijne, Gunnar . 1997. Identification of prokaryotic and eukaryotic signal peptides and prediction of their cleavage sites. *Protein Engineering.* 10:1-6.

- Noble, J.A., M.A. Innis, E.V. Koonin, K.E. Rudd, F. Banuett, and I. Herskowitz. 1993. The *Escherichia coli* hflA locus encodes a putative GTP-binding protein and two membrane proteins, one of which contains a protease-like domain. *Proc Natl Acad Sci U S A.* 90:10866-70.
- O'Hagan, R., M. Chalfie, and M.B. Goodman. 2005. The MEC-4 DEG/ENaC channel of *Caenorhabditis elegans* touch receptor neurons transduces mechanical signals. *Nat Neurosci.* 8:43-50.
- Page, A.J., S.M. Brierley, C.M. Martin, C. Martinez-Salgado, J.A. Wemmie, T.J. Brennan, E. Symonds, T. Omari, G.R. Lewin, M.J. Welsh, and L.A. Blackshaw. 2004. The ion channel ASIC1 contributes to visceral but not cutaneous mechanoreceptor function. *Gastroenterology.* 127:1739-47.
- Patel, A.J., E. Honore, F. Maingret, F. Lesage, M. Fink, F. Duprat, and M. Lazdunski. 1998. A mammalian two pore domain mechano-gated S-like K⁺ channel. *Embo J.* 17:4283-90.
- Perl, E.R. 1968. Myelinated afferent fibres innervating the primate skin and their response to noxious stimuli. *J Physiol.* 197:593-615.
- Persson, L.A., and K. Kristensson. 1979. Uptake of horseradish peroxidase in sensory nerve terminals of mouse trigeminal nerve. *Acta Neuropathol (Berl).* 46:191-6.
- Petrini, J.H. 1999. The mammalian Mre11-Rad50-nbs1 protein complex: integration of functions in the cellular DNA-damage response. *Am J Hum Genet.* 64:1264-9.
- Price, M.P., G.R. Lewin, S.L. McIlwrath, C. Cheng, J. Xie, P.A. Heppenstall, C.L. Stucky, A.G. Mannsfeldt, T.J. Brennan, H.A. Drummond, J. Qiao, C.J. Benson, D.E. Tarr, R.F. Hrstka, B. Yang, R.A. Williamson, and M.J. Welsh. 2000. The mammalian sodium channel BNC1 is required for normal touch sensation. *Nature.* 407:1007-11.
- Price, M.P., S.L. McIlwrath, J. Xie, C. Cheng, J. Qiao, D.E. Tarr, K.A. Sluka, T.J. Brennan, G.R. Lewin, and M.J. Welsh. 2001. The DRASIC cation channel contributes to the detection of cutaneous touch and acid stimuli in mice. *Neuron.* 32:1071-83.
- Price, M.P., R.J. Thompson, J.O. Eshcol, J.A. Wemmie, and C.J. Benson. 2004. Stomatin modulates gating of acid-sensing ion channels. *J Biol Chem.* 279:53886-91.
- Primo-Parmo, S.L., R.C. Sorenson, J. Teiber, and B.N. La Du. 1996. The human serum paraoxonase/arylesterase gene (PON1) is one member of a multigene family. *Genomics.* 33:498-507.

REFERENCES

- Pubols, B.H., Jr. 1990. Slowly adapting type I mechanoreceptor discharge as a function of dynamic force versus dynamic displacement of glabrous skin of raccoon and squirrel monkey hand. *Neurosci Lett.* 110:86-90.
- Rajaram, S., M.M. Sedensky, and P.G. Morgan. 1998. Unc-1: a stomatin homologue controls sensitivity to volatile anesthetics in *Caenorhabditis elegans*. *Proc Natl Acad Sci U S A.* 95:8761-6.
- Reddy, S.T., D.J. Wadleigh, V. Grijalva, C. Ng, S. Hama, A. Gangopadhyay, D.M. Shih, A.J. Lusic, M. Navab, and A.M. Fogelman. 2001. Human paraoxonase-3 is an HDL-associated enzyme with biological activity similar to paraoxonase-1 protein but is not regulated by oxidized lipids. *Arterioscler Thromb Vasc Biol.* 21:542-7.
- Reeh, P.W. 1986. Sensory receptors in mammalian skin in an in vitro preparation. *Neurosci Lett.* 66:141-6.
- Reeh, P.W. 1988. Sensory receptors in a mammalian skin-nerve in vitro preparation. *Prog Brain Res.* 74:271-6.
- Reynolds, M.L., and M. Fitzgerald. 1995. Long-term sensory hyperinnervation following neonatal skin wounds. *J Comp Neurol.* 358:487-98.
- Rossier, B.C., S. Pradervand, L. Schild, and E. Hummler. 2002. Epithelial sodium channel and the control of sodium balance: interaction between genetic and environmental factors. *Annu Rev Physiol.* 64:877-97.
- Roza, C., J.L. Puel, M. Kress, A. Baron, S. Diochot, M. Lazdunski, and R. Waldmann. 2004. Knockout of the ASIC2 channel in mice does not impair cutaneous mechanosensation, visceral mechanonociception and hearing. *J Physiol.* 558:659-69.
- Salzer, U., H. Ahorn, and R. Prohaska. 1993. Identification of the phosphorylation site on human erythrocyte band 7 integral membrane protein: implications for a monotopic protein structure. *Biochim Biophys Acta.* 1151:149-52.
- Salzer, U., and R. Prohaska. 2001. Stomatin, flotillin-1, and flotillin-2 are major integral proteins of erythrocyte lipid rafts. *Blood.* 97:1141-3.
- Sambrook, J., Russell, D.W. 2001 *Molecular Cloning: A Laboratory Manual*, the Third edition, Cold Spring Harbor, New York, 7.84
- Schaeren-Wiemers, Gerfin-Moser 1993. A single protocol to detect transcripts of various types and expression levels in neuronal tissue and cultured cells: in situ hybridization using digoxigenin-labelled cRNA probes. *Histochemistry.* 100(6): 431-40
- Schaible, H.G., and R.F. Schmidt. 1988. Excitation and sensitization of fine articular afferents from cat's knee joint by prostaglandin E2. *J Physiol.* 403:91-104.

- Schofield, J.N., I.N. Day, R.J. Thompson, and Y.H. Edwards. 1995. PGP9.5, a ubiquitin C-terminal hydrolase; pattern of mRNA and protein expression during neural development in the mouse. *Brain Res Dev Brain Res.* 85:229-38.
- Snyers, L., D. Thines-Sempoux, and R. Prohaska. 1997. Colocalization of stomatin (band 7.2b) and actin microfilaments in UAC epithelial cells. *Eur J Cell Biol.* 73:281-5.
- Snyers, L., E. Umlauf, and R. Prohaska. 1998. Oligomeric nature of the integral membrane protein stomatin. *J Biol Chem.* 273:17221-6.
- Snyers, L., E. Umlauf, and R. Prohaska. 1999. Cysteine 29 is the major palmitoylation site on stomatin. *FEBS Lett.* 449:101-4.
- Stewart, G.W., B.E. Hepworth-Jones, J.N. Keen, B.C. Dash, A.C. Argent, and C.M. Casimir. 1992. Isolation of cDNA coding for an ubiquitous membrane protein deficient in high Na⁺, low K⁺ stomatocytic erythrocytes. *Blood.* 79:1593-601.
- Stucky, C.L., T. DeChiara, R.M. Lindsay, G.D. Yancopoulos, and M. Koltzenburg. 1998. Neurotrophin 4 is required for the survival of a subclass of hair follicle receptors. *J Neurosci.* 18:7040-6.
- Stucky, C.L., M. Koltzenburg, M. Schneider, M.G. Engle, K.M. Albers, and B.M. Davis. 1999. Overexpression of nerve growth factor in skin selectively affects the survival and functional properties of nociceptors. *J Neurosci.* 19:8509-16.
- Stucky, C.L., J. Rossi, M.S. Airaksinen, and G.R. Lewin. 2002. GFR alpha2/neurturin signalling regulates noxious heat transduction in isolectin B4-binding mouse sensory neurons. *J Physiol.* 545:43-50.
- Sukharev, S.I., P. Blount, B. Martinac, F.R. Blattner, and C. Kung. 1994. A large-conductance mechanosensitive channel in *E. coli* encoded by *mscL* alone. *Nature.* 368:265-8.
- Suzuki, H., R. Kerr, L. Bianchi, C. Frokjaer-Jensen, D. Slone, J. Xue, B. Gerstbrein, M. Driscoll, and W.R. Schafer. 2003. In vivo imaging of *C. elegans* mechanosensory neurons demonstrates a specific role for the MEC-4 channel in the process of gentle touch sensation. *Neuron.* 39:1005-17.
- Tavernarakis, N., W. Shreffler, S. Wang, and M. Driscoll. 1997. *unc-8*, a DEG/ENaC family member, encodes a subunit of a candidate mechanically gated channel that modulates *C. elegans* locomotion. *Neuron.* 18:107-19.
- Tobin, D.M., and C.I. Bargmann. 2004. Invertebrate nociception: behaviors, neurons and molecules. *J Neurobiol.* 61:161-74.

REFERENCES

- Tuckett, R.P., K.W. Horch, and P.R. Burgess. 1978. Response of cutaneous hair and field mechanoreceptors in cat to threshold stimuli. *J Neurophysiol.* 41:138-49.
- Vallbo, A.B., H. Olausson, J. Wessberg, and N. Kakuda. 1995. Receptive field characteristics of tactile units with myelinated afferents in hairy skin of human subjects. *J Physiol.* 483 (Pt 3):783-95.
- Vallet, V., C. Pfister, J. Loffing, and B.C. Rossier. 2002. Cell-surface expression of the channel activating protease xCAP-1 is required for activation of ENaC in the *Xenopus* oocyte. *J Am Soc Nephrol.* 13:588-94.
- Van Boven, R.W., and K.O. Johnson. 1994. The limit of tactile spatial resolution in humans: grating orientation discrimination at the lip, tongue, and finger. *Neurology.* 44:2361-6.
- Waldmann, R., and M. Lazdunski. 1998. H(+)-gated cation channels: neuronal acid sensors in the NaC/DEG family of ion channels. *Curr Opin Neurobiol.* 8:418-24.
- Walker, R.G., A.T. Willingham, and C.S. Zuker. 2000. A *Drosophila* mechanosensory transduction channel. *Science.* 287:2229-34.
- Wang, D., W.C. Mentzer, T. Cameron, and R.M. Johnson. 1991. Purification of band 7.2b, a 31-kDa integral phosphoprotein absent in hereditary stomatocytosis. *J Biol Chem.* 266:17826-31.
- Wang, Y., and J.S. Morrow. 2000. Identification and characterization of human SLP-2, a novel homologue of stomatin (band 7.2b) present in erythrocytes and other tissues. *J Biol Chem.* 275:8062-71.
- Williams, P.L., and C.P. Wendell-Smith. 1971. Some additional parametric variations between peripheral nerve fibre populations. *J Anat.* 109:505-26.
- Zhang, S., J. Arnadottir, C. Keller, G.A. Caldwell, C.A. Yao, and M. Chalfie. 2004. MEC-2 is recruited to the putative mechanosensory complex in *C. elegans* touch receptor neurons through its stomatin-like domain. *Curr Biol.* 14:1888-96.
- Zhu, Y., C. Paszty, T. Turetsky, S. Tsai, F.A. Kuypers, G. Lee, P. Cooper, P.G. Gallagher, M.E. Stevens, E. Rubin, N. Mohandas, and W.C. Mentzer. 1999. Stomatocytosis is absent in "stomatin"-deficient murine red blood cells. *Blood.* 93:2404-10.