

7. Literaturverzeichnis

- Aller, S.G., Eng, E.T., De Feo, C.J. and Unger, V.M. (2004) Eukaryotic CTR copper uptake transporters require two faces of the third transmembrane domain for helix packing, oligomerization, and function. *J Biol Chem*, 279, 53435-53441.
- Alzheimer, A. (1907) Über eine eigenartige Erkrankung der Hirnrinde. *Allgemeine Zeitschrift für Psychiatrie und Psychisch-Gerichtliche Medizin*, 64, 146-148.
- Anliker, B. and Muller, U. (2006) The functions of mammalian amyloid precursor protein and related amyloid precursor-like proteins. *Neurodegener Dis*, 3, 239-246.
- Annaert, W. and De Strooper, B. (2002) A cell biological perspective on Alzheimer's disease. *Annu Rev Cell Dev Biol*, 18, 25-51.
- Araki, W., Saito, S., Takahashi-Sasaki, N., Shiraishi, H., Komano, H. and Murayama, K.S. (2006) Characterization of APH-1 mutants with a disrupted transmembrane GxxxG motif. *J Mol Neurosci*, 29, 35-43.
- Arispe, N., Rojas, E. and Pollard, H.B. (1993) Alzheimer disease amyloid beta protein forms calcium channels in bilayer membranes: blockade by tromethamine and aluminum. *Proc Natl Acad Sci U S A*, 90, 567-571.
- Arselin, G., Giraud, M.F., Dautant, A., Vaillier, J., Brethes, D., Coulary-Salin, B., Schaeffer, J. and Velours, J. (2003) The GxxxG motif of the transmembrane domain of subunit e is involved in the dimerization/oligomerization of the yeast ATP synthase complex in the mitochondrial membrane. *Eur J Biochem*, 270, 1875-1884.
- Bayer, T.A., Schafer, S., Simons, A., Kemmling, A., Kamer, T., Tepest, R., Eckert, A., Schussel, K., Eikenberg, O., Sturchler-Pierrat, C., Abramowski, D., Staufenbiel, M. and Multhaup, G. (2003) Dietary Cu stabilizes brain superoxide dismutase 1 activity and reduces amyloid Abeta production in APP23 transgenic mice. *Proc Natl Acad Sci U S A*, 100, 14187-14192.
- Bedouelle, H. and Duplay, P. (1988) Production in *Escherichia coli* and one-step purification of bifunctional hybrid proteins which bind maltose. Export of the Klenow polymerase into the periplasmic space. *Eur J Biochem*, 171, 541-549.
- Beher, D., Hesse, L., Masters, C.L. and Multhaup, G. (1996) Regulation of amyloid protein precursor (APP) binding to collagen and mapping of the binding sites on APP and collagen type I. *J Biol Chem*, 271, 1613-1620.
- Beyreuther, K., Multhaup, G. and Masters, C.L. (1996) Alzheimer's disease: genesis of amyloid. *Ciba Found Symp*, 199, 119-127; discussion 127-131.
- Borchardt, T., Camakaris, J., Cappai, R., Masters, C.L., Beyreuther, K. and Multhaup, G. (1999) Copper inhibits beta-amyloid production and stimulates the non-amyloidogenic pathway of amyloid-precursor-protein secretion. *Biochem J*, 344 Pt 2, 461-467.
- Borchelt, D.R., Thinakaran, G., Eckman, C.B., Lee, M.K., Davenport, F., Ratovitsky, T., Prada, C.M., Kim, G., Seekins, S., Yager, D., Slunt, H.H., Wang, R., Seeger, M., Levey, A.I., Gandy, S.E., Copeland, N.G., Jenkins, N.A., Price, D.L., Younkin, S.G. and Sisodia, S.S. (1996) Familial Alzheimer's disease-linked presenilin 1 variants elevate Abeta1-42/1-40 ratio in vitro and in vivo. *Neuron*, 17, 1005-1013.
- Brown, M.S., Ye, J., Rawson, R.B. and Goldstein, J.L. (2000) Regulated intramembrane proteolysis: a control mechanism conserved from bacteria to humans. *Cell*, 100, 391-398.

- Brunkan, A.L. and Goate, A.M. (2005) Presenilin function and gamma-secretase activity. *J Neurochem*, 93, 769-792.
- Bush, A.I., Beyreuther, K. and Masters, C.L. (1992) Beta A4 amyloid protein and its precursor in Alzheimer's disease. *Pharmacol Ther*, 56, 97-117.
- Bush, A.I., Multhaup, G., Moir, R.D., Williamson, T.G., Small, D.H., Rumble, B., Pollwein, P., Beyreuther, K. and Masters, C.L. (1993) A novel zinc(II) binding site modulates the function of the beta A4 amyloid protein precursor of Alzheimer's disease. *J Biol Chem*, 268, 16109-16112.
- Buxbaum, J.D., Gandy, S.E., Cicchetti, P., Ehrlich, M.E., Czernik, A.J., Fracasso, R.P., Ramabhadran, T.V., Unterbeck, A.J. and Greengard, P. (1990) Processing of Alzheimer beta/A4 amyloid precursor protein: modulation by agents that regulate protein phosphorylation. *Proc Natl Acad Sci U S A*, 87, 6003-6006.
- Campion, D., Dumanchin, C., Hannequin, D., Dubois, B., Belliard, S., Puel, M., Thomas-Anterion, C., Michon, A., Martin, C., Charbonnier, F., Raux, G., Camuzat, A., Penet, C., Mesnage, V., Martinez, M., Clerget-Darpoux, F., Brice, A. and Frebourg, T. (1999) Early-onset autosomal dominant Alzheimer disease: prevalence, genetic heterogeneity, and mutation spectrum. *Am J Hum Genet*, 65, 664-670.
- Cao, X. and Sudhof, T.C. (2004) Dissection of amyloid-beta precursor protein-dependent transcriptional transactivation. *J Biol Chem*, 279, 24601-24611.
- Cervantes, S., Gonzalez-Duarte, R. and Marfany, G. (2001) Homodimerization of presenilin N-terminal fragments is affected by mutations linked to Alzheimer's disease. *FEBS Lett*, 505, 81-86.
- Cervantes, S., Saura, C.A., Pomares, E., Gonzalez-Duarte, R. and Marfany, G. (2004) Functional implications of the presenilin dimerization: reconstitution of gamma-secretase activity by assembly of a catalytic site at the dimer interface of two catalytically inactive presenilins. *J Biol Chem*, 279, 36519-36529.
- Chandu, D., Huppert, S.S. and Kopan, R. (2006) Analysis of transmembrane domain mutants is consistent with sequential cleavage of Notch by gamma-secretase. *J Neurochem*, 96, 228-235.
- Chyung, J.H., Raper, D.M. and Selkoe, D.J. (2005) Gamma-secretase exists on the plasma membrane as an intact complex that accepts substrates and effects intramembrane cleavage. *J Biol Chem*, 280, 4383-4392.
- Citron, M., Oltersdorf, T., Haass, C., McConlogue, L., Hung, A.Y., Seubert, P., Vigo-Pelfrey, C., Lieberburg, I. and Selkoe, D.J. (1992) Mutation of the beta-amyloid precursor protein in familial Alzheimer's disease increases beta-protein production. *Nature*, 360, 672-674.
- Citron, M., Vigo-Pelfrey, C., Teplow, D.B., Miller, C., Schenk, D., Johnston, J., Winblad, B., Venizelos, N., Lannfelt, L. and Selkoe, D.J. (1994) Excessive production of amyloid beta-protein by peripheral cells of symptomatic and presymptomatic patients carrying the Swedish familial Alzheimer disease mutation. *Proc Natl Acad Sci U S A*, 91, 11993-11997.
- Cordy, J.M., Hussain, I., Dingwall, C., Hooper, N.M. and Turner, A.J. (2003) Exclusively targeting beta-secretase to lipid rafts by GPI-anchor addition up-regulates beta-site processing of the amyloid precursor protein. *Proc Natl Acad Sci U S A*, 100, 11735-11740.
- Coulson, E.J., Paliga, K., Beyreuther, K. and Masters, C.L. (2000) What the evolution

- of the amyloid protein precursor supergene family tells us about its function. *Neurochem Int*, 36, 175-184.
- Cruts, M., Backhovens, H., Wang, S.Y., Van Gassen, G., Theuns, J., De Jonghe, C.D., Wehnert, A., De Voecht, J., De Winter, G., Cras, P. and et al. (1995) Molecular genetic analysis of familial early-onset Alzheimer's disease linked to chromosome 14q24.3. *Hum Mol Genet*, 4, 2363-2371.
- Curtain, C.C., Ali, F., Volitakis, I., Cherny, R.A., Norton, R.S., Beyreuther, K., Barrow, C.J., Masters, C.L., Bush, A.I. and Barnham, K.J. (2001) Alzheimer's disease amyloid-beta binds copper and zinc to generate an allosterically ordered membrane-penetrating structure containing superoxide dismutase-like subunits. *J Biol Chem*, 276, 20466-20473.
- De Strooper, B. (2007) Loss-of-function presenilin mutations in Alzheimer disease. Talking Point on the role of presenilin mutations in Alzheimer disease. *EMBO Rep*, 8, 141-146.
- De Strooper, B., Saftig, P., Craessaerts, K., Vanderstichele, H., Guhde, G., Annaert, W., Von Figura, K. and Van Leuven, F. (1998) Deficiency of presenilin-1 inhibits the normal cleavage of amyloid precursor protein. *Nature*, 391, 387-390.
- Deibel, M.A., Ehmann, W.D. and Markesberry, W.R. (1996) Copper, iron, and zinc imbalances in severely degenerated brain regions in Alzheimer's disease: possible relation to oxidative stress. *J Neurol Sci*, 143, 137-142.
- Dovey, H.F., Suomensaari-Chrysler, S., Lieberburg, I., Sinha, S. and Keim, P.S. (1993) Cells with a familial Alzheimer's disease mutation produce authentic beta-peptide. *Neuroreport*, 4, 1039-1042.
- Dyrks, T., Dyrks, E., Masters, C. and Beyreuther, K. (1992) Membrane inserted APP fragments containing the beta A4 sequence of Alzheimer's disease do not aggregate. *FEBS Lett*, 309, 20-24.
- Edbauer, D., Winkler, E., Regula, J.T., Pesold, B., Steiner, H. and Haass, C. (2003) Reconstitution of gamma-secretase activity. *Nat Cell Biol*, 5, 486-488.
- Eggert, S., Paliga, K., Soba, P., Evin, G., Masters, C.L., Weidemann, A. and Beyreuther, K. (2004) The proteolytic processing of the amyloid precursor protein gene family members APLP-1 and APLP-2 involves alpha-, beta-, gamma-, and epsilon-like cleavages: modulation of APLP-1 processing by n-glycosylation. *J Biol Chem*, 279, 18146-18156.
- Ehehalt, R., Keller, P., Haass, C., Thiele, C. and Simons, K. (2003) Amyloidogenic processing of the Alzheimer beta-amyloid precursor protein depends on lipid rafts. *J Cell Biol*, 160, 113-123.
- Eriksen, J.L., Sagi, S.A., Smith, T.E., Weggen, S., Das, P., McLendon, D.C., Ozols, V.V., Jessing, K.W., Zavitz, K.H., Koo, E.H. and Golde, T.E. (2003) NSAIDs and enantiomers of flurbiprofen target gamma-secretase and lower Abeta 42 in vivo. *J Clin Invest*, 112, 440-449.
- Esch, F.S., Keim, P.S., Beattie, E.C., Blacher, R.W., Culwell, A.R., Oltersdorf, T., McClure, D. and Ward, P.J. (1990) Cleavage of amyloid beta peptide during constitutive processing of its precursor. *Science*, 248, 1122-1124.
- Esler, W.P., Kimberly, W.T., Ostaszewski, B.L., Diehl, T.S., Moore, C.L., Tsai, J.Y., Rahmati, T., Xia, W., Selkoe, D.J. and Wolfe, M.S. (2000) Transition-state

- analogue inhibitors of gamma-secretase bind directly to presenilin-1. *Nat Cell Biol*, 2, 428-434.
- Ferguson, K.M., Darling, P.J., Mohan, M.J., Macatee, T.L. and Lemmon, M.A. (2000) Extracellular domains drive homo- but not hetero-dimerization of erbB receptors. *Embo J*, 19, 4632-4643.
- Fleming, K.G. and Engelman, D.M. (2001) Specificity in transmembrane helix-helix interactions can define a hierarchy of stability for sequence variants. *Proc Natl Acad Sci U S A*, 98, 14340-14344.
- Furuya, H., Yasuda, M., Terasawa, K.J., Tanaka, K., Murai, H., Kira, J. and Ohyagi, Y. (2003) A novel mutation (L250V) in the presenilin 1 gene in a Japanese familial Alzheimer's disease with myoclonus and generalized convulsion. *J Neurol Sci*, 209, 75-77.
- Gao, Y. and Pimplikar, S.W. (2001) The gamma -secretase-cleaved C-terminal fragment of amyloid precursor protein mediates signaling to the nucleus. *Proc Natl Acad Sci U S A*, 98, 14979-14984.
- Glenner, G.G. and Wong, C.W. (1984) Alzheimer's disease: initial report of the purification and characterization of a novel cerebrovascular amyloid protein. *Biochem Biophys Res Commun*, 120, 885-890.
- Goedert, M. and Spillantini, M.G. (2006) A century of Alzheimer's disease. *Science*, 314, 777-781.
- Goldgaber, D., Lerman, M.I., McBride, O.W., Saffiotti, U. and Gajdusek, D.C. (1987) Characterization and chromosomal localization of a cDNA encoding brain amyloid of Alzheimer's disease. *Science*, 235, 877-880.
- Goutte, C., Tsunozaki, M., Hale, V.A. and Priess, J.R. (2002) APH-1 is a multipass membrane protein essential for the Notch signaling pathway in *Caenorhabditis elegans* embryos. *Proc Natl Acad Sci U S A*, 99, 775-779.
- Gurezka, R. and Langosch, D. (2001) In vitro selection of membrane-spanning leucine zipper protein-protein interaction motifs using POSSYCCAT. *J Biol Chem*, 276, 45580-45587.
- Haass, C., Hung, A.Y., Schlossmacher, M.G., Teplow, D.B. and Selkoe, D.J. (1993) beta-Amyloid peptide and a 3-kDa fragment are derived by distinct cellular mechanisms. *J Biol Chem*, 268, 3021-3024.
- Haass, C., Schlossmacher, M.G., Hung, A.Y., Vigo-Pelfrey, C., Mellon, A., Ostaszewski, B.L., Lieberburg, I., Koo, E.H., Schenk, D., Teplow, D.B. and et al. (1992) Amyloid beta-peptide is produced by cultured cells during normal metabolism. *Nature*, 359, 322-325.
- Hebert, S.S., Godin, C. and Levesque, G. (2003) Oligomerization of human presenilin-1 fragments. *FEBS Lett*, 550, 30-34.
- Hilbich, C., Monning, U., Grund, C., Masters, C.L. and Beyreuther, K. (1993) Amyloid-like properties of peptides flanking the epitope of amyloid precursor protein-specific monoclonal antibody 22C11. *J Biol Chem*, 268, 26571-26577.
- Hoe, H.S. and Rebeck, G.W. (2005) Regulation of ApoE receptor proteolysis by ligand binding. *Brain Res Mol Brain Res*, 137, 31-39.
- Hsia, A.Y., Masliah, E., McConlogue, L., Yu, G.Q., Tatsuno, G., Hu, K., Kholodenko, D., Malenka, R.C., Nicoll, R.A. and Mucke, L. (1999) Plaque-independent disruption

- of neural circuits in Alzheimer's disease mouse models. *Proc Natl Acad Sci U S A*, 96, 3228-3233.
- Huber, O., Kemler, R. and Langosch, D. (1999) Mutations affecting transmembrane segment interactions impair adhesiveness of E-cadherin. *J Cell Sci*, 112 (Pt 23), 4415-4423.
- Ida, N., Hartmann, T., Pantel, J., Schroder, J., Zerfass, R., Forstl, H., Sandbrink, R., Masters, C.L. and Beyreuther, K. (1996) Analysis of heterogeneous A4 peptides in human cerebrospinal fluid and blood by a newly developed sensitive Western blot assay. *J Biol Chem*, 271, 22908-22914.
- Itoh, R. and Fujiki, Y. (2006) Functional domains and dynamic assembly of the peroxin Pex14p, the entry site of matrix proteins. *J Biol Chem*, 281, 10196-10205.
- Iwatsubo, T., Odaka, A., Suzuki, N., Mizusawa, H., Nukina, N. and Ihara, Y. (1994) Visualization of A beta 42(43) and A beta 40 in senile plaques with end-specific A beta monoclonals: evidence that an initially deposited species is A beta 42(43). *Neuron*, 13, 45-53.
- Janssen, J.C., Beck, J.A., Campbell, T.A., Dickinson, A., Fox, N.C., Harvey, R.J., Houlden, H., Rossor, M.N. and Collinge, J. (2003) Early onset familial Alzheimer's disease: Mutation frequency in 31 families. *Neurology*, 60, 235-239.
- Jarrett, J.T. and Lansbury, P.T., Jr. (1993) Seeding „one-dimensional crystallization“ of amyloid: a pathogenic mechanism in Alzheimer's disease and scrapie? *Cell*, 73, 1055-1058.
- Kaether, C., Haass, C. and Steiner, H. (2006a) Assembly, trafficking and function of gamma-secretase. *Neurodegener Dis*, 3, 275-283.
- Kaether, C., Schmitt, S., Willem, M. and Haass, C. (2006b) Amyloid precursor protein and Notch intracellular domains are generated after transport of their precursors to the cell surface. *Traffic*, 7, 408-415.
- Kakuda, N., Funamoto, S., Yagishita, S., Takami, M., Osawa, S., Dohmae, N. and Ihara, Y. (2006) Equimolar production of amyloid beta-protein and amyloid precursor protein intracellular domain from beta-carboxyl-terminal fragment by gamma-secretase. *J Biol Chem*, 281, 14776-14786.
- Kamenetz, F., Tomita, T., Hsieh, H., Seabrook, G., Borchelt, D., Iwatsubo, T., Sisodia, S. and Malinow, R. (2003) APP processing and synaptic function. *Neuron*, 37, 925-937.
- Kamino, K., Orr, H.T., Payami, H., Wijsman, E.M., Alonso, M.E., Pulst, S.M., Anderson, L., O'Dahl, S., Nemens, E., White, J.A. and et al. (1992) Linkage and mutational analysis of familial Alzheimer disease kindreds for the APP gene region. *Am J Hum Genet*, 51, 998-1014.
- Kang, J., Lemaire, H.G., Unterbeck, A., Salbaum, J.M., Masters, C.L., Grzeschik, K.H., Multhaup, G., Beyreuther, K. and Muller-Hill, B. (1987) The precursor of Alzheimer's disease amyloid A4 protein resembles a cell-surface receptor. *Nature*, 325, 733-736.
- Kim, D.Y., Ingano, L.A., Carey, B.W., Pettingell, W.H. and Kovacs, D.M. (2005a) Presenilin/gamma-secretase-mediated cleavage of the voltage-gated sodium channel beta2-subunit regulates cell adhesion and migration. *J Biol Chem*, 280, 23251-23261.

- Kim, D.Y., Ingano, L.A. and Kovacs, D.M. (2002) Nectin-1alpha, an immunoglobulin-like receptor involved in the formation of synapses, is a substrate for presenilin/gamma-secretase-like cleavage. *J Biol Chem*, 277, 49976-49981.
- Kim, S., Jeon, T.J., Oberai, A., Yang, D., Schmidt, J.J. and Bowie, J.U. (2005b) Transmembrane glycine zippers: physiological and pathological roles in membrane proteins. *Proc Natl Acad Sci U S A*, 102, 14278-14283.
- Kimberly, W.T., Xia, W., Rahmati, T., Wolfe, M.S. and Selkoe, D.J. (2000) The transmembrane aspartates in presenilin 1 and 2 are obligatory for gamma-secretase activity and amyloid beta-protein generation. *J Biol Chem*, 275, 3173-3178.
- Kitaguchi, N., Takahashi, Y., Tokushima, Y., Shiojiri, S. and Ito, H. (1988) Novel precursor of Alzheimer's disease amyloid protein shows protease inhibitory activity. *Nature*, 331, 530-532.
- Kleiger, G., Grothe, R., Mallick, P. and Eisenberg, D. (2002) GXXXG and AXXXA: common alpha-helical interaction motifs in proteins, particularly in extremophiles. *Biochemistry*, 41, 5990-5997.
- Klein, W.L., Krafft, G.A. and Finch, C.E. (2001) Targeting small Abeta oligomers: the solution to an Alzheimer's disease conundrum? *Trends Neurosci*, 24, 219-224.
- Kolmar, H., Frisch, C., Kleemann, G., Gotze, K., Stevens, F.J. and Fritz, H.J. (1994) Dimerization of Bence Jones proteins: linking the rate of transcription from an Escherichia coli promoter to the association constant of REIV. *Biol Chem Hoppe Seyler*, 375, 61-70.
- Kornilova, A.Y., Bihel, F., Das, C. and Wolfe, M.S. (2005) The initial substrate-binding site of gamma-secretase is located on presenilin near the active site. *Proc Natl Acad Sci U S A*, 102, 3230-3235.
- Kukar, T., Murphy, M.P., Eriksen, J.L., Sagi, S.A., Weggen, S., Smith, T.E., Ladd, T., Khan, M.A., Kache, R., Beard, J., Dodson, M., Merit, S., Ozols, V.V., Anastasiadis, P.Z., Das, P., Fauq, A., Koo, E.H. and Golde, T.E. (2005) Diverse compounds mimic Alzheimer disease-causing mutations by augmenting Abeta42 production. *Nat Med*, 11, 545-550.
- Lambert, M.P., Barlow, A.K., Chromy, B.A., Edwards, C., Freed, R., Liosatos, M., Morgan, T.E., Rozovsky, I., Trommer, B., Viola, K.L., Wals, P., Zhang, C., Finch, C.E., Krafft, G.A. and Klein, W.L. (1998) Diffusible, nonfibrillar ligands derived from Abeta1-42 are potent central nervous system neurotoxins. *Proc Natl Acad Sci U S A*, 95, 6448-6453.
- Lammich, S., Kojro, E., Postina, R., Gilbert, S., Pfeiffer, R., Jasionowski, M., Haass, C. and Fahrenholz, F. (1999) Constitutive and regulated alpha-secretase cleavage of Alzheimer's amyloid precursor protein by a disintegrin metalloprotease. *Proc Natl Acad Sci U S A*, 96, 3922-3927.
- Langosch, D., Brosig, B., Kolmar, H. and Fritz, H.J. (1996) Dimerisation of the glycophorin A transmembrane segment in membranes probed with the ToxR transcription activator. *J Mol Biol*, 263, 525-530.
- LaVoie, M.J., Fraering, P.C., Ostaszewski, B.L., Ye, W., Kimberly, W.T., Wolfe, M.S. and Selkoe, D.J. (2003) Assembly of the gamma-secretase complex involves early formation of an intermediate subcomplex of Aph-1 and nicastrin. *J Biol Chem*, 278,

- 37213-37222.
- Lee, S.F., Shah, S., Yu, C., Wigley, W.C., Li, H., Lim, M., Pedersen, K., Han, W., Thomas, P., Lundkvist, J., Hao, Y.H. and Yu, G. (2004) A conserved GXXXG motif in APH-1 is critical for assembly and activity of the gamma-secretase complex. *J Biol Chem*, 279, 4144-4152.
- Levine, H., 3rd. (1995) Soluble multimeric Alzheimer beta(1-40) pre-amyloid complexes in dilute solution. *Neurobiol Aging*, 16, 755-764.
- Levy, E., Carman, M.D., Fernandez-Madrid, I.J., Power, M.D., Lieberburg, I., van Duinen, S.G., Bots, G.T., Luyendijk, W. and Frangione, B. (1990) Mutation of the Alzheimer's disease amyloid gene in hereditary cerebral hemorrhage, Dutch type. *Science*, 248, 1124-1126.
- Levy-Lahad, E., Wijsman, E.M., Nemens, E., Anderson, L., Goddard, K.A., Weber, J.L., Bird, T.D. and Schellenberg, G.D. (1995) A familial Alzheimer's disease locus on chromosome 1. *Science*, 269, 970-973.
- Li, Y.M., Lai, M.T., Xu, M., Huang, Q., DiMuzio-Mower, J., Sardana, M.K., Shi, X.P., Yin, K.C., Shafer, J.A. and Gardell, S.J. (2000a) Presenilin 1 is linked with gamma-secretase activity in the detergent solubilized state. *Proc Natl Acad Sci U S A*, 97, 6138-6143.
- Li, Y.M., Xu, M., Lai, M.T., Huang, Q., Castro, J.L., DiMuzio-Mower, J., Harrison, T., Lellis, C., Nadin, A., Neduvvelil, J.G., Register, R.B., Sardana, M.K., Shearman, M.S., Smith, A.L., Shi, X.P., Yin, K.C., Shafer, J.A. and Gardell, S.J. (2000b) Photoactivated gamma-secretase inhibitors directed to the active site covalently label presenilin 1. *Nature*, 405, 689-694.
- Li, Z.W., Stark, G., Gotz, J., Rulicke, T., Gschwind, M., Huber, G., Muller, U. and Weissmann, C. (1996) Generation of mice with a 200-kb amyloid precursor protein gene deletion by Cre recombinase-mediated site-specific recombination in embryonic stem cells. *Proc Natl Acad Sci U S A*, 93, 6158-6162.
- Lichtenthaler, S.F., Beher, D., Grimm, H.S., Wang, R., Shearman, M.S., Masters, C.L. and Beyreuther, K. (2002) The intramembrane cleavage site of the amyloid precursor protein depends on the length of its transmembrane domain. *Proc Natl Acad Sci U S A*, 99, 1365-1370.
- Liu, W., Crocker, E., Zhang, W., Elliott, J.I., Luy, B., Li, H., Aimoto, S. and Smith, S.O. (2005) Structural role of glycine in amyloid fibrils formed from transmembrane alpha-helices. *Biochemistry*, 44, 3591-3597.
- Lue, L.F., Kuo, Y.M., Roher, A.E., Brachova, L., Shen, Y., Sue, L., Beach, T., Kurth, J.H., Rydel, R.E. and Rogers, J. (1999) Soluble amyloid beta peptide concentration as a predictor of synaptic change in Alzheimer's disease. *Am J Pathol*, 155, 853-862.
- Luo, W.J., Wang, H., Li, H., Kim, B.S., Shah, S., Lee, H.J., Thinakaran, G., Kim, T.W., Yu, G. and Xu, H. (2003) PEN-2 and APH-1 coordinately regulate proteolytic processing of presenilin 1. *J Biol Chem*, 278, 7850-7854.
- MacKenzie, K.R., Prestegard, J.H. and Engelman, D.M. (1997) A transmembrane helix dimer: structure and implications. *Science*, 276, 131-133.
- Marambaud, P., Shioi, J., Serban, G., Georgakopoulos, A., Sarner, S., Nagy, V., Baki, L., Wen, P., Efthimiopoulos, S., Shao, Z., Wisniewski, T. and Robakis, N.K. (2002) A presenilin-1/gamma-secretase cleavage releases the E-cadherin intracellular

- domain and regulates disassembly of adherens junctions. *Embo J*, 21, 1948-1956.
- Marchesi, V.T. (2005) An alternative interpretation of the amyloid Abeta hypothesis with regard to the pathogenesis of Alzheimer's disease. *Proc Natl Acad Sci U S A*, 102, 9093-9098.
- Marlow, L., Cain, M., Pappolla, M.A. and Sambamurti, K. (2003) Beta-secretase processing of the Alzheimer's amyloid protein precursor (APP). *J Mol Neurosci*, 20, 233-239.
- Masters, C.L. and Beyreuther, K.T. (1989) The pathology of the amyloid A4 precursor of Alzheimer's disease. *Ann Med*, 21, 89-90.
- Masters, C.L., Simms, G., Weinman, N.A., Multhaup, G., McDonald, B.L. and Beyreuther, K. (1985) Amyloid plaque core protein in Alzheimer disease and Down syndrome. *Proc Natl Acad Sci U S A*, 82, 4245-4249.
- McClain, M.S., Iwamoto, H., Cao, P., Vinion-Dubiel, A.D., Li, Y., Szabo, G., Shao, Z. and Cover, T.L. (2003) Essential role of a GXXXG motif for membrane channel formation by Helicobacter pylori vacuolating toxin. *J Biol Chem*, 278, 12101-12108.
- McGeer, P.L., Schulzer, M. and McGeer, E.G. (1996) Arthritis and anti-inflammatory agents as possible protective factors for Alzheimer's disease: a review of 17 epidemiologic studies. *Neurology*, 47, 425-432.
- McLean, C.A., Cherny, R.A., Fraser, F.W., Fuller, S.J., Smith, M.J., Beyreuther, K., Bush, A.I. and Masters, C.L. (1999) Soluble pool of Abeta amyloid as a determinant of severity of neurodegeneration in Alzheimer's disease. *Ann Neurol*, 46, 860-866.
- Mendrola, J.M., Berger, M.B., King, M.C. and Lemmon, M.A. (2002) The single transmembrane domains of ErbB receptors self-associate in cell membranes. *J Biol Chem*, 277, 4704-4712.
- Menendez-Gonzalez, M., Perez-Pinera, P., Martinez-Rivera, M., Calatayud, M.T. and Blazquez Menes, B. (2005) APP processing and the APP-KPI domain involvement in the amyloid cascade. *Neurodegener Dis*, 2, 277-283.
- Miyahara, M., Nakanishi, H., Takahashi, K., Satoh-Horikawa, K., Tachibana, K. and Takai, Y. (2000) Interaction of nectin with afadin is necessary for its clustering at cell-cell contact sites but not for its cis dimerization or trans interaction. *J Biol Chem*, 275, 613-618.
- Mucke, L., Masliah, E., Yu, G.Q., Mallory, M., Rockenstein, E.M., Tatsuno, G., Hu, K., Kholodenko, D., Johnson-Wood, K. and McConlogue, L. (2000) High-level neuronal expression of abeta 1-42 in wild-type human amyloid protein precursor transgenic mice: synaptotoxicity without plaque formation. *J Neurosci*, 20, 4050-4058.
- Mullan, M., Houlden, H., Windelspecht, M., Fidani, L., Lombardi, C., Diaz, P., Rossor, M., Crook, R., Hardy, J., Duff, K. and et al. (1992) A locus for familial early-onset Alzheimer's disease on the long arm of chromosome 14, proximal to the alpha 1-antichymotrypsin gene. *Nat Genet*, 2, 340-342.
- Multhaup, G. (1994) Identification and regulation of the high affinity binding site of the Alzheimer's disease amyloid protein precursor (APP) to glycosaminoglycans. *Biochimie*, 76, 304-311.
- Multhaup, G., Schlicksupp, A., Hesse, L., Beher, D., Ruppert, T., Masters, C.L. and Beyreuther, K. (1996) The amyloid precursor protein of Alzheimer's disease in the

- reduction of copper(II) to copper(I). *Science*, 271, 1406-1409.
- Munter, L.M., Voigt, P., Harmeier, A., Kaden, D., Gottschalk, K.E., Weise, C., Pipkorn, R., Schaefer, M., Langosch, D. and Multhaup, G. (2007) GxxxG motifs within the amyloid precursor protein transmembrane sequence are critical for the etiology of Abeta42. *Embo J*.
- Murphy, M.P., Hickman, L.J., Eckman, C.B., Uljon, S.N., Wang, R. and Golde, T.E. (1999) gamma-Secretase, evidence for multiple proteolytic activities and influence of membrane positioning of substrate on generation of amyloid beta peptides of varying length. *J Biol Chem*, 274, 11914-11923.
- Ni, C.Y., Murphy, M.P., Golde, T.E. and Carpenter, G. (2001) gamma -Secretase cleavage and nuclear localization of ErbB-4 receptor tyrosine kinase. *Science*, 294, 2179-2181.
- Niimura, M., Issoo, N., Takasugi, N., Tsuruoka, M., Ui-Tei, K., Saigo, K., Morohashi, Y., Tomita, T. and Iwatsubo, T. (2005) Aph-1 contributes to the stabilization and trafficking of the gamma-secretase complex through mechanisms involving intermolecular and intramolecular interactions. *J Biol Chem*, 280, 12967-12975.
- Nilsberth, C., Westlind-Danielsson, A., Eckman, C.B., Condron, M.M., Axelman, K., Forsell, C., Stenh, C., Luthman, J., Teplow, D.B., Younkin, S.G., Naslund, J. and Lannfelt, L. (2001) The 'Arctic' APP mutation (E693G) causes Alzheimer's disease by enhanced Abeta protofibril formation. *Nat Neurosci*, 4, 887-893.
- Oakley, H., Cole, S.L., Logan, S., Maus, E., Shao, P., Craft, J., Guillozet-Bongaarts, A., Ohno, M., Disterhoft, J., Van Eldik, L., Berry, R. and Vassar, R. (2006) Intraneuronal beta-amyloid aggregates, neurodegeneration, and neuron loss in transgenic mice with five familial Alzheimer's disease mutations: potential factors in amyloid plaque formation. *J Neurosci*, 26, 10129-10140.
- Okochi, M., Steiner, H., Fukumori, A., Tanii, H., Tomita, T., Tanaka, T., Iwatsubo, T., Kudo, T., Takeda, M. and Haass, C. (2002) Presenilins mediate a dual intramembranous gamma-secretase cleavage of Notch-1. *Embo J*, 21, 5408-5416.
- Overton, M.C., Chinault, S.L. and Blumer, K.J. (2003) Oligomerization, biogenesis, and signaling is promoted by a glycophorin A-like dimerization motif in transmembrane domain 1 of a yeast G protein-coupled receptor. *J Biol Chem*, 278, 49369-49377.
- Pasternak, S.H., Callahan, J.W. and Mahuran, D.J. (2004) The role of the endosomal/lysosomal system in amyloid-beta production and the pathophysiology of Alzheimer's disease: reexamining the spatial paradox from a lysosomal perspective. *J Alzheimers Dis*, 6, 53-65.
- Pike, C.J., Walencewicz-Wasserman, A.J., Kosmoski, J., Cribbs, D.H., Glabe, C.G. and Cotman, C.W. (1995) Structure-activity analyses of beta-amyloid peptides: contributions of the beta 25-35 region to aggregation and neurotoxicity. *J Neurochem*, 64, 253-265.
- Poirier, J., Davignon, J., Bouthillier, D., Kogan, S., Bertrand, P. and Gauthier, S. (1993) Apolipoprotein E polymorphism and Alzheimer's disease. *Lancet*, 342, 697-699.
- Polgar, O., Robey, R.W., Morisaki, K., Dean, M., Michejda, C., Sauna, Z.E., Ambudkar, S.V., Tarasova, N. and Bates, S.E. (2004) Mutational analysis of ABCG2: role of the GXXXG motif. *Biochemistry*, 43, 9448-9456.
- Qi-Takahara, Y., Morishima-Kawashima, M., Tanimura, Y., Dolios, G., Hirotani, N.,

- Horikoshi, Y., Kametani, F., Maeda, M., Saido, T.C., Wang, R. and Ihara, Y. (2005) Longer forms of amyloid beta protein: implications for the mechanism of intramembrane cleavage by gamma-secretase. *J Neurosci*, 25, 436-445.
- Qiu, W.Q., Ferreira, A., Miller, C., Koo, E.H. and Selkoe, D.J. (1995) Cell-surface beta-amyloid precursor protein stimulates neurite outgrowth of hippocampal neurons in an isoform-dependent manner. *J Neurosci*, 15, 2157-2167.
- Robakis, N.K., Wisniewski, H.M., Jenkins, E.C., Devine-Gage, E.A., Houck, G.E., Yao, X.L., Ramakrishna, N., Wolfe, G., Silverman, W.P. and Brown, W.T. (1987) Chromosome 21q21 sublocalisation of gene encoding beta-amyloid peptide in cerebral vessels and neuritic (senile) plaques of people with Alzheimer disease and Down syndrome. *Lancet*, 1, 384-385.
- Rogaeva, E.A., Fafel, K.C., Song, Y.Q., Medeiros, H., Sato, C., Liang, Y., Richard, E., Rogaev, E.I., Frommelt, P., Sadovnick, A.D., Meschino, W., Rockwood, K., Boss, M.A., Mayeux, R. and St George-Hyslop, P. (2001) Screening for PS1 mutations in a referral-based series of AD cases: 21 novel mutations. *Neurology*, 57, 621-625.
- Rossjohn, J., Cappai, R., Feil, S.C., Henry, A., McKinstry, W.J., Galatis, D., Hesse, L., Multhaup, G., Beyreuther, K., Masters, C.L. and Parker, M.W. (1999) Crystal structure of the N-terminal, growth factor-like domain of Alzheimer amyloid precursor protein. *Nat Struct Biol*, 6, 327-331.
- Saitoh, T., Sundsmo, M., Roch, J.M., Kimura, N., Cole, G., Schubert, D., Oltersdorf, T. and Schenk, D.B. (1989) Secreted form of amyloid beta protein precursor is involved in the growth regulation of fibroblasts. *Cell*, 58, 615-622.
- Sambrook, J., Fritsch, E.F., Maniatis, T. (1989) Molecular cloning: a laboratory manual. New York, Cold Spring Harbor Laboratory Press.
- Sanger, F., Nicklen, S. and Coulson, A.R. (1977) DNA sequencing with chain-terminating inhibitors. *Proc Natl Acad Sci U S A*, 74, 5463-5467.
- Sardi, S.P., Murtie, J., Koirala, S., Patten, B.A. and Corfas, G. (2006) Presenilin-dependent ErbB4 nuclear signaling regulates the timing of astrogenesis in the developing brain. *Cell*, 127, 185-197.
- Saunders, A.M., Strittmatter, W.J., Schmeichel, D., George-Hyslop, P.H., Pericak-Vance, M.A., Joo, S.H., Rosi, B.L., Gusella, J.F., Crapper-MacLachlan, D.R., Alberts, M.J. and et al. (1993) Association of apolipoprotein E allele epsilon 4 with late-onset familial and sporadic Alzheimer's disease. *Neurology*, 43, 1467-1472.
- Schagger, H. and von Jagow, G. (1987) Tricine-sodium dodecyl sulfate-polyacrylamide gel electrophoresis for the separation of proteins in the range from 1 to 100 kDa. *Anal Biochem*, 166, 368-379.
- Scheuermann, S., Hamsch, B., Hesse, L., Stumm, J., Schmidt, C., Beher, D., Bayer, T.A., Beyreuther, K. and Multhaup, G. (2001) Homodimerization of amyloid precursor protein and its implication in the amyloidogenic pathway of Alzheimer's disease. *J Biol Chem*, 276, 33923-33929.
- Schmeichel, A., Strauss, M., Schlicksupp, A., Pipkorn, R., Haass, C., Bayer, T.A. and Multhaup, G. (2004) Human BACE forms dimers and colocalizes with APP. *J Biol Chem*, 279, 39710-39717.
- Schmeichel, A., Zentgraf, H., Scheuermann, S., Fritz, G., Pipkorn, R., Reed, J., Beyreuther, K., Bayer, T.A. and Multhaup, G. (2003) Alzheimer beta-amyloid

- homodimers facilitate A beta fibrillization and the generation of conformational antibodies. *J Biol Chem*, 278, 35317-35324.
- Schroeter, E.H., Ilagan, M.X., Brunkan, A.L., Hecimovic, S., Li, Y.M., Xu, M., Lewis, H.D., Saxena, M.T., De Strooper, B., Coonrod, A., Tomita, T., Iwatsubo, T., Moore, C.L., Goate, A., Wolfe, M.S., Shearman, M. and Kopan, R. (2003) A presenilin dimer at the core of the gamma-secretase enzyme: insights from parallel analysis of Notch 1 and APP proteolysis. *Proc Natl Acad Sci U S A*, 100, 13075-13080.
- Schubert, D., LaCorbiere, M., Saitoh, T. and Cole, G. (1989) Characterization of an amyloid beta precursor protein that binds heparin and contains tyrosine sulfate. *Proc Natl Acad Sci U S A*, 86, 2066-2069.
- Schulz, J.G., Annaert, W., Vandekerckhove, J., Zimmermann, P., De Strooper, B. and David, G. (2003) Syndecan 3 intramembrane proteolysis is presenilin/gamma-secretase-dependent and modulates cytosolic signaling. *J Biol Chem*, 278, 48651-48657.
- Seeger, M., Nordstedt, C., Petanceska, S., Kovacs, D.M., Gouras, G.K., Hahne, S., Fraser, P., Levesque, L., Czernik, A.J., George-Hyslop, P.S., Sisodia, S.S., Thinakaran, G., Tanzi, R.E., Greengard, P. and Gandy, S. (1997) Evidence for phosphorylation and oligomeric assembly of presenilin 1. *Proc Natl Acad Sci U S A*, 94, 5090-5094.
- Selkoe, D.J. (2004) Alzheimer disease: mechanistic understanding predicts novel therapies. *Ann Intern Med*, 140, 627-638.
- Senes, A., Engel, D.E. and DeGrado, W.F. (2004) Folding of helical membrane proteins: the role of polar, GxxxG-like and proline motifs. *Curr Opin Struct Biol*, 14, 465-479.
- Senes, A., Gerstein, M. and Engelman, D.M. (2000) Statistical analysis of amino acid patterns in transmembrane helices: the GxxxG motif occurs frequently and in association with beta-branched residues at neighboring positions. *J Mol Biol*, 296, 921-936.
- Senes, A., Ubarretxena-Belandia, I. and Engelman, D.M. (2001) The Calpha ---H.O hydrogen bond: a determinant of stability and specificity in transmembrane helix interactions. *Proc Natl Acad Sci U S A*, 98, 9056-9061.
- Shah, S., Lee, S.F., Tabuchi, K., Hao, Y.H., Yu, C., LaPlant, Q., Ball, H., Dann, C.E., 3rd, Sudhof, T. and Yu, G. (2005) Nicastrin functions as a gamma-secretase-substrate receptor. *Cell*, 122, 435-447.
- Shen, J., Bronson, R.T., Chen, D.F., Xia, W., Selkoe, D.J. and Tonegawa, S. (1997) Skeletal and CNS defects in Presenilin-1-deficient mice. *Cell*, 89, 629-639.
- Sherrington, R., Rogaev, E.I., Liang, Y., Rogaeva, E.A., Levesque, G., Ikeda, M., Chi, H., Lin, C., Li, G., Holman, K. and et al. (1995) Cloning of a gene bearing missense mutations in early-onset familial Alzheimer's disease. *Nature*, 375, 754-760.
- Shobab, L.A., Hsiung, G.Y. and Feldman, H.H. (2005) Cholesterol in Alzheimer's disease. *Lancet Neurol*, 4, 841-852.
- Siekhaus, A. (2007) Charakterisierung der Enzym-Substrat-Erkennung im g-Sekretase Komplex. Diplomarbeit bei Prof. Dr. Gerd Multhaup, FU Berlin.
- Sisodia, S.S. (1999) Alzheimer's disease: perspectives for the new millennium. *J Clin Invest*, 104, 1169-1170.
- Slack, B.E., Ma, L.K. and Seah, C.C. (2001) Constitutive shedding of the amyloid

- precursor protein ectodomain is up-regulated by tumour necrosis factor-alpha converting enzyme. *Biochem J*, 357, 787-794.
- Small, D.H., Nurcombe, V., Reed, G., Clarris, H., Moir, R., Beyreuther, K. and Masters, C.L. (1994) A heparin-binding domain in the amyloid protein precursor of Alzheimer's disease is involved in the regulation of neurite outgrowth. *J Neurosci*, 14, 2117-2127.
- Snyder, E.M., Nong, Y., Almeida, C.G., Paul, S., Moran, T., Choi, E.Y., Nairn, A.C., Salter, M.W., Lombroso, P.J., Gouras, G.K. and Greengard, P. (2005) Regulation of NMDA receptor trafficking by amyloid-beta. *Nat Neurosci*, 8, 1051-1058.
- Snyder, S.W., Ladror, U.S., Wade, W.S., Wang, G.T., Barrett, L.W., Matayoshi, E.D., Huffaker, H.J., Krafft, G.A. and Holzman, T.F. (1994) Amyloid-beta aggregation: selective inhibition of aggregation in mixtures of amyloid with different chain lengths. *Biophys J*, 67, 1216-1228.
- Soba, P., Eggert, S., Wagner, K., Zentgraf, H., Siehl, K., Kreger, S., Lower, A., Langer, A., Merdes, G., Paro, R., Masters, C.L., Muller, U., Kins, S. and Beyreuther, K. (2006) Homo- and hetero-dimerization of APP family members promotes intercellular adhesion. *Embo J*, 25, 653.
- Spasic, D., Tolia, A., Dillen, K., Baert, V., De Strooper, B., Vrijens, S. and Annaert, W. (2006) Presenilin-1 maintains a nine-transmembrane topology throughout the secretory pathway. *J Biol Chem*, 281, 26569-26577.
- St George-Hyslop, P.H. (2000) Genetic factors in the genesis of Alzheimer's disease. *Ann N Y Acad Sci*, 924, 1-7.
- Stenh, C., Nilsberth, C., Hammarback, J., Engvall, B., Naslund, J. and Lannfelt, L. (2002) The Arctic mutation interferes with processing of the amyloid precursor protein. *Neuroreport*, 13, 1857-1860.
- Struhl, G. and Adachi, A. (2000) Requirements for presenilin-dependent cleavage of notch and other transmembrane proteins. *Mol Cell*, 6, 625-636.
- Suzuki, N., Cheung, T.T., Cai, X.D., Odaka, A., Otvos, L., Jr., Eckman, C., Golde, T.E. and Younkin, S.G. (1994) An increased percentage of long amyloid beta protein secreted by familial amyloid beta protein precursor (beta APP717) mutants. *Science*, 264, 1336-1340.
- Takahashi, Y., Hayashi, I., Tominari, Y., Rikimaru, K., Morohashi, Y., Kan, T., Natsugari, H., Fukuyama, T., Tomita, T. and Iwatsubo, T. (2003) Sulindac sulfide is a noncompetitive gamma-secretase inhibitor that preferentially reduces Abeta 42 generation. *J Biol Chem*, 278, 18664-18670.
- Tang, K., Wang, C., Shen, C., Sheng, S., Ravid, R. and Jing, N. (2003) Identification of a novel alternative splicing isoform of human amyloid precursor protein gene, APP639. *Eur J Neurosci*, 18, 102-108.
- Tanzi, R.E., McClatchey, A.I., Lamperti, E.D., Villa-Komaroff, L., Gusella, J.F. and Neve, R.L. (1988) Protease inhibitor domain encoded by an amyloid protein precursor mRNA associated with Alzheimer's disease. *Nature*, 331, 528-530.
- Tanzi, R.E., Moir, R.D. and Wagner, S.L. (2004) Clearance of Alzheimer's Abeta peptide: the many roads to perdition. *Neuron*, 43, 605-608.
- Thinakaran, G., Regard, J.B., Bouton, C.M., Harris, C.L., Price, D.L., Borchelt, D.R. and Sisodia, S.S. (1998) Stable association of presenilin derivatives and absence of

- presenilin interactions with APP. *Neurobiol Dis*, 4, 438-453.
- Tian, G., Ghanekar, S.V., Aharony, D., Shenvi, A.B., Jacobs, R.T., Liu, X. and Greenberg, B.D. (2003) The mechanism of gamma-secretase: multiple inhibitor binding sites for transition state analogs and small molecule inhibitors. *J Biol Chem*, 278, 28968-28975.
- Troyanovsky, R.B., Sokolov, E. and Troyanovsky, S.M. (2003) Adhesive and lateral E-cadherin dimers are mediated by the same interface. *Mol Cell Biol*, 23, 7965-7972.
- Vassar, R., Bennett, B.D., Babu-Khan, S., Kahn, S., Mendiaz, E.A., Denis, P., Teplow, D.B., Ross, S., Amarante, P., Loeloff, R., Luo, Y., Fisher, S., Fuller, J., Edenson, S., Lile, J., Jarosinski, M.A., Biere, A.L., Curran, E., Burgess, T., Louis, J.C., Collins, F., Treanor, J., Rogers, G. and Citron, M. (1999) Beta-secretase cleavage of Alzheimer's amyloid precursor protein by the transmembrane aspartic protease BACE. *Science*, 286, 735-741.
- Vooijs, M., Schroeter, E.H., Pan, Y., Blandford, M. and Kopan, R. (2004) Ectodomain shedding and intramembrane cleavage of mammalian Notch proteins is not regulated through oligomerization. *J Biol Chem*, 279, 50864-50873.
- Walsh, D.M., Klyubin, I., Fadeeva, J.V., Cullen, W.K., Anwyl, R., Wolfe, M.S., Rowan, M.J. and Selkoe, D.J. (2002) Naturally secreted oligomers of amyloid beta protein potently inhibit hippocampal long-term potentiation in vivo. *Nature*, 416, 535-539.
- Walsh, D.M., Lomakin, A., Benedek, G.B., Condron, M.M. and Teplow, D.B. (1997) Amyloid beta-protein fibrillogenesis. Detection of a protofibrillar intermediate. *J Biol Chem*, 272, 22364-22372.
- Walsh, D.M., Tseng, B.P., Rydel, R.E., Podlisny, M.B. and Selkoe, D.J. (2000) The oligomerization of amyloid beta-protein begins intracellularly in cells derived from human brain. *Biochemistry*, 39, 10831-10839.
- Walter, J., Capell, A., Hung, A.Y., Langen, H., Schnolzer, M., Thinakaran, G., Sisodia, S.S., Selkoe, D.J. and Haass, C. (1997) Ectodomain phosphorylation of beta-amyloid precursor protein at two distinct cellular locations. *J Biol Chem*, 272, 1896-1903.
- Wang, Y. and Ha, Y. (2004) The X-ray structure of an antiparallel dimer of the human amyloid precursor protein E2 domain. *Mol Cell*, 15, 343-353.
- Weggen, S., Eriksen, J.L., Das, P., Sagi, S.A., Wang, R., Pietrzik, C.U., Findlay, K.A., Smith, T.E., Murphy, M.P., Bulter, T., Kang, D.E., Marquez-Sterling, N., Golde, T.E. and Koo, E.H. (2001) A subset of NSAIDs lower amyloidogenic Abeta42 independently of cyclooxygenase activity. *Nature*, 414, 212-216.
- Weggen, S., Eriksen, J.L., Sagi, S.A., Pietrzik, C.U., Ozols, V., Fauq, A., Golde, T.E. and Koo, E.H. (2003) Evidence that nonsteroidal anti-inflammatory drugs decrease amyloid beta 42 production by direct modulation of gamma-secretase activity. *J Biol Chem*, 278, 31831-31837.
- Weidemann, A., Eggert, S., Reinhard, F.B., Vogel, M., Paliga, K., Baier, G., Masters, C.L., Beyreuther, K. and Evin, G. (2002) A novel epsilon-cleavage within the transmembrane domain of the Alzheimer amyloid precursor protein demonstrates homology with Notch processing. *Biochemistry*, 41, 2825-2835.
- Weidemann, A., Konig, G., Bunke, D., Fischer, P., Salbaum, J.M., Masters, C.L. and Beyreuther, K. (1989) Identification, biogenesis, and localization of precursors of Alzheimer's disease A4 amyloid protein. *Cell*, 57, 115-126.

- Wertkin, A.M., Turner, R.S., Pleasure, S.J., Golde, T.E., Younkin, S.G., Trojanowski, J.Q. and Lee, V.M. (1993) Human neurons derived from a teratocarcinoma cell line express solely the 695-amino acid amyloid precursor protein and produce intracellular beta-amyloid or A4 peptides. *Proc Natl Acad Sci U S A*, 90, 9513-9517.
- Westmeyer, G.G., Willem, M., Lichtenthaler, S.F., Lurman, G., Multhaup, G., Assfalg-Machleidt, I., Reiss, K., Saftig, P. and Haass, C. (2004) Dimerization of beta-site beta-amyloid precursor protein-cleaving enzyme. *J Biol Chem*, 279, 53205-53212.
- Wilson, R.S., Mendes De Leon, C.F., Barnes, L.L., Schneider, J.A., Bienias, J.L., Evans, D.A. and Bennett, D.A. (2002) Participation in cognitively stimulating activities and risk of incident Alzheimer disease. *Jama*, 287, 742-748.
- Wolfe, M.S. (2007) When loss is gain: reduced presenilin proteolytic function leads to increased Abeta42/Abeta40. Talking Point on the role of presenilin mutations in Alzheimer disease. *EMBO Rep*, 8, 136-140.
- Wolfe, M.S., Xia, W., Ostaszewski, B.L., Diehl, T.S., Kimberly, W.T. and Selkoe, D.J. (1999) Two transmembrane aspartates in presenilin-1 required for presenilin endoproteolysis and gamma-secretase activity. *Nature*, 398, 513-517.
- Wong, H.K., Sakurai, T., Oyama, F., Kaneko, K., Wada, K., Miyazaki, H., Kurosawa, M., De Strooper, B., Saftig, P. and Nukina, N. (2005) beta Subunits of voltage-gated sodium channels are novel substrates of beta-site amyloid precursor protein-cleaving enzyme (BACE1) and gamma-secretase. *J Biol Chem*, 280, 23009-23017.
- Xu, Y., Shen, J., Luo, X., Zhu, W., Chen, K., Ma, J. and Jiang, H. (2005) Conformational transition of amyloid beta-peptide. *Proc Natl Acad Sci U S A*, 102, 5403-5407.
- Yamasaki, A., Eimer, S., Okochi, M., Smialowska, A., Kaether, C., Baumeister, R., Haass, C. and Steiner, H. (2006) The GxGD motif of presenilin contributes to catalytic function and substrate identification of gamma-secretase. *J Neurosci*, 26, 3821-3828.
- Younkin, S.G. (1998) The role of A beta 42 in Alzheimer's disease. *J Physiol Paris*, 92, 289-292.
- Zhao, G., Cui, M.Z., Mao, G., Dong, Y., Tan, J., Sun, L. and Xu, X. (2005) gamma-Cleavage is dependent on zeta-cleavage during the proteolytic processing of amyloid precursor protein within its transmembrane domain. *J Biol Chem*, 280, 37689-37697.
- Zhao, G., Mao, G., Tan, J., Dong, Y., Cui, M.Z., Kim, S.H. and Xu, X. (2004) Identification of a new presenilin-dependent zeta-cleavage site within the transmembrane domain of amyloid precursor protein. *J Biol Chem*, 279, 50647-50650.
- Zheng, H., Jiang, M., Trumbauer, M.E., Sirinathsinghji, D.J., Hopkins, R., Smith, D.W., Heavens, R.P., Dawson, G.R., Boyce, S., Conner, M.W., Stevens, K.A., Slunt, H.H., Sisoda, S.S., Chen, H.Y. and Van der Ploeg, L.H. (1995) beta-Amyloid precursor protein-deficient mice show reactive gliosis and decreased locomotor activity. *Cell*, 81, 525-531.