
6 REFERENCES

- Airaksinen MS, Koltzenburg M, Lewin GR, Masu Y, Helbig C, Wolf E, Brem G, Toyka KV, Thoenen H, Meyer M (1996) Specific subtypes of cutaneous mechanoreceptors require neurotrophin-3 following peripheral target innervation. *Neuron* 16:287-295.
- Alvarez de la Rosa D, Zhang P, Shao D, White F, Canessa CM (2002) Functional implications of the localization and activity of acid-sensitive channels in rat peripheral nervous system. *Proc Natl Acad Sci U S A* 99:2326-2331.
- Askwith CC, Wemmie JA, Price MP, Rokhlina T, Welsh MJ (2004) ASIC2 modulates ASIC1 H⁺-activated currents in hippocampal neurons. *J Biol Chem*.
- Baranowski R, Lynn B (1985) Receptive field size of hair follicle afferents in rats treated neonatally with capsaicin. *Brain Res* 338:395-397.
- Barnes TM, Jin Y, Horvitz HR, Ruvkun G, Hekimi S (1996) The *Caenorhabditis elegans* behavioral gene *unc-24* encodes a novel bipartite protein similar to both erythrocyte band 7.2 (stomatin) and nonspecific lipid transfer protein. *J Neurochem* 67:46-57.
- Bassilana F, Champigny G, Waldmann R, de Weille JR, Heurteaux C, Lazdunski M (1997) The acid-sensitive ionic channel subunit ASIC and the mammalian degenerin MDEG form a heteromultimeric H⁺-gated Na⁺ channel with novel properties. *J Biol Chem* 272:28819-28822.
- Benson CJ, Xie J, Wemmie JA, Price MP, Henss JM, Welsh MJ, Snyder PM (2002) Heteromultimers of DEG/ENaC subunits form H⁺-gated channels in mouse sensory neurons. *Proc Natl Acad Sci U S A* 99:2338-2343.
- Bessou P, Perl ER (1969) Response of cutaneous sensory units with unmyelinated fibers to noxious stimuli. *J Neurophysiol* 32:1025-1043.
- Bianchi L, Driscoll M (2002) Protons at the gate: DEG/ENaC ion channels help us feel and remember. *Neuron* 34:337-340.
- Birder LA, Perl ER (1994) Cutaneous sensory receptors. *J Clin Neurophysiol* 11:534-552.
- Blount P, Sukharev SI, Moe PC, Martinac B, Kung C (1999) Mechanosensitive channels of bacteria. *Methods Enzymol* 294:458-482.
- Brown AG, Iggo A (1967) A quantitative study of cutaneous receptors and afferent fibres in the cat and rabbit. *J Physiol* 193:707-733.
- Burgess PR, Perl ER (1967) Myelinated afferent fibres responding specifically to noxious stimulation of the skin. *J Physiol* 190:541-562.

-
- Burgess PR, Petit D, Warren RM (1968) Receptor types in cat hairy skin supplied by myelinated fibers. *J Neurophysiol* 31:833-848.
- Cain DM, Khasabov SG, Simone DA (2001) Response properties of mechanoreceptors and nociceptors in mouse glabrous skin: an in vivo study. *J Neurophysiol* 85:1561-1574.
- Canessa CM, Horisberger JD, Rossier BC (1993) Epithelial sodium channel related to proteins involved in neurodegeneration. *Nature* 361:467-470.
- Canessa CM, Schild L, Buell G, Thorens B, Gautschi I, Horisberger JD, Rossier BC (1994) Amiloride-sensitive epithelial Na⁺ channel is made of three homologous subunits. *Nature* 367:463-467.
- Caterina MJ, Leffler A, Malmberg AB, Martin WJ, Trafton J, Petersen-Zeitz KR, Koltzenburg M, Basbaum AI, Julius D (2000) Impaired nociception and pain sensation in mice lacking the capsaicin receptor. *Science* 288:306-313.
- Catton WT (1970) Mechanoreceptor function. *Physiol Rev* 50:297-318.
- Cauna N (1956) Structure and origin of the capsule of Meissner's corpuscle. *Anat Rec* 124:77-93.
- Chalfie M, Au M (1989) Genetic control of differentiation of the *Caenorhabditis elegans* touch receptor neurons. *Science* 243:1027-1033.
- Chalfie M, Wolinsky E (1990) The identification and suppression of inherited neurodegeneration in *Caenorhabditis elegans*. *Nature* 345:410-416.
- Chambers MR, Andres KH, von Duering M, Iggo A (1972) The structure and function of the slowly adapting type II mechanoreceptor in hairy skin. *Q J Exp Physiol Cogn Med Sci* 57:417-445.
- Chelur DS, Ernstrom GG, Goodman MB, Yao CA, Chen L, R OH, Chalfie M (2002) The mechanosensory protein MEC-6 is a subunit of the *C. elegans* touch-cell degenerin channel. *Nature* 420:669-673.
- Chen CC, Zimmer A, Sun WH, Hall J, Brownstein MJ (2002) A role for ASIC3 in the modulation of high-intensity pain stimuli. *Proc Natl Acad Sci U S A* 99:8992-8997.
- Couloigner V, Fay M, Djelidi S, Farman N, Escoubet B, Runembert I, Sterkers O, Friedlander G, Ferrary E (2001) Location and function of the epithelial Na channel in the cochlea. *Am J Physiol Renal Physiol* 280:F214-222.
- Davis KD, Meyer RA, Campbell JN (1993) Chemosensitivity and sensitization of nociceptive afferents that innervate the hairy skin of monkey. *J Neurophysiol* 69:1071-1081.

-
- Djoughri L, Lawson SN (2004) Abeta-fiber nociceptive primary afferent neurons: a review of incidence and properties in relation to other afferent A-fiber neurons in mammals. *Brain Res Brain Res Rev* 46:131-145.
- Drew LJ, Wood JN, Cesare P (2002) Distinct mechanosensitive properties of capsaicin-sensitive and -insensitive sensory neurons. *J Neurosci* 22:RC228.
- Drew LJ, Rohrer DK, Price MP, Blaver K, Cockayne DA, Cesare P, Wood JN (2004) ASIC2 and ASIC3 Do Not Contribute to Mechanically Activated Currents in Mammalian Sensory Neurons. *J Physiol*.
- Driscoll M, Chalfie M (1991) The *mec-4* gene is a member of a family of *Caenorhabditis elegans* genes that can mutate to induce neuronal degeneration. *Nature* 349:588-593.
- Drummond HA, Abboud FM, Welsh MJ (2000) Localization of beta and gamma subunits of ENaC in sensory nerve endings in the rat foot pad. *Brain Res* 884:1-12.
- Drummond HA, Price MP, Welsh MJ, Abboud FM (1998) A molecular component of the arterial baroreceptor mechanotransducer. *Neuron* 21:1435-1441.
- Dubreuil AS, Boukhaddaoui H, Desmadryl G, Martinez-Salgado C, Moshourab R, Lewin GR, Carroll P, Valmier J, Scamps F (2004) Role of T-type calcium current in identified D-hair mechanoreceptor neurons studied in vitro. *J Neurosci* 24:8480-8484.
- Duggan A, Ma C, Chalfie M (1998) Regulation of touch receptor differentiation by the *Caenorhabditis elegans mec-3* and *unc-86* genes. *Development* 125:4107-4119.
- Ernstrom GG, Chalfie M (2002) Genetics of sensory mechanotransduction. *Annu Rev Genet* 36:411-453.
- Eyzaguirre C, Kuffler SW (1955) Processes of excitation in the dendrites and in the soma of single isolated sensory nerve cells of the lobster and crayfish. *J Gen Physiol* 39:87-119.
- Fleischer E, Handwerker HO, Joukhadar S (1983) Unmyelinated nociceptive units in two skin areas of the rat. *Brain Res* 267:81-92.
- French AS (1992) Mechanotransduction. *Annu Rev Physiol* 54:135-152.
- French AS, Torkkeli PH, Seyfarth EA (2002) From stress and strain to spikes: mechanotransduction in spider slit sensilla. *J Comp Physiol A Neuroethol Sens Neural Behav Physiol* 188:739-752.
- Fricke B, Lints R, Stewart G, Drummond H, Dodt G, Driscoll M, von Düring M (2000) Epithelial Na⁺ channels and stomatin are expressed in rat trigeminal mechanosensory neurons. *Cell Tissue Res* 299:327-334.

-
- Fricke B, Stewart GW, Treharne KJ, Mehta A, Knopfle G, Friedrichs N, Muller KM, von Düring M (2003) Stomatin immunoreactivity in ciliated cells of the human airway epithelium. *Anat Embryol (Berl)* 207:1-7.
- Fuller CM, Awaysda MS, Arrate MP, Bradford AL, Morris RG, Canessa CM, Rossier BC, Benos DJ (1995) Cloning of a bovine renal epithelial Na⁺ channel subunit. *Am J Physiol* 269:C641-654.
- Garcia-Anoveros J, Corey DP (1997) The molecules of mechanosensation. *Annu Rev Neurosci* 20:567-594.
- Garcia-Anoveros J, Samad TA, Zúvela-Jelaska L, Woolf CJ, Corey DP (2001) Transport and localization of the DEG/ENaC ion channel BNaC1alpha to peripheral mechanosensory terminals of dorsal root ganglia neurons. *J Neurosci* 21:2678-2686.
- Garell PC, McGillis SL, Greenspan JD (1996) Mechanical response properties of nociceptors innervating feline hairy skin. *J Neurophysiol* 75:1177-1189.
- Gillespie PG, Walker RG (2001) Molecular basis of mechanosensory transduction. *Nature* 413:194-202.
- Goldstein BJ, Kulaga HM, Reed RR (2003) Cloning and characterization of SLP3: a novel member of the stomatin family expressed by olfactory receptor neurons. *J Assoc Res Otolaryngol* 4:74-82.
- Goldstein O, Asher C, Cragoe E, Jr., Kleyman TR, Garty H (1997) An aldosterone regulated chicken intestine protein with high affinity to amiloride. *Comp Biochem Physiol A Physiol* 118:201-208.
- Goodman MB, Ernstrom GG, Chelur DS, O'Hagan R, Yao CA, Chalfie M (2002) MEC-2 regulates *C. elegans* DEG/ENaC channels needed for mechanosensation. *Nature* 415:1039-1042.
- Gu CX, Juranka PF, Morris CE (2001) Stretch-activation and stretch-inactivation of Shaker-IR, a voltage-gated K⁺ channel. *Biophys J* 80:2678-2693.
- Gu G, Caldwell GA, Chalfie M (1996) Genetic interactions affecting touch sensitivity in *Caenorhabditis elegans*. *Proc Natl Acad Sci U S A* 93:6577-6582.
- Handwerker HO, Kilo S, Reeh PW (1991) Unresponsive afferent nerve fibres in the sural nerve of the rat. *J Physiol* 435:229-242.
- Horch KW, Burgess PR (1975) Effect of activation and adaptation on the sensitivity of slowly adapting cutaneous mechanoreceptors. *Brain Res* 98:109-118.
- Howard J, Roberts WM, Hudspeth AJ (1988) Mechano-electrical transduction by hair cells. *Annu Rev Biophys Chem* 17:99-124.

-
- Hu J, Milenkovic N, Lewin GR (2006) The high threshold mechanotransducer: a status report. *Pain* 120:3-7.
- Huang M, Gu G, Ferguson EL, Chalfie M (1995) A stomatin-like protein necessary for mechanosensation in *C. elegans*. *Nature* 378:292-295.
- Iggo A (1960) Cutaneous mechanoreceptors with afferent C fibres. *J Physiol* 152:337-353.
- Iggo A (1976) Is the physiology of cutaneous receptors determined by morphology? *Prog Brain Res* 43:15-31.
- Iggo A, Andres KH (1982) Morphology of cutaneous receptors. *Annu Rev Neurosci* 5:1-31.
- Johansson RS, Vallbo AB (1979) Tactile sensibility in the human hand: relative and absolute densities of four types of mechanoreceptive units in glabrous skin. *J Physiol* 286:283-300.
- Johnson KO (2001) The roles and functions of cutaneous mechanoreceptors. *Curr Opin Neurobiol* 11:455-461.
- Johnson KO, Yoshioka T, Vega-Bermudez F (2000) Tactile functions of mechanoreceptive afferents innervating the hand. *J Clin Neurophysiol* 17:539-558.
- Kellenberger S, Schild L (2002) Epithelial sodium channel/degenerin family of ion channels: a variety of functions for a shared structure. *Physiol Rev* 82:735-767.
- Kernan M, Zuker C (1995) Genetic approaches to mechanosensory transduction. *Curr Opin Neurobiol* 5:443-448.
- Kernan M, Cowan D, Zuker C (1994) Genetic dissection of mechanosensory transduction: mechanoreception-defective mutations of *Drosophila*. *Neuron* 12:1195-1206.
- Koerber HR, Woodbury CJ (2002) Comprehensive phenotyping of sensory neurons using an *ex vivo* somatosensory system. *Physiol Behav* 77:589-594.
- Koltzenburg M, Handwerker HO (1994) Differential ability of human cutaneous nociceptors to signal mechanical pain and to produce vasodilatation. *J Neurosci* 14:1756-1765.
- Koltzenburg M, Stucky CL, Lewin GR (1997) Receptive properties of mouse sensory neurons innervating hairy skin. *J Neurophysiol* 78:1841-1850.
- Kress M, Koltzenburg M, Reeh PW, Handwerker HO (1992) Responsiveness and functional attributes of electrically localized terminals of cutaneous C-fibers *in vivo* and *in vitro*. *J Neurophysiol* 68:581-595.
- Kruger L, Perl ER, Sedivec MJ (1981) Fine structure of myelinated mechanical nociceptor endings in cat hairy skin. *J Comp Neurol* 198:137-154.
- Lawson SN (2002) Phenotype and function of somatic primary afferent nociceptive neurones with C-, Delta- or Aalpha/beta-fibres. *Exp Physiol* 87:239-244.

-
- Leem JW, Willis WD, Chung JM (1993) Cutaneous sensory receptors in the rat foot. *J Neurophysiol* 69:1684-1699.
- Levina N, Totemeyer S, Stokes NR, Louis P, Jones MA, Booth IR (1999) Protection of *Escherichia coli* cells against extreme turgor by activation of MscS and MscL mechanosensitive channels: identification of genes required for MscS activity. *Embo J* 18:1730-1737.
- Lewin GR, McMahon SB (1991) Physiological properties of primary sensory neurons appropriately and inappropriately innervating skin in the adult rat. *J Neurophysiol* 66:1205-1217.
- Lewin GR, Mendell LM (1994) Regulation of cutaneous C-fiber heat nociceptors by nerve growth factor in the developing rat. *J Neurophysiol* 71:941-949.
- Lewin GR, Moshourab R (2004) Mechanosensation and pain. *J Neurobiol* 61:30-44.
- Lewin GR, Ritter AM, Mendell LM (1992) On the role of nerve growth factor in the development of myelinated nociceptors. *J Neurosci* 12:1896-1905.
- Lindemann B (1996) Taste reception. *Physiol Rev* 76:718-766.
- Lingueglia E, de Weille JR, Bassilana F, Heurteaux C, Sakai H, Waldmann R, Lazdunski M (1997) A modulatory subunit of acid sensing ion channels in brain and dorsal root ganglion cells. *J Biol Chem* 272:29778-29783.
- Lockery SR, Goodman MB (1998) Tight-seal whole-cell patch clamping of *Caenorhabditis elegans* neurons. *Methods Enzymol* 293:201-217.
- Loewenstein WR, Skalak R (1966) Mechanical transmission in a Pacinian corpuscle. An analysis and a theory. *J Physiol* 182:346-378.
- Lynn B, Carpenter SE (1982) Primary afferent units from the hairy skin of the rat hind limb. *Brain Res* 238:29-43.
- Macefield G, Gandevia SC, Burke D (1990) Perceptual responses to microstimulation of single afferents innervating joints, muscles and skin of the human hand. *J Physiol* 429:113-129.
- Maingret F, Fosset M, Lesage F, Lazdunski M, Honore E (1999) TRAAK is a mammalian neuronal mechano-gated K⁺ channel. *J Biol Chem* 274:1381-1387.
- Mannsfeldt AG, Carroll P, Stucky CL, Lewin GR (1999) Stomatin, a MEC-2 like protein, is expressed by mammalian sensory neurons. *Mol Cell Neurosci* 13:391-404.
- Mano I, Driscoll M (1999) DEG/ENaC channels: a touchy superfamily that watches its salt. *Bioessays* 21:568-578.
- McCarter GC, Reichling DB, Levine JD (1999) Mechanical transduction by rat dorsal root ganglion neurons in vitro. *Neurosci Lett* 273:179-182.

-
- McDonald FJ, Price MP, Snyder PM, Welsh MJ (1995) Cloning and expression of the beta- and gamma-subunits of the human epithelial sodium channel. *Am J Physiol* 268:C1157-1163.
- McMahon SB, Koltzenburg M (1990) Novel classes of nociceptors: beyond Sherrington. *Trends Neurosci* 13:199-201.
- Meyer RA, Davis KD, Cohen RH, Treede RD, Campbell JN (1991) Mechanically insensitive afferents (MIAs) in cutaneous nerves of monkey. *Brain Res* 561:252-261.
- Munger BL, Ide C (1988) The structure and function of cutaneous sensory receptors. *Arch Histol Cytol* 51:1-34.
- O'Hagan R, Chalfie M (2005) Mechanosensation in *Caenorhabditis elegans*. *Int Rev Neurobiol* 69:169-203.
- O'Hagan R, Chalfie M, Goodman MB (2005) The MEC-4 DEG/ENaC channel of *Caenorhabditis elegans* touch receptor neurons transduces mechanical signals. *Nat Neurosci* 8:43-50.
- Olausson H, Lamarre Y, Backlund H, Morin C, Wallin BG, Starck G, Ekholm S, Strigo I, Worsley K, Vallbo AB, Bushnell MC (2002) Unmyelinated tactile afferents signal touch and project to insular cortex. *Nat Neurosci* 5:900-904.
- Page AJ, Brierley SM, Martin CM, Martinez-Salgado C, Wemmie JA, Brennan TJ, Symonds E, Omari T, Lewin GR, Welsh MJ, Blackshaw LA (2004) The ion channel ASIC1 contributes to visceral but not cutaneous mechanoreceptor function. *Gastroenterology* 127:1739-1747.
- Paoletti P, Ascher P (1994) Mechanosensitivity of NMDA receptors in cultured mouse central neurons. *Neuron* 13:645-655.
- Patel AJ, Honore E, Maingret F, Lesage F, Fink M, Duprat F, Lazdunski M (1998) A mammalian two pore domain mechano-gated S-like K⁺ channel. *Embo J* 17:4283-4290.
- Perl ER (1968) Myelinated afferent fibres innervating the primate skin and their response to noxious stimuli. *J Physiol* 197:593-615.
- Pickles JO, Comis SD, Osborne MP (1984) Cross-links between stereocilia in the guinea pig organ of Corti, and their possible relation to sensory transduction. *Hear Res* 15:103-112.
- Price MP, Snyder PM, Welsh MJ (1996) Cloning and expression of a novel human brain Na⁺ channel. *J Biol Chem* 271:7879-7882.
- Price MP, Thompson RJ, Eshcol JO, Wemmie JA, Benson CJ (2004) Stomatin modulates gating of acid-sensing ion channels. *J Biol Chem* 279:53886-53891.

-
- Price MP, McIlwrath SL, Xie J, Cheng C, Qiao J, Tarr DE, Sluka KA, Brennan TJ, Lewin GR, Welsh MJ (2001) The DRASIC cation channel contributes to the detection of cutaneous touch and acid stimuli in mice. *Neuron* 32:1071-1083.
- Price MP, Lewin GR, McIlwrath SL, Cheng C, Xie J, Heppenstall PA, Stucky CL, Mannsfeldt AG, Brennan TJ, Drummond HA, Qiao J, Benson CJ, Tarr DE, Hrstka RF, Yang B, Williamson RA, Welsh MJ (2000) The mammalian sodium channel BNC1 is required for normal touch sensation. *Nature* 407:1007-1011.
- Pubols BH, Jr. (1990) Slowly adapting type I mechanoreceptor discharge as a function of dynamic force versus dynamic displacement of glabrous skin of raccoon and squirrel monkey hand. *Neurosci Lett* 110:86-90.
- Quilliam TA, Sato M (1955) The distribution of myelin on nerve fibres from Pacinian corpuscles. *J Physiol* 129:167-176.
- Rajaram S, Spangler TL, Sedensky MM, Morgan PG (1999) A stomatin and a degenerin interact to control anesthetic sensitivity in *Caenorhabditis elegans*. *Genetics* 153:1673-1682.
- Reeh PW (1986) Sensory receptors in mammalian skin in an in vitro preparation. *Neurosci Lett* 66:141-146.
- Ritter AM, Lewin GR, Mendell LM (1993) Regulation of myelinated nociceptor function by nerve growth factor in neonatal and adult rats. *Brain Res Bull* 30:245-249.
- Rossier BC, Pradervand S, Schild L, Hummler E (2002) Epithelial sodium channel and the control of sodium balance: interaction between genetic and environmental factors. *Annu Rev Physiol* 64:877-897.
- Sato A, Sato Y, Suzuki H (1985) Aging effects on conduction velocities of myelinated and unmyelinated fibers of peripheral nerves. *Neurosci Lett* 53:15-20.
- Schaible HG, Grubb BD (1993) Afferent and spinal mechanisms of joint pain. *Pain* 55:5-54.
- Schmidt R, Schmelz M, Ringkamp M, Handwerker HO, Torebjork HE (1997) Innervation territories of mechanically activated C nociceptor units in human skin. *J Neurophysiol* 78:2641-2648.
- Schmidt R, Schmelz M, Forster C, Ringkamp M, Torebjork E, Handwerker H (1995) Novel classes of responsive and unresponsive C nociceptors in human skin. *J Neurosci* 15:333-341.
- Shepherd GM, Barres BA, Corey DP (1989) "Bundle blot" purification and initial protein characterization of hair cell stereocilia. *Proc Natl Acad Sci U S A* 86:4973-4977.
- Snyers L, Thines-Sempoux D, Prohaska R (1997) Colocalization of stomatin (band 7.2b) and actin microfilaments in UAC epithelial cells. *Eur J Cell Biol* 73:281-285.

-
- Steen KH, Reeh PW, Anton F, Handwerker HO (1992) Protons selectively induce lasting excitation and sensitization to mechanical stimulation of nociceptors in rat skin, in vitro. *J Neurosci* 12:86-95.
- Stewart GW, Hepworth-Jones BE, Keen JN, Dash BC, Argent AC, Casimir CM (1992) Isolation of cDNA coding for an ubiquitous membrane protein deficient in high Na⁺, low K⁺ stomatocytic erythrocytes. *Blood* 79:1593-1601.
- Stucky CL, DeChiara T, Lindsay RM, Yancopoulos GD, Koltzenburg M (1998) Neurotrophin 4 is required for the survival of a subclass of hair follicle receptors. *J Neurosci* 18:7040-7046.
- Sukharev SI, Blount P, Martinac B, Blattner FR, Kung C (1994) A large-conductance mechanosensitive channel in *E. coli* encoded by *mscL* alone. *Nature* 368:265-268.
- Sutherland SP, Benson CJ, Adelman JP, McCleskey EW (2001) Acid-sensing ion channel 3 matches the acid-gated current in cardiac ischemia-sensing neurons. *Proc Natl Acad Sci U S A* 98:711-716.
- Suzuki H, Kerr R, Bianchi L, Frokjaer-Jensen C, Slone D, Xue J, Gerstbrein B, Driscoll M, Schafer WR (2003a) In vivo imaging of *C. elegans* mechanosensory neurons demonstrates a specific role for the MEC-4 channel in the process of gentle touch sensation. *Neuron* 39:1005-1017.
- Suzuki M, Watanabe Y, Oyama Y, Mizuno A, Kusano E, Hirao A, Ookawara S (2003b) Localization of mechanosensitive channel TRPV4 in mouse skin. *Neurosci Lett* 353:189-192.
- Tavernarakis N, Driscoll M (1997) Molecular modeling of mechanotransduction in the nematode *Caenorhabditis elegans*. *Annu Rev Physiol* 59:659-689.
- Tobin DM, Bargmann CI (2004) Invertebrate nociception: behaviors, neurons and molecules. *J Neurobiol* 61:161-174.
- Tracey WD, Jr., Wilson RI, Laurent G, Benzer S (2003) *painless*, a *Drosophila* gene essential for nociception. *Cell* 113:261-273.
- Vallbo AB, Johansson RS (1984) Properties of cutaneous mechanoreceptors in the human hand related to touch sensation. *Hum Neurobiol* 3:3-14.
- Vallbo AB, Olausson H, Wessberg J, Kakuda N (1995) Receptive field characteristics of tactile units with myelinated afferents in hairy skin of human subjects. *J Physiol* 483 (Pt 3):783-795.
- Waddell PJ, Lawson SN, McCarthy PW (1989) Conduction velocity changes along the processes of rat primary sensory neurons. *Neuroscience* 30:577-584.

-
- Waldmann R, Lazdunski M (1998) H(+)-gated cation channels: neuronal acid sensors in the NaC/DEG family of ion channels. *Curr Opin Neurobiol* 8:418-424.
- Waldmann R, Champigny G, Voilley N, Lauritzen I, Lazdunski M (1996) The mammalian degenerin MDEG, an amiloride-sensitive cation channel activated by mutations causing neurodegeneration in *Caenorhabditis elegans*. *J Biol Chem* 271:10433-10436.
- Waldmann R, Bassilana F, de Weille J, Champigny G, Heurteaux C, Lazdunski M (1997) Molecular cloning of a non-inactivating proton-gated Na⁺ channel specific for sensory neurons. *J Biol Chem* 272:20975-20978.
- Wang D, Mentzer WC, Cameron T, Johnson RM (1991) Purification of band 7.2b, a 31-kDa integral phosphoprotein absent in hereditary stomatocytosis. *J Biol Chem* 266:17826-17831.
- Way JC, Chalfie M (1988) *mec-3*, a homeobox-containing gene that specifies differentiation of the touch receptor neurons in *C. elegans*. *Cell* 54:5-16.
- Weidner C, Schmelz M, Schmidt R, Hansson B, Handwerker HO, Torebjork HE (1999) Functional attributes discriminating mechano-insensitive and mechano-responsive C nociceptors in human skin. *J Neurosci* 19:10184-10190.
- Welsh MJ, Price MP, Xie J (2002) Biochemical basis of touch perception: mechanosensory function of degenerin/epithelial Na⁺ channels. *J Biol Chem* 277:2369-2372.
- Wemmie JA, Chen J, Askwith CC, Hruska-Hageman AM, Price MP, Nolan BC, Yoder PG, Lamani E, Hoshi T, Freeman JH, Jr., Welsh MJ (2002) The acid-activated ion channel ASIC contributes to synaptic plasticity, learning, and memory. *Neuron* 34:463-477.
- Wessberg J, Olausson H, Fernstrom KW, Vallbo AB (2003) Receptive field properties of unmyelinated tactile afferents in the human skin. *J Neurophysiol* 89:1567-1575.
- Wetzel C, Hu J, Riethmacher D, Benckendorff A, Harder L, Eilers A, Moshourab R, Kozlenkov A, Labuz D, Caspani O, Erdmann B, Machelska H, Heppenstall PA, Lewin GR (2006) A stomatin-domain protein essential for touch sensation in the mouse. *Nature*.
- Woodbury CJ, Ritter AM, Koerber HR (2001) Central anatomy of individual rapidly adapting low-threshold mechanoreceptors innervating the "hairy" skin of newborn mice: early maturation of hair follicle afferents. *J Comp Neurol* 436:304-323.
- Xie J, Price MP, Berger AL, Welsh MJ (2002) DRASIC contributes to pH-gated currents in large dorsal root ganglion sensory neurons by forming heteromultimeric channels. *J Neurophysiol* 87:2835-2843.

Zhang S, Arnadottir J, Keller C, Caldwell GA, Yao CA, Chalfie M (2004) MEC-2 is recruited to the putative mechanosensory complex in *C. elegans* touch receptor neurons through its stomatin-like domain. *Curr Biol* 14:1888-1896.