

10 Literature cited

- ALBACH D.C., SOLTIS P.S., SOLTIS D.E. & OLMSTEAD R.G. (2001): Phylogenetic analysis of Asterids based on sequences of four genes. - *Ann. Miss. Bot. Gard.* **88**: 163-212.
- AL-SHEBAZ I.A. (1991): The genera of Boraginaceae in the southeastern United States. - *J. Arnold Arbor., Supplem.* **1**: 1-169.
- AN S.S., FRIEDL T., & HEGEWALD E. (1999): Phylogenetic relationships of *Scenedesmus* and *Scenedesmus*-like coccoid green algae as inferred from ITS-2 rDNA sequence comparison. - *Plant Biol.* **1**: 418-428.
- ASH A., ELLIS B., HICKEY L.J., JOHNSON K., WILF P. & WING S. (eds.) (1999): Manual of leaf architecture – morphological description and categorization of dicotyledonous and net-veined monocotyledonous angiosperms. Leaf Architecture Working Group. Smithsonian Institution. Washington DC.
- AYENSU E.S. (1981): Medicinal plants of the West Indies. Algonac, Michigan, USA: Reference Publications.
- BACON J.D. & BRAGG L.H. (1986): Systematics of *Nama* (Hydrophyllaceae): comparison of seed morphology of sects. *Arachnoidea* and *Cinerascentia* with five species of *Eriodictyon* and *Turricula parryi*. - *Sida* **11**: 271-281.
- BACON J.D. (1987): Systematics of *Nama* (Hydrophyllaceae): seed coat morphology of *Lemmonia californica* and *Nama* species allied with *Nama demissum*. - *Aliso* **11**: 441-450.
- BALDWIN B.G. (1992): Phylogenetic utility of the Internal Transcribed Spacers of nuclear ribosomal DNA in plants: an example from the Compositae. - *Mol. Phyl. Evol.* **1**: 3-16.
- BAMBER R.K. & TER WELLE B.J.H. (1994): Adaptive Trends in the Wood Anatomy of Lianas. - In: IQBAL, M. (ed.): *From Growth Patterns in Vascular Plants*. pp 272-287. Portland, Oregon, USA: Dioscorides Press.
- BARKER R.D. & VESTJENS W.J.M. (1989): *The Food of Australian Birds*. Vol.1. - Lyneham, Australia: CSIRO.
- BARKER R.D. & VESTJENS W.J.M. (1990): *The Food of Australian Birds*. Vol. 2. - Melbourne, Australia: CSIRO.
- BIDER J. (1935): Beiträge zur Pharmakognosie der Boraginaceen und Verbenaceen. Vergleichende Anatomie des Laubblattes. Inaugural-Dissertation Universität Basel.
- BIRECKA H., FROHLICH M.W., HULL L. & CHASKES M.J. (1980): Pyrrolizidine alkaloids of *Heliotropium* from Mexico and adjacent U.S.A. - *Phytochemistry* **19**: 421-426.
- BÖHLE U.-R. & HILGER H.H. (1997): Chloroplast DNA systematics of “Boraginaceae” and related families - a goodbye to the old and familiar concept of five subfamilies. - *Scripta Bot. Belg.* **15**: 30.
- BÖHLE U.-R., HILGER H.H. & MARTIN W.F. (1996): Island colonization and evolution of the insular woody habit in *Echium* L. (Boraginaceae). - *Proc. Natl. Acad. Sci. USA* **93**: 11740-11745.

- BONNEMAIN J.L., BOURQUIN S., RENAULT S., OFFLER C. & FISHER D.G. (1991): Transfer cells : structure and physiology. - In: BONNEMAIN J.L., DELROT S., LUCAS W.L. & DAINTY L. (eds.): Recent advances in phloem transport and assimilate compartmentation. pp. 74-83. Paris, Nantes, France: Ouest Edition, Presses Academiques.
- BOPPRÉ M. (1986): Insects pharmacophagously utilizing defensive plant chemicals (pyrrolizidine alkaloids). - *Naturwissenschaften* **73**: 17-26.
- BOPPRÉ M. (1995): Pharmacophagie: Drogen, Sex und Schmetterlinge. - *Biologie in unserer Zeit* **25**: 8-17.
- BOPPRÉ M. (1999): "Drug-addicted" insects in Africa. - *Metamorphosis* **10**: 3-14.
- BRIESE D. & ZAPATER M. (2001): Biological control of blue Heliotrope. A report for the Rural Industries Research and Development Corporation. No. 01/119, pp.1-18. Kingston, Australia.
- BRITTON D.M. (1951): Cytogenetic studies on the Boraginaceae. - *Brittonia* **7**: 233-266.
- BRUMMIT R.K. (1972): *Heliotropium*. - In: TUTIN T.G. *et al.* (eds.): *Flora Europaea*. **3**: 84-86. Cambridge, Great Britain: Cambridge University Press.
- BULL L.B., CULVENOR C.C.J. & DICK A.T. (1968): The pyrrolizidine alkaloids. – In: NEUBERGER A. & TATUM E.L. (eds.): *Frontiers of Biology*. **9**: 1-293. Amsterdam, Netherlands: North-Holland Publ. Co.
- BURKILL I.H. (1966): A dictionary of the economic products of the Malay Peninsula. Vol 1 and 2. 2nd Ed. Reprint Kuala Lumpur, Malaysia: Ministry of Agriculture and Co. Operatives.
- BURKILL I.H. (1985): The useful plants of west tropical Africa. Vol. 1. Kew, Great Britain: Royal Botanic Garden.
- CARLQUIST S. (1992): Wood anatomy of sympetalous dicotyledon families: a summary, with comments on systematical relationships and evolution of the woody habit. – *Ann. Missouri Bot. Gard.* **79**: 303-332.
- CATALFAMO J.L., MARTIN W.B. & BIRECKA H. (1982): Accumulation of alkaloids and their necines in *Heliotropium curassavicum*, *H. spathulatum* and *H. indicum*. – *Phytochemistry* **21**: 2669-2675.
- CHANCE G.D. & BACON J.D. (1984): Systematic implications of seed coat morphology in *Nama* (Hydrophyllaceae). – *Amer. J. Bot.* **71**: 829-842.
- CHASE M.W., SOLTIS D.E., OLMSTEAD R.G., MORGAN D., LES D.H., MISHLER B.D., DUVALL M.R., PRICE R.A., HILLS H.G., QIU Y.-L., KRON K.A., RETTIG J.H., CONTI E., PALMER J., MANHART J.R., SYTSMA K.J., MICHAELS H.J., KRESS W.J., KAROL K.G., CLARK W.D., HEDRÉN M., GAUT B.S., JANSEN R.K., KIM K.-J., WIMPEE C.F., SMITH J.F., FURNIER G.R., STRAUSS S.H., XIANG Q.-Y., PLUNKETT G.M., SOLTIS P.S., SWENSEN S.M., WILLIAMS S.E., GADEK P.A., QUINN C.J., EGUIARTE L.E., GOLENBERG E., LEARN JR. G.H., GRAHAM S.W., BARRETT S C.H., DAYANANDAN S. & ALBERT V.A. (1993): Phylogenetics of seed plants: an analysis of nucleotide sequences from the plastid gene *rbcl*. – *Ann. Missouri Bot. Gard.* **80** : 528-580.
- CHILTON N.B. & R.B. GASSER (1999): Sequence differences in the Internal Transcribed Spacers of DNA among four species of hookworm (Ancylostomatoidea: *Ancylostoma*). - *Int. J. Parasitol.* **29**: 1971-1977.

- CHUANG T.I. & CONSTANCE L. (1992): Seeds and systematics in Hydrophyllaceae: tribe Hydrophyllae. - Amer. J. Bot. **79**: 257-264.
- COCUCCI, A.E. (1983): New Evidence from embryology in angiosperm classification. – Nord. Bot. J. **3**: 67-73.
- COLEMAN A.W. & MAI J.C. (1997): Ribosomal ITS-1 and ITS-2 sequence comparisons as a tool for predicting genetic relatedness. - J. Mol. Evol. **45**: 168-177.
- COLEMAN A.W., PREPARATA R.M., MEHROTRA B. & MAI J.C. (1998): Derivation of the secondary structure of the ITS-1 transcript in Volvocales and its taxonomic correlations. - Protist **149**: 135-146.
- COSNER M.E., JANSEN R.K. & LAMMERS T.G. (1994): Phylogenetic relationships in the Campanulales based on *rbcL* sequences. - Plant Syst. Evol. **190**: 79-95.
- CRAIG R.M. (1978): Threatened sea-lavender. - In: WARD D.B. (ed.): Rare and endangered biota of Florida. Vol. 5 Plants. Gainesville, Florida, USA: University Press of Florida.
- CRAVEN L.A (1996): A taxonomic revision of *Heliotropium* (Boraginaceae) in Australia. - Austr. Syst. Bot. **9**: 521-657.
- CULVENOR C.C.J. (1973): Alkaloids. - In: BUTLER G.W. & BAILY R.W. (eds.): Chemistry and Biochemistry of Herbage. **1**: 375-446. London, Great Britain: Academic Press.
- CULVENOR C.C.J. (1978): Pyrrolizidine alkaloids – occurrence and systematic importance in angiosperms. - Bot. Notiser **131**: 473-486.
- CZERECHANOV S.K. (1981): Plantae vasculares URSS. pp. 111-119. Leningrad, URSS: Nauka.
- DAHLGREN, R. (1983): General aspects of angiosperm evolution and macrosystematics. – Nord. J. Bot. **3**: 119-149.
- DANDY J.E. (1972): Note on *Argusia* Boehmer (Boraginaceae). - Bot. J. Linn. Soc. **65**: 256.
- DAS V.S.R. & RAGHAVENDRA A.S. (1973): A screening of the dicotyledonous weed flora for the occurrence of C4 dicarboxylic acid pathway of photosynthesis. - Proc. Indian Acad. Sci. **77**: 93-100.
- DECANDOLLE A.P. (1845): CXXXIX. Boraginaceae. Prodomus systematis naturalis regni vegetabilis. Vol. **9**: 466-559. Paris, France: Treuttel & Würtz.
- DIANE N., FÖRTHNER H. & HILGER H.H. (2002a): A systematic analysis of *Heliotropium*, *Tournefortia*, and allied taxa of the Heliotropiaceae (Boraginales) based on ITS1 sequences and morphological data. - Amer. J. Bot. **89**: 287-295.
- DIANE N., HILGER H.H. & GOTTSCHLING M. (2002b): Transfer cells in the seeds of Boraginales. - Bot. J. Linn. Soc. **140**: 155-164.
- DIANE N., FÖRTHNER H., HILGER H.H. & WEIGEND M. (in press a): Heliotropiaceae. - In: KUBITZKI K. (ed.): Families and Genera of the Flowering Plants. Berlin, Germany: Springer.
- DIANE N., JAKOB C. & HILGER H.H. (in press b): Leaf anatomy and foliar trichomes in Heliotropiaceae and their systematic relevance. – Flora **198** (6).

- DiFULVIO T.E. (1967): Recuentos cromosómicos en angiospermas argentinas. - Kurtziana **4**: 87-90.
- DiFULVIO T.E. (1969): Recuentos cromosómicos en *Heliotropium* (Boraginaceae). - Kurtziana **5**: 89-95.
- DiFULVIO T.E. (1978): Sobre la vasculatura floral, embriología y cromosomas de *Ixorhea tschudiana* (Heliotropiaceae). - Kurtziana **11**: 75-105.
- DiFULVIO T.E. (1980): Sobre el polen de *Ixorhea tschudiana* (Heliotropiaceae). - Bol. Soc. Argent. Bot. **19**: 83-90.
- DiFULVIO T.E. (1982): Sobre la anatomía foliar y estomatogénesis de *Ixorhea tschudiana* (Heliotropiaceae). - Kurtziana **15**: 9-18.
- DOCTERS VAN LEEUWEN W.M. (1936) Krakatau 1883-1933. - Ann. Jard. Bot. Buitenzorg: 46-47.
- DOYLE J.J. & J.L. DOYLE. (1990): Isolation of plant DNA from fresh tissue. - Focus **12**: 13-15.
- DUKE J.A. & AYENSU E.S. (1985): Medicinal plants of China. No. 4. Algonac, Michigan, USA: Reference Publications.
- EGERTON-WARBURTON L.M. (1998): A smoke-induced alteration of the sub-testa cuticle in seeds of the post-fire recruiter, *Emmenanthe penduliflora* Benth. (Hydrophyllaceae). - J. Exp. Bot. **49**: 1317-1327.
- ENDLICHER S.L. (1839): Genera plantarum. p. 646. Wien, Austria: Fr.Beck.
- FARQUI S.A. (1961): Cytological studies in *Heliotropium* from West Pakistan. - Caryologia **14**: 313-318.
- FEDOROV A.A. (ed.) (1969): Chromosome numbers of flowering plants. Leningrad, URSS: Nauka.
- FELSENSTEIN J. (1985): Confidence limits on phylogenies: an approach using the bootstrap. - Evolution **39**: 783-791.
- FENZL E. (1886): Vier neue Pflanzenarten Süd-Amerikas. - Verh. K. K. Zoolog.-Bot. Ges. Wien **36**: 287-294.
- FERGUSON D.M. (1998): ITS phylogeny of the 'core' Hydrophyllaceae. GeneBank submission.
- FERGUSON D.M. (1999): Phylogenetic analysis and relationships in Hydrophyllaceae based on *ndhF* sequence data. - Syst. Bot. **23**: 253-268.
- FÖRTHNER H. (1998): Die infragenerische Gliederung der Gattung *Heliotropium* L. und ihre Stellung innerhalb der subfam. Heliotropioideae (Schräd.) Arn. (Boraginaceae). - Sendtnera **5**: 35-241.
- FROHLICH M.W. (1976): Appearance of Vegetation in Ultraviolet Light: Absorbing Flowers, Reflecting Backgrounds. - Science **194**: 839-841.
- FROHLICH M.W. (1978): Systematics of *Heliotropium* section *Orthostachys* in Mexico. Ph. D. Thesis (unpubl.). pp. 1-277. Harvard University, Cambridge, Massachusetts, USA.
- FROHLICH M.W. & COLBURN W.T. (1980): Reports. p. 705. - In: LÖVE Á. (ed.): Chromosome number reports LXIX. Taxon **29**: 703-730.

- FROHLICH M.W. & FERRONE P.J. (1984): Reports pp. 351-352. – In: LÖVE Á (ed.): Chromosome number reports LXXXIII. *Taxon* **33**: 351-354.
- GERSTBERGER P. & LEINS P. (1978): Rasterelektronenmikroskopische Untersuchungen an Blütenknospen von *Physalis philadelphica* (Solanaceae). Anwendung einer neuen Präparationsmethode. – *Ber. Deutsch. Bot. Ges.* **91**: 381-387.
- GOLOBOFF P.A. (1993): Nona Vers. 2.0. Tucumán.
- GOTTSCHLING M., HILGER H.H., WOLF M. & DIANE N. (2001): Secondary structure of the ITS1 transcript and its application in a reconstruction of the phylogeny of Boraginales. – *Plant Biol.* **3** : 629-636.
- GOTTSCHLING M. & HILGER H.H. (2001): Phylogenetic analysis and character evolution of *Ehretia* and *Bourreria* (Ehretiaceae, Boraginales) and their allies based on ITS1 sequences. – *Bot. Jahrb. Syst.* **123**: 249-268.
- GOTTSCHLING M., HILGER H.H., WOLF M. & DIANE N. (2001): Secondary structure of the ITS1 transcript and its application in a reconstruction of the phylogeny of Boraginales. – *Plant Biol.* **3**: 629-636.
- GRAF A.B. (1978): *Tropica. Color cyclopedia of exotic plants and trees from the tropics and subtropics.* East Rutherford, New Jersey, USA: Roehrs Co.
- GRAHAM A. & JARZEN D.M. (1969): Studies in neotropical paleobotany. I. The oligocene communities of Puerto Rico. – *Ann. Missouri Bot. Gard.* **56** : 308-357.
- GRAHAM A. (1976): Studies in neotropical paleobotany. II. The miocene communities of Veracruz, Mexico. – *Ann. Missouri Bot. Gard.* **63** : 787-842.
- GUIGNARD N.L. (1893) : Recherches sur le développement de la graine et en particulier du tégument séminal – Borriginées. – *J. Bot.* **7** : 205-214.
- GUNNING B.E.S. & PATE J.S. (1969): ‘Transfer cells’ plant cells with wall ingrowth, specialized in relation to short distance transport of solutes – their occurrence, structure, and development. – *Protoplasma* **68**: 107-133.
- GÜRKE M. (1893): Borriginaceae. – In: ENGLER A. & PRANTL K. (eds.): *Die natürlichen Pflanzenfamilien.* **4**: 49-96. Leipzig, Germany: W. Engelmann.
- HABERLANDT G. (1896): *Physiologische Pflanzenanatomie.* 2. ed. Leipzig, Germany: W. Engelmann.
- HARRIS J.G. & HARRIS M.W. (1994): *Plant identification terminology.: an illustrated glossary.* Spring Lake, Utah, USA: Spring Lake Publishing.
- HARTWELL J L. (1968): Plants used against cancer. A survey. – *Lloydia* **31**: 71-170.
- HAUMAN L. (1922): Nótula sobre *Oxyosmyles viscosissima* Speg. – *Physis* **5**: 306-307.
- HEINE H. (1976): Borriginaceae. – In : AUBREVILLE A. & LEROY J.-F. (eds.) : *Flora de la Nouvelle Calédonie et dépendances.* **7** : 95-118. Paris, France : Muséum National d’Histoire Naturelle.
- HEPPERLE D. (1997): *Alignment editor align 3.5a.* Heidelberg, Germany.

- HEUBL G.R., GAVIRIA J.C. & WANNER G. (1990): A contribution to the taxonomy and evolution of *Cordia* (Boraginaceae) and allied genera. Chromosome numbers, pollen morphology and crystal pattern in wood. – Bot. Jahrb. Syst. **112**: 129-165.
- HICKEY L.J. (1973): Classification of the architecture of dicotyledonous leaves. – Amer. J. Bot. **60**: 17-33.
- HILGER H.H. (1985): Ontogenie, Morphologie und systematische Bedeutung geflügelter und glochidientragender Cynoglosseae- und Eritrichieae-Früchte (Boraginaceae). – Bot. Jahrb. Syst. **105**: 323-378.
- HILGER H.H. (1987): Fruchtbiologische Untersuchungen an Heliotropioideae (Boraginaceae). I. Die Ontogenie der monospermen Früchte von *Heliotropium supinum* L. – Flora **179**: 291-303.
- HILGER H.H. (1989): Flower and fruit development in the Macaronesian endemic *Ceballosia fruticosa* (syn. *Messerschmidia fruticosa*, Boraginaceae, Heliotropioideae). – Pl. Syst. Evol. **166**: 119-129.
- HILGER H.H. (1992): Morphology of *Heliotropium* (Boraginaceae) dispersal units. – Bot. Acta **105**: 387-393.
- HILGER H.H., HOPPE J.R. & HOFMANN M. (1993): Energiedispersive Röntgenmikroanalyse (EDX) von Boraginaceae subfam. Boraginoideae – Klausenoberflächen (Sind Silicium- und Calcium-Einlagerungen in die Fruchtwand systematisch verwertbare Merkmale?). – Flora **188**: 387-398.
- HILGER H.H. & DIANE N. (submitted): A systematic analysis of Heliotropiaceae (Boraginales) based on *trnL* and ITS1 sequence data. – Syst. Bot. Jahrb.
- HOFMANN M. (1994): Untersuchungen zur Gynoecealentwicklung in der Tribus Eritrichieae (Boraginaceae-Boraginoideae). Inaugural-Dissertation, Freie Universität Berlin (unpublished).
- HOFMANN M. (1999): Flower and fruit development in the genus *Phacelia* (Phacelieae, Hydrophyllaceae): characters of systematic value. – Syst. Geogr. Pl. **68**: 203-212.
- HORNER H.T. & WAGNER B.L. (1992): Association of four different calcium crystals in the anther connective tissue and hypodermal stomium of *Capsicum annuum* (Solanaceae) during microsporogenesis. – Amer. J. Bot. **79**: 531-541.
- HUTCHINSON J. (1959): The families of flowering plants. Oxford, Great Britain: Clarendon Press.
- JAEGER J.A., TURNER D.H. & ZUCKER M. (1989): Improved predictions of secondary structures for RNA. – Proc. Natl. Acad. Sci. USA **86**: 7706-7710.
- JAEGER J.A., TURNER D.H. & ZUCKER M. (1990): Predicting optimal and suboptimal secondary structure for RNA. – Methods Enzymol. **183**: 281-306.
- JOHNSTON I.M. (1928): Studies in the Boraginaceae. VII. 1. The South American species of *Heliotropium*. – Contr. Gray Herb. **81**: 3-83.
- JOHNSTON I.M. (1930): Studies in the Boraginaceae. VIII. 1. 3. Treatment of *Tournefortia*. – Contr. Gray Herb. **92**: 66-89.
- JOHNSTON I.M. (1935a): Studies in the Boraginaceae. XI. 1. The species of *Tournefortia* and *Messerschmidia* in the Old World. – J. Arnold Arbor. **16**: 145 - 168.

- JOHNSTON I.M. (1935b): Studies in the Boraginaceae. X. The Boraginaceae of northeastern South America. - J. Arnold Arbor. **16**: 1-64.
- JOHNSTON I.M. (1949): Studies in the Boraginaceae. XVIII. Boraginaceae of the southeastern West Indies. - J. Arnold Arbor. **30**: 129-133.
- JOHNSTON I.M. (1951): Studies in the Boraginaceae. XX. Representatives of three subfamilies in Eastern Asia. - J. Arnold Arbor. **32**: 118-122.
- JOHNSTON I.M. (1953): Boraginaceae. - In: WILLIAMS R.O. *et al.*, Flora Trinidad and Tobago **2**: 189-209. Trinidad: Port of Spain.
- JOHNSTON I.M. (1964): Boraginaceae. - In: LUNDELL C.L. (ed.): Flora of Texas. **1**: 123-221. Renner, Texas, USA: Found.
- JOHRI B.M. & VASIL I.K. (1956): The embryology of *Ehretia laevis* Roxb. - Phytomorphology **6**: 134-143.
- KHALEEL T.F. (1974): Embryology of *Cynoglossum denticulatum* DC. - Bot. Not. **127**: 193-210.
- KHALEEL T.F. (1977): Embryology of *Ehretia acuminata* R. Br. - Proc. Montana Acad. Sci. **37**: 35-53.
- KHALEEL T.F. (1978): Embryology of *Heliotropium scabrum* and *H. strigosum* (Boraginaceae). - Pl. Syst. Evol. **129**: 45-62.
- Khaleel T.F. (1985): A review of endosperm and the taxonomic position of Boraginaceae. - J. Plant Sci. Res. **1**: 117-133.
- KIMURA M. (1980): A simple method for estimating evolutionary rate of the base substitutions through comparative studies of nucleotide sequences. - J. Mol. Evol. **16**: 111-120.
- KNUTH P. (1899): Handbuch der Blütenbiologie: 92. Familie Boraginaceae. Vol. **2**: 97-125. Leipzig, Germany: W. Engelmann.
- KNUTH P. (1905): Handbuch der Blütenbiologie: 406. *Heliotropium*. Vol. **3**: 64. Leipzig, Germany: W. Engelmann.
- KRAGGE H. (1911): Über die Festigkeit der Blätter der Boraginaceae und verwandter Familien. Inaugural-Dissertaton Universität Kiel. Hamburg, Germany: Petschmann.
- KUMAR K.V. & RAO B.H. (1994): A contribution to the foliar epidermology of *Heliotropium*. - J. Swamy Bot. Cl. **11**: 12-19.
- KUNTZE K.E.O. (1891): Revisio genera plantarum, part 2. Leipzig, Germany: Felix.
- LAKSHMANAN A.J. & SHANMUGASUNDARAM S. (1995): Heliscabine, a pyrrolizidine ester alkaloid from *Heliotropium scabrum*. - Phytochemistry **39**: 473-475.
- LESKO G.L. & WALKER R.B. (1969): Effect of sea water on seed germination in two pacific atoll beach species. - Ecology **50**: 730-734.
- LEWIS W.H. & ELVIN-LEWIS M.P.F. (1977): Medical botany: plant affecting man's health. New York; USA: Wiley.

- LINNÉ C.v. (1753): *Species plantarum*. Stockholm. Facsimile reprint 1957. London, Great Britain: Quaritch.
- LINNÉ C.v. (1767): *Mantissa plantarum, altera*. Stockholm. Facsimile reprint 1961. New York, USA: Wheldon and Wesley, and Hafner Publishing.
- LIU J.-S. & SCHARDL C.L. (1994): A conserved sequence in Internal Transcribed Spacer 1 of plant nuclear rRNA genes. – *Plant Mol. Biol.* **26**: 775-778.
- LUQUE T. (1996): Karyological studies on Spanish Boraginaceae. VII. *Heliotropium* L. - *Willdenowia* **25**: 659-668.
- MANOS P., DOYLE J.J. & NIXON K.C. (1999): Phylogeny, biogeography, and processes of molecular differentiation in *Quercus* subgenus *Quercus* (Fagaceae). - *Mol. Phylogenet. Evol.* **12**: 333-349.
- MARSHALL J.A. (1999): Molecular systematics of *Solanum* section *Lycopersicum* (*Lycopersicon*): nuclear ITS rDNA sequences. Thesis, Department of School of Biosciences, University of Nottingham. GeneBank submission.
- MAYOL M. & ROSSELLÓ J.A. (2001): Why nuclear ribosomal DNA spacers (ITS) tell different stories in *Quercus*. - *Mol. Phylogenet. Evol.* **19**: 167-176.
- METCALFE C.R. & CHALK. L. (1950): Anatomy of the dicotyledons: 195. Boraginaceae. **2**: 945-954. Oxford, Great Britain: Clarendon Press.
- MIERS J. (1868): XI. Observations on some of the Heliotropieae. – *Ann. Mag.Nat. Hist.* **2**: 121-133.
- MULLER J. (1981): Fossil pollen records of extant angiosperms. - *Bot. Rev.* **47**: 1-142.
- NAPP-ZINN K. (1984): Anatomie des Blattes, 2. Blattanatomie der Angiospermen. Handbuch der Anatomie. Berlin-Stuttgart, Germany: Gebrüder Borntraeger.
- NOWICKE J.W. (1969): Boraginaceae. - In: WOODSON R.E., SCHERY R.W. *et al.* (eds.): *Flora of Panama*. *Ann. Missouri Bot. Gard.* **56**: 33-69.
- NOWICKE J.W. & SKVARLA J.J. (1974): A palynological investigation of the genus *Tournefortia* (Boraginaceae). - *Amer. J. Bot.* **61**: 1021-1036.
- NUES R.W.v., RIENTJES J.M.J., SANDE C.A.F.M. VAN DER, ZERP S.F., SLUITER C., VENEMA J., PLANTA R.J. & RAUÉ H.A. (1994): Separate structural elements within Internal Transcribed Spacer 1 of *Saccharomyces cerevisiae* precursor ribosomal RNA direct the formation of 17S and 26S rRNA. - *Nuc. Acids Res.* **22**: 912-919.
- NUTTALL, T. (1837): Collections towards a flora of the territory of Arkansas. - *Trans. Amer. Philos. Soc., n.s.* **5**: 189.
- OLMSTEAD R.G., BREMER B., SCOTT K.M. & PALMER J.D. (1993): A parsimony analysis of the Asteridae sensu lato based on *rbcL* sequences. - *Ann. Missouri Bot. Gard.* **80** : 700-722.
- OLMSTEAD R.G. & FERGUSON D.M. (2001): A molecular phylogeny of the Boraginaceae/ Hydrophyllaceae s.str. (Abstract).
<http://www.2001.botanyconference.org/section21/abstracts/167.shtml>.
- PAL P. (1963) Comparative studies in four Species of *Heliotropium* L. - *Proc. Natl. Inst. Sci. India, B.* **29**: 1-40.

- PARMAR P.J. (1991): 88. Ehretiaceae. - In: SHETTY B.V. & V. SINGH (eds.): Flora of Rajasthan. pp. 500-525. Calcutta, India: Botanical Survey of India.
- PATE J.S., GUNNING B.E.S. & MILLIKEN F.F. (1970): Function of transfer cells in the nodal regions of stems, particularly in relation to the nutrition of young seedlings. - *Protoplasma* **71**: 313-334.
- PATE J.S. & GUNNING B.E.S. (1972): Transfer Cells. - *Annual Rev. Pl. Physiol.* **23**: 173-196.
- PERRY L.M. (1980): Medicinal plants of East and Southeast Asia: attributed properties and uses. Boraginaceae. pp. 60-62. Cambridge, Massachusetts, USA: MIT Press.
- PERVEEN A., QURESHI U.S., & QAISER M. (1995): Pollen flora of Pakistan. 4. Boraginaceae. - *Pak. J. Bot.* **27**: 327-360.
- POPOV M.G. (1953): Boraginaceae. - In: SHISHKIN B.K. (ed.): Flora SSSR **19**:97-691. Leningrad, USSR: Nauka.
- PRAKASH A.S., PEREIRA T.N., REILLY P.E.B. & SEAWRIGHT A.A. (1999) Pyrrolizidine alkaloids in human diet. - *Mutat. Res.* **443**: 53-67.
- QUIROZ-GARCIA D.L. & PALACIOS-CHÁVEZ R. (1985): Catálogo palinológico para la flora Veracruz. No. 27. Familia Boraginaceae. Género *Heliotropium*. - *Biotica* **10**: 341-358.
- QUIROZ-GARCIA D.L. & PALACIOS-CHÁVEZ R. (1986): Catálogo palinológico para la flora Veracruz. No. 32. Familia Boraginaceae. Género *Tournefortia*. - *Biotica* **10**: 55-61.
- QURESHI U.S. (1985): Studies on the pollen morphology of the genus *Heliotropium* L. from Pakistan. - *Pak. J. Bot.* **17**: 107-114.
- RAMBAUT A. (2001): Se-Al. Sequence alignment program v2.0a72. Oxford.
- RECORD J.S. & HESS R.W. (1941): American woods of the family Boraginaceae. - *Tropical Woods* **76**: 19-33.
- RIDLEY N.H. (1930): The dispersal of plants throughout the world. Ashford, Kent, Great Britain: L. Reeve & Co.
- RIEDL H. (1967): Boraginaceae. - In: H.K. RECHINGER (ed.): Flora Iranica, Lfg. **48**: 1-281. Graz, Austria: Akademische Druck- und Verlags-gesellschaft.
- RIEDL H. (1978): Die Gattung *Heliotropium* in Europa. - *Ann. Naturhist. Mus. Wien* **69**: 81-93.
- RITLAND C.E., RITLAND K. & STRAUS N.A. (1993): Variation in the ribosomal internal transcribed spacers (ITS1 and ITS2) among eight taxa of the *Mimulus guttatus* species complex. - *Mol. Biol. Evol.* **10**: 1273-1288.
- ROBINS D.J. (1985) : Pyrrolizidine Alkaloids. - *Nat. Prod. Rep.* **2**: 213-220.
- ROBINS D.J. (1986) : Pyrrolizidine Alkaloids. - *Nat. Prod. Rep.* **3**: 297-305.
- ROEMER J.J. & SCHULTES J.A. (1819): Systema vegetabilium. Vol. 4. Stuttgart, Germany: Cotta..
- ROSANOFF S. (1866): Morphologisch-embryologische Studien. - *Jahrb. Wiss. Bot.* **5**: 72-81.

- SAITOU N. & NEI M. (1987): The neighbor-joining method: a new method for reconstructing phylogenetic trees. - *Mol. Biol. Evol.* **4**: 406-425.
- SANKHLA N., ZIEGLER H., VYAS O.P., STICHLER W. & TRIMBORN P. (1975): Eco-physiological studies on Indian arid zone plants. V. A screening of some species for the C₄-pathway of photosynthetic CO₂-fixation. - *Oecologia*. **21**: 123-129.
- SCHEEL R., YBERT J.-P., BARTH O.M. (1996): Pollen morphology of the Boraginaceae from Santa Catarina State (southern Brazil), with comments on the taxonomy of the family. - *Grana* **35**: 138-153.
- SCHIBLER W. (1887): Beiträge zu einer vergleichend-systematischen Anatomie des Blattes und Stengels der Boragineen. Dissertation, Universität Bern.
- SCHULENBURG J.H.G.V.D., ENGLISCH U. & WÄGELE J.-W. (1999): Evolution of ITS1 rDNA in the Digenea (Platyhelminthes: Trematoda): 3' end sequence conservation and its phylogenetic utility. - *J. Mol. Evol.* **48**: 2-12.
- SCHUMANN K. (1889): Untersuchungen über das Borrageid. - *Ber. Dtsch. Bot. Ges.* **7**: 53-80.
- SCOTT P.E. & MARTIN R.F. (1984): Avian consumers of *Bursea*, *Ficus*, and *Ehretia* fruit in Yucatán. - *Biotropica* **16**: 319-323.
- SEIBERT J. (1978): Fruchtanatomische Untersuchungen an Lithospermeae (Boraginaceae). – *Diss. Bot.* **44**: 1-207.
- SELVI F. & BIGAZZI M. (2001): Leaf surface and anatomy in Boraginaceae tribe Boragineae with respect to ecology and taxonomy. – *Flora* **196**: 269-285.
- SERRATO-VALENTI G., CORNARA L., MODENESI P., PIANA M. & MARIOTTI M.G. (2000): Structure and histochemistry of embryo envelope tissues in the mature dry seed and early germination in *Phacelia tanacetifolia*. – *Ann. Bot.* **85**: 625-634.
- SIMMONS M.P. & OCHOTERENA. H. (2000): Gaps as characters in sequence-based phylogenetic analysis. - *Syst. Biol.* **49**: 369-381.
- SLOTTA T.A.B. (2000): Phylogenetic analysis of *Iliamna* (Malvaceae) using the Internal Transcribed Spacer region. Thesis, Virginia Polytechnic Institute and State University.
- SMITH R.A. & DEPAMPHLIS C.W. (1998): Phylogenetic placement of the holoparasitic family Lennoaceae: preliminary molecular evidence (Abstract). – *Amer. J. Bot. Suppl.* **85**: 157.
- SMITH R.A., FERGUSON D.M., BARKMAN T. J. & DEPAMPHLIS C.W. (2000): Molecular phylogenetic evidence for the origin of Lennoaceae: a case of adelphoparasitism in the angiosperms (Abstract). – *Amer. J. Bot.* **87**: 158.
- SMALL J.K. (1933): Manual of the southeastern flora. p. 1131. New York, USA: University of North Carolina Press.
- SNOW D.W. (1981): Tropical frugivorous birds and their food plants: a world survey. - *Biotropica* **13**: 1-14.
- SOLEREDER H. (1908): Systematische Anatomie der Dikotyledonen. Stuttgart, Germany: Enke.

- SOUÉGES R. (1943): Embryogénie et classification, 3^e fascicule: Essai d'un système embryogénique. - Actualités scientifiques et industrielles **1060**: 1-107.
- SPEGAZZINI C.L. (1901): Plantae novae nonnullae americae australis. - Comun. Mus. Nac. B. Aires **1**: 312-323.
- SPURR A.R. (1969): A low-viscosity epoxy resin embedding medium for electron microscopy. - J. Ultrastruct. Res. **26**: 31-43.
- SVENSSON H.G. (1925): Zur Embryologie der Hydrophyllaceen, Boraginaceen und Heliotropiaceen mit besonderer Rücksicht auf die Endospermibildung. pp. 1-176. Uppsala, Sweden: Uppsala Universitets Årsskrift.
- SWOFFORD D. L. (1998-2002): PAUP*: Phylogenetic analysis using parsimony (*and other methods), Version 4 to 4.0b1. Sunderland, Massachusetts, USA: Sinauer.
- TABERLET P., GIELLY L., PAUTOU G & BOUVET J (1991): Universal primers for amplification of three non-coding regions of chloroplast DNA. - Plant Mol. Biol. **17**: 1105-1109.
- TAKHTAJAN A. (1986): Floristic regions of the world. Berkeley, California, USA: University of California Press.
- TAGO M. (1999): Tokyo Metropolitan University, Makino Herbarium. GeneBank submission.
- THEOBALD W.L., KRAHULIK J.L. & ROLLINS R.C. (1979): Trichome description and classification. - In: METCLAFE C.R. & CHALK, L. (eds.): Anatomy of the dicotyledons, 2nd ed. Vol. **1**: 41-96. Oxford, Great Britain: Clarendon press.
- THOMPSON R.D., HUEROS G., BECKER H.A. & MAITZ M. (2001): Development and functions of seed transfer cells. - Pl. Sci. (Elsevier) **160**: 775-783.
- UPHOF J.C.Th. (1968): Dictionary of economic plants. Lehre, Germany: J. Cramer.
- VAN DE PEER Y. & DE WACHTER R. (1994): TREECON for Windows: a software package for the construction and drawing of evolutionary trees for the Microsoft Windows environment. - Comput. Appl. Biosci. **10**: 569-570.
- VERDCOURT B. (1987): A new genus *Nogalia* (Boraginaceae-Heliotropioideae) from Somaliland and southern Arabia. - Kew Bull. **43**: 431-435.
- VERDCOURT B. (1991): Boraginaceae. - In: POLHILL R.M. (ed.): Flora of tropical East Africa. pp. 1-77. Rotterdam, Brookfield: Balkema.
- YATSKIEVYCH G., MAXON J. & CHARLES T. (1986): A revision of the Lennoaceae. - Syst. Bot. **11**: 531-548.