

## **6. Literaturverzeichnis**

- 1 Chambliss L, Keil U, Dobson A, et al. for the WHO Monica Projekt. Population versus clinical view of case fatality from acute coronary heart disease: Results from the WHO Monica Project 1985 – 1990. *Circulation* 1997; 96:3849-59
- 2 Tunstall-Pedoe H, Kuulasmaa K, Mähönen M, et al. Contribution of trends in survival and coronary event rates to changes in coronary heart disease mortality: 10 years results from 37 WHO-MONICA project populations. *The Lancet* 1999; 355:1547-57
- 3 Ernst Bruckenberger. Heart Report 2002; <http://www.bruckenberger.de>
- 4 Gruppo italiano per lo studio della Streptochinasi nell`infarto miocardico. Long-term effects of intravenous thrombolysis in acute myocardial infarction: Final report of the GISSI Study. *The Lancet* 1987; 871-4
- 5 Randomized trial of intravenous streptokinase, oral aspirin, both or neither among 17 187 cases of suspected acute myocardial infarction: ISIS 2 ( Second International Study of Infarct Survival ). *The Lancet* 1988;349-60
- 6 The GUSTO angiographic investigators: The effects of tissue plasminogen activator, streptokinase or both on coronary-artery patency, ventricular function, and survival after acute myocardial infarction. *New Engl J Med* 1993; 329:1615-22
- 7 Lee KL, Woodlief LH, Topol EJ, et al. for the GUSTO-I Investigators. Predictors of 30-day mortality in the era of reperfusion for acute myocardial infarction. Results from an international trial of 41 021 patients. *Circulation* 1995; 91:1659-68
- 8 De Vreede J, Gorgels A, Verstraaten G, et al. Did prognosis after acute myocardial infarction change during the past 30 years ? A meta analysis. *J Am Coll Cardiol*; 1991: 698-706

- 9 Morrison LJ, Verbeeck PR, Mc Donald AC, et al. Mortality and prehospital thrombolysis for acute myocardial infarction: a meta-analysis. JAMA 2000; 283:2686-92
- 10 Rawles J. 10 years survival of patients with suspected acute myocardial infarction in a randomised comparison of prehospital and hospital thrombolysis ( GREAT ). Heart 2003; 89:563-4
- 11 Bonnefoy E, Lapostolle F, Leizorovicz A, et al. Primary angioplasty versus prehospital fibrinolysis in acute myocardial infarction: a randomised study. A comparison of angioplasty and prehospital thrombolysis in acute myocardial infarction ( CAPTIM ). Lancet 2002; 360:825-9
- 12 Andersen HR, Niesen TT, Rasmussen K, et al. A comparison of coronary angioplasty with fibrinolytic therapy in acute myocardial infarction. N Engl J Med 2003; 349:733-42
- 13 Keely EC, Boura JA, Grines CL, et al. Primary angioplasty versus intravenous thromolytic therapy for acute myocardial infarction: a quantitative review of 23 randomised trials. The Lancet 2003; 361:13-20
- 14 Zahn R, Schiele R, Gitt AK, et al. Impact of prehospital delay on mortality in patients with acute myocardial infarction treated with primary angioplasty and intravenous thrombolysis. Am Heart J 2001; 142:105-11
- 15 Gitt AK, Senges J. The Patient with acute myocardial infarction who does not receive reperfusion treatment. Heart 2001; 86: 243-5
- 16 Antithrombotic Trialists'Collaboration ( 2002 ). Collaborative meta-analysis of randomised trials of antiplatelet therapy for prevention of death, myocardial infarction, and stroke in high risk patients. BMJ 2002; 324:71-86

- 17 Montalescot G, Barragan P, Wittenberg O, et al. Platelet glycoprotein IIb/IIIa inhibition with coronary stenting for acute myocardial infarction. *N Engl J Med* 2000; 344:1895-903
- 18 Stone GW, Grines CL, Cox DA, et al. Comparison of angioplasty with stenting, with or without abciximab, in acute myocardial infarction. *N Engl J Med* 2002; 346:957-66
- 19 Lee DP, Herity NA, Hiatt BL, et al. Adjunctive platelet glycoprotein IIb/IIIa receptor inhibition with Tirofiban before primary angioplasty improves angiographic outcomes: results of the “tirofiban given in the emergency room before primary angioplasty” ( TIGER-PA ) pilot trial. *Circulation* 2003; 107:1497-501
- 20 Antman EM, Louwerenburg HW, Baars HF, et al. Enoxiparin as adjunctive antithrombin therapy for ST-elevation myocardial infarction: results of the ENTIRE-thrombolysis in myocardial Infarction ( TIMI ) 23 trial. *Circulation* 2002; 105:1642-9
- 21 Silvet H, Spencer F, Yarzebski J, et al. Communitywide trends in the use and outcomes associated with β-Blockers in patients with acute myocardial infarction. The Worcester Heart Attack Study. *Arch Intern Med* 2003;163:2175-83
- 22 Flather MD, Yusuf S, Kobe L, et al. Long-term ACE-Inhibitor therapy in patients with heart failure or left-ventricular dysfunction: a systematic overview of data from individual patients. ACE-Inhibitor Myocardial Collaborative Group. *The Lancet* 2000; 355:1575-81
- 23 La Rosa JC, He J, Vupputuri S. Effects of statins on risk of coronary disease: a meta-analysis of randomized controlled trials. *JAMA* 1999; 282:2340-6
- 24 Arbeitsgemeinschaft der Wissenschaftlichen Medizinischen Fachgesellschaften ( AWMF ) 2002. Leitlinien der deutschen Diabetes-Gesellschaft: Diabetes Typ 2.

## AWMF- Leitlinien –Register Nr 057/012K

25 Nissen SE, Tuzcu EM, Schoenhagen P, et al. Effect of intensive compared with moderate lipid-lowering therapy on progression of coronary atherosclerosis. JAMA 2004; 291:1071-180

26 Van den Hoogen PC, Feskens EJ, Nagelkerke NJ, et al. The relation between blood pressure and mortality due to coronary heart disease among men in different parts of the world. Seven countries study research group. N Engl J Med 2000; 342:1-8

27 The Multicenter Postinfarction Research Group. Risk stratification and survival after myocardial infarction. New Engl J Med 309 (1983),331-6

28 Bigger JT, Fleiss JL, Kleiger R et al. The relationships among ventricular arrhythmias, left ventricular dysfunction and mortality in the 2 years after myocardial infarction. Circulation 1984; 69:250-8

29 Hartikainen J, Malik M, Staunton A, et al. Distinction between arrhythmic and non-arrhythmic death after acute myocardial infarction based on heart rate variability, signal averaged electrocardiogram, ventricular arrhythmias and left ventricular ejection fraction. J Am Coll Cardiol 1996;296-304

30 La Rovere MT , Bigger jr J, Marcus FI, et al. Baroreflex sensitivity and heart rate variability in prediction of total cardiac mortality after myocardial infarction . ATRAMI ( Autonomic Tone and Reflexes After Myocardial Infarction ) Investigators. The Lancet 1998; 351: 478-84

31 Rosenbaum DS, Lance E, Jackson LE, et al. Electrical alternans and vulnerability to ventricular arrhythmias. N Engl J Med 1994; 330:235-41

- 32 Buxton AE, Lee KL, Hafley GE, et al. Relation of ejection fraction and inducible ventricular tachycardia to mode of death in patients with coronary artery disease: an analysis of patients enrolled in the multicenter unsustained tachycardia trial. *Circulation* 2002; 106:2466-72
- 33 Buxton AE, Lee KL, Di Carlo L, et al. Electrophysiologic testing to identify patients with coronary artery disease who are at risk for sudden death. Multicenter Unsustained Tachycardia Trial Investigators. *N Engl J Med* 2000; 342:1937-45
- 34 Moss AJ, Hall WJ, Cannom DS, et al. Improved survival with an implanted defibrillator in patients with coronary heart disease at high risk for ventricular arrhythmia. Multicenter Automatic Defibrillator Implantation Trial Investigators. *N Engl J Med* 1996; 335:1933-40
- 35 Moss AJ, Zareba W, Hall WJ, et al. Prophylactic implantation of a defibrillator in patients with myocardial infarction and reduced ejection fraction. Multicenter Automatic Defibrillator Implantation Trial II Investigators. *N Engl J Med* 2002; 342:877-83
- 36 Löwel H, Engel S, Hörmann A, et al. für das MONICA-Augsburg Herzinfarkt-Registerteam. Akuter Herzinfarkt und plötzlicher Herztod aus epidemiologischer Sicht. *Intensivmedizin* 1999; 36: 652-61
- 37 Behrens S, Boecker D, Reek S, et al. The prevention of sudden cardiac death ( PreSCD ) registry: final 18-month-follow-up of a prospective multicenter registry in 5330 patients post myocardial infarction. Abstract for PreSCD 1-Registry; NASPE 2004
- 38 Granger CB, Goldberg RJ, Dabbous O, et al. Predictors of hospital mortality in the global registry of acute coronary events ( GRACE ). *Arch Intern Med.* 2003; 163:2345-53
- 39 Santopinto JJ, Fox KA, Goldberg RJ, et al. Creatinine clearance and adverse

hospital outcomes in patients with acute coronary syndromes: findings from the global registry of acute coronary events ( GRACE ). Heart 2003; 89:1003-08

40 Krumholz HM, Chen J, Chen YT, et al. Predicting one-year mortality among elderly survivors of hospitalisation for an acute myocardial infarction; results from the Cooperative Cardiovascular Project. J Am Coll Cardiol 2001; 38:453-9

41 Newby LK, Bhapkar MV, White HD , et al. for the Symphony and 2<sup>nd</sup> Symphony Investigators. Predictors of 90-day outcome in patients stabilized after acute coronary syndromes. European Heart Journal 2003; 24:172-82

42 Burns RJ, Gibbons RJ, Yi Q, et al. The relationship of left ventricular ejection fraction, endsystolic volume index and infarct size to six-month mortality after hospital discharge following myocardial infarction treated by thrombolysis.

J Am Coll Cardiol 2002;39:30-6

43 Copie X, Hnatkova K, Staunton A, et al. Predictive power of increased heart rate versus depressed left ventricular ejection fraction and heart rate variability for risk stratification after myocardial infarction. Results of a two year follow-up study.

J Am Coll Cardiol 1996; 27:270-6

44 Richards D, Byth K, Ross D, e al. What is the best predictor of spontaneous ventricular tachycardia and sudden death after myocardial infarction ?  
Circulation 1991; 83:756-63

45 Yap Y, Duong T, Bland M, et al. Left ventricular ejection fraction in the thrombolytic era remains a powerful predictor of long-term but not short-term all cause, cardiac and arrhythmic mortality after myocardial infarction - a second meta-analysis of 2828 patients.  
Heart 2000; 83:55

- 46 Hanania G, Cambou JP, Gueret P, et al. Management and in-hospital outcome of patients with acute myocardial infarction admitted to intensive care units at the turn of the century: The French Nationwide USIC 2000 Registry. Heart 2004; 90:1404-10
- 47 Bauer A, Guzik P, Barthel P, et al. Reduced prognostic power of ventricular late potentials in post-infarction patients of the reperfusion era. European Heart Journal 2005; 26:755-61
- 48 Mäkikallio T, Barthel P, Schneider R, et al. Prediction of sudden cardiac death after acute myocardial infarction: The role of holter monitorig in the modern treatment era. European Heart Journal 2005; 26:762-9
- 49 Fathy GJ, Pinski SL, Miller DP, et al. Natural history of isolated bundle branch block Am J Cardiol 1996; 77:1185-90
- 50 Rabkin SW, Matheson FA, Tate RB. Natural history of left bundle branch block. Br Heart J 1980; 43:164-9
- 51 Schneider JF, Thomas Jr HE, Sorlie P, et al. Comparative features of newley acquired left and right bundle branch block in the general Population. The Framingham Study. Am J Cardiol 1981; 47:931-40
- 52 Lee SJ, Mc Culloch C, Mangat I, et al. Isolated bundle branch block and left ventricular dysfunction. J Cardiac Fail 2003; 9:87-92
- 53 Zimetbaum PJ, Buxton AE, Batsford W, et al. Elektrocardiographic predictors of arrhythmic death and total mortality in the multicenter unsustained tachycardia trial Circulation 2004; 110:766-9
- 54 Sternstrand U, Tabrizi F, Lindbäck J, et al. Comorbidity and myocardial dysfunction

are the main explanation for the higher 1 year mortality in acute myocardial infarction with left bundle branch block. Circulation 2004; 110:1896-902

55 Brilakis ES, Wright RS, Kopecky SL, et al. Bundle branch block as a predictor of long term survival after acute myocardial infarction. Am J Cardiol 2001; 88:205-9

56 Newby KH, Pisano E, Krucoff MW, et al. Incidence and clinical relevance of the occurrence of bundle branch block in patients treated with thrombolytic therapy. Circulation 1996; 94:2424-8

57 Balderassoni S, Opasich C, Gorini M, et al. Italian Network on Congestive Heart failure investigators. Left bundle branch block is associated with increased 1-year sudden and total mortality rate in 5517 outpatients with congestive heart failure: a report from the Italian Network on congestive heart failure. Am Heart J 2002; 143:398-405

58 Hesse B, Diaz LA, Snader CE, et al. Complete bundle branch block as an independent predictor of all cause mortality: report of 7073 patients referred for nuclear exercise testing. Am J Med 2001; 110:253-9

59 Pinkau T, Mann JF, Ndrepaa G, et al. Coronary revascularisation in patients with renal insufficiency: restenosis rate and cardiovascular outcomes ( In Process Citation ). Am J Kidney Dis 2004; 44: 627-35

60 Best PJ, Berger PB, Davies BR, et al. Impact of mild or moderate chronic kidney disease on the frequency of restenosis: Results from the PRESTO trial. J Am Coll Cardiol 2004; 44:1786-91

61 Szczech LA, Best PJ, Crowley E, et al. Outcomes of patients with chronic renal insufficiency in the bypass angioplasty revascularisation investigation.

Circulation 2002; 105:2253-8

62 Gibson CM, Pinto DS, Murphy SA, et al. Association of Creatinine and Creatinine clearance on presentation in acute myocardial infarction with subsequent mortality.

J Am Coll Cardiol 2003; 42:1535-43

63 Dumaine R, Collet JP, Tanguy ML, et al. Prognostic significance of renal insufficiency in patients presenting with acute coronary syndrome ( the Prospective Multicenter SYCOMORE study ). Am J Cardiol 2004; 94:1543-7

64 Mc Cullough PA, Soman SS, Sha SS, et al. Risk associated with renal dysfunction in the coronary care unit. J Am Coll Cardiol 2000; 36:679-84

65 Fried LF, Shlipak MG, Crump C, et al. Renal insufficiency as a predictor of cardiovascular outcomes and mortality in elderly individuals. J Am Coll Cardiol 2003; 41:1364-72

66 Mann J, Gerstein HC, Hertzel C, et al. for the HOPE Investigators. Renal insufficiency as a predictor of cardiovascular outcomes and the impact of Ramipril : the HOPE randomized trial. Ann Intern Med 2001; 134:629-36

67 Beattie JN, Somann SS, Sandberg KR, Yee J, et al. Determinants of mortality after myocardial infarction in patients with advanced renal dysfunction. Am J Kidney Dis 2000; 37:1191-200

68 Wright RS, Reeder GS, Herzog CA, et al. Acute myocardial infarction and renal dysfunction: a high risk combination. Ann Intern Med 2002,137:563-70

69 Aronow WS, Ahn C, Mercando AD, et al. Prevalence of coronary artery disease, complex ventricular arrhythmia, and silent myocardial ischemia and incidence of new coronary

events in older persons with chronic renal insufficiency and with normal renal function

Am J Cardiol 2000; 86:1142-3

70 Cerne D, Kaplan-Pavlovic S, Kranjec I, et al. Mildly elevated serum creatinine concentration correlates with the extent of coronary atherosclerosis. Ren Fail 2000; 22:799-808

71 Shlipak MG, Fried LF, Crump C, Bleyer AJ, et al. Cardiovascular disease risk status in elderly persons with renal insufficiency. Kidney Int 2002; 62:997-1004

72 Al Suwaidi J, Reddan DN, Williams K, et al. Prognostic implications of abnormalities in renal function in patients with acute coronary syndromes. Circulation 2002; 106:974-80

73 Herzog CA, Ma JZ, Collins AJ, et al. Poor long term survival after acute myocardial infarction among patients on long-term dialysis. N Engl J Med 1998; 339:799-805

74 Chertow GM, Normand SLT, Silva LR, et al. Survival after acute myocardial infarction among patients on long term dialysis. Results from the cooperative cardiovascular project. Am J Kidney Dis 2000; 35: 1044-51

75 Anavekar NS, Mc Murray JJ, Velazquez EJ, et al. Relation between renal dysfunction and cardiovascular outcomes after myocardial infarction. N Engl J Med 2004; 351:1285-95

76 McCullough PA, Nowak RM, Foreback C, et al. Emergency evaluation of chest pain in patients with advanced kidney disease. Arch Intern Med 2002; 25:2464-8

77 Yusuf S, Reddy S, Ounpuu, et al. Global burden of cardiovascular diseases: Part I: general considerations, the epidemiologic transition, risk factors, and impact of urbanisation. Circulation 2001; 104:2746-53

78 Khot UN, Khot MB, Bajzer CT, et al. Prevalence of conventional risk factors in

patients with coronary heart disease. JAMA 2003; 290:898-904

79 Bartnik M, Malmberg K, Norhammar A, et al. Newly detected abnormal glucose tolerance: an important predictor of long-term outcome after myocardial infarction. European Heart Journal 2004; 25:1990-7

80 Gustafsson I, Hildebrandt P, Seibaek M, et al. Long-term prognosis of diabetic patients with myocardial infarction: Relation to antidiabetic treatment regimen.

The Trace Study Group. European Heart Journal 2000; 21:1937-43

81 Assmann G; Cullen P, Schulte H et al. Simple scoring scheme for calculating the risk of acute coronary events based on a 10-year follow-up of the prospective cardiovascular Munster ( PROCAM ) study. Circulation 2002; 105:310-5

82 Hohnloser SH, Kuck KH, Dorian P et al. Prophylactic use of an implantable cardioverter-defibrillator after acute myocardial infarction. N Engl J Med 2004; 351:2481-8

83 Huikuri HV, Tapanainen JM, Lindgren K, et al. Prediction of sudden cardiac death after myocardial infarction in the beta-blocking era. J Am Coll Cardiol 2003; 42: 652-8

84 ISIS 1 ( First International Study of Infarct Survival ) collaborative group. Randomised trial of intravenous atenolol among 16 027 cases of suspected acute myocardial infarction: ISIS 1. The Lancet 1986; II:57-66

85 Hjalmerson A, Elmfelt D, Herlitz J, et al. Effect on mortality of metoprolol in acute myocardial infarction. A double-blind randomised trial. The Lancet 1981: 823-7

86 The CAPRICORN investigators. Effect of carvedilol on outcome after myocardial infarction in patients with left ventricular dysfunction. The CAPRICORN randomised trial. Lancet 2001; 357:1385-90