

9 LITERATURVERZEICHNIS

- Abraham RT, Wiederrecht GJ. Immunopharmacology of rapamycin. *Ann Rev Immunol* (1996) 14 : 483-510
- Ahmed AR, and Hombal SM. Cyclophosphamide. *J Am Acad Dermatol* 1 (1984): 1115- 1126
- Ahmad W, Pantaleyev A and Christiano AM. The molecular basis of congenital atrichia in humans and mice: mutation in the hairless gene. *J Invest Dermatol* (1999) 4 : 240-2
- Aktories K, Unger C. Mittel zur Behandlung von Tumoren-Tumorchemotherapie. In Aktories K, Förstermann W, Hofmann FB, Starke K. (eds) *Allgemeine und spezielle Pharmakologie und Toxikologie* (9-te Auflage) Urban/ Fischer, München /Jena. (2005) 921-959
- Al Daraji WI, Grant KR, Ryan K, Saxston A, and Reynolds NJ. Localization of calcineurin / NFAT in human skin and psoriasis and inhibition of calcineurin / NFAT activation in human keratinocytes by cyclosporin A. *J Invest Dermatol* (2002) 118 : 779-788
- Allombert-Blaise C, Tamiji S, Mortier L, et al. Terminal differentiation of human epidermal keratinocytes involves mitochondria- and caspasedependent cell death pathway. (2003) *Cell Death Differ.* 10 : 850-2.
- Arck PC, Slominski AA, Theoharides TC et al. Neuroimmunology of stress : skin takes center stage. *J Invest Dermatol* . (2006) *In press*
- Ashkenazi A, Dixit VM. Death receptors : signaling and modulation. *Science* (1998) 281 : 1305-8
- Aurelian L. Cross talk of signaling and apoptotic cascades in the CNS : target for virus modulation. *Front Biosci.* (2005) 10 : 2776-87
- Batchelor D. Hair and cancer chemotherapy : consequences and nursing care a literature study. *Eur J Cancer Care (Engl)* (2001) 10 : 147-63.
- Baker BS, Griffiths CEM, Lambert S, et al. The effects of cyclosporine A on T lymphocyte and dendrite cell subpopulation in psoriasis. *Br J Dermatol* (1987) 116 : 503-510
- Barinov EF, Sulaeva ON. Histophysiology of hair follicles: current concept. *Usp Fiziol Nauk.* (2004) 35 : 65-77.
- Bennet A. Hirsutism and hypertrichosis in adults : investigations and treatment *Ann Dermatol Venereol* 1(2002) 29 : 804-12.
- Bierer BE, Mattila PS, Standaert RF et al. Two distinct signal transmission pathways in T lymphocytes are inhibited by complexes formed between an immunophilin and either FK506 or rapamycin. *Proc Natl Acad Sci USA.* (1990) 87 : 9231-5

- Bleiker TO, Nicolaou N, Traulsen J, Hutchinson PE. Atrophic telogen effluvium from cytotoxic drugs and a randomized controlled trial to investigate the possible protective effect of pretreatment with a topical vitamin D analogue in humans. *Br J Dermatol.* (2005) 153 : 103-12.
- Boatright KM, Salvesen GS. Mechanisms of caspase activation. *Curr Opin Cell Biol* (2003) 15 : 725-31
- Borel JF, Feuer C, Gubler HU, Stähelein H. Biological effects of cyclosporin A, a new anti lymphocytic agent. *Agents Actions* (1976) 6 : 468-475
- Bosse, K. Comparative studies on the physiology and pathology of hair shedding with special consideration of its synchronization. VII. Discussion of personal test results *Hautarzt.* (1967) 18 : 274-83.
- Botchkarev VA, Komarova EA, Siebenhaar F, et al. p53 is essential for chemotherapy-induced hair loss. *Cancer Res.* (2000) 60 : 5002-6
- Botchkarev VA, Komarova EA, Siebenhaar F, et al. p53 Involvement in the control of murine hair follicle regression. *Am J Pathol.* (2001) 158 : 1913-9.
- Botchkarev VA. Molecular mechanisms of chemotherapy-induced hair loss. *J Invest Dermatol Symp Proc.* (2003a) 8 : 72-5.
- Botchkarev VA, Yaar M, Girchlest B and Paus R (p75 Neurotrophin receptor antagonist retards apoptosis-driven hair follicle involuton (catagen). *The Soc Invest Dermatol* (2003b) 120 : 168-169
- Botchkarev VA, Sharov AA. BMP signaling in the control of skin development and hair follicle growth. *Differentiation.* (2004) 72 : 512-26
- Botchkareva NV, Ahluwalia G, Shander D. Apoptosis in the hair follicle. *J Invest Dermatol.* (2006) 126 : 258-64.
- Branzan AL, Landthaler M, Szeimies RM. Skin changes with chemotherapy. *Hautarzt.* (2005) 56 : 591-602
- Braun-Falco, O. Über Veränderungen im Hautbindegewebe während des Haarwachstumszyklus bei der Maus. *Arch. klin exp Derm.* (1959) 209 : 426-432
- Braun-Falco, O. Klinik und Pathomechanismus der Endoxan-Allopecie als Beitrag zum Wesen cytostatischer Alopecien. *Archiv klin exp Dermatol,* (1961a) 212 : 194-216
- Braun-Falco, O. A contribution to a study of falling out of the hair under the influence of chemotherapy. *Dermatol Wochenschr.* (1961b) 23, 144 : 1083-4.
- Braun-Falco, O. Dynamics of normal and pathological hair growth. *Arch klin exp Dermatol.* (1966) 227 : 419-52.

- Butcher, EO.: Development of the pillary system and the replacement of the hair in mammals. Ann NY Acad Sci (1951) 53 : 508-515
- Cai L, Hales BF, Robaire B. Induction of Apoptosis in the germ cells of adult male rats after exposure to cyclophosphamide. Biol Reprod (1998) 56 : 1490-1497
- Cardenas ME, Zhu D, Heitman J. Molecular mechanisms of immunosuppression by cyclosporine, FK506, and rapamycin. Curr Opin Nephrol Hypertens. (1995) 4 : 472-7.
- Chase, HB, Rauch, H, Smith, VW. Critical stages of hair follicle development and pigmentation in the mouse. Physiol Zool (1951) 25 : 1-19
- Chase, HB and Eaton GJ. The Growth of Hair follicles in waves. Ann NY Acad Sci (1959) 83 : 365-368
- Chaturvedi V, Sitailo LA, Bodner B, et al. Defining the caspase-containing apoptotic machinery contributing to cornification in human epidermal equivalents. Exp Dermatol. (2006) 15 : 14-22
- Chen G, Baechle A, Nevins TD, Oh S, et al. Protection against cyclophosphamide-induced alopecia and inhibition of mammary tumor growth by topical 1,25-dihydroxyvitamin D3 in mice. Int J Cancer (1998) 75 : 303-9
- Cece R, Cazzaniga S, Morelli D, et al. Apoptosis of hair follicle cells during doxorubicine-induced alopecia in rats. Lab Invest (1996) 75 : 601-609
- Cerottini JP, Panizzon RG, de Viragh PA. Multifocal alopecia areata during systemic cyclosporine A therapy. Dermatology (1999) 198 : 415-7
- Coggins PR, Ravdin RG, Eisman SH. Clinical evaluation of a new alkylating agent Cytosan, Cyclophosphamide. Cancer (1990) 13 : 1254-1260
- Cohen JJ, Duke RC, Fadok VA, Sellins KS. Apoptosis and programmed cell death. Ann Rev Immunol. (1992) 10 : 267-294
- Colitti M, Allen SP, Price JS. Programmed cell death in the regenerating deer antler. J Anat (2005) 207 : 339-51
- Colvin M, Brondrett RB, Kann HN. Alkylating properties of phosphoramidate mustard. Cancer Res (1976) 36 : 1121-1126
- Comalada M, Valledor AF, Sanchez-Tillo E, et al.. Macrophage colony-stimulating factor-dependent macrophage proliferation is mediated through a calcineurin-dependent but immunophilin-dependent mechanism that mediates the activation of external regulated kinases. Eur J Immunol (2003) 33 : 3091-100

- Connors TA, Cox PJ, Farmer PB. Some studies of the active intermediates formed in the microsomal metabolism of cyclophosphamide and isophosphamide. *Biochem Pharmacol* (1974) 123 :115-129
- Cotsarelis G, Sunn TT, Lavker RM. Label-retaining cells reside in the bulge area of pilosebaceous unit : Implications for follicular stem cells, hair cycle, and skin carcinogenesis. *Cell* 61 : 1329-37
- Cotsarelis G, and Millar S. Towards a molecular understanding of hair loss and its treatment. *Trends Mol Med.* (2001) 7 : 293-301
- Courtois M, Loussouarn G, Hourseau C, Grollier JF. Hair cycle and alopecia. *Skin Pharmacol* (1994) 7 : 84-89
- Crouse RG, Van Scott EJ. Changes in scalp hair roots as a measure of toxicity from cancer chemotherapeutic drugs. *J Invest Dermatol* (1960) 35 : 83-90
- Crow MT, Mani K, Nam YJ, Kitsis RN. The mitochondrial death pathway and cardiac myocyte apoptosis. *Circ Res.* (2004) 95 : 957-70
- Danilenko DM, Ring BD, Pierce GF. Growth factors and cytokines in hair follicle development and cycling : recent insights from animal models and the potential for the clinical therapy. *Mol Med Today* (1996) 2 : 460-46
- Daniel NN, Korsmeyer SJ. Cell death : critical control points. *Cell* (2004) 116 : 205-19
- Dave S and Thappa DM. Peculiar pattern of nail pigmentation following cyclophosphamide therapy. *Dermatology Online Journal* . (2003) 9 (3) : 14
- Davidhizar R, Bartlett D. When your patients lose their hair. *Today's OR Nurse.* (1993) 5 : 39-42
- Davis, BK. Phases of the hair growth cycle. *Nature, Lond.* (1962) 194 : 694-
- Davis ST, Benson BG, Bramson HN, et al. Prevention of chemotherapy-induced alopecia in rats by CDK inhibitors. *Science* (2001) 291 : 134-137
- DeSpain JD. Dermatologic toxicity of Chemotherapy. *Seminars of Oncology.* (1992) 19 : 501-507
- De Paulis A, Stellato C, Cirillo R, et al. Anti-inflammatory effect of FK-506 on human skin mast cells. *J Invest Dermatol* (1992) 99 : 723-740
- Dry FW. Coat of mouse (*mus musculus*). *J Genetics* (1926) 16 : 287-340
- Dryfuss M, Härrä E, Hoffmann H, et al. Cyclosporine A and C: new metabolites from *Trichoderma polysporum* (*Link ex Pers*) Rafai. *European J Appl Microbiol* (1976) 3 : 125-133
- Dudziak, R.: *Lehrbuch der Anästhesiologie.* Stuttgart, New York , (1980) 241-248
- Dumont FJ, Su Q. Mechanism of action of the immunosuppressant rapamycin. *Life Sci* (1996) 58 : 373-95

- Durward, A and Rudall, KM. Hair growth in rat. *J Anat.* (1949) 83 : 325-335
- Ebling FJG, Hale P, Randal VA. Hormones and hair growth. In Goldsmith L (eds) *Biochemistry and Physiology of the Skin.* (2nd edn) NY : Oxford University Press. (1991) 322-350
- Ekert PG, Vaux DL. The mitochondrial death squad : hardened killers or innocent bystanders? *Curr Opin Cell Biol* (2005) 17 : 626-30
- Farmer PB. Metabolism and reactions of alkylating agents. *Pharmacol Ther.* (1987) 35 : 301-8
- Ferri KF, Kroemer G. Organelle-specific initiation of cell death pathways. *Nat Cell Biol* (2001) 3 : E255-63
- Fischer G, Wittmann-Liebold B, Lang K, et al. Cyclophilin and peptidyl-prolyl cis-trans isomerase are probably identical proteins. *Nature.* (1989) 337 : 476-8
- Fisher DE. Apoptosis in cancer therapy : crossing the threshold. *Cell.* (1994) 78 : 539-542
- Fink SL and Coocson BT. Apoptosis, Pyroptosis and Necrosis : Apoptosis, pyroptosis, and necrosis: mechanistic description of dead and dying eukaryotic cells. *Infect Immun* (2005) 73 : 1907-1916.
- Fitzpatrick JE, Hood AF. Histopathologic reactions to chemotherapeutic agents. *Adv Dermatol* (1988) 3 : 161-184
- Fitzpatrick JE. The cutaneous histopathology of chemotherapeutic reactions. *J Cutan Pathol* (1993) 20 : 1-14
- Foitzik K, Paus R. Regulation of human hair growth and therapeutic options. *J Deutsch Dermatol Ges.* (2005) 3 : 901-17.
- Freiser AS. A note on the growth of Rex Angora coats. *J Genet* (1952) 51 : 237-242
- Friedman, J. and Weissman, I. Two cytoplasmic candidates for immunophilin action are revealed by affinity for a new cyclophilin : one in the presence and one in the absence of CsA. *Cell* (1991) 66 : 799-806
- Friedman JS, Lowe SW. Control of apoptosis by p53. *Oncogene* (2003) 22 : 9030-9040
- Furue M, Katz SI. The effect of cyclosporine on epidermal cells. I. Cyclosporine inhibits accessory cell functions of epidermal Langerhans cells in vitro. *J Immunol.* (1988a) 15 : 140 : 4139-43
- Furue M, Gaspari AA, Katz SI. The effect of cyclosporin A on epidermal cells. II. Cyclosporin A inhibits proliferation of normal and transformed keratinocytes. *J Invest Dermatol.* (1988b) 90 : 796-800
- Gothel SF, Marahiel MA. Peptidyl-prolyl cis-trans isomerases, a superfamily of ubiquitous folding catalysts. *Cell Mol Life Sci.* (1999) 55 : 423-36.

- Gandarillas A, Goldsmith LA, Gschmeissner S, et al. Evidence that apoptosis and terminal differentiation of epidermal keratinocytes are distinct processes. *Exp Dermatol.* (1999) 8 : 71-79
- Gavter-Gvili A, Sredni B, Gal R, et al. Cyclosporin A induced hair growth in mice is associated with inhibition of calcineurin-dependent activation of NFAT in follicular keratinocytes. *Am J Physiol Cell Physiol* (2003) 284 : C 1593-C 1603
- Gavter-Gvili A, Kalechman Y, Sredni B, et al. Cyclosporine A-induced hair growth in mice is associated with inhibition of hair follicle regression. *Arch Dermatol Res* (2004) 296 : 265-269
- Giaccia AJ, and Kastan MB. The complexity of p53 modulation : emerging patterns from divergent signals. *Genes Dev* (1998) 12 : 2973-2983
- Goldberg MT, Tackaberry LE, Hardy MH, Noseworthy JH.: Nuclear aberrations in hair follicle cells of patients receiving cyclophosphamide. *Arch Toxicol* (1990) 64 : 116-121
- Goldhirsch A, Kiser J, Joss R, et al. Prevention of cytostatic-related hair loss by hypothermia of a hairy scalp using a cooling cap. *Schweiz Med Wochenschr* (1982) 112 : 568-571
- Grevelman EG, Breed WP. Prevention of chemotherapy-induced hair loss by scalp cooling. *Ann Oncol* (2005) 16 : 352-8
- Gupta AK, Ellis CN, Goldfarb MT, et al Cyclosporine A. *Clin Dermatol.* (1989a) 7 : 98-110.
- Gupta AK, Baadsgaard O, Ellis CN, et al. Lymphocytes and macrophages of the epidermis and dermis in lesiona psoriatic skin, but not epidermal Langerhans cells, are depleted by treatment with cyclosporine A. *Arch Dermatol Res* (1989b) 281 : 219-26
- Hasmall SC, Roberts RA. The perturbation of apoptosis and mitosis by drugs and xenobiotics. *Pharmacol Ther.* (1999) 82 : 63-70.
- Handjiski BK, Eichmüller S, Hofmann U, et al. Alkaline phosphatase activity and localisation during the murine hair cycle. *Br J Dermatol.* (1994)131 : 303-31
- Handjiski B, Mueller-Roever S, Peters E, Paus R. Guidelines and techniques for the accurate classification of chemotherapy-induced hair follicle dystrophy. *J Invest Dermatol* (2001) 117 : 267
- Handschumacher RE, Harding MW, Rice J, Drugge RJ and Speicher DW. Cyclophilin a specific cytosolic bindin protein for cyclosporin A. *Science* (1984) 226 : 544-547
- Hangartner MO. The biochemistry of apoptosis. *Nature* (2000) 407 : 770-776
- Hampton PJ and Reynolds NJ. Calcineurin inhibitors for the treatment of skin disease: how do they work? *Clinical & Experimental Allergy* (2005) 35, 5 : 549-550
- Headington JT. Telogen effluvium. *Arch Dermatol .* (1993) 129 : 356-363

- Hemenway CS, Heitman J. (1999) Calcineurin. Structure, function, and inhibition. *Cell Biochem Biophys.* (1999) 30 : 115-51
- Hendrix S, Handjiski B, Peters EMJ and Paus R. A Guide to Assessing Damage Response Pathways of the Hair Follicle : Lessons from Cyclophosphamide-induced Alopecia in Mice. *J Invest Dermatol* (2005) 125 : 42-51
- Hengartner MO. The biochemistry of apoptosis. *Nature Med* (2000) 407 : 770-776
- Henry LB, Horn TD. Chemotherapy and keratinocytes. *J Cutan Pathol* (2002) 29 : 575-8
- Hoetzenecker W, Meingassner JG, Ecker R et al. Corticosteroids but not Pimecrolimus affect Viability, Maturation and Immune Function of Murine Epidermal Langerhans Cells. *J Invest Dermatol* (2004) 122 : 673-680
- Hibino T, Nishiyama T. Role of TGF- β 2 in the human cycle. *J Dermatol Science* (2004) 35 : 9-18
- Ho S, Clipstone N, Timmermann L, et al. The mechanism of action of cyclosporin A and FK506. *Clin Immunol Immunopathol* (1996) 80 : S40-5.
- Hohorst HJ, Bielicki L, Voelcker G. The enzymatic basis of cyclophosphamide specificity. *Adv Enzyme Regul.* (1986)25 : 99-122
- Hog, H. Apoptosis : no healthy multi-cellular organisms without regulated cell suicide. *Schweiz Rundsch Med Prax.* (2005) 94 : 1215-8
- Hood AF. Cutaneous side effects of cancer chemotherapy. *Med Clin North Am* (1986) 70 : 187-209
- Horn TD. A antineoplastic chemotherapy, sweat, and the skin . *Arch Dermatol* (1997a) 133 : 905-910
- Horn TD. Cutaneous toxicities of drugs. In: Elder D, Elenitsas R, Jaworsky G, Jonson B Jr. (eds.) *Levers histopathology of the skin* (8th edn) NY: Lippincot-Raven. (1997b) 289-301
- Howser DM. Alopecia : the problem of alopecia in cancer. In : Groenewald SL, Frogge MH, Goodman M, (eds) *Cancer Symptom Management.* Jones and Bartlett : Boston ,MA (1996) 261-268
- Huai Q, Kim HY, Liu Y, Zhao Y, et al. Crystal structure of calcineurin-cyclophilin-cyclosporin shows common but distinct recognition of immunophilin-drug complexes. *Proc Natl Acad Sci USA.* (2002) 99 : 12037-12042.
- Hussein AM, Jimenez JJ, McCall CA, Yunis AA. Protection from Chemotherapy-induced Alopecia in a rat model. *Science* (1990) 249 : 1564-1566

- Hussein AM. IL-1 protects against 1- β -D-Arabinofuranosylcytosine-induced alopecia in the newborn rat animal model. *Cancer Res* (1991): 51 : 3329-3330
- Hussein AM, Stuart A, Peters WP. Protection against chemotherapy-induced alopecia by cyclosporin A in the newborn rat animal model. *Dermatology*. (1995) 190 : 192-6
- Iaschke H, Lemasters JJ. Apoptosis vs. oncotic necrosis in hepatic ischemia / reperfusion injury. *Gastroenterology* (2003) 125 : 1246-1257
- Ito M, Sato Y. Dynamic ultrastructural changes of the connective tissue sheath of human hair follicles during hair cycle. *Arch Dermatol Res*. (1990) 282 : 434-41
- Ishibashi M, Moutabarrik A, kameoka H, et al. Effect of FK 506 on generation of activated macrophages invading rejected skin allografts in rats: discrepancy of in vitro versus in vivo effects. *Transplant Proc*. (1991) 23 : 2967-9.
- Ito M, Kizawa K, Hamada K, Cotsarelis G. Hair follicle stem cells in the lower bulge form the secondary germ, a biochemically distinct but functionally equivalent progenitor cell population, at the termination of catagen. *Differentiation*. (2004) 72 : 548-57
- Ivery MTG. Immunophilins : switched on protein binding domains? *Med Res Rev*. (2000) 20 : 452-84.
- Iwabuchi T, Maryama T, Sei Y, Adachi K. Effects of immunosuppressive peptidylpropyl *cis-trans* isomerase (PPIase) inhibitors, cyclosporin A, FK506, ascomycin and rapamycin, on hair growth initiation in mouse immunosuppression is not required for hair growth. *J Dermatol Sci* (1995) 9 : 64-69
- Jahoda CA and Reynolds AJ. Dermal-epidermal interactions. Adult follicle derived cell populations and hair growth. *Dermatol Clin* (1996) 14 : 573-583
- Jessen RT, Straight M, Smith EB. Cutaneous and other complications of cyclophosphamide : a brief review. *Rock Mount Med J* (1978) 75 : 83-90
- Jiang X, Wang Y. Cytochrome C promotes caspase-9 activation by inducing nucleotide binding to Apaf-1. *Biol Chem* . (2000) 275 : 31199-203
- Jiang H, Yamamoto S, Kato R. Induction of anagen in telogen mouse skin by topical application of FK 506, a potent immunosuppressant. *J Invest Dermatol* (1995) 104 : 523-525
- Jimbow K, Fitzpatrick TB, Wick MM. Biochemistry and physiology of melanin pigmentation. *Physiology Bioc NY University Press*, (1991) 545-565
- Jimenez JJ, Alvarez E, Bustamante CD, Yunis AA. Pretreatment with 1,25(OH)₂D₃ protects from Cytoxan-induced alopecia without protecting the leukemic cells from Cytoxan. *Am J Med Sci*. (1995) 310 : 43-7

- Kanduc D, Mittelman A, Serpico R, et al. Cell death : apoptosis vs. necrosis. *Int J Oncol* (2002) 21 : 165-70.
- Karashima T, Hachisuka H, Sasai Y. FK506 and cyclosporin A inhibit growth factor stimulated human keratinocyte proliferation by blocking cells in the G0/G1 phases of the cell cycle. *J Dermatol Sci.* (1996) 12 : 246-54.
- Kato N, Halprin M, Taylor JR. Cyclosporin A does not inhibit epidermal cell growth at therapeutic levels. *J Invest Dermatol* (1987) 88 : 52-54
- Khandke L, Krane JF, Ashinoff R, et al. Cyclosporine in psoriasis treatment. Inhibition of keratinocyte cell-cycle progression in G1 independent of effects on transforming growth factor alpha/epidermal growth factor receptor pathways. *Arch Dermatol.* (1991) 127 : 1172-1179
- Ke H, Huai Q. Crystal structures of cyclophilin and its partners. *Front Biosc* (2004) 9 2285-96
- Kiani A, Rao A, Aramburu J. Manipulating immunoresponses with immunosuppressive agents that target NFAT. *Immunity* (2000) 12 : 359–372
- Kim R. Recent advances in understanding the cell death pathways activated by anticancer therapy. *Cancer.* (2005) 103 : 1551-60
- Kligman, AM. The human hair cycle. *J Invest Dermatol.* (1959) 33 : 307-316
- Kligman, AM. Pathologic dynamics of human hair loss : Telogen effluvium. *Arch Dermatol, Syph.* (1961) 83 : 175-198
- Koppel RA, Boh EE. Cutaneous reactions to chemotherapeutic agents. *Am J Med Sci.* (2003) 321 : 327-35
- Korsmeyer SJ. Programmed cell death and the regulation of homeostasis. *Harvey Lect.* (1999a) 95 : 21-41.
- Korsmeyer SJ. Bcl-2 gene family and the regulation of programmed cell death. *Cancer Res.* (7 Suppl) : (1999b) 1693s-1700s.
- Kostanecki W, Kwiatkowska E, Zborzil J. Die Haarmelanogenese bei der Endoxan Alopezie und deren Beeinflussung durch Corticosteroide. *Archiv klin exp Dermatol* (1966) 226 : 13-20
- Kostanecki W, Kwiatkowska E, Zak R. Der Einfluß des Endoxans auf die Haarmelanogenese und das Haarwachstum bei Mäusen. *Arch klin exp Dermatol* (1967)230 : 396-401
- Krajewski S, Krajewska M, Shabaik A, et al. Immunohistochemical analysis of in vivo patterns of Bcl-X expression. *Cancer Res.* (1994a) 54 5501-7
- Krajewski S, Krajewska M, Shabaik A, et al. Immunohistochemical determination of in vivo distribution of Bax, a dominant inhibitor of Bcl-2. *Am J Pathol.* (1994b) 145 : 1323-36

- Krause K und Foitzik K. Biology of the hair follicle: the basics. *Semin Cutan Med Surg.* (2006) 25 : 2-10
- Kroemer G, Reed JC. Mitochondrial control of cell death. *Nat Med.* (2000) 6 : 513-9.
- Langbein L, Rogers MA, Winter H, et al. The catalog of human hair keratins. I. Expression of the nine type I members in the hair follicle. *J Biol Chem.* (1999) 274 : 19874-84
- Laster SM, Wood JG, Gooding LR. Tumor necrosis factor can induce both apoptotic and necrotic forms of cell lysis. *J Immunol.* (1988) 141 : 2629-34.
- Lavrik IN, Golks A, Krammer PH. Caspases : pharmacological manipulation of cell death. *J Clin Invest.* (2005) 115 : 2665-72
- Lawson MA, Maxfield FR. Ca²⁺ - and calcineurin-dependent recycling of an integrin to the front of migrating neutrophils. *Nature* (1995) 337 : 5-79
- Li H, Kolluri SK, Gu J, et al. Cytochrome C release and apoptosis induced by mitochondrial targeting of nuclear orphan receptor TR3. *Science.* (2000) 18 ; 289 : 1159-64
- Lin KK, Chudova D, Hatfield GW, et al. Identification of hair cycle-associated genes from time-course gene expression profile data by using replicate variance. *Proc Natl Acad Sci U S A.* (2004)101 : 15955-60
- Lindner, G, Botchkarev VA, Botchkareva NV, et al. Analysis of apoptosis during hair follicle regression (catagen) *Am J Pathol* (1997) 151 : 1601-1617
- Lockshin RA, Zakeri Z. Programmed cell death and apoptosis: origins of the theory. *Nat Rev Mol Cell Biol.* (2001) 2 : 545-50
- Lockshin RA, Zakeri Z. Apoptosis, autophagy, and more. *Int J Biochem Cell Biol* (2004) 36 : 2405-19
- Lutz, G. Effects of cyclosporin on hair. *Skin Pharamco* (1994) 7 : 101-4
- Magerl M, Tobin D, Müller-Rover S, et al. Patterns of proliferation and apoptosis during murine hair follicle morphogenesis. *J Invest Dermatol.* (2001) 116 : 947-55
- Majno G, Joris I. Apoptosis, oncosis and necrosis. An overview of cell death. *Am J Pathol* (1995) 146 : 3-15
- Marone G, Genovese A, Granata F, et al. Pharmacological modulation of human mast cells and basophils. *Clin Exp Allergy.* (2002) 32 : 1682-9
- Massey CS. A multicente study to determine the efficacy and patient acceptability of the Paxman Scalp Cooler to prevent hair loss in patients receiving chemotherapy. *Eur J Oncol Nurs* (2004) 8 : 121-130

- Matias JR, Malloy V, Orentreich N. Animal models of androgen-dependent disorders of the pilosebaceous apparatus. 1. The androchronogenetic alopecia (AGA) mouse as a model for male-pattern baldness. *Arch Dermatol Res.* (1989) 281: 247-53
- Maurer M, Handjiski B, Paus R. Hair growth modulation by topical immunophylin ligands : Induction of anagen, inhibition of massive catagen development, and relative protection from chemotherapy-induced alopecia. *Am J Pathol* (1997) 77 : 319-332
- Mayr AC, Nagel GA, Dick HJ, Obrecht JP. Side-effects of cytostatic agents. *Schweiz Med Wochenschr.* (1973) 103 : 1252-5.
- McDuff C, Mackenzie T, Hutcheon A, et al. The effectiveness of scalp cooling in preventing alopecia for patients receiving epirubicin and docetaxel. *Eur J Cancer Care (Engl).* (2003) 12 : 154-61.
- McGregor JM, Berkhout RJ, Rozycka M, et al. p53 mutations implicate sunlight in posttransplant skin cancer irrespective of human papillomavirus status. *Oncogene* (1997) 15 : 1737-40
- Milner Y, Sudnik J, Filippi M, et al. Exogen, shedding phase of the hair growth cycle: characterization of a mouse model. *J Invest Dermatol* (2002) 119 : 639-644
- Mori O, Matsuo K, Hashimoto T. Anticancer drugs induce apoptosis in mouse hair follicles. *Kurume Med J* (2000) 47 : 193-7.
- Muller-Rover S and Paus R. (Topobiology of the hair follicle: adhesion molecules as morphoregulatory signals during hair follicle morphogenesis. In Cheng-Ming Chuong (eds.) *Molecular basis of epithelial appendage morphogenesis.* RG Landes Company (1998) 345-365
- Muller-Rover S, Rossiter H, Paus R, et al. Overexpression of Bcl-2 protects from ultraviolet B-induced apoptosis but promotes hair follicle regression and chemotherapy induced alopecia. *Am J Pathol .* (2000) 156 : 1395-405.
- Müller-Rover S, Handjiski B, Van der Veen C, et al. A comprehensive guide for the accurate classification of murine hair follicles in distinct hair cycle stages. *J Invest Dermatol .* (2001) 117 : 3-15
- Nickoloff BJ, Fisher GJ, Mitra RS, Voorhees JJ. Additive and synergistic antiproliferative effects on cyclosporin A and gamma Interferon on cultured human keratinocytes. *Am J Pathol* (1988) 131 : 12-18
- Ogo M, Suzuki J, Soma T, Takahashi T, et al. Identification of cyclosporin A responsive genes in human keratinocytes. *Exp Dermatol.* (1999) 8 : 357-8.

- Ohnemus U, Unalan M, Handjiski B, Paus R. Topical estrogen accelerates hair regrowth in mice after chemotherapy-induced alopecia by favoring the dystrophic catagen response pathway to damage. *J Invest Dermatol.* (2004) 122 : 7-13.
- Ohta S, Yanagihara K, Nagata K. Mechanism of apoptotic cell death of human gastric carcinoma cells mediated by transforming growth factor beta. *Biochem J.* (1997) 15 ; 324 (P3) : 777-82
- Oltvai ZN and Korsmayer SJ. Checkpoints of dueling dimers foil death wishes. *Cell* (1994) 79 : 189-192
- Orfanos CG, und Gerstein E. Haarausfall nach Zytostatika. *Ärztliche Kosmetologie.* (1976) 6 : 96-100
- Panhans-Gross A, Novak N, Kraft S, Bieber T. Human epidermal Langerhans cells are targets for the immunosuppressive macrolide tacrolimus (FK506). *J Allergy Clin Immunol.* (2001) 107 : 345-52
- Pandaya AG, Guevara IL. Disorders of hyperpigmentation. *Dermatol Clin* (2000) 18 : 91-98
- Panteleyev AA, van der Veen, Rosenbach T, et al. Towards defining the pathogenesis of the hairless phenotype. *J Invest Dermatol* (1998) 110 : 903-907
- Paus R, Stenn KS, Link RE. The induction of anagen hair growth in telogen mouse skin by Cs A administration. *Lab Invest* (1989) 60 : 65-369
- Paus R, Rosenbach T, Haas N, Czarnetzki BM. Patterns of cell death: the significance of apoptosis for dermatology. *Exp Dermatol* (1993): 2 : 3-11
- Paus R, Handjiski B, Eichmüller S and Czarnetzki BM. Chemotherapy-induced Alopecia in Mice: induction by Cyclophosphamide, Inhibition by Cyclosporin A, and Modulation by Dexamethasone. *Am J Pathol* (1994a) 144 : 719-734
- Paus R, Handjiski B, Czarnetzki BM, Eichmuller S. Biology of the hair follicle. *Hautarzt.* (1994b) 45 : 808-25
- Paus R, Handjiski B, Czarnetzki BM, Eichmuller S. A murine model for inducing and manipulating hair follicle regression (catagen): effects of dexamethasone and cyclosporin A. *J Invest Dermatol.* (1994c) 103 :143-7.
- Paus R, Menrad A, Czarnetzki BM. Necrobiology of the skin : apoptosis. *Hautarzt.* (1995) 46 285-303
- Paus R, Schilli MB, Handjiski B, et al. Topical calcitriol enhances normal hair regrowth but does not prevent chemotherapy-induced alopecia in mice. *Cancer Res.* (1996a) 56 : 4438-4443

- Paus R, Böttge JA, Henz BM, Maurer M. Hair growth control by immunosuppression. *Arch Dermatol Res.* (1996b) 288 : 408-10
- Paus R. Control of the hair cycle and hair diseases as cycling disorders. *Curr Opin Dermatol* (1996c) 3 : 248-258
- Paus R and Foitzik K. In search of the hair cycle clock. *Differentiation* (2004) 72 : 489-511
- Parker P, Rocque WJ, Shewchuk L, et al. Prevention of chemotherapy-induced alopecia in rats by CDK inhibitors. *Science.* (2001) 291 :134-137
- Pereira GM, Miller JF, Shevach EM. Mechanism of action of cyclosporine A in vivo. (II) T cell priming in vivo to alloantigen can be mediated by an IL-2-independent cyclosporine A-resistant pathway. *J Immunol* (1990) 144 : 2109-16.
- Peters EM, Foitzik K, Paus R, et al. A new strategy for modulating chemotherapy-induced alopecia, using PTH/PTHrP receptor agonist and antagonist. *J Invest Dermatol* (2001) 117 : 173-8
- Peters EM, Hansen MG, Overall RW, et al. Control of human hair growth by neurotrophins: brain-derived neurotrophic factor inhibits hair shaft elongation, induces catagen, and stimulates follicular transforming growth factor beta 2 expression. *J Invest Dermatol.* (2006) 124 : 675-685
- Petros AM, Olejniczak ET, Fesik W. Structural biology of the Bcl-2 family of proteins. *Biochim Biophys Acta.* (2004) 1644 : 83-94.
- Plonka PM, Handjiski B, Popik M, et al. Zinc as an ambivalent but potent modulator of murine hair growth in vivo-preliminary observations. *Exp Dermatol* (2005) 14 : 844-53
- Powis G, Kooistra KL. Doxorubicin-induced hair loss in the Angora rabbit a study of treatment to protect against the hair loss. *Cancer Chemother Pharmacol.* (1987) 20 : 291-296
- Pratt WB, Toft DO. Steroid receptor interactions with heat shock protein and immunophilin haperones. *Endocr Rev* (1997) 18 : 306-60.
- Randall VA, Sundberg JP, Philpott MP. Animal and in vitro models for the study of hair follicles. *J Investing Dermatol Symp Proc.* (2003) 8 : 39-45.
- Ramirez-Bosca A, Kanitakis J, Thivolet J. Cyclosporin A exerts a cytostatic effect in vivo on human and murine epithelial cells. *Cancer* (1990) 66 : 936-940
- Raj D, Brash DE, Grossman D. Keratinocyte apoptosis in epidermal development and disease. *J Invest Dermatol* (2006) 126 : 243-57.
- Rao A. NF-ATp : a transcription factor required for the co-ordinate induction of cytokine genes. *Immunol Today* (1994) 15 : 274-281

- Rao L, and White E.: Bcl-2 and ICE family of apoptotic regulators: making a connection. *Cur Opin Gen and Development* (1997) 7 : 52-58
- Rebora A, Guarrera M. Kenogen. *Dermatology*, (2002) 205: 108-110
- Rebora A. Pathogenesis of androgenetic alopecia. *J Am Acad Dermatol* (2004) 50 : 777-9
- Rebora A, Guarrera M. Teloptosis and Kenogen : Two new Concepts in Human Trichology. *Arch Dermatol* (2004) 140 : 619-620
- Ridderheim M, Bjurberg M, Gustavsson A. Scalp hypothermia to prevent chemotherapy-induced alopecia is effective and safe : a pilot study of a new digitized scalp cooling system used in 74 patients. *Support Care Cancer*. (2003) 11 : 371-377
- Romais B. *Mikroskopische Technik*. P. Böck.(eds.) (17 Aufl) Urban und Schwarzenberg, München-Wien-Baltimor (1989) 474-475
- Roth SI. Hair and nail. In : *Ultrastructure of normal and Abnormal Skin*. Zellikson BD (eds) Philadelphia : Lea & Febiger, . (1967) 105-107
- Sachs L.: *Angewandte Statistik*. (7th ed.) Springer Verlag Berlin (1992)
- Saitoh M, Uzuka M, Sakamoto M. : Human hair cycle. *J Invest Dermatol*. (1970) 54: 65-81
- Santini MP, Talora C, Seki T, Bolgan L, Dotto GP. Cross talk among calcineurin, Sp1/Sp3, and NFAT in control of p21(WAF1/CIP1) expression in keratinocyte differentiation. (2001) *Proc Natl Acad Sci U S A*. 98 : 9575-80.
- Sato N, Leopold PL, Crystal RG. Effect of adenovirus-mediated expression of sonic hedgehog gene on hair regrowth in mice with chemotherapy-induced alopecia. *J Nat Cancer Inst* (2001) 93 : 1858-1864
- Sawada M, Terada N, Taniguchi H, et al. Cyclosporin stimulate the hair growth in nude mice. *Lab Invest* (1987) 56 684-686
- Schneider P and Tschopp J. Apoptosis induced by death receptors. *Pharm Acta Helv* (2000) 74 : 281-286
- Schilli MB, Paus R, Menrad A. Reduction of intrafollicular apoptosis in chemotherapy-induced alopecia by topical calcitriol analogs. *J Invest Dermatol*. (1998) 111 : 598-604
- Schilli MB (1996) *Persönliche Mitteilung*
- Schreiber SL, Crabtree GR. The mechanismus of action of cyclosporin A and FK 506 *Immunol Today* (1992) 13: 136-142
- Schultz DR, Harrington WJ Jr. Apoptosis: programmed cell death at a molecular level. *Semin Arthritis Rheum*. (2003) 32 : 345-69.

- Sensenbrenner LL, Marini JJ, Colvin M. Comparative effects of cyclophosphamide and phosphoramide mustard on murine hematopoietic and immunocompetent cells. *J Natl Cancer Inst* (1979) 62 : 975-981
- Seilberg M, Marthinuss J, Stenn KF.: Changes in the expression of apoptosis associated genes in skin mark early catagen. *J Invest Dermatol.* (1995) 104 : 78-82
- Selleri S, Arnaboldi F, Vizzotto B, Balsari A, Rumio C. Epitheliummesenchyme compartment interaction and oncosis on chemotherapy-induced hair damage. *Lab Invest.* (2004) 84 : 1404-17
- Sharov AA, Li GZ, Palkina TN, et al. Fas and c-kit are involved in the control of hair ollicle melanocyte apoptosis and migration in chemotherapy-induced hair loss. *J Invest Dermatol.* (2003) 120 : 27-35
- Sharov AA, Siebenhaar F, Sharova TY, et al. Fas signaling is involved in the control of hair follicle response to chemotherapy. *Cancer Res.* (2004) 64 : 6266-70
- Shevach EM. The effects of cyclosporine A on the immune system. *Ann Rev Immunol* (1985) 3 : 397-423
- Shi Y, Sahai BM, Green DR. Cyclosporine A inhibits activation induced cell death in T-cell hybridomas and thymocytes. *Nature* (1989) 339 : 625-626
- Shibasaki F, Price ER, Milan D, McKeon F. Role of kinases and the phosphatase calcineurin in the nuclear shuttling of transcription factor NF-AT4. *Nature.* (1996) 382 :370-3
- Shirai A, Tsunoda H, Tamaoki T, Kamiya T. Topical application of cyclosporin A induces rapid remodeling of damaged anagen hair follicles produced in cyclophosphamide administered mice. *J Dermatol Sci* (2001) 27 : 7-13.
- Silver F, Chase HB. An in vivo method for studying the hair cycle. *Nature.* (1966) 210 : 1051-1060
- Sladek NE. Therapeutic efficacy of cyclophosphamide as a function of inhibition of its metabolites. *Cancer Res* (1973):32: 848-845
- Slominski A, Paus R, Plonka P, et al. Melanogenesis during the anagen-catagen-telogen transformation of the murine hair cycle *J Invest Dermatol* (1994) 102 : 862-9.
- Slominski A, Paus R, Plonka P, et al. Pharmacological disruption of hair follicle pigmentation by cyclophosphamide as a model for studying the melanocyte response to and recovery from cytostatic drug damage *in situ.* *J Invest Dermatol* (1996) 106 : 1203-1211
- Slominski A, Wortsman J, Plonka PM, et al. Hair follicle pigmentation. *J Invest Dermatol.* (2004) 124 :13-21.

- Smith ML, Fornace AJ Jr. Mammalian DNA damage-inducible genes associated with growth arrest and apoptosis. *Mutat Res.* (1996b) 340 : 109-24.
- Soma T, Ogo M, Suzuki J, et al. Analysis of apoptotic cell death in human hair follicles in vivo and in vitro. *J Invest Dermatol* (1998) 111 : 948-954
- Song S, and Lambert PF. Different responses of epidermal and hair follicle cells to radiation correlate with distinct patterns of p53 and p21 induction. *Am J Pathol* (1999) 155 : 1121-1127
- Sperling LC. Hair and systemic disease. *Dermatol Clin.* (2001) 19 : 711-26,
- Staricco RG. The melanocytes in the hair follicle. *J Invest Dermatol* (1960) 35 : 185-194
- Stamnes MA, Zuker CS. Peptidyl-prolyl cis-trans isomerases, cyclophilin, FK506-binding protein, and cyclosporin A : four of a kind. *Curr Opin Cell Biol.* (1990) 2 : 1104-1107
- Stellato C, de Paulis A, Ciccarelli A, et al. Anti-inflammatory effect of cyclosporin A on human skin mast cells. *J Invest Dermatol* (1992) 98 : 800-804
- Stoll E. Evaluation of Cyclophosphamide dosage schedules in breast cancer. *Br J Cancer,* (1974) 24 : 475-450
- Straile WE, Chase HB, Arsenault C. Growth and differentiation of hair follicles between periods of activity and quiescence. *J Exp Zool* (1961) 148 : 205-215
- Straile AW. Possible functions of the external root sheath during growth of the hair follicle. *J Exp Zool* (1962) 150 : 207-23
- Struck RF, Kird MC, Mellet LB, et al. Urinary metabolites of antitumor agent cyclophosphamide. *Mol Pharmacol* (1971) 7 : 519-529
- Sundberg JP, Price VH, King LE Jr. The „hairless“ gene in mouse and man. *Arch Dermatol.* (1999) 135 : 718-720
- Sundberg JP, King LE Jr., Bascom C). Animal models for male pattern (androgenetic) alopecia. *Eur J Dermatol .* (2001) 11: 321-325
- Suzuki S, Ota Y, Ozawa K, Imamura T. Dual mode of hair growth cycle by two GFG-5m gene products. *J Invest Dermatol* (2000) 114 : 456-463
- Szamel M, Bartels F, Resch K. Cyclosporin A inhibits T cell receptor-induced interleukin-2 synthesis of human T lymphocytes by selectively preventing a trans membrane signal transduction pathway leading to sustained activation of a protein kinase C isoenzyme, protein kinase C-beta. *Eur J Immunol.* (1993) 23 : 3072-81
- Taylor M, Ashcroft ATT, Messenger AG. Cyclosporin A prolongs human hair growth in vitro. *J Invest Dermatol* (1993) 100 : 237-239

- Takahashi T, Kimimura A, Shiari A, Yokoo Y. Several selective protein kinase C inhibitors including procyanidins promote hair growth. *Skin Pharmacol Appl Skin Physiol* (2000) 13: 133-142
- Takahashi T, Kimimura A. Cyclosporin a promotes hair epithelial cell proliferation and modulates Protein Kinase C expression and translocation in hair epithelial cells. *J Invest Dermatol.* (2001) 117 : 605-11
- Takahashi K, Reynolds M, Ogawa N, et al. Augmentation of T-cell apoptosis by immunosuppressive agents. *Clin Transplant. Suppl* (2004) 12:72-5.
- Tanaka K, Asanuma M, Ogawa N. Molecular basis of anti-apoptotic effect of immunophilin ligands on hydrogen peroxide-induced apoptosis in human glioma cells. *Neurochem Res.* (2004) 29 : 1529-36.
- Teunissen MBM, De Jager MH, Kapsenberg ML, Bos JD. Inhibitory effect of cyclosporin A on antigen and alloantigen presenting capacity of human epidermal Langerhans cells. *Br J Dermatol* (1991) 25 309-316
- Tierney A, Taylor J, Closs SJ, et al. Hair loss due to a cytostatic chemotherapy : a prospective descriptive study. (1990) *Br J Cancer.* 62 527-528
- Tobin DJ, Slominski A, Botchkarev VA, and Paus R. The fate of hair follicle melanocytes during the hair growth cycle. *J Invest Dermatol Symp Proc* (1999) 4 : 323-332
- Tobin DJ, Gunin A, Magerl M, Handjiski B, Paus R. Plasticity and cytokinetic dynamics of the hair follicle mesenchyme: implications for hair growth control. *J Invest Dermatol.*(2003) 120 : 895-904
- Tollenaar RA, Liefers GJ, Repelaer van Driel OJ, van de Velde CJ. Scalp cooling has no place in the prevention of alopecia in adjuvant chemotherapy for breast cancer. *Eur J Cancer.* (1994) 30A : 1448-53.
- Tossi A, Misciali C, Piraccini BM, et al. Drug-induced hair loss and hair growth. Incidence, management and avoidance. *Drug Saf.* (1994) 10 : 310-
- Trenn G, Taffs R, Hohman R, et al. Biochemical characterization of the inhibitory effect of CsA on cytolytic T lymphocyte effector functions. *J Immunol.* (1989) 142 : 3796-802.
- Trueb RM. Molecular mechanism of androgenetic alopecia. *Experimental Gerontology* (2002) 37 981-990
- Uno, H. Quantitative models for the study of hair growth in vivo. *Ann NY Acad Sci* (1991) 642 : 107-112
- Uno H, Imamura K, Pan H. Androgenetic alopecia in the stump tailed macaque. An important model for investigating the pathology and antiandrogenetic therapy of male pattern

- baldnes. In : Van Neste D, Randal VA, Price V (eds). Hair and its disorders : biology, pathology and management. London : Martin Dunitz (2000) 137-151
- Wagner WL, Bye MG. Body image and patients experiencing alopecia as a result of cancer chemotherapy. *Cancer Nurs* (1979) 2 : 365-369
- Wang P, Heitman J. The cyclophilins. *Genome Biol.* (2005) 6 : 226-256.
- Whitehouse MW, Beck FW, Droge MM, Struck RF. Lymphocyte deactivation by (potential immunosuppressant) alkylating metabolites of cyclophosphamide. *Agents Actions.* (1974) 4 : 117-24
- Whiting DA, Waldstreicher J, Sanchez M, Kaufman KD. Measuring reversal of hair miniaturization in androgenetic alopecia by follicular count in horizontal sections of serial scalp biopsies: results of finasteride 1 mg treatment of men and postmenopausal women. *J Invest Dermatol Symp Proc* (1999) 4 : 282-284
- Whiting DA. Histopathologic Features of Alopecia Areata. *Arch Dermatol*, (2003) 139 1555-1559
- Weedon D, Strutton G. Apoptosis as a mechanism of the involution of hair follicles in catagen transformation. *Acta Derm Venerol* (1990) 61 : 1329-1337
- Wiederrecht G, Lam E, Hung S, et al. The mechanism of action of FK506 and cyclosporin A. *Ann NY Acad Sci.* (1993) 696 : 9-19.
- Wiederrecht GJ, Sabers CJ, Brunn GJ, et al. Mechanism of action of rapamycin : new insights into the regulation of G1 phase progression in eukaryotic cells. *Prog Cell Cycle Res.* (1995) 1 : 53-71
- Wheeler AG, Dansby D, Hawkins HC, et al. A toxicological and haematological evaluations of cyclophosphamide (Cytosan) in experimental animals. *Toxicol. Appl Pharmacol* (1962) 4 : 324-343
- Wysocki GP and Daley TD. Hypertrichosis in patients receiving cyclosporin therapy. *Clin Exp Dermatol* (1987) 12: 191-196
- Wolbach, SB. The hair cycle of the mouse and its importance in the study of sequence of the experimental carcinogenesis. *Ann N.Y. Acad. Sci* (1951) 53 : 517-536,
- Van Gurp M, Festjens N, van Loo G, Saelens X, Vandenabeele P. Mitochondrial intermembrane proteins in cell death. *Biochem Biophys Res Commun.* (2003) 30 : 487-97
- Vogt A, Blume-Peytavi U. Biology of the human hair follicle. New knowledge and the clinical significance. *Hautarzt.* (2003) 54 : 692-8
- Yamamoto T and Nishioka K. Increased expression of p53 and p21 (Waf1/Cip1) in the lesional skin of bleomycin-induced scleroderma. *Arch Dermatol Res.* (2005) 296 : 509-13

- Yarosh DB, Pena AV, Nay SL, et al. Calcineurin Inhibitors Decrease DNA Repair and Apoptosis in Human Keratinocytes Following Ultraviolet B Irradiation. *J Invest Dermatol* (2005) 125 : 1020-1025
- Yu J, Zhang L. Apoptosis in human cancer cells. *Curr Opin Oncol*. (2004) 16 : 19-24.
- Yu J, Zhang L. The transcriptional targets of p53 in apoptosis control. *Biochem Biophys Res Commun*. (2005) 331 : 851-8.
- Zakeri Z, Lockskin RA. Cell death during development. *J Immunol Methods*. (2002) 1: 265 (1-2) : 3-20.
- Zaun H. Pathologische Reaktionen am Haarfollikel. *Annales Universitatis Saraviensis*. (1967) 14 148-223