

## 7 Literaturverzeichnis

1. **Aamodt, A.; Nordsletten, L.; Havelin, L. I. et al.:** Documentation of hip prostheses used in Norway: a critical review of the literature from 1996--2000. *Acta Orthop Scand*, 75(6): 663-76, 2004.
2. **Acklin, Y. P.; Berli, B. J.; Frick, W. et al.:** Nine-year results of Muller cemented titanium Straight Stems in total hip replacement. *Arch Orthop Trauma Surg*, 121(7): 391-8, 2001.
3. **Albrektsson, T.; Branemark, P. I.; Hansson, H. A. et al.:** Osseointegrated titanium implants. Requirements for ensuring a long-lasting, direct bone-to-implant anchorage in man. *Acta Orthop Scand*, 52(2): 155-70, 1981.
4. **Aldinger, G., and Weipert, A.:** [3D-based production of hip joints: the Aldinger system]. *Radiologe*, 31(10): 474-80, 1991.
5. **Aldinger, P. R.; Breusch, S. J.; Lukoschek, M. et al.:** A ten- to 15-year follow-up of the cementless spotorno stem. *J Bone Joint Surg Br*, 85(2): 209-14, 2003.
6. **Aldinger, P. R.; Thomsen, M.; Mau, H. et al.:** Cementless Spotorno tapered titanium stems: excellent 10-15-year survival in 141 young patients. *Acta Orthop Scand*, 74(3): 253-8, 2003.
7. **Arabmotlagh, M.; Hennigs, T.; and Rittmeister, M.:** [Femoral periprosthetic bone remodelling to the proximal femur after implantation of custom made anatomic and standard straight stem hip prostheses]. *Z Orthop Ihre Grenzgeb*, 141(5): 519-25, 2003.
8. **August, A. C.; Aldam, C. H.; and Pynsent, P. B.:** The McKee-Farrar hip arthroplasty. A long-term study. *J Bone Joint Surg Br*, 68(4): 520-7, 1986.
9. **Bailey, A. J., and Mansell, J. P.:** Do subchondral bone changes exacerbate or precede articular cartilage destruction in osteoarthritis of the elderly? *Gerontology*, 43(5): 296-304, 1997.
10. **Ballard, W. T.; Callaghan, J. J.; Sullivan, P. M. et al.:** The results of improved cementing techniques for total hip arthroplasty in patients less than fifty years old. A ten-year follow-up study. *J Bone Joint Surg Am*, 76(7): 959-64, 1994.
11. **Barrack, R. L.; Jasty, M.; Bragdon, C. et al.:** Thigh pain despite bone ingrowth into uncemented femoral stems. *J Bone Joint Surg Br*, 74(4): 507-10, 1992.
12. **Barrack, R. L.; Mulroy, R. D., Jr.; and Harris, W. H.:** Improved cementing techniques and femoral component loosening in young patients with hip arthroplasty. A 12-year radiographic review. *J Bone Joint Surg Br*, 74(3): 385-9, 1992.
13. **Bauer, R., Kerschbaumer F.:** Transglutealer Zugang nach Bauer, in: operative Zugangswege in Orthopädie und Traumatologie, Thieme Verlag. 1986.
14. **Bauer, R., and Russe, W.:** [The transgluteal approach in hip joint arthroplasty]. *Z Orthop Ihre Grenzgeb*, 122(1): 48-9, 1984.
15. **Bellamy, N.:** WOMAC: a 20-year experiential review of a patient-centered self-reported health status questionnaire. *J Rheumatol*, 29(12): 2473-6, 2002.
16. **Bellamy, N.; Buchanan, W. W.; Goldsmith, C. H. et al.:** Validation study of WOMAC: a health status instrument for measuring clinically important patient relevant outcomes to antirheumatic drug therapy in patients with osteoarthritis of the hip or knee. *J Rheumatol*, 15(12): 1833-40, 1988.

17. **Bellamy, N.; Campbell, J.; Hill, J. et al.:** A comparative study of telephone versus onsite completion of the WOMAC 3.0 osteoarthritis index. *J Rheumatol*, 29(4): 783-6, 2002.
18. **Bensmann, G.:** Welcher Werkstoff ist für welche Endoprothese geeignet? In: *Jerosch J, Effenberger H, Fuchs S (Hrsg.): Hüftendoprothetik. Georg Thieme Verlag, Stuttgart New York*, 1996.
19. **Blanchard, C.; Feeny, D.; Mahon, J. L. et al.:** Is the Health Utilities Index responsive in total hip arthroplasty patients? *J Clin Epidemiol*, 56(11): 1046-54, 2003.
20. **Blasius, K.; Cotta, H.; Schneider, U. et al.:** [CLS multicenter study--8-year results]. *Z Orthop Ihre Grenzgeb*, 131(6): 547-52, 1993.
21. **Boardman, D. L.; Dorey, F.; Thomas, B. J. et al.:** The accuracy of assessing total hip arthroplasty outcomes: a prospective correlation study of walking ability and 2 validated measurement devices. *J Arthroplasty*, 15(2): 200-4, 2000.
22. **Bobyn, J. D.; Pilliar, R. M.; Cameron, H. U. et al.:** The optimum pore size for the fixation of porous-surfaced metal implants by the ingrowth of bone. *Clin Orthop Relat Res*, (150): 263-70, 1980.
23. **Bos, I.:** [Tissue reactions around loosened hip joint endoprostheses. A histological study of secondary capsules and interface membranes]. *Orthopade*, 30(11): 881-9, 2001.
24. **Bourne, R. B., and Rorabeck, C. H.:** Assessing the outcomes: what really works? *Orthopedics*, 22(9): 823-5, 1999.
25. **Bragdon, C. R.; Jasty, M.; Muratoglu, O. K. et al.:** Third-body wear testing of a highly cross-linked acetabular liner: the effect of large femoral head size in the presence of particulate poly(methyl-methacrylate) debris. *J Arthroplasty*, 20(3): 379-85, 2005.
26. **Breusch, S. J.; Aldinger, P. R.; Thomsen, M. et al.:** [Anchoring principles in hip endoprostheses. I: Prosthesis stem]. *Unfallchirurg*, 103(11): 918-31, 2000.
27. **Breusch, S. J.; Lukoschek, M.; Kreutzer, J. et al.:** Dependency of cement mantle thickness on femoral stem design and centralizer. *J Arthroplasty*, 16(5): 648-57, 2001.
28. **Breusch, S. J.; Schneider, U.; Kreutzer, J. et al.:** [Effects of the cementing technique on cementing results concerning the coxal end of the femur]. *Orthopade*, 29(3): 260-70, 2000.
29. **Britton, A. R.; Murray, D. W.; Bulstrode, C. J. et al.:** Pain levels after total hip replacement: their use as endpoints for survival analysis. *J Bone Joint Surg Br*, 79(1): 93-8, 1997.
30. **Bulow, J. U.; Scheller, G.; Arnold, P. et al.:** Follow-up (6-9 years) results of the uncemented CLS Spotorno stem. *Arch Orthop Trauma Surg*, 115(3-4): 190-4, 1996.
31. **Bulow, J. U.; Scheller, G.; Arnold, P. et al.:** Uncemented total hip replacement and thigh pain. *Int Orthop*, 20(2): 65-9, 1996.
32. **Callaghan, J. J.; Templeton, J. E.; Liu, S. S. et al.:** Results of Charnley total hip arthroplasty at a minimum of thirty years. A concise follow-up of a previous report. *J Bone Joint Surg Am*, 86-A(4): 690-5, 2004.
33. **Chandler, H. P.; Reineck, F. T.; Wixson, R. L. et al.:** Total hip replacement in patients younger than thirty years old. A five-year follow-up study. *J Bone Joint Surg Am*, 63(9): 1426-34, 1981.
34. **Charnley, J.:** D Total hip replacement. *Jama*, 230(7): 1025-8, 1974.

35. **Charnley, J.:** The long-term results of low-friction arthroplasty of the hip performed as a primary intervention. 1972. *Clin Orthop*, (319): 4-15, 1995.
36. **Charnley, J.:** Low friction arthroplasty of the hip. Springer, Berlin Heidelberg New York. 1979.
37. **Christel, P., and Djian, P.:** Recent advances in adult hip joint surgery. *Curr Opin Rheumatol*, 6(2): 161-71, 1994.
38. **Decking, J.; Rosendahl, T.; and Decking, D.:** [Proximal fixation of hip endoprostheses with porous surface: results of 6 years]. *Z Orthop Ihre Grenzgeb*, 137(2): 108-13, 1999.
39. **Delaunay, C., and Kapandji, A. I.:** Survival analysis of cementless grit-blasted titanium total hip arthroplasties. *J Bone Joint Surg Br*, 83(3): 408-13, 2001.
40. **Dieppe, P.:** Management of hip osteoarthritis. *Bmj*, 311(7009): 853-7, 1995.
41. **Dieppe, P. A., and Lohmander, L. S.:** Pathogenesis and management of pain in osteoarthritis. *Lancet*, 365(9463): 965-73, 2005.
42. **DiFazio, F.; Shon, W. Y.; Salvati, E. A. et al.:** Long-term results of total hip arthroplasty with a cemented custom-designed swan-neck femoral component for congenital dislocation or severe dysplasia: a follow-up note. *J Bone Joint Surg Am*, 84-A(2): 204-7, 2002.
43. **Dorr, L. D.; Luckett, M.; and Conaty, J. P.:** Total hip arthroplasties in patients younger than 45 years. A nine- to ten-year follow-up study. *Clin Orthop Relat Res*, (260): 215-9, 1990.
44. **Ebramzadeh, E.; Sarmiento, A.; McKellop, H. A. et al.:** The cement mantle in total hip arthroplasty. Analysis of long-term radiographic results. *J Bone Joint Surg Am*, 76(1): 77-87, 1994.
45. **Eingartner, C.; Ihm, A.; Maurer, F. et al.:** [Good long term results with a cemented straight femoral shaft prosthesis made of titanium]. *Unfallchirurg*, 105(9): 804-10, 2002.
46. **Ekelund, A.; Rydell, N.; and Nilsson, O. S.:** Total hip arthroplasty in patients 80 years of age and older. *Clin Orthop Relat Res*, (281): 101-6, 1992.
47. **Elke, R.:** [Particle disease. Is tribology a topic in revision surgery?]. *Orthopade*, 30(5): 258-65, 2001.
48. **Engh, C. A.; Bobyn, J. D.; and Glassman, A. H.:** Porous-coated hip replacement. The factors governing bone ingrowth, stress shielding, and clinical results. *J Bone Joint Surg Br*, 69(1): 45-55, 1987.
49. **Engh, C. A.; Zettl-Schaffer, K. F.; Kukita, Y. et al.:** Histological and radiographic assessment of well functioning porous-coated acetabular components. A human postmortem retrieval study. *J Bone Joint Surg Am*, 75(6): 814-24, 1993.
50. **Espehaug, B.; Furnes, O.; Havelin, L. I. et al.:** The type of cement and failure of total hip replacements. *J Bone Joint Surg Br*, 84(6): 832-8, 2002.
51. **Estok, D. M., 2nd, and Harris, W. H.:** Long-term results of cemented femoral revision surgery using second-generation techniques. An average 11.7-year follow-up evaluation. *Clin Orthop Relat Res*, (299): 190-202, 1994.
52. **Frankel, S.; Eachus, J.; Pearson, N. et al.:** Population requirement for primary hip-replacement surgery: a cross-sectional study. *Lancet*, 353(9161): 1304-9, 1999.
53. **Friedebold, G.:** Die Entwicklung des künstlichen Hüftgelenkersatzes. In: Hüftgelenkendoprothetik, R. Rahmanzadeh and M. Faensen, Editors, Springer Verlag: Berlin. 1984.

54. **Friedebold, G.**: Die Entwicklung des künstlichen Hüftgelenkersatzes, in Hüftgelenksendoprothetik, R. Rahmazadeh and M. Faensen, Editors, Springer Verlag: Berlin. 1984.
55. **Garcia-Cimbrelo, E.; Cruz-Pardos, A.; Madero, R. et al.**: Total hip arthroplasty with use of the cementless Zweymüller Alloclassic system. A ten to thirteen-year follow-up study. *J Bone Joint Surg Am*, 85-A(2): 296-303, 2003.
56. **Garellick, G.; Herberts, P.; Stromberg, C. et al.**: Long-term results of Charnley arthroplasty. A 12-16-year follow-up study. *J Arthroplasty*, 9(4): 333-40, 1994.
57. **Garellick, G.; Malchau, H.; and Herberts, P.**: Specific or general health outcome measures in the evaluation of total hip replacement. A comparison between the Harris hip score and the Nottingham Health Profile. *J Bone Joint Surg Br*, 80(4): 600-6, 1998.
58. **Garellick, G.; Malchau, H.; and Herberts, P.**: Survival of hip replacements. A comparison of a randomized trial and a registry. *Clin Orthop Relat Res*, (375): 157-67, 2000.
59. **Gluck, T.**: Die Invaginationsmethode der Osteo- und Arthroplastik, in Aktuelle Probleme in Chirurgie und Orthopädie, M. Ungetüm, Editor. 1978.
60. **Gonzalez, O.; Smith, R. L.; and Goodman, S. B.**: Effect of size, concentration, surface area, and volume of polymethylmethacrylate particles on human macrophages in vitro. *J Biomed Mater Res*, 30(4): 463-73, 1996.
61. **Granchi, D.; Cenni, E.; Trisolino, G. et al.**: Sensitivity to implant materials in patients undergoing total hip replacement. *J Biomed Mater Res B Appl Biomater*, 2005.
62. **Griffiths, H. J.; Priest, D. R.; Kushner, D. M. et al.**: Total hip replacement and other orthopedic hip procedures. *Radiol Clin North Am*, 33(2): 267-87, 1995.
63. **Grubl, A.; Csepan, R.; Delaunay, C. et al.**: [Six to ten year results of use of the alloclassic hip prosthesis--a multicentre survival analysis]. *Z Orthop Ihre Grenzgeb*, 141(3): 303-8, 2003.
64. **Guther, D.; Pap, G.; Bamert, P. et al.**: [Long-term results with the cementless CLS stem in hip replacement]. *Z Orthop Ihre Grenzgeb*, 141(3): 309-15, 2003.
65. **Harris, W. H.**: Osteolysis and particle disease in hip replacement. A review. *Acta Orthop Scand*, 65(1): 113-23, 1994.
66. **Harris, W. H.**: Traumatic arthritis of the hip after dislocation and acetabular fractures: treatment by mold arthroplasty. An end-result study using a new method of result evaluation. *J Bone Joint Surg Am*, 51(4): 737-55, 1969.
67. **Häuselmann, H. J., and Hedbom, E.**: Von der Pathogenese der Arthrose zu therapeutischen Empfehlungen und Knorpelersatz. *Schweiz Med Forum*, Nr. 25: 610- 615, 2002.
68. **Havelin, L. I.; Espehaug, B.; Vollset, S. E. et al.**: The effect of the type of cement on early revision of Charnley total hip prostheses. A review of eight thousand five hundred and seventy-nine primary arthroplasties from the Norwegian Arthroplasty Register. *J Bone Joint Surg Am*, 77(10): 1543-50, 1995.
69. **Healy, W. L.**: Hip implant selection for total hip arthroplasty in elderly patients. *Clin Orthop Relat Res*, (405): 54-64, 2002.
70. **Heisel, C.; Silva, M.; dela Rosa, M. A. et al.**: Short-term in vivo wear of cross-linked polyethylene. *J Bone Joint Surg Am*, 86-A(4): 748-51, 2004.

71. **Hernigou, P.; Intrator, L.; Bahrami, T. et al.**: Interleukin-6 in the blood of patients with total hip arthroplasty without loosening. *Clin Orthop Relat Res*, (366): 147-54, 1999.
72. **Hirakawa, K.; Jacobs, J. J.; Urban, R. et al.**: Mechanisms of failure of total hip replacements: lessons learned from retrieval studies. *Clin Orthop Relat Res*, (420): 10-7, 2004.
73. **Holtzman, J.; Saleh, K.; and Kane, R.**: Effect of baseline functional status and pain on outcomes of total hip arthroplasty. *J Bone Joint Surg Am*, 84-A(11): 1942-8, 2002.
74. **Holtzman, J.; Saleh, K.; and Kane, R.**: Gender differences in functional status and pain in a Medicare population undergoing elective total hip arthroplasty. *Med Care*, 40(6): 461-70, 2002.
75. **Hughes, S. S.; Furia, J. P.; Smith, P. et al.**: Atrophy of the proximal part of the femur after total hip arthroplasty without cement. A quantitative comparison of cobalt-chromium and titanium femoral stems with use of dual x-ray absorptiometry. *J Bone Joint Surg Am*, 77(2): 231-9, 1995.
76. **Hultmark, P.; Karrholm, J.; Stromberg, C. et al.**: Cemented first-time revisions of the femoral component: prospective 7 to 13 years' follow-up using second-generation and third-generation technique. *J Arthroplasty*, 15(5): 551-61, 2000.
77. **Imhof, H.; Czerny, C.; Gahleitner, A. et al.**: [Coxarthrosis]. *Radiologe*, 42(6): 416-31, 2002.
78. **Ingvarsson, T.; Hagglund, G.; and Lohmander, L. S.**: Prevalence of hip osteoarthritis in Iceland. *Ann Rheum Dis*, 58(4): 201-7, 1999.
79. **Jasty, M.; Maloney, W. J.; Bragdon, C. R. et al.**: The initiation of failure in cemented femoral components of hip arthroplasties. *J Bone Joint Surg Br*, 73(4): 551-8, 1991.
80. **Jorring, K.**: Osteoarthritis of the hip. Epidemiology and clinical role. *Acta Orthop Scand*, 51(3): 523-30, 1980.
81. **Joshi, A. B.; Porter, M. L.; Trail, I. A. et al.**: Long-term results of Charnley low-friction arthroplasty in young patients. *J Bone Joint Surg Br*, 75(4): 616-23, 1993.
82. **Judet, R.**: [Total hip endoprosthesis made of porometal without cement anchoring]. *Z Orthop Ihre Grenzgeb*, 113(4): 828-9, 1975.
83. **Kutzer, A., and Löhr, J. F.**: Frühlockerung von Hüftgelenkendoprothesen. *Deutsches Ärzteblatt* 100, 12: A-784/ B-661/ C-621, 2003.
84. **Kawate, K.; Maloney, W. J.; Bragdon, C. R. et al.**: Importance of a thin cement mantle. Autopsy studies of eight hips. *Clin Orthop Relat Res*, (355): 70-6, 1998.
85. **Keener, J. D.; Callaghan, J. J.; Goetz, D. D. et al.**: Long-term function after Charnley total hip arthroplasty. *Clin Orthop Relat Res*, (417): 148-56, 2003.
86. **Knahr, K.; Kryspin-Exner, I.; Jagsch, R. et al.**: [Evaluating the quality of life before and after implantation of a total hip endoprosthesis]. *Z Orthop Ihre Grenzgeb*, 136(4): 321-9, 1998.
87. **Kobayashi, A.; Freeman, M. A.; Bonfield, W. et al.**: Number of polyethylene particles and osteolysis in total joint replacements. A quantitative study using a tissue-digestion method. *J Bone Joint Surg Br*, 79(5): 844-8, 1997.
88. **Konstantoulakis, C.; Anastopoulos, G.; Papaeliou, A. et al.**: Uncemented total hip arthroplasty in the elderly. *Int Orthop*, 23(6): 334-6, 1999.
89. **Krämer, K., Stock, M., Winter, M.**: Klinikleitfaden Orthopädie. 1992.

90. **Kreibich, D. N.; Moran, C. G.; Delves, H. T. et al.:** Systemic release of cobalt and chromium after uncemented total hip replacement. *J Bone Joint Surg Br*, 78(1): 18-21, 1996.
91. **Krismer, M.; Klar, M.; Klestil, T. et al.:** Aseptic loosening of straight- and curved-stem Muller femoral prostheses. *Arch Orthop Trauma Surg*, 110(4): 190-4, 1991.
92. **Lassus, J.; Waris, V.; Xu, J. W. et al.:** Increased interleukin-8 (IL-8) expression is related to aseptic loosening of total hip replacement. *Arch Orthop Trauma Surg*, 120(5-6): 328-32, 2000.
93. **Laupacis, A.; Bourne, R.; Rorabeck, C. et al.:** The effect of elective total hip replacement on health-related quality of life. *J Bone Joint Surg Am*, 75(11): 1619-26, 1993.
94. **Leopold, S. S.; Berger, R. A.; Patterson, L. et al.:** Serum titanium level for diagnosis of a failed, metal-backed patellar component. *J Arthroplasty*, 15(7): 938-43, 2000.
95. **Lieberman, J. R.; Dorey, F.; Shekelle, P. et al.:** Outcome after total hip arthroplasty. Comparison of a traditional disease-specific and a quality-of-life measurement of outcome. *J Arthroplasty*, 12(6): 639-45, 1997.
96. **Lieberman, J. R.; Hawker, G.; and Wright, J. G.:** Hip function in patients >55 years old: population reference values. *J Arthroplasty*, 16(7): 901-4, 2001.
97. **Madey, S. M.; Callaghan, J. J.; Olejniczak, J. P. et al.:** Charnley total hip arthroplasty with use of improved techniques of cementing. The results after a minimum of fifteen years of follow-up. *J Bone Joint Surg Am*, 79(1): 53-64, 1997.
98. **Malchau, H.; Herberts, P.; Eisler, T. et al.:** The Swedish Total Hip Replacement Register. *J Bone Joint Surg Am*, 84-A Suppl 2: 2-20, 2002.
99. **Maloney, W. J.; Galante, J. O.; Anderson, M. et al.:** Fixation, polyethylene wear, and pelvic osteolysis in primary total hip replacement. *Clin Orthop Relat Res*, (369): 157-64, 1999.
100. **Margheritini, F.; Moreschini, O.; and Romanini, L.:** A mould arthroplasty revision: a 43-year follow-up. *Arch Orthop Trauma Surg*, 120(1-2): 110-1, 2000.
101. **McMinn, D., and Treacy, R.:** Birmingham Hip Resurfacing; Geschichte, Entwicklung und klinische Resultate. 3, 2000.
102. **Mittelmeier, H.:** [Fixing endoprostheses without cement using the principle of "weightbearing ribbing" (author's transl)]. *Z Orthop Ihre Grenzgeb*, 112(1): 27-33, 1974.
103. **Mohanty, M.:** Cellular basis for failure of joint prosthesis. *Biomed Mater Eng*, 6(3): 165-72, 1996.
104. **Morscher, E.:** Principles of acetabular fixation in THR with special reference to the "press-fit cup". *Acta Orthop Belg*, 59 Suppl 1: 260-6, 1993.
105. **Morscher, E., and Masar, Z.:** Development and first experience with an uncemented press-fit cup. *Clin Orthop Relat Res*, (232): 96-103, 1988.
106. **Morscher, E. W.:** Der zementfreie Femurschaft. *E. W. Morscher: Endoprothetik*, 1995.
107. **Mulroy, R. D., Jr., and Harris, W. H.:** The effect of improved cementing techniques on component loosening in total hip replacement. An 11-year radiographic review. *J Bone Joint Surg Br*, 72(5): 757-60, 1990.
108. **Mulroy, W. F., and Harris, W. H.:** Acetabular and femoral fixation 15 years after cemented total hip surgery. *Clin Orthop*, (337): 118-28, 1997.

109. **Neumann, L.; Freund, K. G.; and Sorenson, K. H.**: Long-term results of Charnley total hip replacement. Review of 92 patients at 15 to 20 years. *J Bone Joint Surg Br*, 76(2): 245-51, 1994.
110. **Ni, G. X.; Lu, W. W.; Chiu, K. Y. et al.**: Cemented or uncemented femoral component in primary total hip replacement? A review from a clinical and radiological perspective. *J Orthop Surg (Hong Kong)*, 13(1): 96-105, 2005.
111. **Nilsdotter, A. K.; Petersson, I. F.; Roos, E. M. et al.**: Predictors of patient relevant outcome after total hip replacement for osteoarthritis: a prospective study. *Ann Rheum Dis*, 62(10): 923-30, 2003.
112. **Nilsdotter, A. K.; Roos, E. M.; Westerlund, J. P. et al.**: Comparative responsiveness of measures of pain and function after total hip replacement. *Arthritis Rheum*, 45(3): 258-62, 2001.
113. **Oates, K. M.; Barrera, D. L.; Tucker, W. N. et al.**: In vivo effect of pressurization of polymethyl methacrylate bone-cement. Biomechanical and histologic analysis. *J Arthroplasty*, 10(3): 373-81, 1995.
114. **Ochsner, P. E.**: Die Hüfttotalendoprothese: Implantationstechnik und lokale Komplikationen; Hrsg. Peter E. Ochsner.- Berlin; Springer Verlag. 2003.
115. **Okhuisen, S., and Dhert, W.**: Total hip arthroplasty in the Netherlands. *Ned Tojdtschr Geneesk*, 142: 1434-38, 1998.
116. **Older, J.**: Charnley low-friction arthroplasty: a worldwide retrospective review at 15 to 20 years. *J Arthroplasty*, 17(6): 675-80, 2002.
117. **Oparaugo, P. C.; Clarke, I. C.; Malchau, H. et al.**: Correlation of wear debris-induced osteolysis and revision with volumetric wear-rates of polyethylene: a survey of 8 reports in the literature. *Acta Orthop Scand*, 72(1): 22-8, 2001.
118. **Paavolainen, P.; Pukkala, E.; Pulkkinen, P. et al.**: Cancer incidence in Finnish hip replacement patients from 1980 to 1995: a nationwide cohort study involving 31,651 patients. *J Arthroplasty*, 14(3): 272-80, 1999.
119. **Pilliar, R. M.**: Powder metal-made orthopedic implants with porous surface for fixation by tissue ingrowth. *Clin Orthop Relat Res*, (176): 42-51, 1983.
120. **Pingsmann, A.; Muller, R. T.; and Goller, A.**: Cost analysis for total hip arthroplasty by measurement of time and material expenditure. *Arch Orthop Trauma Surg*, 117(8): 421-4, 1998.
121. **Plitz, W.; Veihelmann, A.; and Pellengahr, C.**: [Metal-on-metal pairs for hip prostheses]. *Orthopade*, 32(1): 17-22, 2003.
122. **Poss, R.; Brick, G. W.; Wright, R. J. et al.**: The effects of modern cementing techniques on the longevity of total hip arthroplasty. *Orthop Clin North Am*, 19(3): 591-8, 1988.
123. **R. Berlakovits, G. G.**: Ten years follow up on CLS-system(Spotorno) at a trauma unit. *European Journal of Trauma*, E-Supplement 1/2001: 62-65, 2001.
124. **Rader, C. P.; Hendrich, C.; Low, S. et al.**: [5- to 8-year results of total hip endoprosthesis implantation with the Muller straight shaft prosthesis (cemented TiAlNb shaft)]. *Unfallchirurg*, 103(10): 846-52, 2000.
125. **Revell, P. A., and Jellie, S. E.**: Interleukin 15 production by macrophages in the implant interface membrane of aseptically loosened joint replacements. *J Mater Sci Mater Med*, 9(12): 727-30, 1998.
126. **Riska, E.**: Ceramik endoprosthesis in total hip arthroplasty. *Clin Orthop*: 87-94, 1993.
127. **Robertsson, O., and Dunbar, M. J.**: Patient satisfaction compared with general health and disease-specific questionnaires in knee arthroplasty patients. *J Arthroplasty*, 16(4): 476-82, 2001.

128. **Robinson, R. P.; Lovell, T. P.; and Green, T. M.**: Hip arthroplasty using the cementless CLS stem. A 2--4-year experience. *J Arthroplasty*, 9(2): 177-92, 1994.
129. **Roe, R. J.; Grood, E. S.; Shastri, R. et al.**: Effect of radiation sterilization and aging on ultrahigh molecular weight polyethylene. *J Biomed Mater Res*, 15(2): 209-30, 1981.
130. **Sandhu, J.; Waddell, J. E.; Henry, M. et al.**: The role of T cells in polyethylene particulate induced inflammation. *J Rheumatol*, 25(9): 1794-9, 1998.
131. **Schmalzried, T. P.; Kwong, L. M.; Jasty, M. et al.**: The mechanism of loosening of cemented acetabular components in total hip arthroplasty. Analysis of specimens retrieved at autopsy. *Clin Orthop Relat Res*, (274): 60-78, 1992.
132. **Schramm, M.; Keck, F.; Hohmann, D. et al.**: Total hip arthroplasty using an uncemented femoral component with taper design: outcome at 10-year follow-up. *Arch Orthop Trauma Surg*, 120(7-8): 407-12, 2000.
133. **Schweizer, A.; Riede, U.; Maurer, T. B. et al.**: Ten-year follow-up of primary straight-stem prosthesis (MEM) made of titanium or cobalt chromium alloy. *Arch Orthop Trauma Surg*, 123(7): 353-6, 2003.
134. **Semlitsch, M., and Willert, H. G.**: Clinical wear behaviour of ultra-high molecular weight polyethylene cups paired with metal and ceramic ball heads in comparison to metal-on-metal pairings of hip joint replacements. *Proc Inst Mech Eng [H]*, 211(1): 73-88, 1997.
135. **Semlitsch, M., and Willert, H. G.**: Implant materials for hip endoprostheses: old proofs and new trends. *Arch Orthop Trauma Surg*, 114(2): 61-7, 1995.
136. **Sethi, R. K.; Neavyn, M. J.; Rubash, H. E. et al.**: Macrophage response to cross-linked and conventional UHMWPE. *Biomaterials*, 24(15): 2561-73, 2003.
137. **Siebold, R.; Scheller, G.; Schreiner, U. et al.**: [Long-term results with the cement-free Spotorno CLS shaft]. *Orthopade*, 30(5): 317-22, 2001.
138. **Smith-Petersen, M. N.**: The classic: Evolution of mould arthroplasty of the hip joint by M. N. Smith-Petersen. *J Bone Joint Surg*. 30B:L:59, 1948. *Clin Orthop Relat Res*, (134): 5-11, 1978.
139. **Sochart, D. H., and Porter, M. L.**: The long-term results of Charnley low-friction arthroplasty in young patients who have congenital dislocation, degenerative osteoarthritis, or rheumatoid arthritis. *J Bone Joint Surg Am*, 79(11): 1599-617, 1997.
140. **Soderman, P.**: On the validity of the results from the Swedish National Total Hip Arthroplasty register. *Acta Orthop Scand Suppl*, 71(296): 1-33, 2000.
141. **Soderman, P., and Malchau, H.**: Is the Harris hip score system useful to study the outcome of total hip replacement? *Clin Orthop*, (384): 189-97, 2001.
142. **Soderman, P., and Malchau, H.**: Validity and reliability of Swedish WOMAC osteoarthritis index: a self-administered disease-specific questionnaire (WOMAC) versus generic instruments (SF-36 and NHP). *Acta Orthop Scand*, 71(1): 39-46, 2000.
143. **Soderman, P.; Malchau, H.; and Herberts, P.**: Outcome after total hip arthroplasty: Part I. General health evaluation in relation to definition of failure in the Swedish National Total Hip Arthroplasty register. *Acta Orthop Scand*, 71(4): 354-9, 2000.
144. **Soderman, P.; Malchau, H.; Herberts, P. et al.**: Outcome after total hip arthroplasty: Part II. Disease-specific follow-up and the Swedish National Total Hip Arthroplasty Register. *Acta Orthop Scand*, 72(2): 113-9, 2001.

145. **Spotorno, L.; Romagnoli, S.; Ivaldo, N. et al.**: The CLS system. Theoretical concept and results. *Acta Orthop Belg*, 59 Suppl 1: 144-8, 1993.
146. **Spotorno, L.; Schenk, R. K.; Dietschi, C. et al.**: [Personal experiences with uncemented prostheses]. *Orthopade*, 16(3): 225-38, 1987.
147. **Streicher, R. M., and Thomsen, M.**: [Polyethylene as an implant material]. *Orthopade*, 32(1): 23-31, 2003.
148. **Stucki, G.; Meier, D.; Stucki, S. et al.**: [Evaluation of a German version of WOMAC (Western Ontario and McMaster Universities) Arthritis Index]. *Z Rheumatol*, 55(1): 40-9, 1996.
149. **Sumner, D. R.; Jasty, M.; Jacobs, J. J. et al.**: Histology of porous-coated acetabular components. 25 cementless cups retrieved after arthroplasty. *Acta Orthop Scand*, 64(6): 619-26, 1993.
150. **Sun, Y.; Sturmer, T.; Gunther, K. P. et al.**: [Incidence and prevalence of cox- and gonarthrosis in the general population]. *Z Orthop Ihre Grenzgeb*, 135(3): 184-92, 1997.
151. **Swanson, A. B.; Colwill, J. C.; and de Groot Swanson, G.**: Femoral head intramedullary stemmed implant. A historical prospective. *Clin Orthop*, (281): 107-11, 1992.
152. **Tanzer, M.; Gollish, J.; Leighton, R. et al.**: The effect of adjuvant calcium phosphate coating on a porous-coated femoral stem. *Clin Orthop Relat Res*, (424): 153-60, 2004.
153. **Tauber, C., and Kidron, A.**: Total hip arthroplasty revision using the press-fit CLS Spotorno cementless stem. Twenty-four hips followed between 1987 and 1998. *Arch Orthop Trauma Surg*, 120(3-4): 209-11, 2000.
154. **Thomas, P.**: [Allergic reactions to implant materials]. *Orthopade*, 32(1): 60-4, 2003.
155. **Thomsen, M.; von Strachwitz, B.; Mau, H. et al.**: [Survey of materials in hip endoprosthesis]. *Z Orthop Ihre Grenzgeb*, 133(1): 1-6, 1995.
156. **Thomsen, M., and Willmann, G.**: [Ceramic couplings in orthopedic surgery]. *Orthopade*, 32(1): 11-6, 2003.
157. **Thull, R.**: [Surface modifications to improve biocompatibility and mechanical properties of orthopedic implants]. *Orthopade*, 32(1): 51-9, 2003.
158. **Urban, R. M.; Jacobs, J. J.; Tomlinson, M. J. et al.**: Dissemination of wear particles to the liver, spleen, and abdominal lymph nodes of patients with hip or knee replacement. *J Bone Joint Surg Am*, 82(4): 457-76, 2000.
159. **Urban, R. M.; Tomlinson, M. J.; Hall, D. J. et al.**: Accumulation in liver and spleen of metal particles generated at nonbearing surfaces in hip arthroplasty. *J Arthroplasty*, 19(8 Suppl 3): 94-101, 2004.
160. **Visuri, T.; Pukkala, E.; Paavolainen, P. et al.**: Cancer risk after metal on metal and polyethylene on metal total hip arthroplasty. *Clin Orthop Relat Res*, (329 Suppl): S280-9, 1996.
161. **Warashina, H.; Sakano, S.; Kitamura, S. et al.**: Biological reaction to alumina, zirconia, titanium and polyethylene particles implanted onto murine calvaria. *Biomaterials*, 24(21): 3655-61, 2003.
162. **Wessinghage, D.**: [Historical aspects of joint prosthesis implantation. A bibliographic study]. *Orthopade*, 29(12): 1067-71, 2000.
163. **Willert, H. G.; Bertram, H.; and Buchhorn, G. H.**: Osteolysis in alloarthroplasty of the hip. The role of bone cement fragmentation. *Clin Orthop Relat Res*, (258): 108-21, 1990.

164. **Willert, H. G.; Bertram, H.; and Buchhorn, G. H.**: Osteolysis in alloarthroplasty of the hip. The role of ultra-high molecular weight polyethylene wear particles. *Clin Orthop Relat Res*, (258): 95-107, 1990.
165. **Williamson, O. D.**: Measuring the success of joint replacement surgery. *Med J Aust*, 171(5): 229-30, 1999.
166. **Zanetti, M.; Bruder, E.; Romero, J. et al.**: Bone marrow edema pattern in osteoarthritic knees: correlation between MR imaging and histologic findings. *Radiology*, 215(3): 835-40, 2000.
167. **Zichner, L., and Lindenfeld, T.**: [In-vivo wear of the slide combinations ceramics-polyethylene as opposed to metal-polyethylene]. *Orthopade*, 26(2): 129-34, 1997.