

6. References

Agre P, Saboori AM, Asimos A, and Smith BL. Purification and partial characterization of the Mr 30,000 integral membrane protein associated with the erythrocyte Rh (D) antigen. *J Biol Chem* 262: 17497-17503, 1987

Agre P, Preston GM, Smith BL, Jung JS, Raina S, Moon C, Guggino WB, and Nielsen S. Aquaporin CHIP: the archetypal molecular water channel. *Am J Physiol* 265: F463-F476, 1993

Agre P, Bonhivers M, Borgnia MJ. The aquaporins, blueprints for cellular plumbing systems. *J Biol Chem* 273(24): 14659-14662, 1998

¹Alberts B, Johnson A, Lewis J, Raff M, Roberts K, Walter P. The molecular biology of the cell (4th edition). p 616, *Garland Science*, 2002

²Alberts B, Johnson A, Lewis J, Raff M, Roberts K, Walter P. The molecular biology of the cell (4th edition). p 899, *Garland Science*, 2002

Amlal H, Chen Q, Habo K, Wang Z, Soleimani M. Fasting downregulates renal water channel AQP2 and causes polyuria. *Am J Physiol* 280: F513-F523, 2001

Amlal H, Krane CM, Chen Q, Soleimani M. Early polyuria and urinary concentrating defect in potassium deprivation. *Am J Physiol* 279: F655-F663, 2000

Bagnasco S, Balaban R, Fales HM, Yang YM, Burg M. Predominant osmotically active organic solutes in rat and rabbit renal medullas. *J Biol Chem* 261(13): 5872-7, 1986

Bagnasco SM, Peng T, Nakayama Y, Sands JM. Differential expression of individual UT-A urea transporter isoforms in rat kidney. *J Am Soc Nephrol* 11: 1980-1986, 2000

Beck FX, Dorge A, Rick R, Thurau K. Intra- and extracellular element concentrations of rat renal papilla in antidiuresis. *Kidney Int.* 25: 397-403, 1984

Beck FX, Muller E, Fraek ML, Dorge A, Thurau K. Inner-medullary organic osmolytes and inorganic electrolytes in K depletion [Abstract]. *Pflugers Arch* 439 (4): 471-476, 2000

Bradford MM. A rapid and sensitive method for the quantification of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* 72: 248-254, 1976

¹Capasso JM, Rivard CJ, Berl T. Long-term adaptation of renal cells to hypertonicity: role of MAP kinases and Na-K-ATPase. *Am J Physiol* 280: F768-F776, 2001

²Capasso JM, Rivard C, Berl T. The expression of the γ subunit of Na-K-ATPase is regulated by osmolality via C-terminal Jun kinase and phosphatidylinositol-3 kinase-dependent mechanisms. *PNAS* 98 (23): 13414-13419, 2001

Cha JH, Woo SK, Han KH, Kim YH, Handler JS, Kim J, Kwon HM. Hydration status affects nuclear distribution of transcription factor tonicity responsive enhancer binding protein in rat kidney. *J Am Soc Nephrol* 12: 2221-2230, 2001

Cheng A, Van HA, Yeager M, Verkman AS, Mitra AK. Three-dimensional organization of a human water channel. *Nature* 387: 627-630, 1997

Combet S, Geffroy N, Berthouaud V, Dick B, Teillet L, Verbavatz JM, Corman B, Trinh-Trang-Tan MM. Correction of age-related polyuria by DdAVP: molecular analysis of aquaporins and urea transporters. *Am J Physiol* 284: F199-F208, 2003

Chou CL, Rapko SI, Knepper MA. Phosphoinositide signalling in rat inner medullary collecting duct. *Am J Physiol* 274: F564-F572, 1998

Crawford JD, Doyle AP, Probst H. Service of urea in renal water conservation. *Am J Physiol* 196: 545-548, 1959

Davies SP, Reddy H, Caivano M, Cohen P. Specificity and mechanism of action of some commonly used protein kinase inhibitors. *Biochem J* (351): 95-105, 2000

Deen PM, Verdijk TMAJ, Knoers NVAM, Wieringa B, Monnens LAH, van Os CH, van Oost BA. Requirement of aquaporin-2 for vasopressin-dependent concentration of urine. *Science* 264: 92-95, 1994

Deen PM, Rijss JP, Mulders SM, Errington RJ, van Baal J, van Os CH. Aquaporin-2 transfection of Madin-Darby canine kidney cells reconstitutes vasopressin-regulated transcellular osmotic water transport. *J Am Soc Nephrol* 8(10): 1493-501, 1997

Denker BM, Smith BL, Kuhajda FP, Agre P. Identification, purification, and partial characterization of a novel Mr 28,000 integral membrane protein from erythrocytes and renal tubules. *J Biol Chem* 263(30): 15634-15642, 1988

Earm JH, Christensen BM, Frokiaer J, Marples D, Han JS, Knepper MA, Nielsen S. Decreased aquaporin-2 expression and apical plasma membrane delivery in kidney collecting ducts of polyuric hypercalcemic rats. *J Am Soc Nephrol* 9(12): 2181-2193, 1998

Ecelbarger CA, Nielsen S, Olson BR, Murase T, Baker EA, Knepper MA, Verbalis JG. Role of renal aquaporins in escape from vasopressin-induced antidiuresis in rat. *J Clin Invest.* 99: 1852-1863, 1997

Ecelbarger CA, Kim GH, Terris J, Masilamani S, Mitchell C, Reyes I, Verbalis JG, Knepper MA. Vasopressin-mediated regulation of epithelial sodium channel abundance in rat kidney. *Am J Physiol* 279(1): F46-53, 2000

Ferraris JD, Williams CK, Jung KY, Bedford JJ, Burg MB, Garcia-Perez A. ORE, a eukaryotic minimal essential osmotic response element. The aldose reductase gene in hyperosmotic stress. *J Biol Chem* 271(31): 18318-21, 1996

¹Ferraris JD, Persaud P, Williams CK, Chen Y, MB Burg. cAMP-independent role of PKA in tonicity-induced transactivation of tonicity-responsive enhancer/osmotic response element-binding protein. *PNAS* 99 (26): 16800–16805, 2002

²Ferraris JD, Williams CK, Persaud P, Zhang Z, Chen Y, Burg MB. Activity of the TonEBP/OREBP transactivation domain varies directly with extracellular NaCl concentration. *PNAS* 99(2): 739-744, 2002

Furuno M, Shinichi U, Fumiaki M, Sasaki S. Repressive regulation of the aquaporin-2 gene. *Am J Physiol* 271: F854-F860, 1996

Fujita N, Ishikawa SE, Sasaki S, Fujisawa G, Fushimi K, Marumo F, Saito T. Role of water channel AQP-CD in water retention in SIADH and cirrhotic rats. *Am J Physiol.* 269(6): F926-F931, 1995

Garzia-Perez A and Burg MB. Renal medullary osmolytes. *Physiological reviews: Vol.71, No.4, 1991*

Gonin S, Deschênes G, Roger F, Bens M, Martin PY, Carpentier JL, Vanderwalle A, Doucet A, Féraïlle E. Cyclic AMP increases cell surface expression of functional Na,K-ATPase Units in mammalian cortical collecting duct principal cells. *Mol Biol Cell (13): 255-264, 2001*

Gonzalez GA, Montminy MR. Cyclic AMP stimulates somatostatin gene transcription by phosphorylation of CREB at serine 133. *Cell 59: 675-80, 1989*

Gutsche HU, Peterson LN, Levine DZ. In vivo evidence of impaired solute transport by the thick ascending limb in potassium-depleted rats. *J Clin Invest 73: 908-916, 1984*

¹Handler JS and Kwon JM. Cell and molecular biology of organic osmolyte accumulation in hypertonic renal cells. *Nephron. 87: 106-110, 2001*

²Handler JS and Kwon JM. Transcriptional regulation by changes in tonicity. *Kidney Int 60: 408-411, 2001*

Hanski C, Klußmann E, Wang J, Böhm C, Ogorek D, Hanski ML, Krüger-Krasagakes S, Eberle J, Schmitt-Gräff A, Riecken EO. Fucosyltransferase III and sialyl-Le^x expression correlate in cultured colon carcinoma cells but not in human colon carcinoma tissue. *Glycoconjugate J 13: 1-7, 1996*

Hasler U, Mordasin D, Bens M, Bianchi M, Cluzeaud F, Rousselot M, Vanderwalle A, Féraïlle E, Martin PI. Long term regulation of aquaporin-2 expression in vasopressin-responsive renal collecting duct principal cells. *J Biol Chem 277 (12): 10379-10386, 2002*

Hoffert JD, Leitch V, Agre P, King LS. Hypertonic induction of aquaporin-5 expression through an ERK-dependent pathway. *J Biol Chem 275 (12): 9070-9077, 2000*

- Hozawa S, Holtzman EJ, Ausiello DA. cAMP motifs regulating transcription in the aquaporin 2 gene. *Am J Physiol* 270: C1695-C1702, 1996
- Jo I, Nielsen S, Harris HW. The 17 kD band identified by multiple anti-aquaporin antisera in rat kidney medulla is a histone. *Biochim Biophys Acta* 1324: 91-101, 1997
- Jung JS, Preston GM, Smith BL, Guggino WB, Agre P. Molecular structure of the water channel through aquaporin CHIP. The hourglass model. *J Biol Chem* 269(20): 14648-14654, 1994
- Kamsteeg EJ, Heijnen I, van Os CH, and Deen PMT. The subcellular localization of an Aquaporin-2 tetramer depends on the stoichiometry of phosphorylated and nonphosphorylated monomers. *J Cell Biol* 151 (4) : 919–929, 2000
- Kato A and Sands JM. Active sodium-urea counter-transport is inducible in the basolateral membrane of rat renal initial medullary collecting ducts. *J Clin Invest* 102(5): 1008-1015, 1998
- Klussmann E, Maric K, Wiesner B, Beyermann M, Rosenthal W. Protein kinase A anchoring proteins are required for vasopressin-mediated translocation of aquaporin-2 into cell membranes of renal principal cells. *J Biol Chem* 274: 4934-4938, 1999
- Klussmann E, Maric K, Rosenthal W. The mechanisms of aquaporin control in the renal collecting duct. *Rev Physiol Biochem Pharmacol* 141: 33-95, 2000
- Klussmann E, Tamma G, Lorenz D, Wiesner B, Maric K, Hofmann F, Aktories K, Valenti G, Rosenthal W. An inhibitory role of Rho in the vasopressin-mediated translocation of aquaporin-2 into cell membranes of renal principal cells. *J Biol Chem* 276: 20451-20457, 2001
- Ko BCB, Ruepp B, Bohren KM, Gabbay KH, Chung SSM. Identification of multiple osmotic response sequences in the human aldose reductase gene. *J Biol Chem* 272: 16431-16437, 1997
- Ko BCB, Lam AKM, Kapus A, Fan L, Chung SK, Chung SSM. Fyn and p38 signaling are both required for maximal hypertonic activation of the osmotic response element binding protein/tonicity-responsive enhancer binding protein (OREBP/ TonEBP). *J Biol Chem* 277(48): 46085-46092, 2002

Kültz D, Garcia-Perez A, Ferraris JD, Burg MB. Distinct regulation of osmoprotective genes in yeast and mammals. Aldose reductase osmotic response element is induced independent of p38 and stress-activated protein kinase/Jun N-terminal kinase in rabbit kidney cells. *J Biol Chem* 272(20): 13165-13170, 1997

Kültz D, Chakravarty D. Hyperosmolality in the form of elevated NaCl but not urea causes DNA damage in murine kidney cells. *PNAS* 98: 1999-2004, 2001

Kwon HM, Itoh T, Rim JS, and Handler JS. The MAP kinase cascade is not essential for transcriptional stimulation of osmolyte transporter genes. *Biochem Biophys Res Commun* 213: 975-979, 1995

Laemmli UK. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 227: 680-685, 1970

Lee SD, Colla E, Sheen ME, Na KY, Kwon HM, Multiple domains of TonEBP cooperate to stimulate transcription in response to hypertonicity. *JBC Papers in Press as Manuscript: M308795200*, 2003

Leitch V, Agre P, Landon SK. Altered ubiquitination and stability of aquaporin-1 in hypertonic stress. *PNAS* 98 (5): 2894-2898, 2000

Levinsky NG, Berliner RW. The role of urea in the urine concentration mechanism. *J Clin Invest* 38: 741-748, 1959

Li H, Lee S, Jap BK. Molecular design of aquaporin-1 water channel as revealed by electron crystallography. *Nat Struct Biol* 4: 263-265, 1997

Liebenhoff U, Rosenthal W. Identification of Rab3-, Rab5a- and synaptobrevin II-like proteins in a preparation of rat kidney vesicles containing the vasopressin-regulated water channel. *FEBS* 365: 209-213, 1995

López-Rodríguez C, Aramburu J, Rakeman AS, Rao A. NFAT5, a constitutively nuclear NFAT protein that does not cooperate with Fos and Jun. *PNAS* 96: 7214-7219, 1999

- López-Rodríguez C, Aramburu J, Jin L, Rakeman AS, Michino M, Rao A. Bridging the NFAT and NF-kappaB families: NFAT5 dimerization regulates cytokine gene transcription in response to osmotic stress. *Immunity* 15(1): 47-58, 2001
- Lorenz D, Krylov A, Hahm D, Hagen V, Rosenthal W, Pohl P, Maric K. Cyclic AMP is sufficient for triggering the exocytic recruitment of aquaporin-2 in renal epithelial cells. *EMBO Rep* 4(1): 88-93, 2003
- Maric K, Oksche A, Rosenthal W. Aquaporin-2 expression in primary cultured rat inner medullary collecting duct cells. *Am J Physiol* 275(5 Pt 2): F796-801, 1998
- Maric K, Wiesner B, Lorenz D, Klussmann E, Betz T, Rosenthal W. Cell volume kinetics of adherent epithelial cells measured by laser scanning reflection microscopy: determination of water permeability changes of renal principal cells. *Biophys J* 80(4): 1783-1790, 2001
- Marples D, Christensen S, Christensen EI, Ottosen PD, Nielsen S. Lithium-induced downregulation of aquaporin-2 water channel expression in rat kidney medulla. *J Clin Invest* 95(4): 1838-1845, 1995
- Marples D, Froekiaer J, Dorup J, Knepper MA, Nielsen S. Hypokalemia-induced downregulation of aquaporin-2 water channel expression in rat kidney medulla and cortex. *J Clin Invest* 97(8): 1960-8, 1996
- Marples D, Moenster Christensen B, Froekiaer J, Knepper MA, Nielsen S. Dehydration reverses vasopressin antagonist-induced diuresis and aquaporin-2 downregulation in rats. *Am J Physiol* 275(4): F400-F409, 1998
- Matsumura Y, Uchida S, Rai T, Sasaki S, Marumo Fs. Transcriptional regulation of aquaporin-2 water channel gene by cAMP. *J Am Soc Nephrol* 8: 861-867, 1997
- Matsuzaki T, Suzuki T, Takata K. Hypertonic induction of aquaporin 3 in MDCK cells. *Am J Physiol* (281): C55-C63, 2001

- Michea L, Ferguson DR, Peters EM, Andrews PM, Kirby MR, Burg MB. Cell cycle delay and apoptosis are induced by high salt and urea in renal medullary cells. *Am J Physiol* 278: F209-F218, 2000
- Miyai A, Yamauchi A, Moriyama T, Kaneko T, Takenaka M, Sugiura T, Kitamura H, Ando A, Tohyama M, Shimada S, Imai E, Kamada T. Expression of betaine mRNA: its unique localization and rapid regulation in rat kidney (Abstract). *Kidney Int* 50 (3): 819-827, 1996
- Miyakawa H, Woo SK, Chen CP, Dahl SC, Handler JS, Kwon HM. Cis- and trans-acting factors regulating transcription of the BGT1 gene in response to hypertonicity. *Am J Physiol* 274: F753-61, 1998
- Miyakawa H, Woo SK, Dahl SC, Handler JS, Kwon HM. Tonicity-responsive enhancer binding protein, a Rel-like protein that stimulates transcription in response to hypertonicity. *PNAS* 96: 2538-2542, 1999
- Moriyama T, Garcia-Perez A, Burg MB. Osmotic regulation of aldose reductase protein synthesis in renal medullary cells. *J Biol Chem* 264(28): 16810-16814, 1989
- Müller E, Neuhofer W, Ohno A, Rucker S, Thureau K, Beck FX. Heat shock proteins HSP25, HSP60, HSP72, HSP73, in isoosmotic cortex and hyperosmotic medulla of rat kidney. *Pflugers Arch* 431: 608-617, 1996
- Müller E, Neuhofer W, Burger-Kentischer A, Ohno A, Thureau K, Beck FX. Effects of long-term changes in medullary osmolality on heat shock proteins HSP25, HSP60, HSP72, and HSP73. *Pflugers Arch* 435: 705-712, 1998
- Murata K, Mitsuoka K, Hirai T, Walz T, Agre P, Heymann JB, Engel A, Fujioishi Y. Structural determinants of water permeation through Aquaporin-1. *Nature* 407: 599-605, 2000
- Muto S. Potassium transport in the mammalian collecting duct. *Physiol Reviews* 81 (1): 85-106, 2001
- Nakayama Y, Peng T, Sands JM, Bagnasco SM. The TonE/TonEBP pathway mediates tonicity-responsive regulation of UT-A transporter expression. *J Biol Chem* 275: 38275-38280, 2000

Nelson RD, Stricklett P, Gustafson C, Stevens A, Ausiello D, Brown D, Kohan DE. Expression of an AQP2 Cre recombinase transgene in kidney and male reproductive system of transgenic mice. *Am J Physiol* 275 (4): C216-C226, 1998

Neuhofer W, Müller E, Burger-Kentischer A, Fraek ML, Thureau K, Beck FX. Inhibition of NaCl-induced heat shock protein 72 expression renders MDCK cells susceptible to high urea concentrations. *Pflugers Arch* 437: 611-616, 1999

Neuhofer W, Lugmayr K, Fraek ML, Beck FX. Regulated overexpression of HSP72 protects Madin-Darby canine kidney cells from the detrimental effects of high urea concentrations. *J Am Soc Nephrol* 12: 2565-2571, 2001

Neuhofer W, Woo SK, Na KY, Grünbein R, Park WK, Nahm O, Beck FX, Kwon HM. Regulation of TonEBP transcriptional activator in MDCK cells following changes in ambient tonicity. *Am J Physiol* 283: C1604-1611, 2002

Nielsen S, DiGiovanni SR, Christensen EI, Knepper MA, Harris HW. Cellular and subcellular immunolocalization of vasopressin-regulated water channel in rat kidney. *PNAS* 90: 11663-11667, 1993

Nielsen S, Chou CL, Marples D, Christensen EI, Kishore BK, Knepper MA. Vasopressin increases water permeability of kidney collecting duct by inducing translocation of aquaporin-CD water channels to plasma membrane. *PNAS* 92(4): 1013-1017, 1995

Nielsen S, Frokiaer J, Marples D, Kwon TH, Agre P, Knepper MA. Aquaporins in the kidney: From molecules to medicine. *Physiol Rev.*28: 205-244, 2002

Ohara M, Martin PY, Xu DL, St John J, Pattison TA, Kim JK, Schrier RW. Upregulation of aquaporin 2 water channel expression in pregnant rats. *J Clin Invest* 101(5): 1076-1083, 1998

Ohtaka A, Muto S, Kawakami K, Nagano K, Asano Y. Hyperosmolality stimulates Na-K-ATPase gene expression in inner medullary collecting duct cells. *Am J Physiol* 270 (5): F728-F738, 1996

Peet M, Pratt JP. Lithium. Current status in psychiatric disorders. *Drugs* 46(1): 7-17, 1993

Pennell JP, Sanjana V, Frey NR, Jamison RL. The effect of urea infusion on the urinary concentrating mechanism in protein-depleted rats. *J Clin Invest* 55(2): 399-409, 1975

Preisser L, Teillet L, Aliotti S, Gobin R, Berthouaud V, Chevalier J, Corman B, Verbavatz JM. Downregulation of aquaporin-2 and -3 in aging kidney is independent of V(2) vasopressin receptor. *Am J Physiol Renal Physiol* 279(1): F144-52, 2000

Preston GM, Agre P. Isolation of the cDNA for erythrocyte integral membrane protein of 28 kilodaltons: member of an ancient channel family. *PNAS* 88(24): 11110-11114, 1991

Preston GM, Carroll TP, Guggino WB, Agre P. Appearance of water channels in *Xenopus* oocytes expressing red cell CHIP28 protein. *Science* 256(5055): 385-387, 1992

Promeneur D, Kwon TH, Yasui M, Kim GH, Froekiaer J, Knepper MA, Agre P, Nielsen S. Regulation of AQP6 mRNA and protein in response to altered acid-base or water balance. *Am J Physiol* 279 (6): F1014-F1026, 2000

Puliyanda DP, Ward DT, Baum MA, Hammond TG, Harris Jr. HW. Calpain-mediated AQP2 proteolysis in inner medullary collecting duct. *Bioch Biophys Res Comm* 303: 52-58, 2003

Rai T, Uchida S, Marumo F, Sasaki S. Cloning of rat and mouse aquaporin-2 gene promoters and identification of a negative cis-regulatory element. *Am J Physiol* 273: F264-F273, 1997

Rim JS, Atta MG, Dahl SC, Berry GT, Handler JS, Kwon HM. Transcription of the Sodium/*myo*-Inositol Cotransporter Gene Is Regulated by Multiple Tonicity-responsive Enhancers Spread over 50 Kilobase Pairs in the 5'-Flanking Region. *J Biol Chem* 273 (32): 20615-20621, 1998

Rosenthal W, Seibold A, Antaramian A, Lonergan M, Arthus MF, Hendy GN, Birnbaumer M, Bichet DG. Molecular identification of the gene responsible for congenital nephrogenic diabetes insipidus. *Nature* 359(6392): 233-235, 1992

Rossato M, Balercia G, Lucarelli G, Foresta C, Mantero F. Role of seminal osmolarity in the reduction of human sperm motility. *Int J Androl* 25(4): 230-235, 2002

Sambrook J, Fritsch EF, Maniatis T. Molecular Cloning 2nd edition. *Cold Spring Harbort Laboratory, 1989*

Sands JM, Naruse M, Jacobs JD, Wilcox JN, Klein JD. Changes in aquaporin-2 protein contribute to the urine concentrating defect in rats fed a low-protein diet. *J Clin Invest 97(12): 2807-14, 1996*

Sands JM, Naruse M, Baum M, Jo I, Hebert SC, Brown EM, Harris HW. Apical extracellular calcium/polyvalent cation-sensing receptor regulates vasopressin-elicited water permeability in rat kidney inner medullary collecting duct. *J Clin Invest 99(6): 1399-405, 1997*

Sands JM. Regulation of urea transporter proteins in kidney and liver. *Mount Sinai J of Med 67 (2): 112-119, 2000*

Schmidt-Nielsen B, Barrett JM, Graves B, Crossley B. Physiological and morphological responses of the rat kidney to reduced dietary protein. *Am J Physiol 248(2): F31-F42, 1985*

Sheikh-Hammad D, Di Mari J, Suki WN, Safirstein R, Watts BA III, and Rouse D. p38 kinase activity is essential for osmotic induction of mRNAs for HSP 70 and transporter for organic solute betaine in Madin-Darby canine kidney cells. *J Biol Chem 273: 1832-1837, 1998*

Sone M, Ohno A, Albrecht GJ, Thureau K, Beck FX. Restoration of urine concentrating ability and accumulation of medullary osmolytes after chronic diuresis. *Am J Physiol 269 (4): F480-490, 1995*

Stevens AL, Breton S, Gustafson CE, Bouley R, Nelson RD, Kohan DE, Brown D. Aquaporin 2 is a vasopressin-independent, constitutive apical membrane protein in rat vas deferens. *Am J Physiol 278: C791–C802, 2000*

Storm R. Diplom thesis. *FMP-Berlin / Freie Universität Berlin, 1999*

Storm R, Maric K, Klußmann E, Lorenz D, Rosenthal W. NHE proteins involved in the long-term regulation of aquaporin2 in primary cultured IMCD cells. *Molecular Biology and Physiology of Water and Solute Transport – MIP 2000; 1st – 5th July; Göteborg/Sweden (Abstract/Poster), 2000*

- Storm R, Klussmann E, Geelhaar A, Rosenthal W, Maric K. Osmolality and solute composition are strong regulators of AQP2 expression in renal principal cells. *Am J Physiol* 284(1): F189-198, 2003
- Stroud JC, López-Rodríguez C, Rao A, Chen L. Structure of a TonEBP-DNA complex reveals DNA encircled by a transcription factor. *Nat Struct Biol* 2: 90-94, 2002
- Sugiyama Y, Ota Y, Hara M, Inoue S. Osmotic stress up-regulates aquaporin-3 gene expression in cultured human keratinocytes. *Biochim Biophys A* 1522: 82-88, 2001
- Tamarappoo BK and Verkman AS. Defective aquaporin-2 trafficking in Nephrogenic Diabetes Insipidus and correction by chemical chaperones. *J Clin Invest* 101: 2257-2267, 1998
- Tamma G, Klussmann E, Maric K, Aktories K, Svelto M, Rosenthal W, Valenti G. Rho inhibits cAMP-induced translocation of aquaporin-2 into the apical membrane of renal cells. *Am J Physiol* 281: F1092-F1101, 2001
- Terris J, Ecelbarger CA, Nielsen S, Knepper MA. Long-term regulation of four renal aquaporins in rats. *Am J Physiol* 271: F414-422, 1996
- Tian W, Cohen DM. Urea inhibits hypertonicity-inducible TonEBP expression and action. *Am J Physiol* 280(5): F904-912, 2001
- Trama J, Lu Q, Hawley RG, Ho SN. The NFAT-related protein NFATL1 (TonEBP/NFAT5) is induced upon T cell activation in a calcineurin-dependent manner. *J Immunol* 165(9): 4884-4894, 2000
- Umenishi F and Schrier RW. Identification and characterization of a novel hypertonicity-responsive element in the human aquaporin-1 gene. *Biochem Biophys Res Comm* 292: 711-775, 2002
- Umenishi F and Schrier RW. Hypertonicity-induced aquaporin-1 expression is mediated by the activation of MAPK pathways and hypertonicity-responsive element in the AQP1 gene. *J Biol Chem* 278 (18): 15765-15770, 2003

Valtin H. Sequestration of urea and nonurea solutes in renal tissues of rats with hereditary hypothalamic Diabetes insipidus: effects of vasopressin and dehydration on the countercurrent mechanism. *J Clin Invest* 45: 337-345, 1966

Vinciguerra M, Dêschenes G, Hasler U, Mordasini D, Rousselot M, Doucet A, Vanderwalle A, Martin PY, Féraille E. Intracellular Na⁺ controls cell surface expression of Na,K-ATPase via a cAMP-independent PKA pathway in mammalian kidney collecting duct cells. *Mol Biol Cell* 14: 2677-2688, 2003

Watts BA 3rd, Good DW. Hypoosmolality stimulates apical membrane Na⁽⁺⁾/H⁽⁺⁾ exchange and HCO₃⁽⁻⁾ absorption in renal thick ascending limb. *J Clin Invest* 104(11): 1593-602; 1999

Wojtaszek PA, Heasley LE, Berl T. In vivo regulation of MAP kinases in Ratus norvegicus renal papilla by water-loading and restriction. *J Clin Invest* 102: 1874-1881, 1998

¹Woo SK, Dahl SC, Handler JS, Kwon HM. Bidirectional regulation of tonicity-responsive enhancer binding protein in response to changes in tonicity. *Am J Physiol* 278: F1006-F1012, 2000

²Woo SK, Maouyo D, Handler JS, Kwon HM. Nuclear redistribution of tonicity-responsive enhancer binding protein requires proteasome activity. *Am J Physiol* 278(2): C323-330, 2000

Woo SK, Handler JS, Kwon HM. TonEBP stimulates transcription of a HSP70 gene [abstract]. *FASEB J* 15: A851, 2001

Woo SK, Lee SD, Kwon HM. TonEBP transcriptional activator in the cellular response to increased osmolality. *Pflugers Arch - Eur J Physiol* 444: 579-585, 2002

Xu DL, Martin PY, Ohara M, St John J, Pattison T, Meng X, Morris K, Kim JK, Schrier RW. Upregulation of aquaporin-2 water channel expression in chronic heart failure rat. *J Clin Invest* 99(7): 1500-1505, 1997

Yamaki M, Kusano E, Tetsuka T, Takeda S, Homma S, Murayama N, Asano Y. Cellular mechanism of lithium-induced nephrogenic diabetes insipidus in rats. *Am J Physiol* 261(2): F505-511, 1991

Yamauchi A, Miyai A, Shimada S, Minami Y, Tohyama M, Imai E, Kamada T, Ueda N. Localization and rapid regulation of Na⁺/myo-inositol cotransporter in rat kidney (Abstract). *J Clin Invest* 96 (3): 1195-1201, 1995

Yancey PH, Clark ME, Hand SC. Living with water stress: evolution of osmolyte systems. *Science* 217: 1214-1222, 1982

Zhao H, Tian W, and Cohen DM. Rottlerin inhibits tonicity-dependent expression and action of TonEBP in a PKC δ -independent fashion. *Am J Physiol* 282: F710-F717, 2002

Zhuang S, Hirai SI, Ohno S. Hyperosmolality induces activation of cPKC and nPKC, a requirement for ERK1/2 activation in NIH/3T3 cells. *Am J Physiol* 278: C102-C109, 2000