

9 Literaturverzeichnis

- [1] Abriél H, Loffing J, Rebhun JF, Pratt JH, Schild L, Horisberger JD, Rotin D, Staub O. Defective regulation of the epithelial Na⁺ channel by Nedd4 in Liddle's syndrome. *J Clin Invest* 1999; 103:667-673.
- [2] Agrawal R, Lichtenstein I, Siebe H, Hierholzer K. In vitro 6 beta-hydroxylation of progesterone in human renal tissue. *Wien Klin Wochenschr* 1997; 109:493-496.
- [3] Albiston AL, Obeyesekere VR, Smith RE, Krozowski ZS. Cloning and tissue distribution of the human 11 beta-hydroxysteroid dehydrogenase type 2 enzyme. *Mol Cell Endocrinol* 1994; 105:R11-R17.
- [4] Allolio B, Oremus M, Reincke M, Schaeffer HJ, Winkelmann W, Heck G, Schulte HM. High-dose progesterone infusion in healthy males: evidence against antiglucocorticoid activity of progesterone. *Eur J Endocrinol* 1995; 133(6):696-700.
- [5] Anderson RA, Baillie TA, Axelson M, Cronholm T, Sjovall K, Sjovall J. Stable isotope studies on steroid metabolism and kinetics: sulfates of 3alpha-hydroxy-5 alpha-pregnane derivatives in human pregnancy. *Steroids* 1990; 55:443-457.
- [6] Armanini D, Endres S, Kuhnle U, Weber PC. Parallel determination of mineralocorticoid and glucocorticoid receptors in T- and B-lymphocytes of human spleen. *Acta Endocrinol (Copenh)* 1988; 118:479-482.
- [7] Arriza JL, Weinberger C, Cerelli G, Glaser TM, Handelin BL, Housman DE, Evans RM. Cloning of human mineralocorticoid receptor complementary DNA: structural and functional kinship with the glucocorticoid receptor. *Science* 1987; 237:268-275.
- [8] Auchus RJ, Lee TC, Miller WL. Cytochrome b5 augments the 17,20-lyase activity of human P450c17 without direct electron transfer. *J Biol Chem* 1998; 273(6):3158-3165.
- [9] Auzou G, Fagart J, Souque A, Hellal-Levy C, Wurtz JM, Moras D, Rafestin-Oblin ME. A single amino acid mutation of ala-773 in the mineralocorticoid receptor confers agonist properties to 11beta-substituted spiro lactones. *Mol Pharmacol* 2000; 58:684-691.
- [10] Baillie TA, Curstedt T, Sjovall K, Sjovall J. Production rates and metabolism of sulphates of 3 beta-hydroxy-5 alpha-pregnane derivatives in pregnant women. *J Steroid Biochem* 1980; 13:1473-1488.
- [11] Ballard PL, Carter JP, Graham BS, Baxter JD. A radioreceptor assay for evaluation of the plasma glucocorticoid activity of natural and synthetic steroids in man. *J Clin Endocrinol Metab* 1975; 41:290-304.

- [12] Bartter FC, Henkin RI, Bryan GT. Aldosterone hypersecretion in "non-salt-losing" congenital adrenal hyperplasia. *J Clin Invest* 1968; 47:1742-1752.
- [13] Beavan S, Horner A, Bord S, Ireland D, Compston J. Colocalization of glucocorticoid and mineralocorticoid receptors in human bone. *J Bone Miner Res* 2001; 16:1496-1504.
- [14] Belelli D, Bolger MB, Gee KW. Anticonvulsant profile of the progesterone metabolite 5 alpha-pregnan-3alpha-ol-20-one. *Eur J Pharmacol* 1989; 166:325-329.
- [15] Binart N, Lombes M, Rafestin-Oblin ME, Baulieu EE. Characterization of human mineralocorticosteroid receptor expressed in the baculovirus system. *Proc Natl Acad Sci U S A*. 1991; 88:10681-10685.
- [16] Bird CE, Wiqvist N, Diczfalusy E, Solomon S. Metabolism of progesterone by the perfused previable human fetus. *J Clin Endocrinol Metab* 1966; 26:1144-1154.
- [17] Brown RD, Strott CA, Liddle GW. Plasma deoxycorticosterone in normal and abnormal pregnancy. *J Clin Endocrinol Metab* 1972; 35:736-742.
- [18] Bumke-Vogt C, Bähr V, Diederich S, Herrmann SM, Anagnostopoulos I, Oelkers W, Quinkler M. Expression of the progesterone receptor and progesterone-metabolising enzymes in the female and male human kidney. *J Endocrinol* 2002; 175:349-364.
- [19] Burton G, Galigniana M, De Lavallaz S, Brachet-Cota AL, Sproviero EM, Ghini AA, Lantos CP, Damasco MC. Sodium-retaining activity of some natural and synthetic 21-deoxysteroids. *Mol Pharmacol* 1995; 47:535-543.
- [20] Buster JE, Chang RJ, Preston DL, Elashoff RM, Cousins LM, Abraham GE, Hobel CJ, Marshall JR. Interrelationships of circulating maternal steroid concentrations in third trimester pregnancies. I. C21 steroids: progesterone, 16alpha-hydroxyprogesterone, 17 alpha-hydroxyprogesterone, 20alpha-dihydroprogesterone, delta 5-pregnenolone, delta 5-pregnenolone sulfate, and 17-hydroxy delta 5-pregnenolone. *J Clin Endocrinol Metab* 1979; 48:133-138.
- [21] Butkus A, Congiu M, Scoggins BA, Coghlan JP. The affinity of 17 alpha-hydroxyprogesterone and 17 alpha, 20alpha-dihydroxyprogesterone for classical mineralocorticoid or glucocorticoid receptors. *Clin Exp Pharmacol Physiol* 1982; 9:157-163.
- [22] Casey ML, Howell ML, Winkel CA, Simpson ER, MacDonald PC. Deoxycorticosterone sulfate biosynthesis in human fetal kidney. *J Clin Endocrinol Metab* 1981; 53:990-996.
- [23] Casey ML, MacDonald PC. Formation of deoxycorticosterone from progesterone in extraadrenal tissues: demonstration of steroid 21-hydroxylase activity in human aorta. *J Clin Endocrinol Metab* 1982; 55(4):804-806.

- [24] Casey ML, Winkel CA, MacDonald PC. Conversion of progesterone to deoxycorticosterone in the human fetus: steroid 21-hydroxylase activity in fetal tissues. *J Steroid Biochem.* 1983; 18:449-452.
- [25] Chany E, Slaunwhine WK, Sandberg AA. Biliary and urinary metabolites of 4-C¹⁴ progesterone in human subjects. *J Clin Endocrinol Metab* 1960; 20:1568
- [26] Chapman AB, Zamudio S, Woodmansee W et al. Systemic and renal hemodynamic changes in the luteal phase of the menstrual cycle mimic early pregnancy. *Am J Physiol* 1997; 273(5 Pt 2):F777-F782.
- [27] Chen SY, Bhargava A, Mastroberardino L, Meijer OC, Wang J, Buse P, Firestone GL, Verrey F, Pearce D. Epithelial sodium channel regulated by aldosterone-induced protein sgk. *Proc Natl Acad Sci U S A* 1999; 96:2514-2519.
- [28] Cheng Y, Prusoff WH. Relationship between the inhibition constant (K₁) and the concentration of inhibitor which causes 50 per cent inhibition (I₅₀) of an enzymatic reaction. *Biochem Pharmacol* 1973; 22:3099-3108.
- [29] Couette B, Jalaguier S, Hellal-Levy C, Lupo B, Fagart J, Auzou G, Rafestin-Oblin ME. Folding requirements of the ligand-binding domain of the human mineralocorticoid receptor. *Mol Endocrinol* 1998; 12:855-863.
- [30] Couette B, Le Ricousse S, Fortin D, Rafestin-Oblin ME, Richard-Foy H. The establishment of the long terminal repeat of the mouse mammary tumor virus into CV-1 cells allows a functional analysis of steroid receptors. *Biochim Biophys Acta* 1994; 1219:607-612.
- [31] Couzinet B, Meduri G, Lecce MG et al. The postmenopausal ovary is not a major androgen-producing gland. *J Clin Endocrinol Metab* 2001; 86(10):5060-6.
- [32] Crawley JN, Glowa JR, Majewska MD, Paul SM. Anxiolytic activity of an endogenous adrenal steroid. *Brain Res* 1986; 398:382-385.
- [33] Cunningham FG, MacDonald PC, Gant NF. *Williams Obstetrics.* 1989 Appleton & Lange, Norwalk, CT, p 155
- [34] Demura T, Driscoll WJ, Strott CA. Nuclear progesterone-binding protein in the guinea pig adrenal cortex: distinction from the classical progesterone receptor. *Endocrinology* 1989; 124:2200-2207.
- [35] Diederich S, Quinkler M, Hanke B, Bähr V, Oelkers W. 11 beta-hydroxysteroid dehydrogenases: key enzymes in the action of mineralocorticoids and glucocorticoids. *Dtsch Med Wochenschr* 1999; 124:51-55.
- [36] Dinudom A, Harvey KF, Komwatana P, Young JA, Kumar S, Cook DI. Nedd4 mediates control of an epithelial Na⁺ channel in salivary duct cells by cytosolic Na⁺. *Proc Natl Acad Sci U S A* 1998; 95:7169-7173.

- [37] Dunn JF, Nisula BC, Rodbard D. Transport of steroid hormones: binding of 21 endogenous steroids to both testosterone-binding globulin and corticosteroid-binding globulin in human plasma. *J Clin Endocrinol Metab* 1981; 53:58-68.
- [38] Edelman IS. Mechanism of action of aldosterone: energetic and permeability factors. *J Endocrinol* 1979; 81:49P-53P.
- [39] Epple HJ, Schulzke JD, Schmitz H, Fromm M. Enzyme- and mineralocorticoid receptor-controlled electrogenic Na⁺ absorption in human rectum in vitro. *Am J Physiol* 1995; 269:G42-G48.
- [40] Evans RM. The steroid and thyroid hormone receptor superfamily. *Science* 1988; 240:889-895.
- [41] Everett RB, Worley RJ, MacDonald PC, Gant NF. Modification of vascular responsiveness to angiotensin II in pregnant women by intravenously infused 5alpha-dihydroprogesterone. *Am J Obstet Gynecol* 1978; 131:352-7.
- [42] Fagart J, Couette B, Souque A, Davioud E, Marquet A, Rafestin-Oblin ME. Photoaffinity labelling of the human mineralocorticoid receptor with steroids having a reactive group at position 3, 18 or 21. *Biochim Biophys Acta* 1998; 1388:35-44.
- [43] Fagart J, Wurtz JM, Souque A, Hellal-Levy C, Moras D, Rafestin-Oblin ME. Antagonism in the human mineralocorticoid receptor. *EMBO J* 1998; 17:3317-3325.
- [44] Farman N, Rafestin-Oblin ME. Multiple aspects of mineralocorticoid selectivity. *Am J Physiol Renal Physiol* 2001; 280:F181-F192.
- [45] Frey FJ, Escher G, Frey BM. Pharmacology of 11 beta-hydroxysteroid dehydrogenase. *Steroids* 1994; 59:74-9.
- [46] Frindt G, Palmer LG. Regulation of Na channels in the rat cortical collecting tubule: effects of cAMP and methyl donors. *Am J Physiol* 1996; 271:F1086-F1092.
- [47] Fronius M, Rehn M, Eckstein-Ludwig U, Clauss W. Inhibitory non-genomic effects of progesterone on Na⁺ absorption in epithelial cells from *Xenopus* kidney (A6). *J Comp Physiol [B]* 2001; 171(5):377-86.
- [48] Fukushima K, Sasano H, Nagura H, Sasaki I, Matsuno S, Krozowski ZS. Immunohistochemical localization of mineralocorticoid receptor in human gut. *Tohoku J Exp Med* 1991; 165:155-163.
- [49] Fukushima K, Sasano H, Sasaki I, Nagura H, Funayama Y, Matsuno S. Increased expression of mineralocorticoid receptor in human ileum after total colectomy: immunohistochemical and immunoblotting studies. *Tohoku J Exp Med* 1994; 173:383-390.

- [50] Fuller PJ, Funder JW. Mineralocorticoid and glucocorticoid receptors in human kidney. *Kidney Int* 1976; 10:154-157.
- [51] Funder JW, Pearce PT, Smith R, Smith AI. Mineralocorticoid action: target tissue specificity is enzyme, not receptor, mediated. *Science* 1988; 242:583-585.
- [52] Funder JW. How can aldosterone act as a mineralocorticoid? *Endocr Res* 1989;15:227-238.
- [53] Galigniana MD, Vicent GP, Piwien-Pilipuk G, Burton G, Lantos CP. Mechanism of action of the potent sodium-retaining steroid 11,19-oxidoprogesterone. *Mol Pharmacol* 2000; 58:58-70.
- [54] Garty H, Palmer LG. Epithelial sodium channels: function, structure, and regulation. *Physiol Rev* 1997; 77:359-396.
- [55] Geller DS, Farhi A, Pinkerton N, Fradley M, Moritz M, Spitzer A, Meinke G, Tsai FT, Sigler PB, Lifton RP. Activating mineralocorticoid receptor mutation in hypertension exacerbated by pregnancy. *Science*. 2000; 289:119-123.
- [56] Gellersen B, Kempf R, Telgmann R, DiMattia GE. Nonpituitary human prolactin gene transcription is independent of Pit-1 and differentially controlled in lymphocytes and in endometrial stroma. *Mol Endocrinol*. 1994; 8:356-373.
- [57] Granick S, Kappas A. Steroid induction of porphyrin synthesis in liver cell culture. I. Structural basis and possible physiological role in the control of heme formation. *J Biol Chem* 1967; 242:4587-4593.
- [58] Greig M, Macnaughton MC. Radioactive metabolites in the liver and adrenals of the human foetus after administration of [4-14C]progesterone. *J Endocrinol* 1967; 39:153-62.
- [59] Gyermek L. Effects of pregnanolone and progesterone. *Lancet* 1968; 2:1195.
- [60] Hellal-Levy C, Couette B, Fagart J, Souque A, Gomez-Sanchez C, Rafestin-Oblin M. Specific hydroxylations determine selective corticosteroid recognition by human glucocorticoid and mineralocorticoid receptors. *FEBS Lett* 1999; 464:9-13.
- [61] Hellal-Levy C, Fagart J, Souque A, Rafestin-Oblin ME. Mechanistic aspects of mineralocorticoid receptor activation. *Kidney Int* 2000; 57:1250-1255.
- [62] Hellal-Levy C, Fagart J, Souque A, Wurtz JM, Moras D, Rafestin-Oblin ME. Crucial role of the H11-H12 loop in stabilizing the active conformation of the human mineralocorticoid receptor. *Mol Endocrinol* 2000; 14:1210-1221.
- [63] Hirasawa G, Sasano H, Takahashi K, Fukushima K, Suzuki T, Hiwatashi N, Toyota T, Krozowski ZS, Nagura H. Colocalization of 11 beta-hydroxysteroid dehydrogenase type II and mineralocorticoid receptor in human epithelia. *J Clin Endocrinol Metab* 1997; 82:3859-3863.

- [64] Hirasawa G, Takeyama J, Sasano H, Fukushima K, Suzuki T, Muramatu Y, Darnel AD, Kaneko C, Hiwatashi N, Toyota T, Nagura H, Krozowski ZS. 11Beta-hydroxysteroid dehydrogenase type II and mineralocorticoid receptor in human placenta. *J Clin Endocrinol Metab* 2000; 85:1306-1309.
- [65] Holzbauer M. Physiological aspects of steroids with anaesthetic properties. *Med Biol* 1976; 54:227-42.
- [66] Iswari S, Colas AE, Karavolas HJ. Binding of 5 alpha-dihydroprogesterone and other progestins to female rat anterior pituitary nuclear extracts. *Steroids* 1986; 47:189-203.
- [67] Jamieson PM, Chapman KE, Edwards CR, Seckl JR. 11 beta-hydroxysteroid dehydrogenase is an exclusive 11 beta- reductase in primary cultures of rat hepatocytes: effect of physicochemical and hormonal manipulations. *Endocrinology* 1995; 136:4754-4761.
- [68] Jensen FC et al. Infection of human and simian tissue cultures with Rous Sarcoma Virus. *Proc Natl Acad Sci U S A* 1964; 52: 53-59.
- [69] Johansson ED, Jonasson LE. Progesterone levels in amniotic fluid and plasma from women. I. Levels during normal pregnancy. *Acta Obstet Gynecol Scand* 1971; 50:339-343.
- [70] Kamynina E, Staub O. Concerted action of ENaC, Nedd4-2, and Sgk1 in transepithelial Na(+) transport. *Am J Physiol Renal Physiol* 2002; 283:F377-387.
- [71] Kappas A, Soybel W, Glickman P. Fever-producing steroids of endogenous origin in men. *Arch Intern Med* 1960; 105:701
- [72] Kato K, Sasano H, Ohara S, Sekine H, Mochizuki S, Mune T, Yasuda K, Nagura H, Shimosegawa T, Toyota T, Krozowski Z. Coexpression of mineralocorticoid receptors and 11beta-hydroxysteroid dehydrogenase 2 in human gastric mucosa. *J Clin Endocrinol Metab* 1999; 84:2568-2573.
- [73] Kenouch S, Lombes M, Delahaye F, Eugene E, Bonvalet JP, Farman N. Human skin as target for aldosterone: coexpression of mineralocorticoid receptors and 11 beta-hydroxysteroid dehydrogenase. *J Clin Endocrinol Metab* 1994; 79:1334-1341.
- [74] Kobayashi T, Cohen P. Activation of serum- and glucocorticoid-regulated protein kinase by agonists that activate phosphatidylinositide 3-kinase is mediated by 3-phosphoinositide-dependent protein kinase-1 (PDK1) and PDK2. *Biochem J* 1999; 339:319-328.
- [75] Kuhnle U, Land M, Ulick S. Evidence for the secretion of an antimineralocorticoid in congenital adrenal hyperplasia. *J Clin Endocrinol Metab* 1986; 62:934-940.

- [76] Kumar S, Tomooka Y, Noda M. Identification of a set of genes with developmentally down-regulated expression in the mouse brain. *Biochem Biophys Res Commun* 1992; 185:1155-1161.
- [77] Laatikainen T, Karjalainen O. Excretion of conjugates of neutral steroids in human bile during late pregnancy. *Acta Endocrinol (Copenh)* 1972; 69:775-788.
- [78] Laidlaw JC, Ruse JL, Gornall AG. The influence of estrogen and progesterone on aldosterone excretion. *J Clin Endocrinol Metab* 1962; 22:161-71.
- [79] Landau RL, Bergenstal DM, Lugibihl K, Kascht ME. The metabolic effects of progesterone in man. *J Clin Endocrinol Metab* 1955; 15: 1194-1215.
- [80] Landau RL, Lugibihl K, Bergenstal DM, Dimick DF. The metabolic effects of progesterone in man: Dose response relationships. *J Lab Clin Med* 1957; 50:613-620.
- [81] Landau RL, Lugibihl K. Inhibition of the sodium-retaining influence of aldosterone by progesterone. *J Clin Endocrinol Metab* 1958; 18:1237-1245.
- [82] Latif SA, Sheff MF, Ribeiro CE, Morris DJ. Selective inhibition of sheep kidney 11 beta-hydroxysteroid dehydrogenase isoform 2 activity by 5 alpha-reduced (but not 5 beta) derivatives of adrenocorticosteroids. *Steroids* 1997; 62:230-237.
- [83] Lee A, Ellis G. Serum 17 alpha-hydroxyprogesterone in infants and children as measured by a direct radioimmunoassay kit. *Clin Biochem* 1991; 24(6):505-511.
- [84] Lim-Tio SS, Keightley MC, Fuller PJ. Determinants of specificity of transactivation by the mineralocorticoid or glucocorticoid receptor. *Endocrinology* 1997; 138:2537-2543.
- [85] Lisboa BP, Gustafsson JA. Biosynthesis of 16-alpha-hydroxyprogesterone by human fetal liver microsomes. *Steroids* 1968; 12:249-60.
- [86] Liu W, Wang J, Sauter NK, Pearce D. Steroid receptor heterodimerization demonstrated in vitro and in vivo. *Proc Natl Acad Sci U S A* 1995; 92:12480-12484.
- [87] Lombes M, Alfaidy N, Eugene E, Lessana A, Farman N, Bonvalet JP. Prerequisite for cardiac aldosterone action. Mineralocorticoid receptor and 11 beta-hydroxysteroid dehydrogenase in the human heart. *Circulation* 1995;92:175-182.
- [88] Lombes M, Binart N, Oblin ME, Joulin V, Baulieu EE. Characterization of the interaction of the human mineralocorticosteroid receptor with hormone response elements. *Biochem J* 1993; 292:577-583.

- [89] Lombes M, Farman N, Oblin ME, Baulieu EE, Bonvalet JP, Erlanger BF, Gasc JM. Immunohistochemical localization of renal mineralocorticoid receptor by using an anti-idiotypic antibody that is an internal image of aldosterone. *Proc Natl Acad Sci U S A* 1990; 87:1086-1088.
- [90] Lombes M, Kenouch S, Souque A, Farman N, Rafestin-Oblin ME. The mineralocorticoid receptor discriminates aldosterone from glucocorticoids independently of the 11 beta-hydroxysteroid dehydrogenase. *Endocrinology* 1994; 135:834-840.
- [91] MacDonald PC, Dombroski RA, Casey ML. Recurrent secretion of progesterone in large amounts: an endocrine/metabolic disorder unique to young women? *Endocr Rev* 1991; 12:372-401.
- [92] Maeyama M, Matuoka H, Tuchida Y, Hashimoto Y. Metabolism of neutral steroids in the human fetus. I. Distribution of radioactive material in the previable human fetus after administration of progesterone-4-¹⁴C. *Steroids* 1969;14:144-150.
- [93] Majewska MD, Harrison NL, Schwartz RD, Barker JL, Paul SM. Steroid hormone metabolites are barbiturate-like modulators of the GABA receptor. *Science* 1986; 232:1004-1007.
- [94] Marver D, Edelman IS. Dihydrocortisol: a potential mineralocorticoid. *J Steroid Biochem* 1978; 9:1-7.
- [95] Matulich DT, Spindler BJ, Schambelan M, Baxter JD. Mineralocorticoid receptors in human kidney. *J Clin Endocrinol Metab* 1976; 43:1170-1174.
- [96] McShane PM, Fencl MD. Conversion of progesterone to corticosteroids by the midterm fetal adrenal and kidney. *Steroids* 1983; 42(3):299-310.
- [97] Mercer J, Funder JW. The affinity of hydroxylated progesterone derivatives for classical steroid receptors. *J Steroid Biochem* 1978; 9:33-37.
- [98] Milewich L, Gomez-Sanchez C, Madden JD, MacDonald PC. Isolation and characterization of 5alpha-pregnane-3,20-dione and progesterone in peripheral blood of pregnant women. measurement throughout pregnancy. *Gynecol Invest* 1975; 6:291-306.
- [99] Mirshahi M, Mirshahi S, Golestaneh N, Mishal Z, Nicolas C, Hecquet C, Agarwal MK. Demonstration of the mineralocorticoid hormone receptor and action in human leukemic cell lines. *Leukemia* 2000;14:1097-1104.
- [100] Mirshahi M, Mirshahi S, Golestaneh N, Nicolas C, Mishal Z, Lounes KC, Hecquet C, Dagonet F, Pouliquen Y, Agarwal MK. Mineralocorticoid hormone signaling regulates the 'epithelial sodium channel' in fibroblasts from human cornea. *Ophthalmic Res* 2001; 33:7-19.

- [101] Mizoguchi H, Levere RD. Enhancement of heme and globin synthesis in cultured human marrow by certain 5-H steroid metabolites. *J Exp Med* 1971; 134:1501-1512.
- [102] Morris DJ. The biological significance of reduced metabolites of aldosterone. *Hum Pathol* 1979; 10:128-131.
- [103] Motulsky HJ. *Analyzing Data with GraphPad Prism*. 1999, GraphPad Software Inc., San Diego CA
- [104] Mujais SK, Nora NA, Chen Y. Regulation of the renal Na:K pump: role of progesterone. *J Am Soc Nephrol* 1993; 3(8):1488-1495.
- [105] Murakami T, Watanabe Ogura E, Tanaka Y, Yamamoto M. High blood pressure lowered by pregnancy. *Lancet* 2000; 356:1980.
- [106] Myles K, Funder JW. Progesterone binding to mineralocorticoid receptors: in vitro and in vivo studies. *Am J Physiol* 1996; 270:E601-E607.
- [107] Myslobodsky MS, Kofman O. Regular and lasting neocortical spiking produced by systemic administration of a steroid derivative in the rat. *Neuropharmacology* 1983; 22:157-164.
- [108] Nolten WE, Lindheimer MD, Oparil S, Ehrlich EN. Desoxycorticosterone in normal pregnancy. I. Sequential studies of the secretory patterns of desoxycorticosterone, aldosterone, and cortisol. *Am J Obstet Gynecol* 1978; 132:414-420.
- [109] Oelkers W, Schöneshofer M, Blumel A. Effects of progesterone and four synthetic progestagens on sodium balance and the renin-aldosterone system in man. *J Clin Endocrinol Metab* 1974; 39:882-890.
- [110] Oelkers WK. Effects of estrogens and progestogens on the renin-aldosterone system and blood pressure. *Steroids* 1996; 61:166-171.
- [111] Oparil S, Ehrlich EN, Lindheimer MD. Effect of progesterone on renal sodium handling in man: relation to aldosterone excretion and plasma renin activity. *Clin Sci Mol Med* 1975; 49(2):139-147.
- [112] Paillard J, Baviera E, Agarwal MK. Gluco- and mineralocorticoid receptors in human liver and kidney. *Biochem Med* 1980; 24:201-209.
- [113] Park J, Leong ML, Buse P, Maiyar AC, Firestone GL, Hemmings BA. Serum and glucocorticoid-inducible kinase (SGK) is a target of the PI3-kinase-stimulated signaling pathway. *EMBO J* 1999; 18:3024-3033.
- [114] Parker CR Jr, Winkel CA, Rush AJ Jr, Porter JC, MacDonald PC. Plasma concentrations of 11-deoxycorticosterone in women during the menstrual cycle. *Obstet Gynecol* 1981; 58:26-30.

- [115] Pearson Murphy BE, Allison CM. Determination of progesterone and some of its neuroactive ring A-reduced metabolites in human serum. *J Steroid Biochem Mol Biol* 2000; 74:137-142.
- [116] Pearson Murphy BE, Steinberg SI, Hu FY, Allison CM. Neuroactive ring A-reduced metabolites of progesterone in human plasma during pregnancy: elevated levels of 5alpha-dihydroprogesterone in depressed patients during the latter half of pregnancy. *J Clin Endocrinol Metab* 2001; 86(12):5981-5987.
- [117] Quinkler M, Johanssen S, Bumke-Vogt C, Oelkers W, Bähr V, Diederich S. Enzyme-mediated protection of the mineralocorticoid receptor against progesterone in the human kidney. *Mol Cell Endocrinol* 2001; 171:21-24.
- [118] Quinkler M, Johanssen S, Grossmann C, Bähr V, Müller M, Oelkers W, Diederich S. Progesterone metabolism in the human kidney and inhibition of 11beta-hydroxysteroid dehydrogenase type 2 by progesterone and its metabolites. *J Clin Endocrinol Metab* 1999; 84:4165-4171.
- [119] Rafestin-Oblin ME, Lombes M, Michiel JB, Michaud A, Claire M. Mineralocorticoid receptors in the epithelial cells of human colon and ileum. *J Steroid Biochem*. 1984; 20:311-315.
- [120] Ramirez VD, Dluzen D. Is progesterone a pre-hormone in the CNS? *J Steroid Biochem* 1987;27:589-598.
- [121] Rogerson FM, Dimopoulos N, Sluka P, Chu S, Curtis AJ, Fuller PJ. Structural determinants of aldosterone binding selectivity in the mineralocorticoid receptor. *J Biol Chem* 1999; 274:36305-36311.
- [122] Rogerson FM, Fuller PJ. Mineralocorticoid action. *Steroids* 2000; 65:61-73.
- [123] Rosenthal HE, Slaunwhite WR Jr, Sandberg AA. Transcortin: a corticosteroid-binding protein of plasma. X. Cortisol and progesterone interplay and unbound levels of these steroids in pregnancy. *J Clin Endocrinol Metab* 1969; 29:352-367.
- [124] Rundle SE, Smith AI, Stockman D, Funder JW. Immunocytochemical demonstration of mineralocorticoid receptors in rat and human kidney. *J Steroid Biochem* 1989; 33:1235-1242.
- [125] Rupperecht R, Arriza JL, Spengler D, Reul JM, Evans RM, Holsboer F, Damm K. Transactivation and synergistic properties of the mineralocorticoid receptor: relationship to the glucocorticoid receptor. *Mol Endocrinol* 1993; 7:597-603.
- [126] Rupperecht R, Reul JM, van Steensel B, Spengler D, Soder M, Berning B, Holsboer F, Damm K. Pharmacological and functional characterization of human mineralocorticoid and glucocorticoid receptor ligands. *Eur J Pharmacol* 1993; 247:145-154.

- [127] Saitoh H, Hirato K, Yanaihara T, Nakayama T. A study of 5 alpha-reductase in human fetal brain. *Endocrinol Jpn* 1982; 29:461-467.
- [128] Sano Y, Okinaga S, Arai K. Metabolism of [14C]-C21 steroids in the human fetal ovaries. *J Steroid Biochem* 1982; 16:721-724.
- [129] Sasano H, Frost AR, Saitoh R, Matsunaga G, Nagura H, Krozowski ZS, Silverberg SG. Localization of mineralocorticoid receptor and 11 beta-hydroxysteroid dehydrogenase type II in human breast and its disorders. *Anticancer Res* 1997; 17:2001-2007.
- [130] Sasano H, Fukushima K, Sasaki I, Matsuno S, Nagura H, Krozowski ZS. Immunolocalization of mineralocorticoid receptor in human kidney, pancreas, salivary, mammary and sweat glands: a light and electron microscopic immunohistochemical study. *J Endocrinol* 1992; 132:305-310.
- [131] Scatchard G. The attractions of proteins for small molecules and ions. *Ann N Y Acad Sci* 1948; 51:664-672.
- [132] Schmidt RF, Thews G(Hrsg.): *Physiologie des Menschen*. 27. Auflage. Springer-Verlag 1997, Berlin. Seite 748
- [133] Seckl JR, Dickson KL, Yates C, Fink G. Distribution of glucocorticoid and mineralocorticoid receptor messenger RNA expression in human postmortem hippocampus. *Brain Res* 1991; 561:332-337.
- [134] Sekihara H, Island DP, Liddle GW. New mineralocorticoids: 5alpha-dihydroaldosterone and 5alpha-dihydro-11-deoxycorticosterone. *Endocrinology* 1978; 103:1450-1452.
- [135] Selye H. Anesthetic effect of steroid hormones. *Proc Soc Exp Biol Med* 1941; 46:116
- [136] Shimkets RA, Lifton R, Canessa CM. In vivo phosphorylation of the epithelial sodium channel. *Proc Natl Acad Sci U S A* 1998; 95:3301-3305.
- [137] Soucy P, Luu-The V. Conversion of pregnenolone to DHEA by human 17alpha-hydroxylase/17, 20-lyase (P450c17). Evidence that DHEA is produced from the released intermediate, 17alpha-hydroxypregnenolone. *Eur J Biochem* 2000; 267(11):3243-7.
- [138] Souque A, Fagart J, Couette B, Davioud E, Sobrio F, Marquet A, Rafestin-Oblin ME. The mineralocorticoid activity of progesterone derivatives depends on the nature of the C18 substituent. *Endocrinology* 1995; 136:5651-5658.
- [139] Staub O, Yeager H, Plant PJ, Kim H, Ernst SA, Rotin D. Immunolocalization of the ubiquitin-protein ligase Nedd4 in tissues expressing the epithelial Na⁺ channel (ENaC). *Am J Physiol* 1997; 272:C1871-C1880.

- [140] Stewart PM, Murry BA, Mason JJ. Human kidney 11 beta-hydroxysteroid dehydrogenase is a high affinity nicotinamide adenine dinucleotide-dependent enzyme and differs from the cloned type I isoform. *J Clin Endocrinol Metab* 1994; 79:480-484.
- [141] Stockand JD. New ideas about aldosterone signaling in epithelia. *Am J Physiol Renal Physiol* 2002; 282(4):F559-F576.
- [142] Stokes J, Noble J, Brett L, Phillips C, Seckl JR, O'Brien C, Andrew R. Distribution of glucocorticoid and mineralocorticoid receptors and 11beta-hydroxysteroid dehydrogenases in human and rat ocular tissues. *Invest Ophthalmol Vis Sci* 2000;41:1629-1638.
- [143] Swinney DC. Progesterone metabolism in hepatic microsomes. Effect of the cytochrome P-450 inhibitor, ketoconazole, and the NADPH 5 alpha-reductase inhibitor, 4-MA, upon the metabolic profile in human, monkey, dog, and rat. *Drug Metab Dispos* 1990; 18:859-65.
- [144] Technical Bulletin TB239. Promega GmbH, Mannheim, Deutschland
- [145] Trapp T, Holsboer F. Ligand-induced conformational changes in the mineralocorticoid receptor analysed by protease mapping. *Biochem Biophys Res Commun* 1995; 215:286-291.
- [146] Tsai MJ, O'Malley BW. Molecular mechanisms of action of steroid/thyroid receptor superfamily members. *Annu Rev Biochem* 1994; 63:451-486.
- [147] Verrey F. Early aldosterone effects. *Exp Nephrol* 1998; 6:294-301.
- [148] Vilee CA, Loring JM. 20alpha-hydroxysteroid dehydrogenase and 3alpha-hydroxysteroid dehydrogenase in human fetal liver. *Endocrinology* 1963; 72:824-828
- [149] Voice MW, Seckl JR, Edwards CR, Chapman KE. 11 beta-hydroxysteroid dehydrogenase type 1 expression in 2S FAZA hepatoma cells is hormonally regulated: a model system for the study of hepatic glucocorticoid metabolism. *Biochem J* 1996; 317:621-625.
- [150] Wambach G, Higgins JR, Kem DC, Kaufmann W. Interaction of synthetic progestagens with renal mineralocorticoid receptors. *Acta Endocrinol (Copenh)* 1979; 92:560-567.
- [151] Wambach G, Higgins JR. Antimineralocorticoid action of progesterone in the rat: correlation of the effect on electrolyte excretion and interaction with renal mineralocorticoid receptors. *Endocrinology* 1978; 102:1686-1693.
- [152] Warnock DG. Liddle syndrome: an autosomal dominant form of human hypertension. *Kidney Int* 1998; 53:18-24.

- [153] Watzka M, Beyenburg S, Blumcke I, Elger CE, Bidlingmaier F, Stoffel-Wagner B. Expression of mineralocorticoid and glucocorticoid receptor mRNA in the human hippocampus. *Neurosci Lett* 2000; 290:121-124.
- [154] White PC, Speiser PW. Congenital adrenal hyperplasia due to 21-hydroxylase deficiency. *Endocr Rev* 2000; 21:245-291.
- [155] Whorwood CB, Mason JI, Ricketts ML, Howie AJ, Stewart PM. Detection of human 11 beta-hydroxysteroid dehydrogenase isoforms using reverse-transcriptase-polymerase chain reaction and localization of the type 2 isoform to renal collecting ducts. *Mol Cell Endocrinol* 1995; 110:R7-R12.
- [156] Wilson JD, Gloyna RE. The intranuclear metabolism of testosterone in the accessory organs of reproduction. *Recent Prog Horm Res* 1970; 26:309-336.
- [157] Wilson R, Bird CE, Wiqvist N, Solomon S, Diczfalusy E. Metabolism of progesterone by the perfused adrenalectomized human fetus. *J Clin Endocrinol Metab* 1966; 26:1155-1159.
- [158] Winkel CA, Casey ML, Simpson ER, MacDonald PC. Deoxycorticosterone biosynthesis from progesterone in kidney tissue of the human fetus. *J Clin Endocrinol Metab* 1981; 53(1):10-15.
- [159] Winkel CA, Milewich L, Parker CR Jr, Gant NF, Simpson ER, MacDonald PC. Conversion of plasma progesterone to deoxycorticosterone in men, nonpregnant and pregnant women, and adrenalectomized subjects. *J Clin Invest* 1980; 66:803-812.
- [160] Winkel CA, Parker CR Jr, Simpson ER, MacDonald PC. Production rate of deoxycorticosterone in women during the follicular and luteal phases of the ovarian cycle: the role of extraadrenal 21-hydroxylation of circulating progesterone in deoxycorticosterone production. *J Clin Endocrinol Metab* 1980; 51:1354-1358.
- [161] Winkel CA, Simpson ER, Milewich L, MacDonald PC. Deoxycorticosterone biosynthesis in human kidney: potential for formation of a potent mineralocorticosteroid in its site of action. *Proc Natl Acad Sci U S A* 1980; 77(12):7069-7073.
- [162] Wudy SA, Hartmann M, Homoki J. Hormonal diagnosis of 21-hydroxylase deficiency in plasma and urine of neonates using benchtop gas chromatography-mass spectrometry. *J Endocrinol* 2000; 165:679-683.
- [163] Wurtz JM, Bourguet W, Renaud JP, Vivat V, Chambon P, Moras D, Gronemeyer H. A canonical structure for the ligand-binding domain of nuclear receptors. *Nat Struct Biol* 1996; 3:87-94.

- [164] Zanisi M, Martini L. Effects on gonadotropin secretion of 5-alpha-pregnane-3 alpha, 20 alpha-diol, a physiological progesterone derivative. Proc Soc Exp Biol Med 1979; 161:66-69.