

## 1 Literatur

BOLL, M., MARKOVICH, D., WEBER, W.-M., KORTE, H., DANIEL, H. und MURER, H. (1994)

Expression cloning of a cDNA from rabbit small intestine related to proton-coupled transport of peptides,  $\beta$ -lactam antibiotics and ACE-inhibitors  
Pflügers Arch. 429, 146-149

BOLL, M., HERGET, M., WAGENER, M., WEBER, W.M., MARKOVICH, D., BIBER, J., CLAUSS, W., MURER, H. und DANIEL, H. (1996)

Expression cloning and functional characterisation of the kidney cortex high-affinity proton coupled peptide transporter  
Proc. Natl. Acad. Sci. USA 93, 284-289

BRODERICK, G. und CRAIG, W. M. (1989)

Metabolism of peptides and amino acids during in vitro protein degradation by mixed rumen organisms.  
J. Dairy Sci. 72, 2540- 2548

BRODERICK, G.A. und WALLACE, R.J. (1988)

Effects of nitrogen dietary source on concentrations of ammonia, free amino acids and fluorescamine-reactive peptides in the sheep rumen.  
J. Anim. Sci. 66, 2233-2238

BRODERICK, G.A., WALLACE, R.J., ÖRSKOV, E.R. (1991)

Control of rate and extend of protein degradation  
Physiological aspects of digestion and metabolism in ruminants:  
Proceedings of the seventh international symposium in ruminant physiology  
Academic Press, Inc.

CHEESEMAN, C.I., (1980)

The transport of dipeptides by the small intestine.  
Can. J. Physiol. Pharmacol. 58, 1326-1333

CHEN ,G., RUSSEL, J.B. und SNIFFEN, C.J. (1987)

A procedure for measuring peptides in rumen fluid and evidence that peptide uptake can be the rate limiting step in ruminal protein degradation.  
J. Dairy Sci. 70, 1211-1219

CHEN, G. und RUSSEL, J.B. (1991)

Effect of monensin and a protonophore on protein degradation, peptide accumulation and deamination by mixed ruminal microorganisms in vitro.

J. Anim. Sci. 69, 2196 - 2203

CHEN, G., SNIFFEN, C.G. und RUSSEL, B. ( 1987)

Concentration and estimated flow of peptides from the rumen of dairy cattle : Effects of protein quantity, protein solubility, and feeding frequency.

J. Dairy Sci. 70, 983 – 992

DANIEL, H. (1996)

Topical Review: Function and molecular structure of brush border membrane peptide/H<sup>+</sup> Symporters

J. Membr. Biol. 154, 197 – 203

DANIEL, H. und HERGET, M.

Transport of Amino-Acids and Peptides in intestine and kidney: from humans and animals to single molecules.

Institute of Nutritional Sciences, Justus- Liebig- Universität Giessen. Germany

DANIEL, H. und HERGET, M., (1997)

Invited Review: Cellular and molecular mechanisms of renal peptide transport  
Am. Physiol. Society, 0363-6127/97

DANILSON, D.A., WEBB, K.E.Jr. und HERBEIN, J.H.(1987)

Transport and hindlimb exchange of peptide and serum protein amino acids in calves fed soy-or urea-based purified diets

J. Anim. Sci. 64, 1852-1857

DIRKSEN, G., LIEBICH, H.G., BROSI, G., HAGEMEISTER, H., MAYER, E. (1984)

Morphologie der Pansenschleimhaut und Fettsäurereresorption beim Rind- bedeutende Faktoren für Gesundheit und Leistung.

Zbl. Vet. Med. A, 414-430

Verlag Paul Parey, Berlin und Hamburg

ISSN 0721-0981 / Inter Code: ZVRAAX

DOREAU, M., FERCHAL und E., BECKERS, Y. (1997)

Effects of level of intake and of available volatile fatty acids on the absorptive capacity of sheep rumen.

Small Ruminant Research 25, 99-105

DYER, J., BEECHY, R.B., GORVEL, J.P., SMITH, R.T., WOOTON, R. und SHIRAZI-BEECHEY, S.P. (1990)

Glycyl-L-proline transport in rabbit enterocyte basolateral-membrane vesicles.

Biochem. J. 269, 565-571

FEI, Y.J., KANAI, Y., NUSSBERGER, S., GANAPATHY, V., LEIBACH, F.H., ROMERO, M.F., SINGH, S., BORON, W.F. und HEDIGER, M.A. (1994)

Expression cloning of a mammalian proton-coupled oligopeptide transporter  
Nature 386, 563-566

GANAPATHY, V., MENDICINO, J.F. und LEIBACH, F.H (1981)

Transport of Glycyl-Lproline into intestinal and renal brush border vesicles from rabbit.  
J. Biol. Chem. 256, No1, pp.118-124

GANAPATHY, V., RHADHAKRISHNAN, A.N. (1980)

Sodium-dependent inhibition of amino acid and dipeptide transport by harmaline in  
monkey small intestine.  
Biochem. Pharmacol. 29, 713-716

HARRISON, F.A., KEYNES, R.D., RANKIN, J.C. und ZURICH, L. (1975)

The effect of ouabain on ion transport across isolated sheep rumen epithelium.  
J. Physiol. 249, 669-677

KOELN, L.L., SCHLAGHECK, T.G. und WEBB, K.E. Jr. (1993)

Amino acid flux across the gastrointestinal tract and liver of calves  
J. Dairy Sci. 76, 2275-2285.

KURKOWSKI, S. und MARTENS, H., (1998)

Auswirkungen eines erhöhten osmotischen Druckes auf die Gewebeleitfähigkeit und den  
Na- Transport des isolierten Pansenepithels von Schafen.  
Proc. Soc. Nutr. Physiol. 7, 67

LANG, I., (1997)

In-vitroUntersuchungen zur Charakterisierung eines spannungsabhängigen Kationenkanals  
in der luminalen Membran des Pansenepithels beim Schaf.  
Diss., Berlin, Freie Universität

LEBZIEN P., VOIGT, J., GABEL, M. und GÄDEKEN, D. (1996)

Zur Schätzung der Menge an nutzbarem Rohprotein am Duodenum von Milchkühen.  
J. Anim. Physiol. A. Anim. Nutr. 76: 218-223

LEONHARD-MAREK, S. und MARTENS, H. (1996)

Effects of potassium on magnesium transport across rumen epithelium.  
Am. J. Physiol. G1034-G1038

MARTENS, H., GÄBEL, G., und STROZYK, B. (1991)

Mechanism of electrically silent Na and Cl transport across the rumen epithelium of sheep.  
Exp. Physiol. 76, 103-114

MARTENS, H., HARMEYER, J.und TEINKHAO, M.(1974)

Modellversuch über die Passage von Aminosäuren durch die Pansenwand in vivo

MARTENS, H. und RAYSSIGUIER, Y., (1980)

Magnesium metabolism and hypomagnesaemia.  
In: Digestive Physiologie and metabolism in Ruminants  
Clermont-Ferrand, 447-466

MATTHEWS, D.M. und PAYNE, J.W. (1980)

Transmembrane transport of small peptides.  
Curr. Top. Membr. Transp. 14, 331-425

MATTHEWS, D.M. (1975)

Intestinal absorption of peptides.  
Physiol. Rev. 55, 537-608

MATTHEWS, J.C. und WEBB, K.E., Jr.(1995):

Absorption of L-Carnosine, L-Methionine, and L-Methionylglycine by isolated Sheep ruminal and omasal epithelial tissue  
J. Anim. Sci. 73, 3464-3475

MATTHEWS, J.C., WONG, E.A., BENDER, P.K., BLOOMQUIST, J.R. und WEBB, K.E.,Jr. (1996)

Demonstration and characterization of dipeptide transport system activity in sheep omasal epithelium by expression of mRNA in *Xenopus laevis* Oocytes  
J. Anim. Sci. 74, 1720-1727

MC CORMICK, M.E. und WEBB, K.E.Jr. (1982)

Plasmafree, erythrocytefree and plasma peptide aminoacid exchange of calves in steady state and fasting metabolism  
J. Nutr. 112, No.2

MC COLLUM, M.Q. und WEBB, K.E. (1998)

Glycyl-L-Sarcosine absorption across ovine omasal epithelium during coincubation with other peptide substrates and volatile fatty acids  
J. Anim. Sci. 76, 2706-2711

MEREDITH, D. und BOYD, C.A.R. (1995)

Topical Review: Oligopeptide transport by epithelial cells  
J. Membr. Biol. 145, 1-12

PAN, Y.X., WONG, E.A., BLOOMQUIST, J.R., WEBB, K.E.Jr. (1997)

Poly(A)<sup>+</sup> RNA from sheep omasal epithelium induces expression of a peptide transport protein in *Xenopus laevis* oocytes  
J. Anim. Sci. 75, 3323-3330

POWELL, D.W. (1981)

Barrier function of epithelia  
Am. J. Physiol. 241, G275-G288.

RUBINO, A., FIELD, M., und SHWACHMAN, H. (1971)

Intestinal transport of amino acid residues of dipeptides.  
J. Biol. Chem. 246, No.11, 3542-3548

RUSSEL, J.B., SNIFFEN, C.J. und VAN SOEST, P.J. (1983)

Effect of carbohydrate limitation on degradation and utilization of casein by mixed rumen bacteria  
J. Dairy Sci. 66, 763-775

SCHEFFLER, A. (1984)

Charakterisierung der Wirkung von Pharmaka auf die elektrophysiologischen Parameter gastrointestinaler Epithelgewebe mit einer computergesteuerten Strom und Spannungsklemmeneinrichtung.  
Diss., Achen, Med. Fak. der TU

SCHULTHEISS, G. (1995)

Calcium-sensitiver, elektogener Natriumtransport des Psalterepithels von Schafen.  
Diss., Berlin, Freie Universität

SEAL, C.J. und PARKER, D.S. (1996)

Effect of intraruminal propionic acid infusion on metabolism of mesenteric- and portal-drained viscera in growing steers fed a forage diet :II. ammonia, urea, amino acids, and peptides  
J. Anim. Sci. 74, 245-256

STEVENS, C.E., STETTLER, B.K. (1965)

Factors affecting the transport of volatile fatty acids across rumen epithelium.  
Am. J. Physiol. 210 (2), 365-372

STEWART, A.K., BOYD, C.A.R. und VAUGHAN-JONES, R.D. (1999)

A novel role for carbonic anhydrase: cytoplasmic pH gradient dissipation in mouse small intestinal enterocytes  
J. Physiol. 516 (1) 209-217

TERADA, T., SAWADA, K., SAITO, H., HASHIMOTO, Y., INUI, K. (1999)

Funktional characteristics of basolateral peptide transporter in human intestinal cell line Caco-2.  
Am. Phys. Society 0193-1857/99, G1435-G1441

THAMOTHARAN, M., BAWANI, S., ZHOU, X. und ADIBI, S.A.(1999)

Hormonal regulation of oligopeptide transporter Pept-1 in a human intestinal cell line.  
Am. J. Physiol. C821-C626

THWAITES, D.T., BROWN, C.D.A., HIRST, B.H., SIMMONS, N.L. (1993)

H<sup>+</sup>-coupled dipeptide (glycylsarcosine) transport across apical and basal borders of human intestinal Caco-2 cell monolayers display distinctive characteristics.  
Biochim. Biophys. Acta 1151, 237-245

USSING, H.H. (1949)

The active ion transport through the isolated frog skin in the light of tracer studies  
Acta physiol. Scand. 17, 1-37

WALKER, D., THWAITES, D.T., SIMMONS, N.L., GILBERT, H.J., HIRST, B.H. (1998)

Substrate upregulation of the human small intestinal peptide transporter, hPepT1.  
J. Physiol. 507 (3), 697-706

WALLACE, R.J., NEWBOLD, C.J., WATT, N.D., BUCHANAN, V. und BROWN, D.S. (1993)

Amino acid composition of degradation-resistant peptides in extracellular rumen fluid of sheep.  
J. Agric. Sci. 120, 129 - 133

WEBB, K.E. JR., DI RIENZO, B.D. und MATTHEWS, J.C. (1993)

Recent Developments in Gastrointestinal Absorption and Tissue Utilization of Peptides: A Review  
J. Dairy Sci. 76, 351-361

WEBB, K.E. JR., MATTHEWS, J.C. und DI RIENZO, B.D., (1992)

Peptide absorption : A review of current concepts and future perspectives  
J. Anim. Sci. 70, 3248-3257

WOLFFRAM, S., GRENAKER, B. und SCHARRER, E. (1998)

H<sup>+</sup>-coupled uphill transport of the dipeptide Glycyl-Sarcosine by bovine intestinal brush-border membrane vesicles  
J. Dairy Sci. 81, 2595-2603

