

Literaturverzeichnis

- ¹ Arnett FC, Edworthy SM, Bloch DA, McShane DJ, Fries JF, Cooper NS, Healey LA, Kaplan SR, Liang MH, Luthra HS, Medsger Jr TA, Mitchell DM, Neustadt DH, Pinals RS, Schaller JG, Sharp JT, Wilder RL, Hunder GG. The American Rheumatism Association 1987 revised criteria for the classification of rheumatoid arthritis. *Arthritis Rheum* 1988; 31: 315-24
- ² Bläß St, Specker Ch, Lakomek HJ, Schneider EM, Schwochau M. Novel 68k autoantigen detected by rheumatoid arthritis specific antibodies. *Ann Rheum Dis* 1995; 54: 355-60
- ³ Wernick R, Merryman P, Jaffe I, Ziff M. IgG and IgM rheumatoid factor in rheumatoid arthritis. *Arthritis Rheum* 1983; 26: 593
- ⁴ Jones MG, Shipley ME, Hearn JP, Hay FC. Specificity of rheumatoid factors in relation to the disease state in rheumatoid arthritis. *Arthritis Rheum* 1990; 49: 757-62
- ⁵ Vaughan JH. Pathogenetic concepts and origins of rheumatoid factors in rheumatoid arthritis. *Arthritis Rheum* 1993; 36: 1-6
- ⁶ Klareskog L, Forsum U, Kabelitz D, Ploen L, Sundstrom C, Nilsson K, Wigren A, Wigzell H. Immune functions of human synovial cells. Phenotypic and T cell regulatory properties of macrophages-line cells that express HLA-DR. *Arthritis Rheum* 1982; 25: 488
- ⁷ Naujoks C. Die volkswirtschaftlichen Kosten von Rheuma in der Bundesrepublik Deutschland. Merkblätter Deutsche Rheuma-Liga 1990 Nr.6.7
- ⁸ Jones M, Symmons D, Finn J, Wolfe F. Does exposure to immunosuppressive therapy increase the 10 year malignancy and mortality risks in rheumatoid arthritis? A matched cohort study. *Br J Rheumatol* 1996; 35: 738-45
- ⁹ Goronzy J, Weyand CM, Fathman CG. Shared T cell recognition sites on human histocompatibility leukocyte antigen class II molecules of patients with seropositive rheumatoid arthritis. *J Clin Invest* 1986; 77: 1042-9
- ¹⁰ Nelson JL, Mickelson E, Masewicz S, Barrington R, Dugowson C, Koepsell T, Hansen JA. Dw14 (DRB1*0404) is a Dw4-dependent risk factor for rheumatoid arthritis: Rethinking the "shared epitope" hypothesis. *Tissue Antigens* 1991; 38: 145-51
- ¹¹ Wraith DC, McDevitt HO, Steinman L, Acha-Orbea H. T cell recognition as the target for immune intervention in autoimmune disease. *Cell* 1989; 57: 709-15
- ¹² Lang B, Melchers I, Peter H-H. Das HLA-System und die rheumatoide Arthritis. *Z Rheumatol* 1990; 49: 197-207
- ¹³ Bell JI, Denney JrD, Foster L, Belt T, Todd JA, McDevitt HO. Allelic variation in DR subregion of the human major histocompatibility complex. *Proc Natl Acad Sci USA* 1987; 84: 6234-8
- ¹⁴ Bell JI, Todd JA, McDevitt HO. The molecular basis of HLA-disease association. In Harris H, HirschhornK. Advances in human genetics. Plenum Press New York 1989; 1-4
- ¹⁵ Lanchbury JSS, Sakkas LI, Stephens HAF, Welsch KI, Panay GS. Susceptibility to rheumatoid arthritis is associated with specific HLA-DR4B1 gene sequences. *Arthritis Rheum* 1989; 32 (Suppl.): 39
- ¹⁶ Weyand CM, Goronzy JJ. Function of the shared third hypervariable region (HVR3) of the HLA-DR 1-chain in rheumatoid arthritis-associated alleles; lacking evidence for crossreactive antigen-specific T cell responses. *Arthritis Rheum* 1989; 32 (Suppl.): 40

-
- ¹⁷ Gregersen PK., Silver J., Winchester RJ. The shared epitope hypothesis. An approach to understanding the molecular genetics of susceptibility to rheumatoid arthritis. *Arthritis Rheum* 1987; 30: 1205-13
- ¹⁸ Hall FC, Weeks DE, Camilleri JP, Williams LA, Amos N, Darke C, Gibson K, Pile K, Wordsworth BP, Jessop JD. Influence of the HLA-DRB1 locus on susceptibility and severity in rheumatoid arthritis. *Qjm* 1996; 89: 821-9
- ¹⁹ Boki KA, Panayi GS, Vaughan RW, Dross AA, Moutsopoulos HM, Lanchbury JS. HLA class II sequence polymorphisms and susceptibility to rheumatoid arthritis in Greeks. *Arthritis Rheum* 1992; 35: 749-55
- ²⁰ Smolen JS, Steiner GS. Are autoantibodies active players or epiphenomena? *Curr Opin Rheum* 1998; 18: 201-206
- ²¹ Arnett FC, Edworthy SM, Bloch DA, McShane DJ, Fries JF, Cooper NS, Healey LA, Kaplan SR, Liang MH, Luthra HS, Medsger TA jr, Mitchell DM, Neustadt DH, Pinals RS, Schaller JG, Sharp JT, Wilder RL, Hunder GG. The American Rheumatism Association 1987 revised criteria for the classification of rheumatoid arthritis. *Arthritis Rheum* 1988; 31: 315-24
- ²² Bläß S, Engel JM, Burmester GR. Der Immunologische Homunculus der RA. *Z Rheumatol* 2001; 60: 1-16
- ²³ Weyand CM, Goronzy JJ. HLA polymorphisms and T cells in rheumatoid arthritis. *Int Rev Immunol* 1999; 18: 37-59
- ²⁴ Parry SL, Hall FC, Olson J, Kamradt T, Sonderstrup G. Autoreactivity versus autoaggression: a different perspective on human autoantigens. *Curr Opin Immunol* 1998; 10: 663-8
- ²⁵ Schmidt D, Goronzy JJ, Weyand CM. CD4+CD7-CD28- T cells are expanded in rheumatoid arthritis and are characterized by autoreactivity. *J Clin Invest* 1996; 97: 2027-37
- ²⁶ Cohen IR, Young DB. Autoimmunity, microbial immunity and the immunological homunculus. *Immunol Today* 1991; 12: 105-10
- ²⁷ Bläß S, Engel JM, Burmester GR. The immunologic homunculus in rheumatoid arthritis. *Arth. Rheum.* 1999; 42(12): 2499-2506
- ²⁸ Goff Le P, Saraux A, Youinou P. New autoantibodies in rheumatoid arthritis. *Rev Rheum [Engl. Ed.]* 1997; 64: 638-644
- ²⁹ Saraux A, Valls I, Voisin V, Koreichi A, Baron D, Youinou P, Le Goff P. How useful are tests for rheumatoid factors, antiperinuclear factors, antikeratin antibody and the HLA-DR4 antigen for the diagnosis of rheumatoid arthritis? *Rev Rheum [Eng. Ed.]* 1995; 62: 16-20
- ³⁰ Nogueira L, Sebbag M, Vincent C, Arnaud M, Fournie B, Cantagrel A, Jolivet M, Serre G. Performance of two ELISA for antifilaggrin autoantibodies, using either affinity purified or deiminated recombinant human filaggrin, in the diagnosis of rheumatoid arthritis. *Ann Rheum Dis* 2001; 60: 882-887
- ³¹ Bizzaro N, Mazzanti G, Tonutti E, Villalta D, Tozzoli R. Diagnostic accuracy of the anti-citrulline antibody assay for rheumatoid arthritis. *Clin Chem* 2001; 1089-1093
- ³² Goldbach-Mansky R, Lee J, McCoy A, Hoxworth J, Yarboro C, Smolen JS, Steiner G, Rosen A, Zhang C, Menard HA, Zhou ZJ, Palosuo T, Van Venrooij WJ, Wilder RL, Klippel JH, Schumacher HR Jr, El-Gabalawy HS. Rheumatoid arthritis associated autoantibodies in patients with synovitis of recent onset. *Arthritis Res* 2000; 2: 236-243
- ³³ Feldmann M, Brennan FM, Field M, Maini RN. Pathogenesis of rheumatoid arthritis. Cellular and cytokine interactions. In Smolen JS, Kalden JR, Maini RN. *Rheumatoid Arthritis - recent research advances*. Springer-Verlag 1992; 41-54

-
- ³⁴ Salazar-Onfray F, Petersson M, Franksson L, Matsuda M, Blankenstein T, Karre K, Kiessling R. IL-10 converts mouse lymphoma cells to a CTL-resistant, NK-sensitive phenotype with low but peptide-inducible MHC class I expression. *J Immunol* 1995; 154: 6291-8
- ³⁵ Klimiuk PA, Goronzy JJ, Bjornsson J, Beckenbaugh RD, Weyand CM. Tissue cytokine patterns distinguish variants of rheumatoid synovitis. *Am J Pathol* 1997; 151 (5): 1311-9
- ³⁶ Potocnik AJ, Kinne R, Menninger H, Zacher J, Emmrich F, Krocze RA. Expression of activation antigens on T cells in rheumatoid arthritis patients. *Scand J Immunol* 1990; 31: 213-24
- ³⁷ Kohem CL, Brezinschek RI, Wisbey H, Tortorella C, Lipsky PE, Oppenheimer-Marks N. Enrichment of differentiated CD45RBdim, CD27-memory T cells in the peripheral blood, synovial fluid and synovial tissue of patients with rheumatoid arthritis. *Arthritis Rheum* 1996; 39: 844-54
- ³⁸ Ali M, Ponchel F, Wilson K-E, Francis M-J, Wu X, Verhoef A, Boylston A-W, Veale D-J, Emery P, Markham A-F, Lamb J-R, Isaacs J-D. Rheumatoid arthritis synovial T cells regulate transcription of several genes associated with antigen-induced anergy. *J Clin Invest*. 2001; 107(4): 519-28
- ³⁹ Carson DA. Unconventional T-cell activation by IL-15 in rheumatoid arthritis. *Nat Med*. 1997; 3 (2): 148-9
- ⁴⁰ Shimizu Y, Newman W, Tanaka Y, Shaw S. Lymphocyte interaction with endothelial cells. *Immunol Today* 1992; 13: 106
- ⁴¹ Hain N, Stuhlmüller B, Hahn GR, Kalden JR, Deutzmann R, Burmester GR. Characterisation and microsequencing of a 205k synovial protein stimulatory for T cells and reactive with rheumatoid factor containing sera. *J Immunol* 1996; 157: 1773-80
- ⁴² Bas S, Perneger TV, Mikhnevitch E, Seitz M, Tiercy, J-M, Roux-Lombard P, Guerne PA. Association of rheumatoid factors and anti.filaggrin antibodies with severity of erosions in rheumatoid arthritis. *Rheumatology* 2000; 39: 1082-1088
- ⁴³ Lapoint E, Dery U, Vaillancourt F, Menard HA, Sherbrooke TS. Rheumatoid sera potentially recognize all citrullinated proteins. *Arthritis Rheum* 1999; 42(Suppl): 86
- ⁴⁴ Nienhuis R, Mandema E. A new serum factor in patients with rheumatoid arthritis. The antiperinuclear factor. *Ann Rheum Dis* 1964; 23: 302-305
- ⁴⁵ Mediawake R, Isenberg DA, Schellekens GA, van Venrooij WJ. Use of anti-citrullinated peptide and anti-RA33 antibodies in distinguishing erosive arthritis in patients with systemic lupus erythematosus and rheumatoid arthritis. *Ann-Rheum-Dis*. 2001; 60 (1): 67-8
- ⁴⁶ Smolen JS, Steiner G. Are autoantibodies active players or epiphrenomena? *Current Opinion in Rheumatology* 1998; 10: 201-206
- ⁴⁷ Bas S, Perneger TV, Seitz M, Tiercy JM, Roux-Lombard P, Guerne PA. Diagnostic tests for rheumatoid arthritis: comparison of anti-cyclic citrullinated peptide antibodies, anti-keratin antibodies and IgM rheumatoid factors. *Rheumatology* 2002; 41 (7): 809-14
- ⁴⁸ Bizzaro N, Mazzanti G, Tonutti E, Villalta D, Tozzoli R. Diagnostic accuracy of the anti-citrulline antibody assay for rheumatoid arthritis. *Clin Chem* 2001; 1089-1093
- ⁴⁹ Nogueira L, Sebbag M, Vincent C, Arnaud M, Fournie B, Cantagrel A, Jolivet M, Serre G. Performance of two ELISA for antifilaggrin autoantibodies, using either affinity purified or deiminated recombinant human filaggrin, in the diagnosis of rheumatoid arthritis. *Ann Rheum Dis* 2001; 60: 882-887
- ⁵⁰ Schellekens GA, de Jong BA, van den Hoogen FH, van de Putte LB, van Venrooij WJ. Citrulline is an essential constituent of antigenic determinants recognized by rheumatoid arthritis-specific autoantibodies. *J Clin Invest* 1998; 101: 273-281

-
- ⁵¹ Jethmalani SM, Henle KJ, Kaushal GP. Heat shock-induced prompt glycosylation. Identification of P-SG67 as calreticulin. *J Biol Chem* 1994; 269(38): 23603-23609
- ⁵² Llewellyn DH, Johnson S, Eggleton P. Calreticulin comes of age. *Trends Cell Biol* 2000; 10: 399-402
- ⁵³ Meldolesi J, Krause K-H, Michalak M. Calreticulin: how many functions in how many cellular components? *Cell Calcium* 1996; 20: 83-86
- ⁵⁴ Nauseef WM, Mc Cormick SJ, Clark RA. Calreticulin functions as a molecular chaperone in the biosynthesis of myeloperoxidase. *J Biol Chem* 1995; 270(9): 4741-4747
- ⁵⁵ Eggleton P, Ward FJ, Johnson J et al. Fine specificity of autoantibodies to calreticulin: epitope mapping and characterization. *Clin Exp Immunol* 2000; 120: 381-391
- ⁵⁶ Verreck FA, Elferink D, Vermeulen CJ, Amons R, Breedveld F, de Vries RR, Koning F. DR4Dw4/DR53 molecules contain a peptide from the autoantigen calreticulin. *Tissue Antigens* 1995; 45: 270-5
- ⁵⁷ Wang KK, Posmantur R, Nadimpalli R, Nath R, Mohan P, Nixon RA, Talanian RV, Keegan M, Herzog L, Allen H. Caspase-mediated fragmentation of calpain inhibitor protein calpastatin during apoptosis. *Arch-Biochem-Biophys* 1998; 356(2): 187-96
- ⁵⁸ Cong J, Thompson VF, Goll DE. Effect of monoclonal antibodies specific for the 28-kDa submit on catalytic properties of the calpains. *J Biol Chem* 1993; 268: 25740
- ⁵⁹ Lane RD, Allan DM, Mellgren RL. A comparison of the intracellular distribution of mu-calpain, m-calpain, and calpastain in proliferating human A431 cells. *Exper Cell Res* 1992; 203: 5
- ⁶⁰ Geesink GH, Nonneman D, Koohmaraie M. An improved purification protocol for heart and skeletal muscle calpastatin reveals two isoforms resulting from alternative splicing. *Arch-Biochem-Biophys* 1998; 356(1): 19-24
- ⁶¹ Yamamoto S, Shimizu K, Shimizu K, Suzuki K, Nakagawa Y, Yamamuro T. Calcium-dependent cysteine proteinase (calpain) in human arthritic synovial joints. *Arthritis Rheum* 1992; 35: 1309-17
- ⁶² Després N, Talbot G, Plouffe B, Boire G, Ménard HA. Detection and expression of a cDNA clone that encodes a polypeptide containing two inhibitory domains of human calpastatin and its recognition by rheumatoid arthritis sera. *J Clin Invest* 1995; 95: 1891-6
- ⁶³ Mimori T, Suganuma K, Tanami Y, Jojima T, Matsumura M, Fujii T, Yoshizawa T, Suzuki K, Akizuki M. Autoantibodies to calpastatin (an endogenous inhibitor for calcium-dependent neutral protease, calpain) in systemic rheumatic diseases. *Med Sci* 1995; 92: 7267-71
- ⁶⁴ Schlosser U, Lackner KJ, Scheckenhofer C, Spannagl M, Spengel FA, Hahn G, Lang B, Schmitz G. Autoantibodies against the protease inhibitor calpastatin: a new risk factor for venous thrombosis? *Thromb Haemost* 1997; 77: 11-3
- ⁶⁵ Takada R, Matsumoto M, Yosida M, Nojima T, Hirakata M, Ohosone Y, Mimori T. Detection of isotype-specific autoantibodies to calpastatin in sera from patients with rheumatic diseases using heat treated HeLa-cell extract as an antigen source for immunoblotting. *Nihon-Rinsho-Meneki-Gakkai-Kaishi* 1998; 21(4): 150-B
- ⁶⁶ Toth E, Bruin JP, Heinsbroek RP, Joosten RN. Spatial learning and memory in calpastatin-deficient rats. *Neurobiol Learn mem* 1996; 66: 230-235
- ⁶⁷ Despres N, Boire G, Lopez-Longo FJ, Menard HA. The Sa-system: a novel antibody system specific for rheumatoid arthritis. *J Rheumatol* 1994; 21: 1027-1033
- ⁶⁸ Mimori T, Suganuma K, Tanami Y, Nojima T, Matsumura M, Fujii T, Yoshizawa T, Suzuki K, Akizuki M. Autoantibodies to calpastatin (an endogenous inhibitor for calcium dependent neutral protease, calpain) in systemic rheumatic diseases. *Proc Natl Acad Sci* 1995; 92: 7267-7271

-
- ⁶⁹ Munro S, Pelham HRB. A Hsp70 like protein in th ER: identity with the 78-kd glucose-regulated protein and immunoglobulin heavy chain binding protein. *Cell* 1986; 46: 291-300
- ⁷⁰ Blobel G, Dobberstein B. Transfer to proteins across membranes. II. Reconstitution of functional rough microsomes from heterologous components. *J Cell Biol* 1975; 67: 852-862
- ⁷¹ Bläß St, Meier Ch, Vohr HW, Schwochau M, Specker Ch, Burmester GR. The p68 autoantigen characteristic of rheumatoid arthritis is reactive with carbohydrate epitope specific autoantibodies. *Ann Rheum Dis* 1998; 57: 220-25
- ⁷² Baldwin GS, Chandler R, Seet KL, Weinstock J, Grego B, Rubira M, Moritz RL, Simpson RJ. Structural studies on a 75kDa glycoprotein isolated from porcine gastric mucosal membranes: close homology with the 78kDa glucose-regulated family of proteins. *Protein Seq Data Anal* 1987; 1: 7-12
- ⁷³ Chang SC, Wooden SK, Nakaki T, Kim YK, Lin AY, Kung, L, Attenello JW, Lee AS. Rat gene encoding the 78kDa glucose-regulated protein Grp78: its regulatory sequences and the effect of protein glycosylation on its expression. *Proc Natl Acad Sci* 1987; 84: 680-684
- ⁷⁴ Delpino A, Piselli P, Vismara D, Vendetti S, Colizzi V. Cell surface localization of the 78kD glucose regulated protein (Grp78) induced by thapsigargin. *Mol Membr Biol* 1998; 15: 21-26
- ⁷⁵ Bläß S, Schneider SM, Specker C et al. Nover 68 Kda autoantigen detected by rheumatoid arthritis specific antibodies. *Ann Rheum Dis* 1995; 54: 355-60
- ⁷⁶ Bläß S, Haferkamp C, Specker C et al. Rheumatoid arthritis: Autoreactive T-cells directed against a novel 68k autoantigen. *Ann Rheum Dis* 1997; 55: 317-22
- ⁷⁷ Corrigall VM, Bodman-Smith MD, Fife MS, Canas B, Myers LK, Wooley PH, Soh C, Staines SH, Pappin DJC, Berlo SE, van Eden W, van der Zee R, Lanchbury JS., Panayi GS. The Human Endoplasmic Reticulum Molecular Chaperone BiP Is an Autoantigen for Rheumatoid Arthritis and Prevents the Induction of Experimental Arthritis. *J Immunol* 2001; 166: 1492-98
- ⁷⁸ Hassfeld W, Steiner G, Studnicka-Benke A, Skriner K, Graninger W, Fischer I, Smolen JS. Autoimmune response to the splicosome. An immunologic link between rheumatoid arthritis, mixed connective tissue disease and systemic lupus erythematosus. *Arthritis Rheum* 1995; 38: 777-85
- ⁷⁹ Steiner G. The A2 protein of the heterogeneous nuclear ribonucleoprotein (hnRNP-A2)/RA33. In: van Venrooij WJ, Maini RN.:Manual of Biological Markers of Diseases. Kluwer Academic Publishers, Dordrecht, Boston, London (1996)
- ⁸⁰ Meyer O, Tauxe F, Fabregas D et al. Anti-RA33 antinuclear autoantibody in rheumatoid arthritis and mixed connective tissue disease: comparison with antikeratin and antiperinuclear antibodies. *Clin Ex Rheumatol* 1993; 11: 473-478
- ⁸¹ Lal RB, Edison LJ, Chused TM. Fixation and long-term storage of human lymphocytes for surface marker analysis by flow cytometry. *Cytometry* 1988; 9: 213-9
- ⁸² Dolbeare F, Beisker W, Pallavicini MG, Vanderlaan M, Gray JW. Cytochemistry for bromodeoxyuridine/DNA analysis: Stoichiometry and sensitivity. *Cytometry* 1985; 8: 521-30
- ⁸³ Gey GD, Coffman WD, Kubicek MD. Tissue culture studies of the proliferative capacity of cervical carcinoma and normal epithelium. *Cancer Re* 1952; 12: 264-265
- ⁸⁴ Puck TT, Marcus PI, Cieciura SJ. Clonal growth of mammalian cells in vitro. *J Exp Med* 1956; 103: 273-284
- ⁸⁵ Laemmli UK. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 1970; 227: 680-5

-
- ⁸⁶ Dunn MJ, Burghes AHM. High resolution two-dimensional polyacrylamide gel electrophoresis. I. Methodological procedures. *Electrophoresis* 1983; 4: 97-116
- ⁸⁷ Bradford MM. A rapid and sensitive method for the quantization of microgram quantities of protein using the principle of protein-dye-binding. *Anal Biochem* 1976; 72: 248-54
- ⁸⁸ Hain N, Stuhlmüller B, Hahn GR, Kalden JR, Deutzmann R, Burmester GR. Characterisation and microsequencing of a 205k synovial protein stimulatory for T cells and reactive with rheumatoid factor containing sera. *J Immunol* 1996; 157: 1773-80
- ⁸⁹ Priem S, Rittig MG, Kamradt T, Burmester GR, Krause A. An optimized PCR leads to rapid and highly sensitive detection of *Borrelia burgdorferi* in patients with lyme borreliosis. *J Clin Microbiol* 1997; 35: 685-90
- ⁹⁰ Bläß S, Union A, Raymackers J, Schumann F, Ungethüm U, Müller-Steinbach S, De Kayser F, Burmester GR: The stress protein BiP is overexpressed and is a major B- and T-cell target in rheumatoid arthritis, *Arthritis Rheum.* 2001; 44
- ⁹¹ Bläß S, Schumann F, Hain NAK, Engel JM, Stuhlmüller B, Burmester GR. p205 is a major target of autoreactive T cells in rheumatoid arthritis. *Arthritis Rheum.* 1999; 42: 971-980
- ⁹² Panayi GS, Lanchbury JS, Kingsley GH. The importance of the T cell in initiating and maintaining the chronic synovitis in rheumatoid arthritis. *Arthritis Rheum.* 1992; 35: 729-735
- ⁹³ Gans HA, Moldonado Y, Yasukawa LL, Beeler J, Audett S, Rinki MM, DeHovitz R, Arvin AM. IL-12, IFN- and T cell proliferation to measles in immunized infants. *J. Immunol.* 1999; 162: 5569-5575
- ⁹⁴ Dupuis M, Schaefer E, Krause KH, Tschopp J. The calcium binding protein calreticulin is a major constituent of lytic granules in cytolytic T lymphocytes. *J. Exp. Med.* 1993; 177: 1-7
- ⁹⁵ Ali M, Ponchel F, Wilson KE, Francis MJ, Wu X, Verhoef A, Boylston AW, Veale DJ, Emery P, Markham AF, Lamb JR, Isaacs JD. Rheumatoid arthritis synovial T cells regulate transcription of several genes associated with antigen-induced anergy. *J. Clin. Invest.* 2001 Feb; 107(4): 519-28
- ⁹⁶ Kim HJ, Berek C. B cells in rheumatoid arthritis. *Arthritis-research* 2000; 2(2): 126-31
- ⁹⁷ David-Ameline J, Lim A, Davodeau F, Peyrat MA, Berthelot JM, Semana G, Pannetier C, Gaschet J, Vie H, Even J, Bonneville M. Selection of T cells reactive against autologous B lymphoblastoid cells during chronic rheumatoid arthritis. *J Immunol* 1996; 157: 4697-706
- ⁹⁸ Fox DA. The role of T cells in the immunopathogenesis of rheumatoid arthritis. *Arthritis Rheum* 1997; 40: 598-609
- ⁹⁹ Tak PP, van der Lubbe PA, Cauli A, Daha MR, Smeets TJM, Kluij PM, Meinders AE, Yanni G, Panayi GS, Breedveld FC. Reduction of synovial inflammation after anti-CD4 monoclonal antibody treatment in early rheumatoid arthritis. *Arthritis Rheum* 1995; 38: 1457-65
- ¹⁰⁰ Kishore U, Sontheimer RD, Sastry et.al. The systemic lupus erythematosus disease autoantigen-calreticulin can inhibit C1q association with immune complexes. *Clin. Exp. Immunol.* 1997; 108: 181-190
- ¹⁰¹ Routsias JG, Tzioufas AG, Sakarellos C, Moutsopoulos HM. Calreticulin synthetic peptide analogues: anti-peptide antibodies in autoimmune rheumatic diseases. *Clin. Exp. Immunol.* 1993; 91: 437-441
- ¹⁰² Moyes SP, Brown CM, Scott BB, Maini RN, Mayed RA. Analysis of V kappa genes in rheumatoid arthritis synovial B lymphocytes provides evidence for both polyclonal activation and antigen-driven selection. *Clin Exp Immunol* 1996; 105: 89-98
- ¹⁰³ Scott GK et.al. Hirudin-sensitive proteinase. *Bio Chemistry* (has been accepted for publication)
- ¹⁰⁴ Sakaguchi S. Regulatory T cells key controllers of immunologic self-tolerance. *Cell* 2000; 101: 455-458

-
- ¹⁰⁵ Shevach EM. Regulatory T cells in autoimmunity. *Ann. Rev. Immunol.* 2000; 18: 423-49
- ¹⁰⁶ Groux H, Bigler M, O'Garra A, Rouleau M, Antonenka S, de Vries JE, Roncarolo MG. A CD4⁺ T cell subset inhibits antigen-specific T cell responses and prevents colitis. *Nature* 1997; 389: 737
- ¹⁰⁷ Bloom BR, Modlin RL, Salgame P. Stigma variations: observations on suppressor T cells and leprosy. *Ann. Rev. Immunol.* 1992; 10: 453-488
- ¹⁰⁸ Fievet N, Moussa M, Tami G, Maubert B, Cot M, Deleron P, Chaouat G. Plasmodium falciparum induces TH1/TH2 disequilibrium, favoring the TH1 type pathway in the human placenta. *JID* 2001; 183: 1530-1534
- ¹⁰⁹ Joss A, Akdis M, Faith A, Blaser K, Akdis CA. IL-10 directly acts on T cells by specifically altering the CD28 co-stimulation pathway. *Eur. J. Immunol.* 2000; 30: 1683-1690
- ¹¹⁰ Combe B, Pope RM, Fischbach M, Darnell B, Baron S, Talal N. Interleukin 2 in rheumatoid arthritis. Production and response to interleukin 2 in rheumatoid synovial fluid, synovial tissue and peripheral blood. *Clin. Exp. Immunol.* 1985; 59: 520-528
- ¹¹¹ Firestein GS, Zvaifler NJ. Peripheral blood and synovial fluid monocyte activation in inflammatory arthritis II. Low levels of synovial fluid and synovial tissue interferon suggests that gamma-interferon is not the primary macrophage activating factor. *Arthritis Rheum.* 1987; 30: 864-871
- ¹¹² Nathan CF, Kaplan G, Levis WR et.al. Local and systemic effects of intradermal interferon-gamma in patients with lepromatous leprosy. *N. Eng. J. Med.* 1986; 315: 6-15
- ¹¹³ Alsalameh S, Kalden JR, Burmester GR. Die Rolle von Zytokinen und Wachstumsfaktoren bei der rheumatoide Gelenkdestruktion. *Z. Rheumatol.* 1991; 50: 347-359
- ¹¹⁴ Alvaro-Gracia GM, Zvaifler NJ, Firestein GM. Cytokines in chronic inflammatory arthritis V. Mutual antagonism between interferon-gamma and tumor necrosis factor-alpha on HLA-DR expression , proliferation, collagenase production and GMCSF-production in rheumatoid arthritis synoviocytes. *J. Clin. Invest.* 1990; 86: 1790-1798
- ¹¹⁵ Klimiuk PA, Yang H, Goronzy J, Weyand CM. Production of cytokines and metalloproteinases in rheumatoid synovitis is T cell dependent. *Clinical Immunology* 1999; 90: 65-78
- ¹¹⁶ Wagner AD, Bjornsson J, Barthley GB, Goronzy JJ, Weyand CM. Interferon-gamma producing T cells in giant cell vasculitis represent a minority of tissue-infiltrating cells and are located in the distant from the site of pathology. *Am. J. Pathol.* 1996; 148: 1925-1933
- ¹¹⁷ Coyle AI, Lloyd C, Tian J, Nguyen T, Eriksson C, Wang L, Ottoson P, Person P, Delaney T, Lehar S, Lin S, Poison L, Meisel C, Kamradt T, Bjerke T, Levinson D, Gutierrez-Ramos J. Crucial role of the interleukin 1 receptor family member T1/ST2 in the T helper cell type 2-mediated lung mucosal immune responses. *J. Exp. Med* 1999; 190: 895-902
- ¹¹⁸ Yan XM, Guo J, Pichurin P, Tanaka K, Jaume JC, Rapaport B, McLauchlan SM. Cytokines, IgG subclasses and costimulation in mouse model of thyroid autoimmunity induced by injection of fibroblasts co-expressing MHC class II and thyroid antigens. *Clin. Exp. Immunol.* 2000; 122: 170-179
- ¹¹⁹ Miossec P, Naviliat M, Dupuy d'Angeac A, Sany J, Banchereau J. Low levels of Interleukin 4 and high levels of TGF- β in rheumatoid arthritis. *Arthritis Rheum.* 1990; 145: 2514-2517
- ¹²⁰ McInnes IB, Liew FY. Interleukin15: a proinflammatory role in rheumatoid synovitis. *Immunology Today* 1998; 19: 75-79
- ¹²¹ Agostini C, Trentin L, Facco M. Role of IL-15, IL-2, and their receptors in the development of T cell alveolitis in pulmonary sarcoidosis. *J. Immunol.* 1996; 157: 910-918

-
- ¹²² Jullien D, Sieling PA, Uyemura K, Mar ND, Rea TH, Modlin RL. IL-15, an immunomodulator of T cell responses in intracellular infections. *J. Immunol.* 1997; 158: 800-806
- ¹²³ Kirman I, Nielsen OH. Increased numbers of interleukin-15-expressing cells in active ulcerative colitis. *Am. J. Gastroenterology* 1996; 91: 1789-1794
- ¹²⁴ Dooms H, Van Belle T, Desmedt M, Roittiers P, Grotten J. IL-15 redirects the outcome of a tolerizing T cell stimulus from apoptosis to anergy. *Blood* 2000; 96: 1006-1012
- ¹²⁵ Kelleher P, Maroof A, Knight SC. Retrovally induced switch from production of IL-12 to IL-4 in dendritic cells. *Eur. J. Immunol.* 1999; 29: 2309-2318
- ¹²⁶ de Waal Malefyt R, Haanem J, Spits H, Roncarollo MG, te Velde A, de Vries JE et. al. IL-10 and viral IL-10 strongly reduce antigen-specific human T cell proliferation by diminishing the antigen presenting capacity of monocytes via downregulation of class II MHC expression. *J. Exp. Med.* 1991; 174: 915-921
- ¹²⁷ de Waal Malefyt R, Abrams J, de Vries JE et.al. IL-10 inhibits cytokine synthesis by human monocytes: an autoregulatory role of IL-10 produced by monocytes. *J. Exp. Med.* 1991; 174: 1209-1214
- ¹²⁸ Gerosa F, Paganin C, Trinchieri G et.al. Interleukin 12 primes human CD4 and CD8 T cell clones for high production of both interferon-gamma and interleukin-10. *J. Exp. Med.* 1996; 183: 2559-2565
- ¹²⁹ Aarvak T, Chabaud M, Thoen J, Miossec P, Natvig JB. Changes in the TH1 or TH2 cytokine dominance in the synovium of rheumatoid arthritis: a kinetic study of the TH subsets in one unusual RA patient. *Rheumatology* 2000 39: 513-522
- ¹³⁰ Hopkins SJ, Meager A. Cytokines in synovial fluid II. The presence of tumor necrosis factor and interferon. *Clin. Exp. Immunol.* 1988; 73: 88-92
- ¹³¹ Laouar Y, Crispe IN. Functional flexibility in T cells: independent regulation of CD4+ T cell proliferation and effector function in vivo. *Immunity* 2000; 13: 291-301