

8 Literatur

- [1] Arbeitsgemeinschaft Bevölkerungsbezogener Krebsregister in Deutschland. Krebsregister . 2004.
- [2] Häring R. Chirurgie. Walter de Gruyter Berlin, 1992.
- [3] Sarli L, Costi R, Sansebastiano G, Trivelli M, Roncoroni L Prospective randomized trial of low-pressure pneumoperitoneum for reduction of shoulder-tip pain following laparoscopy. *Br J Surg* 2000; 87(9):1161-5.
- [4] Murdock CM, Wolff AJ, Van Geem T. Risk factors for hypercarbia, subcutaneous emphysema, pneumothorax, and pneumomediastinum during laparoscopy. *Obstet Gynecol* 2000; 95(5):704-9.
- [5] Galizia G, Prizio G, Lieto E et al. Hemodynamic and pulmonary changes during open, carbon dioxide pneumoperitoneum and abdominal wall-lifting cholecystectomy. A prospective, randomized study. *Surg Endosc* 2001; 15(5):477-83.
- [6] Dexter SP, Vucevic M, Gibson J, McMahon MJ. Hemodynamic consequences of high- and low-pressure capnoperitoneum during laparoscopic cholecystectomy. *Surg Endosc* 1999; 13(4):376-81.
- [7] Cunningham AJ, Brull SJ. Laparoscopic cholecystectomy: anesthetic implications. *Anesth Analg* 1993; 76(5):1120-33.
- [8] Volpino P, Cangemi V, D'Andrea N, Cangemi B, Piat G. Hemodynamic and pulmonary changes during and after laparoscopic cholecystectomy. A comparison with traditional surgery. *Surg Endosc* 1998; 12(2):119-23.
- [9] Chmielnicki Z, Noras K, Boldys S, Celarek B. [Effect of CO₂ insufflation into the peritoneal cavity on selected indices of respiratory system function during laparoscopic cholecystectomy]. *Wiad Lek* 1994; 47(13-14):503-5.
- [10] Allendorf JD, Bessler M, Whelan RL et al. Better preservation of immune function after laparoscopic-assisted vs. open bowel resection in a murine model. *Dis Colon Rectum* 1996; 39(10 Suppl):S67-S72.

- [11] Jacobi CA, Ordemann J, Zieren HU *et al.* Increased systemic inflammation after laparotomy vs laparoscopy in an animal model of peritonitis. *Arch Surg* 1998; 133(3):258-62.
- [12] Karayiannakis AJ, Makri GG, Mantzioka A, Karousos D, Karatzas G. Systemic stress response after laparoscopic or open cholecystectomy: a randomized trial. *Br J Surg* 1997; 84(4):467-71.
- [13] Ordemann J, Jacobi CA, Schwenk W, Stosslein R, Muller JM. Cellular and humoral inflammatory response after laparoscopic and conventional colorectal resections. *Surg Endosc* 2001; 15(6):600-8.
- [14] Perttila J, Salo M, Ovaska J *et al.* Immune response after laparoscopic and conventional Nissen fundoplication. *Eur J Surg* 1999; 165(1):21-8.
- [15] McMahon AJ, O'Dwyer PJ, Cruikshank AM *et al.* Comparison of metabolic responses to laparoscopic and minilaparotomy cholecystectomy. *Br J Surg* 1993; 80(10):1255-8.
- [16] Ortega AE, Peters JH, Incarbone R *et al.* A prospective randomized comparison of the metabolic and stress hormonal responses of laparoscopic and open cholecystectomy. *J Am Coll Surg* 1996; 183(3):249-56.
- [17] Milsom JW, Bohm B, Hammerhofer KA *et al.* A prospective, randomized trial comparing laparoscopic versus conventional techniques in colorectal cancer surgery: a preliminary report. *J Am Coll Surg* 1998; 187(1):46-54.
- [18] Nilsson G, Larsson S, Johnsson F. Randomized clinical trial of laparoscopic versus open fundoplication: blind evaluation of recovery and discharge period. *Br J Surg* 2000; 87(7):873-8.
- [19] Squirrell DM, Majeed AW, Troy G *et al.* A randomized, prospective, blinded comparison of postoperative pain, metabolic response, and perceived health after laparoscopic and small incision cholecystectomy. *Surgery* 1998; 123(5):485-95.
- [20] Lacy AM, Garcia-Valdecasas JC, Pique JM *et al.* Short-term outcome analysis of a randomized study comparing laparoscopic vs open colectomy for colon cancer. *Surg Endosc* 1995; 9(10):1101-5.

- [21] Schwenk W, Bohm B, Witt C *et al.* Pulmonary function following laparoscopic or conventional colorectal resection: a randomized controlled evaluation. *Arch Surg* 1999; 134(1):6-12.
- [22] Poulin EC, Mamazza J, Schlachta CM, Gregoire R, Roy N. Laparoscopic resection does not adversely affect early survival curves in patients undergoing surgery for colorectal adenocarcinoma. *Ann Surg* 1999; 229(4):487-92.
- [23] Lechaux D, Trebuchet G, Le Calve JL. Five-year results of 206 laparoscopic left colectomies for cancer. *Surg Endosc* 2002; 16(10):1409-12.
- [24] Fleshman JW, Wexner SD, Anvari M *et al.* Laparoscopic vs. open abdominoperineal resection for cancer. *Dis Colon Rectum* 1999; 42(7):930-9.
- [25] Kockerling F, Schneider C, Reymond MA *et al.* Early results of a prospective multicenter study on 500 consecutive cases of laparoscopic colorectal surgery. Laparoscopic Colorectal Surgery Study Group (LCSSG). *Surg Endosc* 1998; 12(1):37-41.
- [26] Stocchi L, Nelson H. Laparoscopic colectomy for colon cancer: trial update. *J Surg Oncol* 1998; 68(4):255-67.
- [27] Franklin ME, Kazantsev GB, Abrego D *et al.* Laparoscopic surgery for stage III colon cancer: long-term follow-up. *Surg Endosc* 2000; 14(7):612-6.
- [28] Lacy AM, Delgado S, Garcia-Valdecasas JC *et al.* Port site metastases and recurrence after laparoscopic colectomy. A randomized trial. *Surg Endosc* 1998; 12(8):1039-42.
- [29] Hahnloser D, Hetzer FH. [Laparoscopy resection for colon cancer. The new standard?]. *Schweiz Rundsch Med Prax* 2004; 93(36):1447-53.
- [30] A comparison of laparoscopically assisted and open colectomy for colon cancer *N Engl J Med* 2004; 350(20):2050-9.
- [31] Berends FJ, Kazemier G, Bonjer HJ, Lange JF. Subcutaneous metastases after laparoscopic colectomy. *Lancet* 1994; 344(8914):58.

- [32] Schaeff B, Paolucci V, Thomopoulos J. Port site recurrences after laparoscopic surgery. A review. *Dig Surg* 1998; 15(2):124-34.
- [33] Neuhaus SJ, Texler M, Hewett PJ, Watson DI. Port-site metastases following laparoscopic surgery. *Br J Surg* 1998; 85(6):735-41.
- [34] Ziprin P, Ridgway PF, Peck DH, Darzi AW. The theories and realities of port-site metastases: a critical appraisal. *J Am Coll Surg* 2002; 195(3):395-408.
- [35] Martinez J, Targarona EM, Balague C, Pera M, Trias M. Port site metastasis. An unresolved problem in laparoscopic surgery. A review. *Int Surg* 1995; 80(4):315-21.
- [36] Wexner SD, Cohen SM. Port site metastases after laparoscopic colorectal surgery for cure of malignancy. *Br J Surg* 1995; 82(3):295-8.
- [37] Schiedeck TH, Schwandner O, Baca I *et al*. Laparoscopic surgery for the cure of colorectal cancer: results of a German five-center study. *Dis Colon Rectum* 2000; 43(1):1-8.
- [38] Curet MJ, Putrakul K, Pitcher DE, Josloff RK, Zucker KA. Laparoscopically assisted colon resection for colon carcinoma: perioperative results and long-term outcome. *Surg Endosc* 2000; 14(11):1062-6.
- [39] Hartley JE, Mehigan BJ, MacDonald AW, Lee PW, Monson JR. Patterns of recurrence and survival after laparoscopic and conventional resections for colorectal carcinoma. *Ann Surg* 2000; 232(2):181-6.
- [40] Reilly WT, Nelson H, Schroeder G *et al*. Wound recurrence following conventional treatment of colorectal cancer. A rare but perhaps underestimated problem. *Dis Colon Rectum* 1996; 39(2):200-7.
- [41] Fusco MA, Paluzzi MW. Abdominal wall recurrence after laparoscopically assisted colectomy for adenocarcinoma of the colon. Report of a case. *Dis Colon Rectum* 1993; 36(9):858-61.
- [42] Clair DG, Lautz DB, Brooks DC. Rapid development of umbilical metastases after laparoscopic cholecystectomy for unsuspected gallbladder carcinoma. *Surgery* 1993; 113(3):355-8.

- [43] Lee SW, Whelan RL, Southall JC, Bessler M. Abdominal wound tumor recurrence after open and laparoscopic-assisted splenectomy in a murine model. *Dis Colon Rectum* 1998; 41(7):824-31.
- [44] Kim SH, Milsom JW, Gramlich TL *et al.* Does laparoscopic vs. conventional surgery increase exfoliated cancer cells in the peritoneal cavity during resection of colorectal cancer? *Dis Colon Rectum* 1998; 41(8):971-8.
- [45] Whalen GF, Ingber DE. Inhibition of tumor-cell attachment to extracellular matrix as a method for preventing tumor recurrence in a surgical wound. *Ann Surg* 1989; 210(6):758-64.
- [46] Gutt CN, Riemer V, Kim ZG *et al.* Impact of laparoscopic colonic resection on tumour growth and spread in an experimental model. *Br J Surg* 1999; 86(9):1180-4.
- [47] Jones DB, Guo LW, Reinhard MK *et al.* Impact of pneumoperitoneum on trocar site implantation of colon cancer in hamster model. *Dis Colon Rectum* 1995; 38(11):1182-8.
- [48] Zmora O, Gervaz P, Wexner SD. Trocar site recurrence in laparoscopic surgery for colorectal cancer. *Surg Endosc* 2001; 15(8):788-93.
- [49] Bouvy ND, Marquet RL, Jeekel H, Bonjer HJ. Impact of gas(less) laparoscopy and laparotomy on peritoneal tumor growth and abdominal wall metastases. *Ann Surg* 1996; 224(6):694-700.
- [50] Neuhaus SJ, Watson DI, Ellis T *et al.* Wound metastasis after laparoscopy with different insufflation gases. *Surgery* 1998; 123(5):579-83.
- [51] Dorrance HR, Oien K, O'Dwyer PJ. Effects of laparoscopy on intraperitoneal tumor growth and distant metastases in an animal model. *Surgery* 1999; 126(1):35-40.
- [52] Farrell TM, Metreveli RE, Johnson AB, Smith CD, Hunter JG. Choice of insufflating gas influences on wound metastasis. *Surg Endosc* 2000; 14(11):1047-9.
- [53] Jacobi CA, Sabat R, Bohm B *et al.* Pneumoperitoneum with carbon dioxide stimulates growth of malignant colonic cells. *Surgery* 1997; 121(1):72-8.

- [54] Gutt CN, Kim ZG, Schmadera T, Paolucci V, Lorenz M. Carbon dioxide pneumoperitoneum is associated with increased liver metastases in a rat model. *Surgery* 2000; 127(5):566-70.
- [55] Ishida H, Murata N, Yamada H *et al.* Pneumoperitoneum with carbon dioxide enhances liver metastases of cancer cells implanted into the portal vein in rabbits. *Surg Endosc* 2000; 14(3):239-42.
- [56] Jacobi CA, Ordemann J, Bohm B *et al.* Inhibition of peritoneal tumor cell growth and implantation in laparoscopic surgery in a rat model. *Am J Surg* 1997; 174(3):359-63.
- [57] Gutt CN, Kim ZG, Hollander D, Bruttel T, Lorenz M. CO₂ environment influences the growth of cultured human cancer cells dependent on insufflation pressure. *Surg Endosc* 2001; 15(3):314-8.
- [58] Ishida H, Murata N, Yokoyama M *et al.* The influence of different insufflation pressures during carbon dioxide pneumoperitoneum on the development of pulmonary metastasis in a mouse model. *Surg Endosc* 2000; 14(6):578-81.
- [59] Wittich P, Steyerberg EW, Simons SH, Marquet RL, Bonjer HJ. Intraperitoneal tumor growth is influenced by pressure of carbon dioxide pneumoperitoneum. *Surg Endosc* 2000; 14(9):817-9.
- [60] Jacobi CA, Wenger FA, Ordemann J *et al.* Experimental study of the effect of intra-abdominal pressure during laparoscopy on tumour growth and port site metastasis. *Br J Surg* 1998; 85(10):1419-22.
- [61] Basson MD, Yu CF, Herden-Kirchoff O *et al.* Effects of increased ambient pressure on colon cancer cell adhesion. *J Cell Biochem* 2000; 78(1):47-61.
- [62] Volz J, Koster S, Spacek Z, Paweletz N. Characteristic alterations of the peritoneum after carbon dioxide pneumoperitoneum. *Surg Endosc* 1999; 13(6):611-4.
- [63] Buck RC. Walker 256 tumor implantation in normal and injured peritoneum studied by electron microscopy, scanning electron microscopy, and autoradiography. *Cancer Res* 1973; 33(12):3181-8.

- [64] Rovensky YA, Gvichiya AS, Vasiliev JM. SEM study of the attachment of mouse ascitic hepatoma cells to various substrata. *Scan Electron Microsc* 1980;(3):71-8.
- [65] Eleftheriadis E, Kotzampassi K, Papanotas K, Heliadis N, Sarris K. Gut ischemia, oxidative stress, and bacterial translocation in elevated abdominal pressure in rats. *World J Surg* 1996; 20(1):11-6.
- [66] Wu L, Mustoe TA. Effect of ischemia on growth factor enhancement of incisional wound healing. *Surgery* 1995; 117(5):570-6.
- [67] Jacobi CA, Peter FJ, Wenger FA, Ordemann J, Muller JM. New therapeutic strategies to avoid intra- and extraperitoneal metastases during laparoscopy: results of a tumor model in the rat. *Dig Surg* 1999; 16(5):393-9.
- [68] Wildbrett P, Oh A, Naundorf D, Volk T, Jacobi CA. Impact of laparoscopic gases on peritoneal microenvironment and essential parameters of cell function. *Surg Endosc* 2003; 17(1):78-82.
- [69] Kopernik G, Avinoach E, Grossman Y *et al*. The effect of a high partial pressure of carbon dioxide environment on metabolism and immune functions of human peritoneal cells-relevance to carbon dioxide pneumoperitoneum. *Am J Obstet Gynecol* 1998; 179(6 Pt 1):1503-10.
- [70] Jacobi CA, Ordemann J, Halle E, Volk HD, Muller JM. Impact of laparoscopy with carbon dioxide versus helium on local and systemic inflammation in an animal model of peritonitis. *J Laparoendosc Adv Surg Tech A* 1999; 9(3):305-12.
- [71] Watson DI, Mathew G, Ellis T *et al*. Gasless laparoscopy may reduce the risk of port-site metastases following laparoscopic tumor surgery. *Arch Surg* 1997; 132(2):166-8.
- [72] Neuhaus SJ, Watson DI, Ellis T *et al*. Influence of gases on intraperitoneal immunity during laparoscopy in tumor-bearing rats. *World J Surg* 2000; 24(10):1227-31.
- [73] Mathew G, Watson DI, Ellis TS, Jamieson GG, Rofe AM. The role of peritoneal immunity and the tumour-bearing state on the development of wound and peritoneal metastases after laparoscopy. *Aust N Z J Surg* 1999;

69(1):14-8.

- [74] Mathew G, Watson DI, Rofe AM *et al.* Wound metastases following laparoscopic and open surgery for abdominal cancer in a rat model. *Br J Surg* 1996; 83(8):1087-90.
- [75] Vittimberga FJ, Jr., Foley DP, Meyers WC, Callery MP. Laparoscopic surgery and the systemic immune response. *Ann Surg* 1998; 227(3):326-34.
- [76] Watson RW, Redmond HP, McCarthy J, Burke PE, Bouchier-Hayes D. Exposure of the peritoneal cavity to air regulates early inflammatory responses to surgery in a murine model. *Br J Surg* 1995; 82(8):1060-5.
- [77] Bedrosian I, Sofia RD, Wolff SM, Dinarello CA. Taurolidine, an analogue of the amino acid taurine, suppresses interleukin 1 and tumor necrosis factor synthesis in human peripheral blood mononuclear cells. *Cytokine* 1991; 3(6):568-75.
- [78] Goldstein DS, Lu ML, Hattori T *et al.* Inhibition of peritoneal tumor-cell implantation: model for laparoscopic cancer surgery. *J Endourol* 1993; 7(3):237-41.
- [79] Gorman SP, McCafferty DF, Woolfson AD, Jones DS. Reduced adherence of micro-organisms to human mucosal epithelial cells following treatment with Taurolin, a novel antimicrobial agent. *J Appl Bacteriol* 1987; 62(4):315-20.
- [80] Skipper D, Jeffrey MJ, Cooper AJ, Alexander P, Taylor I. Enhanced growth of tumour cells in healing colonic anastomoses and laparotomy wounds. *Int J Colorectal Dis* 1989; 4(3):172-7.
- [81] Ruoslahti E, Pierschbacher MD. New perspectives in cell adhesion: RGD and integrins. *Science* 1987; 238(4826):491-7.
- [82] Castronovo V, Taraboletti G, Sobel ME. Laminin receptor complementary DNA-deduced synthetic peptide inhibits cancer cell attachment to endothelium. *Cancer Res* 1991; 51(20):5672-8.
- [83] Ruoslahti E. The Walter Herbert Lecture. Control of cell motility and tumour invasion by extracellular matrix interactions. *Br J Cancer* 1992; 66(2):239-42.

- [84] Hirabayashi Y, Yamaguchi K, Shiraishi N *et al.* Development of port-site metastasis after pneumoperitoneum. *Surg Endosc* 2002; 16(5):864-8.
- [85] Tseng LN, Berends FJ, Wittich P *et al.* Port-site metastases. Impact of local tissue trauma and gas leakage. *Surg Endosc* 1998; 12(12):1377-80.
- [86] Aoki Y, Shimura H, Li H *et al.* A model of port-site metastases of gallbladder cancer: the influence of peritoneal injury and its repair on abdominal wall metastases. *Surgery* 1999; 125(5):553-9.
- [87] Aznavoorian S, Murphy AN, Stetler-Stevenson WG, Liotta LA. Molecular aspects of tumor cell invasion and metastasis. *Cancer* 1993; 71(4):1368-83.
- [88] Stetler-Stevenson WG, Aznavoorian S, Liotta LA. Tumor cell interactions with the extracellular matrix during invasion and metastasis. *Annu Rev Cell Biol* 1993; 9:541-73.
- [89] Lanfrancone L, Boraschi D, Ghiara P *et al.* Human peritoneal mesothelial cells produce many cytokines (granulocyte colony-stimulating factor [CSF], granulocyte-monocyte-CSF, macrophage-CSF, interleukin-1 [IL-1], and IL-6) and are activated and stimulated to grow by IL-1. *Blood* 1992; 80(11):2835-42.
- [90] Yanagisawa M, Imai H, Fukushima Y *et al.* Effects of tumour necrosis factor alpha and interleukin 1 beta on the proliferation of cultured glomerular epithelial cells. *Virchows Arch* 1994; 424(6):581-6.
- [91] Zetter BR. Adhesion molecules in tumor metastasis. *Semin Cancer Biol* 1993; 4(4):219-29.
- [92] Honn KV, Tang DG. Adhesion molecules and tumor cell interaction with endothelium and subendothelial matrix. *Cancer Metastasis Rev* 1992; 11(3-4):353-75.
- [93] Kinsella AR, Green B, Lepts GC *et al.* The role of the cell-cell adhesion molecule E-cadherin in large bowel tumour cell invasion and metastasis. *Br J Cancer* 1993; 67(5):904-9.
- [94] Mulder JW, Wielenga VJ, Pals ST, Offerhaus GJ. p53 and CD44 as clinical markers of tumour progression in colorectal carcinogenesis. *Histochem J* 1997; 29(6):439-52.

- [95] Wielenga VJ, van d, V, Mulder JW *et al.* CD44 splice variants as prognostic markers in colorectal cancer. *Scand J Gastroenterol* 1998; 33(1):82-7.
- [96] Haier J, Nasralla M, Nicolson GL. Cell surface molecules and their prognostic values in assessing colorectal carcinomas. *Ann Surg* 2000; 231(1):11-24.
- [97] Jiang WG. Cell adhesion molecules in the formation of liver metastasis. *J Hepatobiliary Pancreat Surg* 1998; 5(4):375-82.
- [98] Jiang WG. E-cadherin and its associated protein catenins, cancer invasion and metastasis. *Br J Surg* 1996; 83(4):437-46.
- [99] Kinsella AR, Lepts GC, Hill CL, Jones M. Reduced E-cadherin expression correlates with increased invasiveness in colorectal carcinoma cell lines. *Clin Exp Metastasis* 1994; 12(4):335-42.
- [100] Weg-Remers S, Anders M, von Lampe B *et al.* Decreased expression of CD44 splicing variants in advanced colorectal carcinomas. *Eur J Cancer* 1998; 34(10):1607-11.
- [101] Gumbiner BM. Cell adhesion: the molecular basis of tissue architecture and morphogenesis. *Cell* 1996; 84(3):345-57.
- [102] Birchmeier W, Hulsken J, Behrens J. E-cadherin as an invasion suppressor. *Ciba Found Symp* 1995; 189:124-36; discussion 136-41, 174-6.:124-36.
- [103] Takeichi M. Cadherins in cancer: implications for invasion and metastasis. *Curr Opin Cell Biol* 1993; 5(5):806-11.
- [104] Kim ZG FAU, Mehl CF, Lorenz MF, Gutt CN. Impact of laparoscopic CO₂-insufflation on tumor-associated molecules in cultured colorectal cancer cells. *Surg Endosc* 2002;16(8):1182-6.
- [105] Cai KL, Wang GB, Xiong LJ. Effects of carbon dioxide and nitrogen on adhesive growth and expressions of E-cadherin and VEGF of human colon cancer cell CCL-228. *World J Gastroenterol* 2003; 9(7):1594-7.
- [106] Ponta H, Sherman L, Herrlich PA. CD44: from adhesion molecules to signalling regulators. *Nat Rev Mol Cell Biol* 2003; 4(1):33-45.

- [107] Wielenga VJ, Heider KH, Offerhaus GJ *et al.* Expression of CD44 variant proteins in human colorectal cancer is related to tumor progression. *Cancer Res* 1993; 53(20):4754-6.
- [108] Ropponen KM, Eskelinan MJ, Lipponen PK, Alhava E, Kosma VM. Expression of CD44 and variant proteins in human colorectal cancer and its relevance for prognosis. *Scand J Gastroenterol* 1998; 33(3):301-9.
- [109] Heider KH, Hofmann M, Hors E *et al.* A human homologue of the rat metastasis-associated variant of CD44 is expressed in colorectal carcinomas and adenomatous polyps. *J Cell Biol* 1993; 120(1):227-33.
- [110] Gunthert U, Hofmann M, Rudy W *et al.* A new variant of glycoprotein CD44 confers metastatic potential to rat carcinoma cells. *Cell* 1991; 65(1):13-24.
- [111] Mulder JW, Kruyt PM, Sewnath M *et al.* Difference in expression of CD44 splice variants between proximal and distal adenocarcinoma of the large bowel. *Br J Surg* 1995; 82(11):1468-70.
- [112] Shibuya YF, Okabayashi TF, Oda KF, Tanaka N. Ratio of CD44 epithelial variant to CD44 hematopoietic variant is a useful prognostic indicator in gastric and colorectal carcinoma. *Jpn J Clin Oncol* 1998;(10):609-14.
- [113] Mulder JW, Kruyt PM, Sewnath M *et al.* Colorectal cancer prognosis and expression of exon-v6-containing CD44 proteins. *Lancet* 1994; 344(8935):1470-2.
- [114] Hynes RO. Integrins: versatility, modulation, and signaling in cell adhesion. *Cell* 1992; 69(1):11-25.
- [115] Howe A, Aplin AE, Alahari SK, Juliano RL. Integrin signaling and cell growth control. *Curr Opin Cell Biol* 1998; 10(2):220-31.
- [116] Varner JA, Cheresh DA. Integrins and cancer. *Curr Opin Cell Biol* 1996; 8(5):724-30.
- [117] Felding-Habermann B. Integrin adhesion receptors in tumor metastasis. *Clin Exp Metastasis* 2003; 20(3):203-13.

- [118] Neuhaus SJ, Ellis T, Jamieson GG, Watson DI. Experimental study of the effect of intraperitoneal heparin on tumour implantation following laparoscopy. *Br J Surg* 1999; 86(3):400-4.
- [119] Jacobi CA, Sabat R, Ordemann J *et al.* [Peritoneal instillation of taurolidine and heparin for preventing intraperitoneal tumor growth and trocar metastases in laparoscopic operations in the rat model]. *Langenbecks Arch Chir* 1997; 382(4 Suppl 1):S31-S36.
- [120] Neuhaus SJ, Watson DI, Ellis T *et al.* Efficacy of cytotoxic agents for the prevention of laparoscopic port-site metastases. *Arch Surg* 1998; 133(7):762-6.
- [121] Balli JE, Franklin ME, Almeida JA *et al.* How to prevent port-site metastases in laparoscopic colorectal surgery. *Surg Endosc* 2000; 14(11):1034-6.
- [122] Teschner M. Taurolidin in der septischen Chirurgie. 1996. Georg Thieme Verlag Stuttgart.
- [123] Braumann C, Ordemann J, Kilian M, Wenger FA, Jacobi CA. Local and systemic chemotherapy with taurolidine and taurolidine/heparin in colon cancer-bearing rats undergoing laparotomy. *Clin Exp Metastasis* 2003; 20(5):387-94.
- [124] Braumann C, Ordemann J, Wildbrett P, Jacobi CA. Influence of intraperitoneal and systemic application of taurolidine and taurolidine/heparin during laparoscopy on intraperitoneal and subcutaneous tumour growth in rats. *Clin Exp Metastasis* 2000; 18(7):547-52.
- [125] Opitz I, Veen HC, Braumann C *et al.* The influence of adhesion prophylactic substances and taurolidine/heparin on local recurrence and intraperitoneal tumor growth after laparoscopic-assisted bowel resection of colon carcinoma in a rat model. *Surg Endosc* 2003; 17:1098-1104.
- [126] McCourt M, Wang JH, Sookhai S, Redmond HP. Taurolidine inhibits tumor cell growth in vitro and in vivo. *Ann Surg Oncol* 2000; 7(9):685-91.
- [127] Victor FC, Gottlieb AB. TNF-alpha and apoptosis: implications for the pathogenesis and treatment of psoriasis. *J Drugs Dermatol* 2002; 1(3):264-

- [128] Volz J, Volz-Koster S, Kanis S *et al*. Modulation of tumor-induced lethality after pneumoperitoneum in a mouse model. *Cancer* 2000; 89(2):262-6.
- [129] Calabresi P, Goulette FA, Darnowski JW. Taurolidine: cytotoxic and mechanistic evaluation of a novel antineoplastic agent. *Cancer Res* 2001; 61(18):6816-21.
- [130] Han Z, Ribbizi I, Pantazis P *et al*. The antibacterial drug taurolidine induces apoptosis by a mitochondrial cytochrome c-dependent mechanism. *Anticancer Res* 2002; 22(4):1959-64.
- [131] Petrovic L, Schlegel KA, Ries J *et al*. [In vitro effect of taurolidine on squamous cell carcinoma in the oral cavity]. *Mund Kiefer Gesichtschir* 2003; 7(2):102-7.
- [132] Stendel R, Scheurer L, Stoltenburg-Didinger G, Brock M, Mohler H. Enhancement of Fas-ligand-mediated programmed cell death by taurolidine. *Anticancer Res* 2003; 23(3B):2309-14.
- [133] Braumann C, Henke W, Jacobi CA, Dubiel W. The tumor-suppressive reagent taurolidine is an inhibitor of protein biosynthesis. *Int J Cancer* 2004; 112(2):225-30.
- [134] Blenkharn JI. Sustained anti-adherence activity of taurolidine (Taurolin) and noxythiolin (Noxyflex S) solutions. *J Pharm Pharmacol* 1988; 40(7):509-11.
- [135] Knight BI, Skellern GG, Browne MK, Pfirrmann RW. Peritoneal absorption of the antibacterial and antiendotoxin taurolin in peritonitis. *Br J Clin Pharmacol* 1981; 12(5):695-9.
- [136] Ribizzi I, Darnowski JW, Goulette FA *et al*. Taurolidine: preclinical evaluation of a novel, highly selective, agent for bone marrow purging. *Bone Marrow Transplant* 2002; 29(4):313-9.
- [137] Da Costa ML, Redmond P, Bouchier-Hayes DJ. The effect of laparotomy and laparoscopy on the establishment of spontaneous tumor metastases. *Surgery* 1998; 124(3):516-25.

- [138] Sawada T, Nishizawa H, Nishio E, Kadouaki M. Postoperative adhesion prevention with an oxidized regenerated cellulose adhesion barrier in infertile women. *J Reprod Med* 2000; 45(5):387-9.
- [139] Szabo A, Haj M, Waxsman I, Eitan A. Evaluation of seprafilm and amniotic membrane as adhesion prophylaxis in mesh repair of abdominal wall hernia in rats. *Eur Surg Res* 2000; 32(2):125-8.
- [140] Tzianabos AO, Cisneros RL, Gershkovich J *et al.* Effect of surgical adhesion reduction devices on the propagation of experimental intra-abdominal infection. *Arch Surg* 1999; 134(11):1254-9.
- [141] Underwood RA, Wu JS, Wright MP *et al.* Sodium hyaluronate carboxymethylcellulose-based bioresorbable membrane (Seprafilm)--does it affect tumor implantation at abdominal wound sites? *Dis Colon Rectum* 1999; 42(5):614-8.
- [142] Wiseman DM, Gottlick-Iarkowski L, Kamp L. Effect of different barriers of oxidized regenerated cellulose (ORC) on cecal and sidewall adhesions in the presence and absence of bleeding. *J Invest Surg* 1999; 12(3):141-6.
- [143] Wiseman DM, Kamp LF, Saferstein L *et al.* Improving the efficacy of INTERCEED Barrier in the presence of blood using thrombin, heparin or a blood insensitive barrier, modified INTERCEED (nTC7). *Prog Clin Biol Res* 1993; 381:205-12.
- [144] Wiseman DM, Trout JR, Franklin RR, Diamond MP. Metaanalysis of the safety and efficacy of an adhesion barrier (Interceed TC7) in laparotomy. *J Reprod Med* 1999; 44(4):325-31.
- [145] Johns DB, Keyport GM, Hoehler F, diZerega GS. Reduction of postsurgical adhesions with Intergel adhesion prevention solution: a multicenter study of safety and efficacy after conservative gynecologic surgery. *Fertil Steril* 2001; 76(3):595-604.
- [146] Johns DB, Rodgers KE, Donahue WD, Kiorpis TC, diZerega GS. Reduction of adhesion formation by postoperative administration of ionically cross-linked hyaluronic acid. *Fertil Steril* 1997; 68(1):37-42.

- [147] Thornton MH, Johns DB, Campeau JD, Hoehler F, diZerega GS. Clinical evaluation of 0.5% ferric hyaluronate adhesion prevention gel for the reduction of adhesions following peritoneal cavity surgery: open-label pilot study. *Hum Reprod* 1998; 13(6):1480-5.
- [148] Sekiba K. Use of Interceed(TC7) absorbable adhesion barrier to reduce postoperative adhesion reformation in infertility and endometriosis surgery. The Obstetrics and Gynecology Adhesion Prevention Committee. *Obstet Gynecol* 1992; 79(4):518-22.
- [149] Prevention of postsurgical adhesions by INTERCEED(TC7), an absorbable adhesion barrier: a prospective randomized multicenter clinical study. INTERCEED(TC7) Adhesion Barrier Study Group *Fertil Steril* 1989; 51(6):933-8.
- [150] Azziz R. Microsurgery alone or with INTERCEED Absorbable Adhesion Barrier for pelvic sidewall adhesion re-formation. The INTERCEED (TC7) Adhesion Barrier Study Group II. *Surg Gynecol Obstet* 1993; 177(2):135-9.
- [151] Franklin RR. Reduction of ovarian adhesions by the use of Interceed. Ovarian Adhesion Study Group. *Obstet Gynecol* 1995; 86(3):335-40.
- [152] Mais V, Ajossa S, Piras B *et al.* Prevention of de-novo adhesion formation after laparoscopic myomectomy: a randomized trial to evaluate the effectiveness of an oxidized regenerated cellulose absorbable barrier. *Hum Reprod* 1995; 10(12):3133-5.
- [153] Jacobi CA, Wildbrett P, Volk T, Muller JM. Influence of different gases and intraperitoneal instillation of antiadherent or cytotoxic agents on peritoneal tumor cell growth and implantation with laparoscopic surgery in a rat model. *Surg Endosc* 1999; 13(10):1021-5.
- [154] Franklin ME, Jr., Rosenthal D, Abrego-Medina D *et al.* Prospective comparison of open vs. laparoscopic colon surgery for carcinoma. Five-year results. *Dis Colon Rectum* 1996; 39(10 Suppl):S35-S46.
- [155] Falk PM, Beart RW, Jr., Wexner SD *et al.* Laparoscopic colectomy: a critical appraisal. *Dis Colon Rectum* 1993; 36(1):28-34.

- [156] Hoffman GC, Baker JW, Doxey JB *et al.* Minimally invasive surgery for colorectal cancer. Initial follow-up. *Ann Surg* 1996; 223(6):790-6.
- [157] Fleshman JW, Nelson H, Peters WR *et al.* Early results of laparoscopic surgery for colorectal cancer. Retrospective analysis of 372 patients treated by Clinical Outcomes of Surgical Therapy (COST) Study Group. *Dis Colon Rectum* 1996; 39(10 Suppl):S53-S58.
- [158] Kruitwagen RF, Swinkels BM, Keyser KG, Doesburg WH, Schijf CP. Incidence and effect on survival of abdominal wall metastases at trocar or puncture sites following laparoscopy or paracentesis in women with ovarian cancer. *Gynecol Oncol* 1996; 60(2):233-7.
- [159] Nduka CC, Monson JR, Menzies-Gow N, Darzi A. Abdominal wall metastases following laparoscopy. *Br J Surg* 1994; 81(5):648-52.
- [160] Larsson B. Efficacy of Interceed in adhesion prevention in gynecologic surgery: a review of 13 clinical studies. *J Reprod Med* 1996; 41(1):27-34.
- [161] Pagidas K, Tulandi T. Effects of Ringer's lactate, Interceed(TC7) and Gore-Tex Surgical Membrane on postsurgical adhesion formation. *Fertil Steril* 1992; 57(1):199-201.
- [162] Maxson WS, Herbert CM, Oldfield EL, Hill GA. Efficacy of a modified oxidized cellulose fabric in the prevention of adhesion formation. *Gynecol Obstet Invest* 1988; 26(2):160-5.
- [163] Best CL, Rittenhouse D, Vasquez C *et al.* Evaluation of interceed(TC7) for reduction of postoperative adhesions in rabbits. *Fertil Steril* 1992; 58(4):817-20.
- [164] Haney AF, Doty E. Expanded-polytetrafluoroethylene but not oxidized regenerated cellulose prevents adhesion formation and reformation in a mouse uterine horn model of surgical injury. *Fertil Steril* 1993; 60(3):550-8.
- [165] West JL, Chowdhury SM, Sawhney AS *et al.* Efficacy of adhesion barriers. Resorbable hydrogel, oxidized regenerated cellulose and hyaluronic acid. *J Reprod Med* 1996; 41(3):149-54.

- [166] Shushan A, Mor-Yosef S, Avgar A, Laufer N. Hyaluronic acid for preventing experimental postoperative intraperitoneal adhesions. *J Reprod Med* 1994; 39(5):398-402.
- [167] Diamond MP. Reduction of de novo postsurgical adhesions by intraoperative precoating with Sepracoat (HAL-C) solution: a prospective, randomized, blinded, placebo-controlled multicenter study. The Sepracoat Adhesion Study Group. *Fertil Steril* 1998; 69(6):1067-74.
- [168] Burns JW, Skinner K, Colt MJ *et al.* A hyaluronate based gel for the prevention of postsurgical adhesions: evaluation in two animal species. *Fertil Steril* 1996; 66(5):814-21.
- [169] De Iaco PA, Stefanetti M, Pressato D *et al.* A novel hyaluronan-based gel in laparoscopic adhesion prevention: preclinical evaluation in an animal model. *Fertil Steril* 1998; 69(2):318-23.
- [170] Yung S, Coles GA, Williams JD, Davies M. The source and possible significance of hyaluronan in the peritoneal cavity. *Kidney Int* 1994; 46(2):527-33.
- [171] Asplund T, Versnel MA, Laurent TC, Heldin P. Human mesothelioma cells produce factors that stimulate the production of hyaluronan by mesothelial cells and fibroblasts. *Cancer Res* 1993; 53(2):388-92.
- [172] Kocak I, Unlu C, Akcan Y, Yakin K. Reduction of adhesion formation with cross-linked hyaluronic acid after peritoneal surgery in rats. *Fertil Steril* 1999; 72(5):873-8.
- [173] Tan B, Wang JH, Wu QD, Kirwan WO, Redmond HP. Sodium hyaluronate enhances colorectal tumour cell metastatic potential in vitro and in vivo. *Br J Surg* 2001; 88(2):246-50.
- [174] Ropponen K, Tammi M, Parkkinen J *et al.* Tumor cell-associated hyaluronan as an unfavorable prognostic factor in colorectal cancer. *Cancer Res* 1998; 58(2):342-7.
- [175] Fujisaki TF, Tanaka YF, Fujii KF *et al.* CD44 stimulation induces integrin-mediated adhesion of colon cancer cell lines to endothelial cells by up-regulation of integrins and c-Met and activation of integrins. *Cancer Res*

1999; 59(17):4427-34.

- [176] Xu YF, Yu Q. E-cadherin negatively regulates CD44-hyaluronan interaction and CD44-mediated tumor invasion and branching morphogenesis. *J Biol Chem* 2003; 278(10):8661-8.
- [177] Sneath RJ, Mangham DC. The normal structure and function of CD44 and its role in neoplasia. *Mol Pathol* 1998; 51(4):191-200.
- [178] Lokeshwar VB, Iida N, Bourguignon LY. The cell adhesion molecule, GP116, is a new CD44 variant (ex14/v10) involved in hyaluronic acid binding and endothelial cell proliferation. *J Biol Chem* 1996; 271(39):23853-64.
- [179] Turley EA, Austen L, Vandeligt K, Clary C. Hyaluronan and a cell-associated hyaluronan binding protein regulate the locomotion of ras-transformed cells. *J Cell Biol* 1991; 112(5):1041-7.
- [180] Tanabe KK, Saya H. The CD44 adhesion molecule and metastasis. *Crit Rev Oncog* 1994; 5(2-3):201-12.
- [181] Korell M. Modern adhesion prophylaxis in gynecological operations. *Gynäkologe* 2002; 35:1218-23.
- [182] See WA, Chapman PH. Heparin prevention of tumor cell adherence and implantation on injured urothelial surfaces. *J Urol* 1987; 138(1):182-6.
- [183] Loo BM, Kreuger J, Jalkanen M, Lindahl U, Salmivirta M. Binding of heparin/heparan sulfate to fibroblast growth factor receptor 4. *J Biol Chem* 2001; 276(20):16868-76.
- [184] Dudas J, Ramadori G, Knittel T *et al.* Effect of heparin and liver heparan sulphate on interaction of HepG2-derived transcription factors and their cis-acting elements: altered potential of hepatocellular carcinoma heparan sulphate. *Biochem J* 2000; 350 Pt 1:245-51.
- [185] Watson RW, Redmond HP, Mc CJ, Bouchier-Hayes D. Taurolidine, an antilipopolysaccharide agent, has immunoregulatory properties that are mediated by the amino acid taurine. *J Leukoc Biol* 1995; 58(3):299-306.

- [186] Pidgeon GP, Harmey JH, Kay E *et al.* The role of endotoxin/lipopolysaccharide in surgically induced tumour growth in a murine model of metastatic disease. *Br J Cancer* 1999; 81(8):1311-7.
- [187] Kerbel RS. Tumor angiogenesis: past, present and the near future. *Carcinogenesis* 2000; 21(3):505-15.
- [188] Norrby K. Angiogenesis: new aspects relating to its initiation and control. *APMIS* 1997; 105(6):417-37.
- [189] Wordemann M, Fandrey J, Jelkmann W. Tumor necrosis factor-alpha production by human hepatoma cell lines is resistant to drugs that are inhibitory to macrophages. *J Interferon Cytokine Res* 1998; 18(12):1069-75.
- [190] Eigler A, Loher F, Endres S. [Suppression of synthesis of tumor necrosis factor]. *Internist (Berl)* 2001; 42(1):28-34.
- [191] Van Antwerp DJ, Martin SJ, Verma IM, Green DR. Inhibition of TNF-induced apoptosis by NF-kappa B. *Trends Cell Biol* 1998; 8(3):107-11.
- [192] Stendel R, Stoltenburg-Didinger G, Al Keikh CL, Wattrodt M, Brock M. The effect of taurolidine on brain tumor cells. *Anticancer Res* 2002; 22(2A):809-14.
- [193] Harrington KJ, Syrigos KN, Harrington KJ. The role of E-cadherin-catenin complex: more than an intercellular glue? *Ann Surg Oncol* 2000; 7(10):783-8.
- [194] Birchmeier WF, Hulskens JF, Behrens J. Adherens junction proteins in tumour progression. *Cancer Surv* 1995; 24:129-40.
- [195] Wijnhoven BP, Dinjens WN, Pignatelli M. E-cadherin-catenin cell-cell adhesion complex and human cancer. *Br J Surg* 2000; 87(8):992-1005.
- [196] Dorudi S, Sheffield JP, Poulsom R, Northover JM, Hart IR. E-cadherin expression in colorectal cancer. An immunocytochemical and in situ hybridization study. *Am J Pathol* 1993; 142(4):981-6.
- [197] Nigam AK, Savage FJ, Boulos PB *et al.* Loss of cell-cell and cell-matrix adhesion molecules in colorectal cancer. *Br J Cancer* 1993; 68(3):507-14.

- [198] Shiozaki HF, Oka HF, Inoue MF, Tamura S, Monden M. E-cadherin mediated adhesion system in cancer cells. *Cancer* 1996; 77(8 Suppl):1605-13.
- [199] Ikeguchi MF, Makino MF, Kaibara N. Clinical significance of E-cadherin-catenin complex expression in metastatic foci of colorectal carcinoma. *J Surg Oncol* 2001; 77(3):201-7.
- [200] Keller SH, Nigam SK. Biochemical processing of E-cadherin under cellular stress. *Biochem Biophys Res Commun* 2003; 307(2):215-23.
- [201] Steinhusen U, Weiske J, Badock V *et al.* Cleavage and shedding of E-cadherin after induction of apoptosis. *J Biol Chem* 2001; 276(7):4972-80.
- [202] Choi SH, Takahashi K, Eto H, Yoon SS, Tanabe KK. CD44s expression in human colon carcinomas influences growth of liver metastases. *Int J Cancer* 2000; 85(4):523-6.
- [203] Streit MF, Schmidt RF, Hilgenfeld RU, Thiel E, Kreuser ED. Adhesion receptors in malignant transformation and dissemination of gastrointestinal tumors. *Recent Results Cancer Res* 1996; 142:19-50.
- [204] Washington K, Gottfried MR, Telen MJ. Expression of the cell adhesion molecule CD44 in gastric adenocarcinomas. *Hum Pathol* 1994; 25(10):1043-9.
- [205] Yamaguchi A, Urano T, Goi T *et al.* Expression of a CD44 variant containing exons 8 to 10 is a useful independent factor for the prediction of prognosis in colorectal cancer patients. *J Clin Oncol* 1996; 14(4):1122-7.
- [206] Dall P, Heider KH, Sinn HP *et al.* Comparison of immunohistochemistry and RT-PCR for detection of CD44v-expression, a new prognostic factor in human breast cancer. *Int J Cancer* 1995; 60(4):471-7.
- [207] Kaufmann M, Heider KH, Sinn HP *et al.* CD44 variant exon epitopes in primary breast cancer and length of survival. *Lancet* 1995; 345(8950):615-9.
- [208] Seiter S, Arch R, Reber S *et al.* Prevention of tumor metastasis formation by anti-variant CD44. *J Exp Med* 1993; 177(2):443-55.

- [209] Trochon V, Mabilat C, Bertrand P *et al.* Evidence of involvement of CD44 in endothelial cell proliferation, migration and angiogenesis in vitro. *Int J Cancer* 1996; 66(5):664-8.
- [210] Barshishat M, Ariel A, Cahalon L *et al.* TNFalpha and IL-8 regulate the expression and function of CD44 variant proteins in human colon carcinoma cells. *Clin Exp Metastasis* 2002; 19(4):327-37.
- [211] Agrez MV, Bates RC. Colorectal cancer and the integrin family of cell adhesion receptors: current status and future directions. *Eur J Cancer* 1994; 30A(14):2166-70.
- [212] Schaller MD, Parsons JT. Focal adhesion kinase and associated proteins. *Curr Opin Cell Biol* 1994; 6(5):705-10.
- [213] Christofori G. Changing neighbours, changing behaviour: cell adhesion molecule-mediated signalling during tumour progression. *EMBO J* 2003; 22(10):2318-23.
- [214] Chan BM, Matsuura N, Takada Y, Zetter BR, Hemler ME. In vitro and in vivo consequences of VLA-2 expression on rhabdomyosarcoma cells. *Science* 1991; 251(5001):1600-2.
- [215] Zutter MM, Santoro SA, Staatz WD, Tsung YL. Re-expression of the alpha 2 beta 1 integrin abrogates the malignant phenotype of breast carcinoma cells. *Proc Natl Acad Sci U S A* 1995; 92(16):7411-5.
- [216] Hangan D, Uniyal S, Morris VL *et al.* Integrin VLA-2 (alpha2beta1) function in postextravasation movement of human rhabdomyosarcoma RD cells in the liver. *Cancer Res* 1996; 56(13):3142-9.
- [217] Varner JA, Emerson DA, Juliano RL. Integrin alpha 5 beta 1 expression negatively regulates cell growth: reversal by attachment to fibronectin. *Mol Biol Cell* 1995; 6(6):725-40.
- [218] Zhang Z, Vuori K, Reed JC, Ruoslahti E. The alpha 5 beta 1 integrin supports survival of cells on fibronectin and up-regulates Bcl-2 expression. *Proc Natl Acad Sci U S A* 1995; 92(13):6161-5.

- [219] Koukoulis GK FAU, Patriarca CF, Gould VE. Adhesion molecules and tumor metastasis. *Hum Pathol* 1998;(9):889-92.
- [220] Stallmach A, von Lampe B, Orzechowski HD, Matthes H, Riecken EO. Increased fibronectin-receptor expression in colon carcinoma-derived HT 29 cells decreases tumorigenicity in nude mice. *Gastroenterology* 1994; 106(1):19-27.
- [221] Juliano RL. The role of beta 1 integrins in tumors. *Semin Cancer Biol* 1993; 4(5):277-83.
- [222] Wang JH, Manning BJ, Wu QD *et al.* Endotoxin/lipopolysaccharide activates NF-kappa B and enhances tumor cell adhesion and invasion through a beta 1 integrin-dependent mechanism. *J Immunol* 2003; 170(2):795-804.
- [223] Wu JS, Brasfield EB, Guo LW *et al.* Implantation of colon cancer at trocar sites is increased by low pressure pneumoperitoneum. *Surgery* 1997; 122(1):1-7.
- [224] Rosman C, Westerveld GJ, van Oeveren W, Kooi K, Bleichrodt RP. Effect of intraperitoneal antimicrobials on the concentration of bacteria, endotoxin, and tumor necrosis factor in abdominal fluid and plasma in rats. *Eur Surg Res* 1996; 28(5):351-60.