

8. Literatur

- Agell,N., Jaumot,M., Rodriguez-Vilarrupla,A., Brun,S., Abella,N., Canela,N., Estanyol,J.M., and Bachs,O. (2006). The Diverging Roles of Calmodulin and PKC in the Regulation of p21 Intracellular Localization. *Cell Cycle*, **5**, 3-6.
- Aho,T.L., Sandholm,J., Peltola,K.J., Mankonen,H.P., Lilly,M., and Koskinen,P.J. (2004). Pim-1 kinase promotes inactivation of the pro-apoptotic Bad protein by phosphorylating it on the Ser112 gatekeeper site. *FEBS Lett.*, **571**, 43-49.
- Asada,M., Yamada,T., Ichijo,H., Delia,D., Miyazono,K., Fukumuro,K., and Mizutani,S. (1999). Apoptosis inhibitory activity of cytoplasmic p21(Cip1/WAF1) in monocytic differentiation. *EMBO J.*, **18**, 1223-1234.
- Bachmann,M., Hennemann,H., Xing,P.X., Hoffmann,I., and Moroy,T. (2004). The oncogenic serine/threonine kinase Pim-1 phosphorylates and inhibits the activity of Cdc25C-associated kinase 1 (C-TAK1): a novel role for Pim-1 at the G2/M cell cycle checkpoint. *J. Biol. Chem.*, **279**, 48319-48328.
- Bachmann,M., Kosan,C., Xing,P.X., Montenarh,M., Hoffmann,I., and Moroy,T. (2006). The oncogenic serine/threonine kinase Pim-1 directly phosphorylates and activates the G2/M specific phosphatase Cdc25C. *Int. J. Biochem. Cell Biol.*, **38**, 430-443.
- Bachmann,M. and Moroy,T. (2005). The serine/threonine kinase Pim-1. *Int. J. Biochem. Cell Biol.*, **37**, 726-730.
- Bar-Nun,S., Shneyour,Y., and Beckmann,J.S. (1983). G-418, an elongation inhibitor of 80 S ribosomes. *Biochim. Biophys. Acta*, **741**, 123-127.
- Beaudouin,J., Gerlich,D., Daigle,N., Eils,R., and Ellenberg,J. (2002). Nuclear envelope breakdown proceeds by microtubule-induced tearing of the lamina. *Cell*, **108**, 83-96.

- Bhattacharya,N., Wang,Z., Davitt,C., McKenzie,I.F., Xing,P.X., and Magnuson,N.S. (2002). Pim-1 associates with protein complexes necessary for mitosis. *Chromosoma*, **111**, 80-95.
- Birnboim,H.C. and Doly,J. (1979). A rapid alkaline extraction procedure for screening recombinant plasmid DNA. *Nucleic Acids Res.*, **7**, 1513-1523.
- Bissonnette,N. and Hunting,D.J. (1998). p21-induced cycle arrest in G1 protects cells from apoptosis induced by UV-irradiation or RNA polymerase II blockage. *Oncogene*, **16**, 3461-3469.
- Blagosklonny,M.V. (2002). Are p27 and p21 cytoplasmic oncoproteins? *Cell Cycle*, **1**, 391-393.
- Blake,M.S., Johnston,K.H., Russell-Jones,G.J., and Gotschlich,E.C. (1984). A rapid, sensitive method for detection of alkaline phosphatase-conjugated anti-antibody on Western blots. *Anal. Biochem.*, **136**, 175-179.
- Booher,R.N., Holman,P.S., and Fattaey,A. (1997). Human Myt1 is a cell cycle-regulated kinase that inhibits Cdc2 but not Cdk2 activity. *J. Biol. Chem.*, **272**, 22300-22306.
- Bullock,A.N., Debreczeni,J., Amos,A.L., Knapp,S., and Turk,B.E. (2005). Structure and substrate specificity of the Pim-1 kinase. *J. Biol. Chem.*, **280**, 41675-41682.
- Chalfie,M., Tu,Y., Euskirchen,G., Ward,W.W., and Prasher,D.C. (1994). Green fluorescent protein as a marker for gene expression. *Science*, **263**, 802-805.
- Child,E.S. and Mann,D.J. (2006). The Intricacies of p21 Phosphorylation: Protein/Protein Interactions, Subcellular Localization and Stability. *Cell Cycle*, **5**.
- Clute,P. and Pines,J. (1999). Temporal and spatial control of cyclin B1 destruction in metaphase. *Nat. Cell Biol.*, **1**, 82-87.
- Coqueret,O. (2003). New roles for p21 and p27 cell-cycle inhibitors: a function for each cell compartment? *Trends Cell Biol.*, **13**, 65-70.

- Cuypers,H.T., Selten,G., Quint,W., Zijlstra,M., Maandag,E.R., Boelens,W., van Wezenbeek,P., Melief,C., and Berns,A. (1984). Murine leukemia virus-induced T-cell lymphomagenesis: integration of proviruses in a distinct chromosomal region. *Cell*, **37**, 141-150.
- Dash,B.C. and El Deiry,W.S. (2005). Phosphorylation of p21 in G2/M promotes cyclin B-Cdc2 kinase activity. *Mol. Cell Biol.*, **25**, 3364-3387.
- Delavaine,L. and La Thangue,N.B. (1999). Control of E2F activity by p21Waf1/Cip1. *Oncogene*, **18**, 5381-5392.
- Dhanasekaran,S.M., Barrette,T.R., Ghosh,D., Shah,R., Varambally,S., Kurachi,K., Pienta,K.J., Rubin,M.A., and Chinnaiyan,A.M. (2001). Delineation of prognostic biomarkers in prostate cancer. *Nature*, **412**, 822-826.
- Dotto,G.P. (2000). p21(WAF1/Cip1): more than a break to the cell cycle? *Biochim. Biophys. Acta*, **1471**, M43-M56.
- Downs,T.R. and Wilfinger,W.W. (1983). Fluorometric quantification of DNA in cells and tissue. *Anal. Biochem.*, **131**, 538-547.
- Doxsey,S., Zimmerman,W., and Mikule,K. (2005). Centrosome control of the cell cycle. *Trends Cell Biol.*, **15**, 303-311.
- Dyson,N. (1998). The regulation of E2F by pRB-family proteins. *Genes Dev.*, **12**, 2245-2262.
- Edamatsu,M. (2001). The molecular mechanism of targeted vesicle transport in cytokinesis. *Cell Struct. Funct.*, **26**, 567-570.
- Fattaey,A. and Booher,R.N. (1997). Myt1: a Wee1-type kinase that phosphorylates Cdc2 on residue Thr14. *Prog. Cell Cycle Res.*, **3**, 233-240.
- Felgner,P.L., Gadek,T.R., Holm,M., Roman,R., Chan,H.W., Wenz,M., Northrop,J.P., Ringold,G.M., and Danielsen,M. (1987). Lipofection: a highly efficient, lipid-

- mediated DNA-transfection procedure. *Proc. Natl. Acad. Sci. U. S. A.*, **84**, 7413-7417.
- Firestone,G.L. and Winguth,S.D. (1990). Immunoprecipitation of proteins. *Methods Enzymol.*, **182**, 688-700.
- Friedmann,M., Nissen,M.S., Hoover,D.S., Reeves,R., and Magnuson,N.S. (1992). Characterization of the proto-oncogene pim-1: kinase activity and substrate recognition sequence. *Arch. Biochem. Biophys.*, **298**, 594-601.
- Gartel,A.L. and Radhakrishnan,S.K. (2005). Lost in transcription: p21 repression, mechanisms, and consequences. *Cancer Res.*, **65**, 3980-3985.
- Gartel,A.L. and Tyner,A.L. (1999). Transcriptional regulation of the p21((WAF1/CIP1)) gene. *Exp. Cell Res.*, **246**, 280-289.
- Girard,F., Fernandez,A., and Lamb,N. (1995). Delayed cyclin A and B1 degradation in non-transformed mammalian cells. *J. Cell Sci.*, **108 (Pt 7)**, 2599-2608.
- Gomez-Lazaro,M., Fernandez-Gomez,F.J., and Jordan,J. (2004). [The role of protein p53 in neurodegenerative processes throughout the 25 years of its history]. *Rev. Neurol.*, **39**, 243-250.
- Gulbis,J.M., Kelman,Z., Hurwitz,J., O'Donnell,M., and Kuriyan,J. (1996). Structure of the C-terminal region of p21(WAF1/CIP1) complexed with human PCNA. *Cell*, **87**, 297-306.
- Hannon,G.J. (2002). RNA interference. *Nature*, **418**, 244-251.
- Harper,J.W. and Elledge,S.J. (1996). Cdk inhibitors in development and cancer. *Curr. Opin. Genet. Dev.*, **6**, 56-64.
- Hoefnagel,J.J., Dijkman,R., Basso,K., Jansen,P.M., Hallermann,C., Willemze,R., Tensen,C.P., and Vermeer,M.H. (2005). Distinct types of primary cutaneous large B-cell lymphoma identified by gene expression profiling. *Blood*, **105**, 3671-3678.

- Ionov,Y., Le,X., Tunquist,B.J., Sweetenham,J., Sachs,T., Ryder,J., Johnson,T., Lilly,M.B., and Kraft,A.S. (2003). Pim-1 protein kinase is nuclear in Burkitt's lymphoma: nuclear localization is necessary for its biologic effects. *Anticancer Res.*, **23**, 167-178.
- Kaldis,P. (1999). The cdk-activating kinase (CAK): from yeast to mammals. *Cell Mol. Life Sci.*, **55**, 284-296.
- Kaneuchi,M., Yamashita,T., Shindoh,M., Segawa,K., Takahashi,S., Furuta,I., Fujimoto,S., and Fujinaga,K. (1999). Induction of apoptosis by the p53-273L (Arg -> Leu) mutant in HSC3 cells without transactivation of p21Waf1/Cip1/Sdi1 and bax. *Mol. Carcinog.*, **26**, 44-52.
- Katayama,K., Fujita,N., and Tsuruo,T. (2005). Akt/protein kinase B-dependent phosphorylation and inactivation of WEE1Hu promote cell cycle progression at G2/M transition. *Mol. Cell Biol.*, **25**, 5725-5737.
- Kelly,T.J. and Brown,G.W. (2000). Regulation of chromosome replication. *Annu. Rev. Biochem.*, **69**, 829-880.
- King,R.W., Glotzer,M., and Kirschner,M.W. (1996). Mutagenic analysis of the destruction signal of mitotic cyclins and structural characterization of ubiquitinated intermediates. *Mol. Biol. Cell*, **7**, 1343-1357.
- Konietzko,U., Kauselmann,G., Scafidi,J., Staubli,U., Mikkers,H., Berns,A., Schweizer,M., Waltereit,R., and Kuhl,D. (1999). Pim kinase expression is induced by LTP stimulation and required for the consolidation of enduring LTP. *EMBO J.*, **18**, 3359-3369.
- Laskey,R.A. (1980). The use of intensifying screens or organic scintillators for visualizing radioactive molecules resolved by gel electrophoresis. *Methods Enzymol.*, **65**, 363-371.
- Lei,M. and Tye,B.K. (2001). Initiating DNA synthesis: from recruiting to activating the MCM complex. *J. Cell Sci.*, **114**, 1447-1454.

- Leone,G., DeGregori,J., Sears,R., Jakoi,L., and Nevins,J.R. (1997). Myc and Ras collaborate in inducing accumulation of active cyclin E/Cdk2 and E2F. *Nature*, **387**, 422-426.
- Li,Y., Dowbenko,D., and Lasky,L.A. (2002). AKT/PKB phosphorylation of p21Cip/WAF1 enhances protein stability of p21Cip/WAF1 and promotes cell survival. *J. Biol. Chem.*, **277**, 11352-11361.
- Liang,H., Hittelman,W., and Nagarajan,L. (1996). Ubiquitous expression and cell cycle regulation of the protein kinase PIM-1. *Arch. Biochem. Biophys.*, **330**, 259-265.
- Liang,J. and Slingerland,J.M. (2003). Multiple roles of the PI3K/PKB (Akt) pathway in cell cycle progression. *Cell Cycle*, **2**, 339-345.
- Lilly,M., Sandholm,J., Cooper,J.J., Koskinen,P.J., and Kraft,A. (1999). The PIM-1 serine kinase prolongs survival and inhibits apoptosis-related mitochondrial dysfunction in part through a bcl-2-dependent pathway. *Oncogene*, **18**, 4022-4031.
- Lipinski,M.M. and Jacks,T. (1999). The retinoblastoma gene family in differentiation and development. *Oncogene*, **18**, 7873-7882.
- Lundgren,K., Walworth,N., Booher,R., Dembski,M., Kirschner,M., and Beach,D. (1991). mik1 and wee1 cooperate in the inhibitory tyrosine phosphorylation of cdc2. *Cell*, **64**, 1111-1122.
- Maller,J., Gautier,J., Langan,T.A., Lohka,M.J., Shenoy,S., Shalloway,D., and Nurse,P. (1989). Maturation-promoting factor and the regulation of the cell cycle. *J. Cell Sci. Suppl*, **12**, 53-63.
- Malumbres,M. and Barbacid,M. (2005). Mammalian cyclin-dependent kinases. *Trends Biochem. Sci.*, **30**, 630-641.
- Massague,J. (2004). G1 cell-cycle control and cancer. *Nature*, **432**, 298-306.

- Matsumoto,Y. (2001). Molecular mechanism of PCNA-dependent base excision repair. *Prog. Nucleic Acid Res. Mol. Biol.*, **68**, 129-138.
- Merdes,A., Heald,R., Samejima,K., Earnshaw,W.C., and Cleveland,D.W. (2000). Formation of spindle poles by dynein/dynactin-dependent transport of NuMA. *J. Cell Biol.*, **149**, 851-862.
- Mochizuki,T., Kitanaka,C., Noguchi,K., Muramatsu,T., Asai,A., and Kuchino,Y. (1999). Physical and functional interactions between Pim-1 kinase and Cdc25A phosphatase. Implications for the Pim-1-mediated activation of the c-Myc signaling pathway. *J. Biol. Chem.*, **274**, 18659-18666.
- Morgan,D.O. (1997). Cyclin-dependent kinases: engines, clocks, and microprocessors. *Annu. Rev. Cell Dev. Biol.*, **13**, 261-291.
- Morris,E.J. and Dyson,N.J. (2001). Retinoblastoma protein partners. *Adv. Cancer Res.*, **82**, 1-54.
- Moustakas,A., Pardali,K., Gaal,A., and Heldin,C.H. (2002). Mechanisms of TGF-beta signaling in regulation of cell growth and differentiation. *Immunol. Lett.*, **82**, 85-91.
- Nilsson,I. and Hoffmann,I. (2000). Cell cycle regulation by the Cdc25 phosphatase family. *Prog. Cell Cycle Res.*, **4**, 107-114.
- Obata,T., Yaffe,M.B., Leparc,G.G., Piro,E.T., Maegawa,H., Kashiwagi,A., Kikkawa,R., and Cantley,L.C. (2000). Peptide and protein library screening defines optimal substrate motifs for AKT/PKB. *J. Biol. Chem.*, **275**, 36108-36115.
- Ohata,M., Nakamura,S., Fujita,H., and Isemura,M. (2005). Prognostic implications of p21 (Waf1/Cip1) immunolocalization in multiple myeloma. *Biomed. Res.*, **26**, 91-98.
- Okaichi,K., Wang,L.H., Sasaki,J., Saya,H., Tada,M., and Okumura,Y. (1999). A point mutation of human p53, which was not detected as a mutation by a yeast functional assay, led to apoptosis but not p21Waf1/Cip1/Sdi1 expression in

- response to ionizing radiation in a human osteosarcoma cell line, Saos-2. *Int. J. Radiat. Oncol. Biol. Phys.*, **45**, 975-980.
- Pardali,K., Kowanetz,M., Heldin,C.H., and Moustakas,A. (2005). Smad pathway-specific transcriptional regulation of the cell cycle inhibitor p21(WAF1/Cip1). *J. Cell Physiol*, **204**, 260-272.
- Pardali,K., Kurisaki,A., Moren,A., ten Dijke,P., Kardassis,D., and Moustakas,A. (2000). Role of Smad proteins and transcription factor Sp1 in p21(Waf1/Cip1) regulation by transforming growth factor-beta. *J. Biol. Chem.*, **275**, 29244-29256.
- Pavletich,N.P. (1999). Mechanisms of cyclin-dependent kinase regulation: structures of Cdks, their cyclin activators, and Cip and INK4 inhibitors. *J. Mol. Biol.*, **287**, 821-828.
- Pepper,D.S., Upton,T.M., Ladha,M.H., Neuman,E., Zalvide,J., Bernards,R., DeCaprio,J.A., and Ewen,M.E. (1997). Ras signalling linked to the cell-cycle machinery by the retinoblastoma protein. *Nature*, **386**, 177-181.
- Pei,X.H. and Xiong,Y. (2005). Biochemical and cellular mechanisms of mammalian CDK inhibitors: a few unresolved issues. *Oncogene*, **24**, 2787-2795.
- Prasher,D.C. (1995). Using GFP to see the light. *Trends Genet.*, **11**, 320-323.
- Prasher,D.C., Eckenrode,V.K., Ward,W.W., Prendergast,F.G., and Cormier,M.J. (1992). Primary structure of the Aequorea victoria green-fluorescent protein. *Gene*, **111**, 229-233.
- Qian,K.C., Wang,L., Hickey,E.R., Studts,J., Barringer,K., Peng,C., Kronkaitis,A., Li,J., White,A., Mische,S., and Farmer,B. (2005). Structural basis of constitutive activity and a unique nucleotide binding mode of human Pim-1 kinase. *J. Biol. Chem.*, **280**, 6130-6137.
- Rodriguez-Vilarrupla,A., Jaumot,M., Abella,N., Canela,N., Brun,S., Diaz,C., Estanyol,J.M., Bachs,O., and Agell,N. (2005). Binding of calmodulin to the carboxy-terminal region of p21 induces nuclear accumulation via inhibition of

- protein kinase C-mediated phosphorylation of Ser153. *Mol. Cell Biol.*, **25**, 7364-7374.
- Rossig,L., Jadidi,A.S., Urbich,C., Badorff,C., Zeiher,A.M., and Dimmeler,S. (2001). Akt-dependent phosphorylation of p21(Cip1) regulates PCNA binding and proliferation of endothelial cells. *Mol. Cell Biol.*, **21**, 5644-5657.
- Sancar,A., Lindsey-Boltz,L.A., Unsal-Kacmaz,K., and Linn,S. (2004). Molecular mechanisms of mammalian DNA repair and the DNA damage checkpoints. *Annu. Rev. Biochem.*, **73**, 39-85.
- Saramaki,A., Banwell,C.M., Campbell,M.J., and Carlberg,C. (2006). Regulation of the human p21(waf1/cip1) gene promoter via multiple binding sites for p53 and the vitamin D3 receptor. *Nucleic Acids Res.*, **34**, 543-554.
- Scott,M.T., Morrice,N., and Ball,K.L. (2000). Reversible phosphorylation at the C-terminal regulatory domain of p21(Waf1/Cip1) modulates proliferating cell nuclear antigen binding. *J. Biol. Chem.*, **275**, 11529-11537.
- Selten,G., Cuypers,H.T., and Berns,A. (1985). Proviral activation of the putative oncogene Pim-1 in MuLV induced T-cell lymphomas. *EMBO J.*, **4**, 1793-1798.
- Smits,V.A. and Medema,R.H. (2001). Checking out the G(2)/M transition. *Biochim. Biophys. Acta*, **1519**, 1-12.
- Sparks,C.A., Fey,E.G., Vidair,C.A., and Doxsey,S.J. (1995). Phosphorylation of NUMA occurs during nuclear breakdown and not mitotic spindle assembly. *J. Cell Sci.*, **108 (Pt 11)**, 3389-3396.
- Srinivas,P.R., Srivastava,S., Hanash,S., and Wright,G.L., Jr. (2001). Proteomics in early detection of cancer. *Clin. Chem.*, **47**, 1901-1911.
- Srinivas,P.R., Verma,M., Zhao,Y., and Srivastava,S. (2002). Proteomics for cancer biomarker discovery. *Clin. Chem.*, **48**, 1160-1169.

- Stark,G.R. and Taylor,W.R. (2004). Analyzing the G2/M checkpoint. *Methods Mol. Biol.*, **280**, 51-82.
- Stevenson-Lindert,L.M., Fowler,P., and Lew,J. (2003). Substrate specificity of CDK2-cyclin A. What is optimal? *J. Biol. Chem.*, **278**, 50956-50960.
- Stout,B.A., Bates,M.E., Liu,L.Y., Farrington,N.N., and Bertics,P.J. (2004). IL-5 and granulocyte-macrophage colony-stimulating factor activate STAT3 and STAT5 and promote Pim-1 and cyclin D3 protein expression in human eosinophils. *J. Immunol.*, **173**, 6409-6417.
- Suzuki,A., Kawano,H., Hayashida,M., Hayasaki,Y., Tsutomi,Y., and Akahane,K. (2000). Procaspsase 3/p21 complex formation to resist fas-mediated cell death is initiated as a result of the phosphorylation of p21 by protein kinase A. *Cell Death. Differ.*, **7**, 721-728.
- Suzuki,A., Tsutomi,Y., Yamamoto,N., Shibutani,T., and Akahane,K. (1999). Mitochondrial regulation of cell death: mitochondria are essential for procaspsase 3-p21 complex formation to resist Fas-mediated cell death. *Mol. Cell Biol.*, **19**, 3842-3847.
- Tsurimoto,T. (1999). PCNA binding proteins. *Front Biosci.*, **4**, D849-D858.
- Ubersax,J.A., Woodbury,E.L., Quang,P.N., Paraz,M., Blethrow,J.D., Shah,K., Shokat,K.M., and Morgan,D.O. (2003). Targets of the cyclin-dependent kinase Cdk1. *Nature*, **425**, 859-864.
- Valdman,A., Fang,X., Pang,S.T., Ekman,P., and Egevad,L. (2004). Pim-1 expression in prostatic intraepithelial neoplasia and human prostate cancer. *Prostate*, **60**, 367-371.
- Waga,S., Hannon,G.J., Beach,D., and Stillman,B. (1994). The p21 inhibitor of cyclin-dependent kinases controls DNA replication by interaction with PCNA. *Nature*, **369**, 574-578.

- Wagman,G.H., Testa,R.T., Marquez,J.A., and Weinstein,M.J. (1974). Antibiotic G-418, a new Micromonospora-produced aminoglycoside with activity against protozoa and helminths: fermentation, isolation, and preliminary characterization. *Antimicrob. Agents Chemother.*, **6**, 144-149.
- Walworth,N.C. (2001). DNA damage: Chk1 and Cdc25, more than meets the eye. *Curr. Opin. Genet. Dev.*, **11**, 78-82.
- Wang,Z., Bhattacharya,N., Mixter,P.F., Wei,W., Sedivy,J., and Magnuson,N.S. (2002). Phosphorylation of the cell cycle inhibitor p21Cip1/WAF1 by Pim-1 kinase. *Biochim. Biophys. Acta*, **1593**, 45-55.
- Wang,Z., Bhattacharya,N., Weaver,M., Petersen,K., Meyer,M., Gapter,L., and Magnuson,N.S. (2001). Pim-1: a serine/threonine kinase with a role in cell survival, proliferation, differentiation and tumorigenesis. *J. Vet. Sci.*, **2**, 167-179.
- Warbrick,E. (1998). PCNA binding through a conserved motif. *Bioessays*, **20**, 195-199.
- Warbrick,E. (2000). The puzzle of PCNA's many partners. *Bioessays*, **22**, 997-1006.
- Winters,Z.E., Hunt,N.C., Bradburn,M.J., Royds,J.A., Turley,H., Harris,A.L., and Norbury,C.J. (2001). Subcellular localisation of cyclin B, Cdc2 and p21(WAF1/CIP1) in breast cancer. association with prognosis. *Eur. J. Cancer*, **37**, 2405-2412.
- Woo,R.A. and Poon,R.Y. (2003). Cyclin-dependent kinases and S phase control in mammalian cells. *Cell Cycle*, **2**, 316-324.
- Xia,W., Chen,J.S., Zhou,X., Sun,P.R., Lee,D.F., Liao,Y., Zhou,B.P., and Hung,M.C. (2004). Phosphorylation/cytoplasmic localization of p21Cip1/WAF1 is associated with HER2/neu overexpression and provides a novel combination predictor for poor prognosis in breast cancer patients. *Clin. Cancer Res.*, **10**, 3815-3824.
- Xie,Y., Xu,K., Dai,B., Guo,Z., Jiang,T., Chen,H., and Qiu,Y. (2006). The 44 kDa Pim-1 kinase directly interacts with tyrosine kinase Etk/BMX and protects human

- prostate cancer cells from apoptosis induced by chemotherapeutic drugs. *Oncogene*, **25**, 70-78.
- Xu,Y., Zhang,T., Tang,H., Zhang,S., Liu,M., Ren,D., and Niu,Y. (2005). Overexpression of PIM-1 is a potential biomarker in prostate carcinoma. *J. Surg. Oncol.*, **92**, 326-330.
- Yan,B., Zemskova,M., Holder,S., Chin,V., Kraft,A., Koskinen,P.J., and Lilly,M. (2003). The PIM-2 kinase phosphorylates BAD on serine 112 and reverses BAD-induced cell death. *J. Biol. Chem.*, **278**, 45358-45367.
- Yanai,I., Benjamin,H., Shmoish,M., Chalifa-Caspi,V., Shklar,M., Ophir,R., Bar-Even,A., Horn-Saban,S., Safran,M., Domany,E., Lancet,D., and Shmueli,O. (2005). Genome-wide midrange transcription profiles reveal expression level relationships in human tissue specification. *Bioinformatics*, **21**, 650-659.
- Zhivotovsky,B., Burgess,D.H., Schlegel,J., Porn,M.I., Vanags,D., and Orrenius,S. (1997). Proteases in Fas-mediated apoptosis. *J. Cell Biochem*, **64**, 43-49.
- Zhou,B.P., Liao,Y., Xia,W., Spohn,B., Lee,M.H., and Hung,M.C. (2001). Cytoplasmic localization of p21Cip1/WAF1 by Akt-induced phosphorylation in HER-2/neu-overexpressing cells. *Nat. Cell Biol.*, **3**, 245-252.
- Zhu,L. (2005). Tumour suppressor retinoblastoma protein Rb: a transcriptional regulator. *Eur. J. Cancer*, **41**, 2415-2427.