

## 6. Literaturverzeichnis

- Adler, A.I., Stevens, R.J., Manley, S.E., Bilous, R.W., Cull, C.A. und Holman, R.R. (2003). "Development and progression of nephropathy in type 2 diabetes: the United Kingdom Prospective Diabetes Study (UKPDS 64)." Kidney Int **63**(1): 225-32.
- Agrawal, B., Berger, A., Wolf, K. und Luft, F.C. (1996). "Microalbuminuria screening by reagent strip predicts cardiovascular risk in hypertension." J Hypertens **14**(2): 223-8.
- Ahmad, J., Siddiqui, M.A. und Ahmad, H. (1997). "Effective postponement of diabetic nephropathy with enalapril in normotensive type 2 diabetic patients with microalbuminuria." Diabetes Care **20**(10): 1576-81.
- Alvestrand, A., Gutierrez, A., Bucht, H. und Bergstrom, J. (1988). "Reduction of blood pressure retards the progression of chronic renal failure in man." Nephrol Dial Transplant **3**(5): 624-31.
- BAYER (2003). "Microalbustix<sup>TM</sup> - Reagent Strips for Determining Albumin and Creatinine in Urine." Gebrauchsanleitung Microalbustix<sup>TM</sup>.
- Bender, R. und Lange, S. (2001). "Die Vierfeldertafel." DMW **126**(15): T36-T38.
- Berson, S.A., Yalow, R.S., Bauman, A., Rothschild, M.A. und Newerly, K. (1956). "Insulin-I131 metabolism in human subjects: demonstration of insulin binding globulin in the circulation of insulin treated subjects." J Clin Invest **35**(2): 170-90.
- Bianchi, S., Bigazzi, R., Caiazza, A. und Campese, V.M. (2003). "A controlled, prospective study of the effects of atorvastatin on proteinuria and progression of kidney disease." Am J Kidney Dis **41**(3): 565-70.
- Bilous, R. (1995). "Diabetes: should you be screening for microalbuminuria?" Practitioner **239**(1550): 343-5.
- Boero, R., Prodi, E., Elia, F., Porta, L., Martelli, S., Ferraro, L. und Quarello, F. (2003). "How well are hypertension and albuminuria treated in type II diabetic patients?" J Hum Hypertens **17**(6): 413-8.

Bohrer, M.P., Baylis, C., Humes, H.D., Glasscock, R.J., Robertson, C.R. und Brenner, B.M. (1978). "Permselectivity of the glomerular capillary wall. Facilitated filtration of circulating polycations." J Clin Invest **61**(1): 72-8.

Borch-Johnsen, K., Nissen, H., Salling, N., Henriksen, E., Kreiner, S., Deckert, T. und Nerup, J. (1987). "The natural history of insulin-dependent diabetes in Denmark: 2. Long-term survival--who and why." Diabet Med **4**(3): 211-6.

Bramlage, P., Wittchen, H.U., Pittrow, D., Dikow, R., Kirch, W., Lehnert, H. und Ritz, E. (2003). "[Microalbuminuria is an early marker for increased morbidity and mortality]." Fortschr Med Orig **121 Suppl 1**: 28-32.

Brazy, P.C., Stead, W.W. und Fitzwilliam, J.F. (1989). "Progression of renal insufficiency: role of blood pressure." Kidney Int **35**(2): 670-4.

Brenner, B.M., Cooper, M.E., de Zeeuw, D., Keane, W.F., Mitch, W.E., Parving, H.H., Remuzzi, G., Snapinn, S.M., Zhang, Z. und Shahinfar, S. (2001). "Effects of losartan on renal and cardiovascular outcomes in patients with type 2 diabetes and nephropathy." N Engl J Med **345**(12): 861-9.

Briganti, E.M., Branley, P., Chadban, S.J., Shaw, J.E., McNeil, J.J., Welborn, T.A. und Atkins, R.C. (2002). "Smoking is associated with renal impairment and proteinuria in the normal population: the AusDiab kidney study. Australian Diabetes, Obesity and Lifestyle Study." Am J Kidney Dis **40**(4): 704-12.

Chachati, A., von Frenckell, R., Foidart-Willems, J., Godon, J.P. und Lefebvre, P.J. (1987). "Variability of albumin excretion in insulin-dependent diabetics." Diabet Med **4**(5): 441-5.

Chase, H.P., Garg, S.K., Marshall, G., Berg, C.L., Harris, S., Jackson, W.E. und Hamman, R.E. (1991). "Cigarette smoking increases the risk of albuminuria among subjects with type I diabetes." Jama **265**(5): 614-7.

Clausen, P., Jensen, J.S., Jensen, G., Borch-Johnsen, K. und Feldt-Rasmussen, B. (2001). "Elevated urinary albumin excretion is associated with impaired arterial dilatory capacity in clinically healthy subjects." Circulation **103**(14): 1869-74.

Comper, W.D., Jerums, G. und Osicka, T.M. (2003). "Deficiency in the detection of microalbuminuria by urinary dipstick in diabetic patients." Diabetes Care **26**(11): 3195-6.

Cordonnier, D.J., Pinel, N., Barro, C., Maynard, M., Zaoui, P., Halimi, S., Hurault de Ligny, B., Reznic, Y., Simon, D. und Bilous, R.W. (1999). "Expansion of cortical interstitium is limited by converting enzyme inhibition in type 2 diabetic patients with glomerulosclerosis. The Diabiopsies Group." J Am Soc Nephrol **10**(6): 1253-63.

Damsgaard, E.M., Froland, A., Jorgensen, O.D. und Mogensen, C.E. (1990). "Microalbuminuria as predictor of increased mortality in elderly people." Bmj **300**(6720): 297-300.

de Jong, P.E. und Brenner, B.M. (2004). "From secondary to primary prevention of progressive renal disease: the case for screening for albuminuria." Kidney Int **66**(6): 2109-18.

Deckert, T., Feldt-Rasmussen, B., Borch-Johnsen, K., Jensen, T. und Kofoed-Enevoldsen, A. (1989). "Albuminuria reflects widespread vascular damage. The Steno hypothesis." Diabetologia **32**(4): 219-26.

Dornblüth, O., Pschyrembel, W. und Zink, C. (1990). "Pschyrembel Klinisches Wörterbuch." **256**.

Feldt-Rasmussen, B., Dinesen, B. und Deckert, M. (1985). "Enzyme immunoassay: an improved determination of urinary albumin in diabetics with incipient nephropathy." Scand J Clin Lab Invest **45**(6): 539-44.

Fernandez Fernandez, I., Paez Pinto, J.M., Hermosin Bono, T., Vazquez Garijo, P., Ortiz Camunez, M.A. und Tarilonte Delgado, M.A. (1998). "Rapid screening test evaluation for microalbuminuria in diabetes mellitus." Acta Diabetol **35**(4): 199-202.

Gall, M.A., Rossing, P., Skott, P., Damsbo, P., Vaag, A., Bech, K., Dejgaard, A., Lauritzen, M., Lauritzen, E., Hougaard, P. und et al. (1991). "Prevalence of micro- and macroalbuminuria, arterial hypertension, retinopathy and large vessel disease in European type 2 (non-insulin-dependent) diabetic patients." Diabetologia **34**(9): 655-61.

Gerstein, H.C., Mann, J.F., Yi, Q., Zinman, B., Dinneen, S.F., Hoogwerf, B., Halle, J.P., Young, J., Rashkow, A., Joyce, C., Nawaz, S. und Yusuf, S. (2001). "Albuminuria and risk of cardiovascular events, death, and heart failure in diabetic and nondiabetic individuals." *Jama* **286**(4): 421-6.

Gilbert, R.E., Akdeniz, A. und Jerums, G. (1997). "Detection of microalbuminuria in diabetic patients by urinary dipstick." *Diabetes Res Clin Pract* **35**(1): 57-60.

Hansson, L., Zanchetti, A., Carruthers, S.G., Dahlof, B., Elmfeldt, D., Julius, S., Menard, J., Rahn, K.H., Wedel, H. und Westerling, S. (1998). "Effects of intensive blood-pressure lowering and low-dose aspirin in patients with hypertension: principal results of the Hypertension Optimal Treatment (HOT) randomised trial. HOT Study Group." *Lancet* **351**(9118): 1755-62.

Hasslacher, C., Danne, T., Sawicki, P.T. und Walter, H. (1999). "Frühdiagnose der diabetischen Nephropathie." *Deutsches Ärzteblatt* **96**(1-2): A-51/B-47/C-47.

Herold, G. (2003). "Nephrologie." *Innere Medizin*: 506ff.

Hillege, H.L., Fidler, V., Diercks, G.F., van Gilst, W.H., de Zeeuw, D., van Veldhuisen, D.J., Gans, R.O., Janssen, W.M., Grobbee, D.E. und de Jong, P.E. (2002). "Urinary albumin excretion predicts cardiovascular and noncardiovascular mortality in general population." *Circulation* **106**(14): 1777-82.

Hunsicker, L.G., Shearer, T.P. und Shaffer, S.J. (1981). "Acute reversible proteinuria induced by infusion of the polycation hexadimethrine." *Kidney Int* **20**(1): 7-17.

Jeffers, B.W., Estacio, R.O., Raynolds, M.V. und Schrier, R.W. (1997). "Angiotensin-converting enzyme gene polymorphism in non-insulin dependent diabetes mellitus and its relationship with diabetic nephropathy." *Kidney Int* **52**(2): 473-7.

Jensen, J.S., Feldt-Rasmussen, B., Strandgaard, S., Schroll, M. und Borch-Johnsen, K. (2000). "Arterial hypertension, microalbuminuria, and risk of ischemic heart disease." *Hypertension* **35**(4): 898-903.

Jones, C.A., Francis, M.E., Eberhardt, M.S., Chavers, B., Coresh, J., Engelgau, M., Kusek, J.W., Byrd-Holt, D., Narayan, K.M., Herman, W.H., Jones, C.P., Salive, M. und Agodoa, L.Y. (2002). "Microalbuminuria in the US population: third National Health and Nutrition Examination Survey." Am J Kidney Dis **39**(3): 445-59.

K/DOQI (2002). "Clinical practice guidelines for chronic kidney disease: evaluation, classification, and stratification. National Kidney Foundation, Kidney Disease Outcome Quality Initiative. ."

Kanwar, Y.S., Veis, A., Kimura, J.H. und Jakubowski, M.L. (1984). "Characterization of heparan sulfate-proteoglycan of glomerular basement membranes." Proc Natl Acad Sci U S A **81**(3): 762-6.

Kjellen, L., Bielefeld, D. und Hook, M. (1983). "Reduced sulfation of liver heparan sulfate in experimentally diabetic rats." Diabetes **32**(4): 337-42.

Klausen, K., Borch-Johnsen, K., Feldt-Rasmussen, B., Jensen, G., Clausen, P., Scharling, H., Appleyard, M. und Jensen, J.S. (2004). "Very low levels of microalbuminuria are associated with increased risk of coronary heart disease and death independently of renal function, hypertension, and diabetes." Circulation **110**(1): 32-5.

Klinke, R. und Silbernagel, S. (1996). "Die Funktion der Nieren." Lehrbuch der Physiologie 2: 294.

Kraft, S.K., Lazaridis, E.N., Qiu, C., Clark, C.M., Jr. und Marrero, D.G. (1999). "Screening and treatment of diabetic nephropathy by primary care physicians." J Gen Intern Med **14**(2): 88-97.

Leoncini, G., Sacchi, G., Ravera, M., Viazzi, F., Ratto, E., Vettoretti, S., Parodi, D., Bezante, G.P., Del Sette, M., Deferrari, G. und Pontremoli, R. (2002). "Microalbuminuria is an integrated marker of subclinical organ damage in primary hypertension." J Hum Hypertens **16**(6): 399-404.

Levey, A.S., Coresh, J., Balk, E., Kausz, A.T., Levin, A., Steffes, M.W., Hogg, R.J., Perrone, R.D., Lau, J. und Eknayan, G. (2003). "National Kidney Foundation practice guidelines for chronic kidney disease: evaluation, classification, and stratification." Ann Intern Med **139**(2): 137-47.

Lewis, E.J., Hunsicker, L.G., Clarke, W.R., Berl, T., Pohl, M.A., Lewis, J.B., Ritz, E., Atkins, R.C., Rohde, R. und Raz, I. (2001). "Renoprotective effect of the angiotensin-receptor antagonist irbesartan in patients with nephropathy due to type 2 diabetes." N Engl J Med **345**(12): 851-60.

Liese, A.D., Hense, H.W., Doring, A., Stieber, J. und Keil, U. (2001). "Microalbuminuria, central adiposity and hypertension in the non-diabetic urban population of the MONICA Augsburg survey 1994/95." J Hum Hypertens **15**(11): 799-804.

Mainous, A.G., 3rd und Gill, J.M. (2001). "The lack of screening for diabetic nephropathy: evidence from a privately insured population." Fam Med **33**(2): 115-9.

Mathiesen, E.R., Rønn, B., Jensen, T., Storm, B. und Deckert, T. (1990). "Relationship between blood pressure and urinary albumin excretion in development of microalbuminuria." Diabetes **39**(2): 245-9.

Matthews, C.M. (1957). "The theory of tracer experiments with  $^{131}\text{I}$ -labelled plasma proteins." Phys Med Biol **2**(1): 36-53.

McKenna, K. und Thompson, C. (1997). "Microalbuminuria: a marker to increased renal and cardiovascular risk in diabetes mellitus." Scott Med J **42**(4): 99-104.

Misiani, R., Marchesi, D., Tiraboschi, G., Gualandris, L., Pagni, R., Goglio, A., Amuso, G., Muratore, D., Bertuletti, P. und Massazza, M. (1991). "Urinary albumin excretion in normal pregnancy and pregnancy-induced hypertension." Nephron **59**(3): 416-22.

Mitzner, S.R. (1999). "MARS<sup>®</sup> (Molecular Adsorbents Recirculating System)-ein neues Blutentgiftungsverfahren." Dialyse aktuell **3**: 6-7.

Mogensen, C.E. (1971). "Urinary albumin excretion in early and long-term juvenile diabetes." Scand J Clin Lab Invest **28**(2): 183-93.

Mogensen, C.E. (1982). "Long-term antihypertensive treatment inhibiting progression of diabetic nephropathy." Br Med J (Clin Res Ed) **285**(6343): 685-8.

Mogensen, C.E. (2005). "Zielblutdruck und Screening bei Diabetikern mit Nephropathie." Deutsches Ärzteblatt **102**(16).

Mogensen, C.E., Chachati, A., Christensen, C.K., Close, C.F., Deckert, T., Hommel, E., Kastrup, J., Lefebvre, P., Mathiesen, E.R., Feldt-Rasmussen, B. und et al. (1985). "Microalbuminuria: an early marker of renal involvement in diabetes." *Uremia Invest* **9**(2): 85-95.

Mogensen, C.E., Vestbo, E., Poulsen, P.L., Christiansen, C., Damsgaard, E.M., Eiskjaer, H., Froland, A., Hansen, K.W., Nielsen, S. und Pedersen, M.M. (1995). "Microalbuminuria and potential confounders. A review and some observations on variability of urinary albumin excretion." *Diabetes Care* **18**(4): 572-81.

Mogensen, C.E., Viberti, G.C., Peheim, E., Kutter, D., Hasslacher, C., Hofmann, W., Renner, R., Bojestig, M., Poulsen, P.L., Scott, G., Thoma, J., Kuefer, J., Nilsson, B., Gambke, B., Mueller, P., Steinbiss, J. und Willamowski, K.D. (1997). "Multicenter evaluation of the Micral-Test II test strip, an immunologic rapid test for the detection of microalbuminuria." *Diabetes Care* **20**(11): 1642-6.

Molitch, M.E., DeFronzo, R.A., Franz, M.J., Keane, W.F., Mogensen, C.E., Parving, H.H. und Steffes, M.W. (2004). "Nephropathy in diabetes." *Diabetes Care* **27 Suppl 1**: S79-83.

Orth, S.R., Ritz, E. und Schrier, R.W. (1997). "The renal risks of smoking." *Kidney Int* **51**(6): 1669-77.

Parving, H.H., Ranek, L. und Lassen, N.A. (1977). "Increased transcapillary escape rate of albumin in patients with cirrhosis of the liver." *Scand J Clin Lab Invest* **37**(7): 643-8.

Parving, H.H. und Rossing, N. (1973). "Simultaneous determination of the transcapillary escape rate of albumin and IgG in normal and long-term juvenile diabetic subjects." *Scand J Clin Lab Invest* **32**(3): 239-44.

Pedrinelli, R., Dell'Omo, G., Di Bello, V., Pontremoli, R. und Mariani, M. (2002). "Microalbuminuria, an integrated marker of cardiovascular risk in essential hypertension." *J Hum Hypertens* **16**(2): 79-89.

Pedrinelli, R., Giampietro, O., Carmassi, F., Melillo, E., Dell'Omo, G., Catapano, G., Matteucci, E., Talarico, L., Morale, M., De Negri, F. und et al. (1994). "Microalbuminuria and endothelial dysfunction in essential hypertension." *Lancet* **344**(8914): 14-8.

Pedrini, M.T., Levey, A.S., Lau, J., Chalmers, T.C. und Wang, P.H. (1996). "The effect of dietary protein restriction on the progression of diabetic and nondiabetic renal diseases: a meta-analysis." Ann Intern Med **124**(7): 627-32.

Peterson, J.C., Adler, S., Burkart, J.M., Greene, T., Hebert, L.A., Hunsicker, L.G., King, A.J., Klahr, S., Massry, S.G. und Seifter, J.L. (1995). "Blood pressure control, proteinuria, and the progression of renal disease. The Modification of Diet in Renal Disease Study." Ann Intern Med **123**(10): 754-62.

Pontremoli, R. (1996). "Microalbuminuria in essential hypertension--its relation to cardiovascular risk factors." Nephrol Dial Transplant **11**(11): 2113-5.

Pontremoli, R., Leoncini, G., Ravera, M., Viazzi, F., Vettoretti, S., Ratto, E., Parodi, D., Tomillo, C. und Deferrari, G. (2002). "Microalbuminuria, cardiovascular, and renal risk in primary hypertension." J Am Soc Nephrol **13 Suppl 3**: S169-72.

Poulsen, P.L. und Mogensen, C.E. (1998). "Clinical evaluation of a test for immediate and quantitative determination of urinary albumin-to-creatinine ratio. A brief report." Diabetes Care **21**(1): 97-8.

Ravid, M., Brosh, D., Levi, Z., Bar-Dayan, Y., Ravid, D. und Rachmani, R. (1998). "Use of enalapril to attenuate decline in renal function in normotensive, normoalbuminuric patients with type 2 diabetes mellitus. A randomized, controlled trial." Ann Intern Med **128**(12 Pt 1): 982-8.

Reiber, H. (1994). "Flow rate of cerebrospinal fluid (CSF)--a concept common to normal blood-CSF barrier function and to dysfunction in neurological diseases." J Neurol Sci **122**(2): 189-203.

Rohrbach, R. (1986). "Reduced content and abnormal distribution of anionic sites (acid proteoglycans) in the diabetic glomerular basement membrane." Virchows Arch B Cell Pathol Incl Mol Pathol **51**(2): 127-35.

Romundstad, S., Holmen, J., Hallan, H., Kvenild, K. und Ellekjaer, H. (2003). "Microalbuminuria and all-cause mortality in treated hypertensive individuals: does sex matter? The Nord-Trondelag Health Study (HUNT), Norway." Circulation **108**(22): 2783-9.

Rossing, N. (1978). "Intra- and extravascular distribution of albumin and immunoglobulin in man." Lymphology **11**(4): 138-42.

Rothschild, M.A., Oratz, M. und Schreiber, S.S. (1988). "Serum albumin." Hepatology **8**(2): 385-401.

Russo, L.M., Bakris, G.L. und Comper, W.D. (2002). "Renal handling of albumin: a critical review of basic concepts and perspective." Am J Kidney Dis **39**(5): 899-919.

Scherberich, J. (1998). "Nichtinvasive Diagnostik von Nierenerkrankungen: Differentielle klinische Bewertung der Proteinurieformen." Der Bayerische Internist **18**: 80-88.

Schiffl, H., Lang, S.M. und Fischer, R. (2002). "Stopping smoking slows accelerated progression of renal failure in primary renal disease." J Nephrol **15**(3): 270-4.

Schmidt, R.F. und Thews, G. (1997). "Blutplasma." Physiologie des Menschen **27**: 412ff.

Schrier, R.W., Estacio, R.O., Esler, A. und Mehler, P. (2002). "Effects of aggressive blood pressure control in normotensive type 2 diabetic patients on albuminuria, retinopathy and strokes." Kidney Int **61**(3): 1086-97.

Selby, J.V., FitzSimmons, S.C., Newman, J.M., Katz, P.P., Sepe, S. und Showstack, J. (1990). "The natural history and epidemiology of diabetic nephropathy. Implications for prevention and control." Jama **263**(14): 1954-60.

Stow, J.L., Sawada, H. und Farquhar, M.G. (1985). "Basement membrane heparan sulfate proteoglycans are concentrated in the laminae rarae and in podocytes of the rat renal glomerulus." Proc Natl Acad Sci U S A **82**(10): 3296-300.

UKPDSG (1998). "Intensive blood-glucose control with sulphonylureas or insulin compared with conventional treatment and risk of complications in patients with type 2 diabetes (UKPDS 33). UK Prospective Diabetes Study (UKPDS) Group." Lancet **352**(9131): 837-53.

van der Woude, F.J. und van Det, N.F. (1997). "Heparan sulphate proteoglycans and diabetic nephropathy." Exp Nephrol **5**(3): 180-8.

Viberti, G.C., Hill, R.D., Jarrett, R.J., Argyropoulos, A., Mahmud, U. und Keen, H. (1982). "Microalbuminuria as a predictor of clinical nephropathy in insulin-dependent diabetes mellitus." Lancet **1**(8287): 1430-2.

Volhard, F. (1923). "Der arterielle Blutdruck." Verh Dtsch Inn Med **135**: 134-184.

Weber, M.H., Scholz, P., Stibbe, W. und Scheler, F. (1985). "[Alpha 1-microglobulin in the urine and serum in proteinuria and kidney insufficiency]." Klin Wochenschr **63**(15): 711-7.

Weigand, K. (1980). "[Synthesis, distribution and importance of serum albumin]." Klin Anasthesiol Intensivther **21**: 52-64.

Weir, M.R. (2004). "Microalbuminuria in type 2 diabetics: an important, overlooked cardiovascular risk factor." J Clin Hypertens (Greenwich) **6**(3): 134-41; quiz 142-3.

Wenzel, U.O. (2001). "Angiotensin-converting enzyme inhibitors and progression of renal disease: evidence from clinical studies." Contrib Nephrol(135): 200-11.

Zelmanovitz, T., Gross, J.L., Oliveira, J.R., Paggi, A., Tatsch, M. und Azevedo, M.J. (1997). "The receiver operating characteristics curve in the evaluation of a random urine specimen as a screening test for diabetic nephropathy." Diabetes Care **20**(4): 516-9.

Zheng, Y.L., Liao, J., Takeda, Y., Io, H., Kobata, M., Kanamaru, Y., Maeda, A., Shou, I., Igari, J. und Tomino, Y. (1999). "Determination of sensitivity and specificity of the Micral-Test II strip for detection of microalbuminuria in diabetic and nondiabetic patients." Nephron **81**(4): 455.

Zimmerhackl, L.B. (1993). "Evaluation of nephrotoxicity with renal antigens in children: role of Tamm-Horsfall protein." Eur J Clin Pharmacol **44 Suppl 1**: S39-42.

