

## Literaturverzeichnis

- Aaltonen, L. A., Peltomaki, P., Mecklin, J. P., Jarvinen, H., Jass, J. R., Green, J. S., Lynch, H. T., Watson, P., Tallqvist, G., Juhola, M. und et al. (1994) Replication errors in benign and malignant tumors from hereditary nonpolyposis colorectal cancer patients. *Cancer Res* **54**: 1645-8.
- Agoff, S. N., Brentnall, T. A., Crispin, D. A., Taylor, S. L., Raaka, S., Haggitt, R. C., Reed, M. W., Afonina, I. A., Rabinovitch, P. S., Stevens, A. C., Feng, Z. und Bronner, M. P. (2000) The role of cyclooxygenase 2 in ulcerative colitis-associated neoplasia. *Am J Pathol* **157**: 737-45.
- Allgayer, H., Deschryver, K. und Stenson, W. F. (1989) Treatment with 16,16'-dimethyl prostaglandin E2 before and after induction of colitis with trinitrobenzenesulfonic acid in rats decreases inflammation. *Gastroenterology* **96**: 1290-300.
- Ambs, S., Merriam, W. G., Bennett, W. P., Felley-Bosco, E., Ogunfusika, M. O., Oser, S. M., Klein, S., Shields, P. G., Billiar, T. R. und Harris, C. C. (1998) Frequent nitric oxide synthase-2 expression in human colon adenomas: implication for tumor angiogenesis and colon cancer progression. *Cancer Res* **58**: 334-41.
- Andoh, A., Tsujikawa, T. und Fujiyama, Y. (2003) Role of dietary fiber and short-chain fatty acids in the colon. *Curr Pharm Des* **9**: 347-58.
- Andus, T., Daig, R., Vogl, D., Aschenbrenner, E., Lock, G., Hollerbach, S., Kollinger, M., Scholmerich, J. und Gross, V. (1997) Imbalance of the interleukin 1 system in colonic mucosa-association with intestinal inflammation and interleukin 1 receptor antagonist [corrected] genotype 2. *Gut* **41**: 651-7.
- Appleton, I., Tomlinson, A. und Willoughby, D. A. (1996) Induction of cyclo-oxygenase and nitric oxide synthase in inflammation. *Adv Pharmacol* **35**: 27-78.
- Araki, Y., Okamura, S., Hussain, S. P., Nagashima, M., He, P., Shiseki, M., Miura, K. und Harris, C. C. (2003) Regulation of cyclooxygenase-2 expression by the Wnt and ras pathways. *Cancer Res* **63**: 728-34.
- Asp, N. G. (1992) Preface: Resistant starch. Proceedings of the second plenary meeting of EURESTA: European Flair Concerted action No. 11 (Cost 911) on Physiological implications of the consumption of resistant starch in man. *Eur. J. Clin. Nutr.* **46, Suppl 2**: 1.
- Babbs, C. F. (1990) Free radicals and the etiology of colon cancer. *Free Radic Biol Med* **8**: 191-200.
- Barshishat, M., Polak-Charcon, S. und Schwartz, B. (2000) Butyrate regulates E-cadherin transcription, isoform expression and intracellular position in colon cancer cells. *Br J Cancer* **82**: 195-203.
- Behrens, J., von Kries, J. P., Kuhl, M., Bruhn, L., Wedlich, D., Grosschedl, R. und Birchmeier, W. (1996) Functional interaction of beta-catenin with the transcription factor LEF-1. *Nature* **382**: 638-42.
- Bhatia, B., Maldonado, C. J., Tang, S., Chandra, D., Klein, R. D., Chopra, D., Shappell, S. B., Yang, P., Newman, R. A. und Tang, D. G. (2003) Subcellular localization and tumor-suppressive functions of 15-lipoxygenase 2 (15-LOX2) and its splice variants. *J Biol Chem* **278**: 25091-100.
- Bird, R. P. (1995) Role of aberrant crypt foci in understanding the pathogenesis of colon cancer. *Cancer Lett* **93**: 55-71.
- Bishop, J. M. (1991) Molecular themes in oncogenesis. *Cell* **64**: 235-48.
- Bissonnette, M., Khare, S., von Lintig, F. C., Wali, R. K., Nguyen, L., Zhang, Y., Hart, J., Skarosi, S., Varki, N., Boss, G. R. und Brasitus, T. A. (2000) Mutational and nonmutational activation of p21ras in rat colonic azoxymethane-induced tumors: effects on mitogen-activated protein kinase, cyclooxygenase-2, and cyclin D1. *Cancer Res* **60**: 4602-9.
- Blottiere, H. M., Buecher, B., Galmiche, J. P. und Cherbut, C. (2003) Molecular analysis of the effect of short-chain fatty acids on intestinal cell proliferation. *Proc Nutr Soc* **62**: 101-6.

- Blum, C. A., Xu, M., Orner, G. A., Fong, A. T., Bailey, G. S., Stoner, G. D., Horio, D. T. und Dashwood, R. H. (2001) beta-Catenin mutation in rat colon tumors initiated by 1,2-dimethylhydrazine and 2-amino-3-methylimidazo[4,5-f]quinoline, and the effect of post-initiation treatment with chlorophyllin and indole-3-carbinol. *Carcinogenesis* **22**: 315-20.
- Boese-Landgraf, J. (1998) Epidemiologie, Vorstufen und Pathogenese des kolorektalen Karzinoms. *Der Onkologe* **4**: 2-4.
- Boirivant, M., Pica, R., DeMaria, R., Testi, R., Pallone, F. und Strober, W. (1996) Stimulated human lamina propria T cells manifest enhanced Fas-mediated apoptosis. *J Clin Invest* **98**: 2616-22.
- Bondeson, J., Browne, K. A., Brennan, F. M., Foxwell, B. M. und Feldmann, M. (1999) Selective regulation of cytokine induction by adenoviral gene transfer of IkappaBalpha into human macrophages: lipopolysaccharide-induced, but not zymosan-induced, proinflammatory cytokines are inhibited, but IL-10 is nuclear factor-kappaB independent. *J Immunol* **162**: 2939-45.
- Bonen, D. K. und Cho, J. H. (2003) The genetics of inflammatory bowel disease. *Gastroenterology* **124**: 521-36.
- Bos, J. L., Fearon, E. R., Hamilton, S. R., Verlaan-de Vries, M., van Boom, J. H., van der Eb, A. J. und Vogelstein, B. (1987) Prevalence of ras gene mutations in human colorectal cancers. *Nature* **327**: 293-7.
- Boughton-Smith, N. K., Evans, S. M., Hawkey, C. J., Cole, A. T., Balsitis, M., Whittle, B. J. und Moncada, S. (1993) Nitric oxide synthase activity in ulcerative colitis and Crohn's disease. *Lancet* **342**: 338-40.
- Brash, A. R., Boeglin, W. E. und Chang, M. S. (1997) Discovery of a second 15S-lipoxygenase in humans. *Proc Natl Acad Sci U S A* **94**: 6148-52.
- Brash, A. R. (1999) Lipoxygenases: occurrence, functions, catalysis, and acquisition of substrate. *J Biol Chem* **274**: 23679-82.
- Brigelius-Flohé, R. (1999) Tissue-specific functions of individual glutathione peroxidases. *Free Radic Biol Med* **27**: 951-65.
- Brigelius-Flohé, R., Maurer, S., Lotzer, K., Bol, G., Kallionpaa, H., Lehtolainen, P., Viita, H. und Yla-Herttuala, S. (2000) Overexpression of PHGPx inhibits hydroperoxide-induced oxidation, NFkappaB activation and apoptosis and affects oxLDL-mediated proliferation of rabbit aortic smooth muscle cells. *Atherosclerosis* **152**: 307-16.
- Brigelius-Flohé, R., Muller, C., Menard, J., Florian, S., Schmehl, K. und Wingler, K. (2001) Functions of GI-GPx: lessons from selenium-dependent expression and intracellular localization. *Biofactors* **14**: 101-6.
- Brinckmann, R., Schnurr, K., Heydeck, D., Rosenbach, T., Kolde, G. und Kuhn, H. (1998) Membrane translocation of 15-lipoxygenase in hematopoietic cells is calcium-dependent and activates the oxygenase activity of the enzyme. *Blood* **91**: 64-74.
- Bronner, C. E., Baker, S. M., Morrison, P. T., Warren, G., Smith, L. G., Lescoe, M. K., Kane, M., Earabino, C., Lipford, J., Lindblom, A. und et al. (1994) Mutation in the DNA mismatch repair gene homologue hMLH1 is associated with hereditary non-polyposis colon cancer. *Nature* **368**: 258-61.
- Bulow, S. (1986) Clinical features in familial polyposis coli. Results of the Danish Polyposis Register. *Dis Colon Rectum* **29**: 102-7.
- Butzner, J. D., Parmar, R., Bell, C. J. und Dalal, V. (1996) Butyrate enema therapy stimulates mucosal repair in experimental colitis in the rat. *Gut* **38**: 568-73.
- Cameron, I. L., Garza, J. und Hardman, W. E. (1996) Distribution of lymphoid nodules, aberrant crypt foci and tumours in the colon of carcinogen-treated rats. *Br J Cancer* **73**: 893-8.
- Caspari, R., Jungck, M., Lamberti, C., Schmidt-Wolf, I. G., Friedl, W. und Propping, P. (2000) [The diagnosis of hereditary colorectal carcinomas]. *Dtsch Med Wochenschr* **125**: 153-8.
- Cassand, P., Maziere, S., Champ, M., Meflah, K., Bornet, F. und Narbonne, J. F. (1997) Effects of resistant starch- and vitamin A-supplemented diets on the promotion of precursor lesions of colon cancer in rats. *Nutr Cancer* **27**: 53-9.

- Chandrasekharan, N. V., Dai, H., Roos, K. L., Evanson, N. K., Tomsik, J., Elton, T. S. und Simmons, D. L. (2002) COX-3, a cyclooxygenase-1 variant inhibited by acetaminophen and other analgesic/antipyretic drugs: cloning, structure, and expression. *Proc Natl Acad Sci U S A* **99**: 13926-31.
- Chapple, K. S., Cartwright, E. J., Hawcroft, G., Tisbury, A., Bonifer, C., Scott, N., Windsor, A. C., Guillou, P. J., Markham, A. F., Coletta, P. L. und Hull, M. A. (2000) Localization of cyclooxygenase-2 in human sporadic colorectal adenomas. *Am J Pathol* **156**: 545-53.
- Chapple, K. S., Scott, N., Guillou, P. J., Coletta, P. L. und Hull, M. A. (2002) Interstitial cell cyclooxygenase-2 expression is associated with increased angiogenesis in human sporadic colorectal adenomas. *J Pathol* **198**: 435-41.
- Chazotte-Aubert, L., Hainaut, P. und Ohshima, H. (2000) Nitric oxide nitrates tyrosine residues of tumor-suppressor p53 protein in MCF-7 cells. *Biochem Biophys Res Commun* **267**: 609-13.
- Cho, J. H., Nicolae, D. L., Gold, L. H., Fields, C. T., LaBuda, M. C., Rohal, P. M., Pickles, M. R., Qin, L., Fu, Y., Mann, J. S., Kirschner, B. S., Jabs, E. W., Weber, J., Hanauer, S. B., Bayless, T. M. und Brant, S. R. (1998) Identification of novel susceptibility loci for inflammatory bowel disease on chromosomes 1p, 3q, and 4q: evidence for epistasis between 1p and IBD1. *Proc Natl Acad Sci U S A* **95**: 7502-7.
- Chu, F. F., Doroshow, J. H. und Esworthy, R. S. (1993) Expression, characterization, and tissue distribution of a new cellular selenium-dependent glutathione peroxidase, GSHPx-GI. *J Biol Chem* **268**: 2571-6.
- Chu, F. F., Esworthy, R. S., Ho, Y. S., Bermeister, M., Swiderek, K. und Elliott, R. W. (1997) Expression and chromosomal mapping of mouse Gpx2 gene encoding the gastrointestinal form of glutathione peroxidase, GPX-GI. *Biomed Environ Sci* **10**: 156-62.
- Clancy, R., Varenika, B., Huang, W., Ballou, L., Attur, M., Amin, A. R. und Abramson, S. B. (2000) Nitric oxide synthase/COX cross-talk: nitric oxide activates COX-1 but inhibits COX-2-derived prostaglandin production. *J Immunol* **165**: 1582-7.
- Crew, T. E., Elder, D. J. und Paraskeva, C. (2000) A cyclooxygenase-2 (COX-2) selective non-steroidal anti-inflammatory drug enhances the growth inhibitory effect of butyrate in colorectal carcinoma cells expressing COX-2 protein: regulation of COX-2 by butyrate. *Carcinogenesis* **21**: 69-77.
- Csordas, A. (1996) Butyrate, aspirin and colorectal cancer. *Eur J Cancer Prev* **5**: 221-31.
- Cuzzocrea, S., Mazzon, E., Serraino, I., Dugo, L., Centorrino, T., Ciccolo, A., Sautebin, L. und Caputi, A. P. (2001) Celecoxib, a selective cyclo-oxygenase-2 inhibitor reduces the severity of experimental colitis induced by dinitrobenzene sulfonic acid in rats. *Eur J Pharmacol* **431**: 91-102.
- DGE. (2000) Referenzwerte für die Nährstoffzufuhr, (eds. Deutsche Gesellschaft für Ernährung (DGE), Österreichische Gesellschaft für Ernährung (ÖGE), Schweizerische Gesellschaft für Ernährungsforschung (SGE) und (SVE), S. V. f. E.), pp. 59-63, Umschau/Braus, Frankfurt am Main.
- Dijkstra, G., Moshage, H., van Dullemen, H. M., de Jager-Krikken, A., Tiebosch, A. T., Kleibeuker, J. H., Jansen, P. L. und van Goor, H. (1998) Expression of nitric oxide synthases and formation of nitrotyrosine and reactive oxygen species in inflammatory bowel disease. *J Pathol* **186**: 416-21.
- Dikopoulos, N., Nussler, A. K., Liptay, S., Bachem, M., Reinshagen, M., Stiegler, M., Schmid, R. M., Adler, G. und Weidenbach, H. (2001) Inhibition of nitric oxide synthesis by aminoguanidine increases intestinal damage in the acute phase of rat TNB-colitis. *Eur J Clin Invest* **31**: 234-9.
- Dohi, T., Fujihashi, K., Kiyono, H., Elson, C. O. und McGhee, J. R. (2000) Mice deficient in Th1- and Th2-type cytokines develop distinct forms of hapten-induced colitis. *Gastroenterology* **119**: 724-33.
- Duchmann, R., Hoffmann, J., Marth, T., Schneider, T., Stallmach, A., Zeitz, M. (1999) Mukosales Immunsystem im Darm - Immunologische Grundlagen chronisch entzündlicher Darmerkrankungen und der HIV-induzierten Enteropathie. *Magazin Forschung* **1**: 47-64.

- Eberhart, C. E., Coffey, R. J., Radhika, A., Giardiello, F. M., Ferrenbach, S. und DuBois, R. N. (1994) Up-regulation of cyclooxygenase 2 gene expression in human colorectal adenomas and adenocarcinomas. *Gastroenterology* **107**: 1183-8.
- Ekbom, A., Daszak, P., Kraaz, W. und Wakefield, A. J. (1996) Crohn's disease after in-utero measles virus exposure. *Lancet* **348**: 515-7.
- el-Deiry, W. S., Harper, J. W., O'Connor, P. M., Velculescu, V. E., Canman, C. E., Jackman, J., Pietenpol, J. A., Burrell, M., Hill, D. E., Wang, Y. und et al. (1994) WAF1/CIP1 is induced in p53-mediated G1 arrest and apoptosis. *Cancer Res* **54**: 1169-74.
- Ell, C. und Raithel, M. (1998) [Diagnostic standard in chronic inflammatory bowel diseases]. *Internist (Berl)* **39**: 1013-23.
- Elson, C. O., Sartor, R. B., Tennyson, G. S. und Riddell, R. H. (1995) Experimental models of inflammatory bowel disease. *Gastroenterology* **109**: 1344-67.
- Elson, C. O., Beagley, K. W., Sharmanov, A. T., Fujihashi, K., Kiyono, H., Tennyson, G. S., Cong, Y., Black, C. A., Ridwan, B. W. und McGhee, J. R. (1996) Hapten-induced model of murine inflammatory bowel disease: mucosa immune responses and protection by tolerance. *J Immunol* **157**: 2174-85.
- Englyst, H. N., Kingman, S. M. und Cummings, J. H. (1992) Classification and measurement of nutritionally important starch fractions. *Eur J Clin Nutr* **46 Suppl 2**: S33-50.
- Esworthy, R. S., Swiderek, K. M., Ho, Y. S. und Chu, F. F. (1998) Selenium-dependent glutathione peroxidase-GI is a major glutathione peroxidase activity in the mucosal epithelium of rodent intestine. *Biochim Biophys Acta* **1381**: 213-26.
- Esworthy, R. S., Aranda, R., Martin, M. G., Doroshow, J. H., Binder, S. W. und Chu, F. F. (2001) Mice with combined disruption of Gpx1 and Gpx2 genes have colitis. *Am J Physiol Gastrointest Liver Physiol* **281**: G848-55.
- Esworthy, R. S., Binder, S. W., Doroshow, J. H. und Chu, F. F. (2003) Microflora trigger colitis in mice deficient in selenium-dependent glutathione peroxidase and induce Gpx2 gene expression. *Biol Chem* **384**: 597-607.
- Evans, D. G., Walsh, S., Jeacock, J., Robinson, C., Hadfield, L., Davies, D. R. und Kingston, R. (1997) Incidence of hereditary non-polyposis colorectal cancer in a population-based study of 1137 consecutive cases of colorectal cancer. *Br J Surg* **84**: 1281-5.
- Fasano, A. (2002) Toxins and the gut: role in human disease. *Gut* **50 Suppl 3**: III9-14.
- Fearnhead, N. S., Britton, M. P. und Bodmer, W. F. (2001) The ABC of APC. *Hum Mol Genet* **10**: 721-33.
- Femia, A. P., Luceri, C., Dolara, P., Giannini, A., Biggeri, A., Salvadori, M., Clune, Y., Collins, K. J., Paglierani, M. und Caderni, G. (2002) Antitumorigenic activity of the prebiotic inulin enriched with oligofructose in combination with the probiotics *Lactobacillus rhamnosus* and *Bifidobacterium lactis* on azoxymethane-induced colon carcinogenesis in rats. *Carcinogenesis* **23**: 1953-60.
- Fink, S. P., Reddy, G. R. und Marnett, L. J. (1997) Mutagenicity in *Escherichia coli* of the major DNA adduct derived from the endogenous mutagen malondialdehyde. *Proc Natl Acad Sci U S A* **94**: 8652-7.
- Fiocchi, C. (1998) Inflammatory bowel disease: etiology and pathogenesis. *Gastroenterology* **115**: 182-205.
- Fishel, R., Lescoe, M. K., Rao, M. R., Copeland, N. G., Jenkins, N. A., Garber, J., Kane, M. und Kolodner, R. (1993) The human mutator gene homolog MSH2 and its association with hereditary nonpolyposis colon cancer. *Cell* **75**: 1027-38.
- Fishel, R. und Kolodner, R. D. (1995) Identification of mismatch repair genes and their role in the development of cancer. *Curr Opin Genet Dev* **5**: 382-95.
- Flohé, L., Günzler, W. Schock, H.H. (1973) Glutathion peroxidase: A selenoenzyme. *FEBS Lett* **32**: 132-134.
- Florian, S., Wingler, K., Schmehl, K., Jacobasch, G., Kreuzer, O. J., Meyerhof, W. und Brigelius-Flohé, R. (2001) Cellular and subcellular localization of gastrointestinal glutathione peroxidase in normal and malignant human intestinal tissue. *Free Radic Res* **35**: 655-63.

- Foitzik, T., Kruschewski, M., Kroesen, A. und Buhr, H. J. (1999) Does microcirculation play a role in the pathogenesis of inflammatory bowel diseases? Answers from intravital microscopic studies in animal models. *Int J Colorectal Dis* **14**: 29-34.
- Fürstenberger, G., Hagedorn, H., Jacobi, T., Besemfelder, E., Stephan, M., Lehmann, W. D. und Marks, F. (1991) Characterization of an 8-lipoxygenase activity induced by the phorbol ester tumor promoter 12-O-tetradecanoylphorbol-13-acetate in mouse skin in vivo. *J Biol Chem* **266**: 15738-45.
- Fuss, I. J., Neurath, M., Boirivant, M., Klein, J. S., de la Motte, C., Strong, S. A., Fiocchi, C. und Strober, W. (1996) Disparate CD4+ lamina propria (LP) lymphokine secretion profiles in inflammatory bowel disease. Crohn's disease LP cells manifest increased secretion of IFN-gamma, whereas ulcerative colitis LP cells manifest increased secretion of IL-5. *J Immunol* **157**: 1261-70.
- Futami, S., Aoyama, N., Honsako, Y., Tamura, T., Morimoto, S., Nakashima, T., Ohmoto, A., Okano, H., Miyamoto, M., Inaba, H. und et al. (1995) HLA-DRB1\*1502 allele, subtype of DR15, is associated with susceptibility to ulcerative colitis and its progression. *Dig Dis Sci* **40**: 814-8.
- Garewal, H., Ramsey, L., Fass, R., Hart, N. K., Payne, C. M., Bernstein, H. und Bernstein, C. (2003) Perils of immunohistochemistry: variability in staining specificity of commercially available COX-2 antibodies on human colon tissue. *Dig Dis Sci* **48**: 197-202.
- Geboes, K. und Dalle, I. (2002) Influence of treatment on morphological features of mucosal inflammation. *Gut* **50 Suppl 3**: III37-42.
- Gendler, S. J. und Spicer, A. P. (1995) Epithelial mucin genes. *Annu Rev Physiol* **57**: 607-34.
- Ghosh, D. K. und Stuehr, D. J. (1995) Macrophage NO synthase: characterization of isolated oxygenase and reductase domains reveals a head-to-head subunit interaction. *Biochemistry* **34**: 801-7.
- Gibson, G. R. und Roberfroid, M. B. (1995) Dietary modulation of the human colonic microbiota: introducing the concept of prebiotics. *J Nutr* **125**: 1401-12.
- Glasgow, W. C., Afshari, C. A., Barrett, J. C. und Eling, T. E. (1992) Modulation of the epidermal growth factor mitogenic response by metabolites of linoleic and arachidonic acid in Syrian hamster embryo fibroblasts. Differential effects in tumor suppressor gene (+) and (-) phenotypes. *J Biol Chem* **267**: 10771-9.
- Glasgow, W. C., Hui, R., Everhart, A. L., Jayawickreme, S. P., Angerman-Stewart, J., Han, B. B. und Eling, T. E. (1997) The linoleic acid metabolite, (13S)-hydroperoxyoctadecadienoic acid, augments the epidermal growth factor receptor signaling pathway by attenuation of receptor dephosphorylation. Differential response in Syrian hamster embryo tumor suppressor phenotypes. *J Biol Chem* **272**: 19269-76.
- Goodwin, D. C., Landino, L. M. und Marnett, L. J. (1999) Effects of nitric oxide and nitric oxide-derived species on prostaglandin endoperoxide synthase and prostaglandin biosynthesis. *Faseb J* **13**: 1121-36.
- Groden, J., Thliveris, A., Samowitz, W., Carlson, M., Gelbert, L., Albertsen, H., Joslyn, G., Stevens, J., Spirio, L., Robertson, M. und et al. (1991) Identification and characterization of the familial adenomatous polyposis coli gene. *Cell* **66**: 589-600.
- Gurler, A., Boyvat, A., Turksen, U. (1997) Clinical Manifestations of Behcet's Disease: An Analysis of 2147 Patients. *Yonsei Med J* **38**: 423-427.
- Gustafson-Svard, C., Lilja, I., Hallbook, O. und Sjodahl, R. (1996) Cyclooxygenase-1 and cyclooxygenase-2 gene expression in human colorectal adenocarcinomas and in azoxymethane induced colonic tumours in rats. *Gut* **38**: 79-84.
- Gyde, S. N., Prior, P., Allan, R. N., Stevens, A., Jewell, D. P., Truelove, S. C., Lofberg, R., Brostrom, O. und Hellers, G. (1988) Colorectal cancer in ulcerative colitis: a cohort study of primary referrals from three centres. *Gut* **29**: 206-17.
- Halter, F., Tarnawski, A. S., Schmassmann, A. und Peskar, B. M. (2001) Cyclooxygenase 2-implications on maintenance of gastric mucosal integrity and ulcer healing: controversial issues and perspectives. *Gut* **49**: 443-53.
- Hao, X. P., Pretlow, T. G., Rao, J. S. und Pretlow, T. P. (2001) Inducible nitric oxide synthase (iNOS) is expressed similarly in multiple aberrant crypt foci and colorectal tumors from the same patients. *Cancer Res* **61**: 419-22.

- Herlin, T., Fogh, K., Hansen, E. S., Andreasen, A., Knudsen, V., Henriksen, T. B., Bunger, C. und Kragballe, K. (1990) 15-HETE inhibits leukotriene B4 formation and synovial cell proliferation in experimental arthritis. *Agents Actions* **29**: 52-3.
- Herrlinger, K., Fellermann, K. und Stange, E. F. (2002) Standard therapy of ulcerative colitis. Evidence- and consensus-based recommendations. *Internist (Berl)* **43**: 1367-8, 1371-5.
- Hightower, L. E. (1991) Heat shock, stress proteins, chaperones, and proteotoxicity. *Cell* **66**: 191-7.
- Hoffmann, J. C. und Zeitz, M. (2002) Standard therapy of Crohn disease. Consensus-based recommendations. *Internist (Berl)* **43**: 1376, 1379-85.
- Hofseth, L. J., Saito, S., Hussain, S. P., Espey, M. G., Miranda, K. M., Araki, Y., Jhappan, C., Higashimoto, Y., He, P., Linke, S. P., Quezado, M. M., Zurer, I., Rotter, V., Wink, D. A., Appella, E. und Harris, C. C. (2003) Nitric oxide-induced cellular stress and p53 activation in chronic inflammation. *Proc Natl Acad Sci U S A* **100**: 143-8.
- Holtmann, M., Mudter, J., Galle, P.R., Neurath, M. F. (2002) Das mukosale Immunsystem. Wie klar ist die Pathophysiologie? *Der Internist* **43**: 1343-1353.
- Hsi, L. C., Angerman-Stewart, J. und Eling, T. E. (1999) Introduction of full-length APC modulates cyclooxygenase-2 expression in HT-29 human colorectal carcinoma cells at the translational level. *Carcinogenesis* **20**: 2045-9.
- Hsi, L. C., Wilson, L., Nixon, J. und Eling, T. E. (2001) 15-lipoxygenase-1 metabolites down-regulate peroxisome proliferator-activated receptor gamma via the MAPK signaling pathway. *J Biol Chem* **276**: 34545-52.
- Huang, W. C., Chen, J. J., Inoue, H. und Chen, C. C. (2003) Tyrosine Phosphorylation of I-kappaB Kinase alpha/beta by Protein Kinase C-Dependent c-Src Activation Is Involved in TNF-alpha-Induced Cyclooxygenase-2 Expression. *J Immunol* **170**: 4767-75.
- Hugot, J. P., Zouali, H., Lesage, S. und Thomas, G. (1999) Etiology of the inflammatory bowel diseases. *Int J Colorectal Dis* **14**: 2-9.
- Hugot, J. P., Chamaillard, M., Zouali, H., Lesage, S., Cezard, J. P., Belaiche, J., Almer, S., Tysk, C., O'Morain, C. A., Gassull, M., Binder, V., Finkel, Y., Cortot, A., Modigliani, R., Laurent-Puig, P., Gower-Rousseau, C., Macry, J., Colombel, J. F., Sahbatou, M. und Thomas, G. (2001) Association of NOD2 leucine-rich repeat variants with susceptibility to Crohn's disease. *Nature* **411**: 599-603.
- Hussain, S. P., Amstad, P., Raja, K., Ambs, S., Nagashima, M., Bennett, W. P., Shields, P. G., Ham, A. J., Swenberg, J. A., Marrogi, A. J. und Harris, C. C. (2000) Increased p53 mutation load in noncancerous colon tissue from ulcerative colitis: a cancer-prone chronic inflammatory disease. *Cancer Res* **60**: 3333-7.
- Huycke, M. M., Abrams, V. und Moore, D. R. (2002) Enterococcus faecalis produces extracellular superoxide and hydrogen peroxide that damages colonic epithelial cell DNA. *Carcinogenesis* **23**: 529-36.
- Ikawa, H., Kamitani, H., Calvo, B. F., Foley, J. F. und Eling, T. E. (1999) Expression of 15-lipoxygenase-1 in human colorectal cancer. *Cancer Res* **59**: 360-6.
- Ilyas, M., Straub, J., Tomlinson, I. P. und Bodmer, W. F. (1999) Genetic pathways in colorectal and other cancers. *Eur J Cancer* **35**: 1986-2002.
- Inan, M. S., Rasoulpour, R. J., Yin, L., Hubbard, A. K., Rosenberg, D. W. und Giardina, C. (2000) The luminal short-chain fatty acid butyrate modulates NF-kappaB activity in a human colonic epithelial cell line. *Gastroenterology* **118**: 724-34.
- Ionov, Y., Peinado, M. A., Malkhosyan, S., Shibata, D. und Perucho, M. (1993) Ubiquitous somatic mutations in simple repeated sequences reveal a new mechanism for colonic carcinogenesis. *Nature* **363**: 558-61.
- Jackson, L. M., Wu, K. C., Mahida, Y. R., Jenkins, D. und Hawkey, C. J. (2000) Cyclooxygenase (COX) 1 and 2 in normal, inflamed, and ulcerated human gastric mucosa. *Gut* **47**: 762-70.
- Jacobasch, G., Schmiedl, D., Kruschewski, M. und Schmehl, K. (1999) Dietary resistant starch and chronic inflammatory bowel diseases. *Int J Colorectal Dis* **14**: 201-11.
- Jacobasch, G. und Dongowski, G. (2000) Ballaststoffe mit spezifischen Wirkungen. In *Praxishandbuch Functional Food*, (eds. Erbersdobler, H. F. und Meyer, A. H.), pp. 1-34, Behr's Verlag, Hamburg.

- Jacobasch, G. und Schmiedl, D. (2002) Die Bedeutung resistenter Stärken für eine gesundheitsorientierte Ernährung. *Ernährungs-Umschau* **49**: 4-9.
- Jaiswal, M., LaRusso, N. F., Burgart, L. J. und Gores, G. J. (2000) Inflammatory cytokines induce DNA damage and inhibit DNA repair in cholangiocarcinoma cells by a nitric oxide-dependent mechanism. *Cancer Res* **60**: 184-90.
- Jaiswal, M., LaRusso, N. F. und Gores, G. J. (2001) Nitric oxide in gastrointestinal epithelial cell carcinogenesis: linking inflammation to oncogenesis. *Am J Physiol Gastrointest Liver Physiol* **281**: G626-34.
- Jakob, U., Gaestel, M., Engel, K. und Buchner, J. (1993) Small heat shock proteins are molecular chaperones. *J Biol Chem* **268**: 1517-20.
- Jenab, M., Chen, J. und Thompson, L. U. (2001) Sialomucin production in aberrant crypt foci relates to degree of dysplasia and rate of cell proliferation. *Cancer Lett* **165**: 19-25.
- Jenkins, D. C., Charles, I. G., Thomsen, L. L., Moss, D. W., Holmes, L. S., Baylis, S. A., Rhodes, P., Westmore, K., Emson, P. C. und Moncada, S. (1995) Roles of nitric oxide in tumor growth. *Proc Natl Acad Sci U S A* **92**: 4392-6.
- Kadmon, M., Tandara, A. und Herfarth, C. (2001) Duodenal adenomatosis in familial adenomatous polyposis coli. A review of the literature and results from the Heidelberg Polyposis Register. *Int J Colorectal Dis* **16**: 63-75.
- Kamitani, H., Geller, M. und Eling, T. (1998) Expression of 15-lipoxygenase by human colorectal carcinoma Caco-2 cells during apoptosis and cell differentiation. *J Biol Chem* **273**: 21569-77.
- Kankuri, E., Vaali, K., Knowles, R. G., Lahde, M., Korpela, R., Vapaatalo, H. und Moilanen, E. (2001a) Suppression of acute experimental colitis by a highly selective inducible nitric-oxide synthase inhibitor, N-[3-(aminomethyl)benzyl]acetamidine. *J Pharmacol Exp Ther* **298**: 1128-32.
- Kankuri, E., Vaali, K., Korpela, R., Paakkari, I., Vapaatalo, H. und Moilanen, E. (2001b) Effects of a COX-2 preferential agent nimesulide on TNBS-induced acute inflammation in the gut. *Inflammation* **25**: 301-10.
- Kayanoki, Y., Fujii, J., Islam, K. N., Suzuki, K., Kawata, S., Matsuzawa, Y. und Taniguchi, N. (1996) The protective role of glutathione peroxidase in apoptosis induced by reactive oxygen species. *J Biochem (Tokyo)* **119**: 817-22.
- Kelner, M. J., Bagnell, R. D., Montoya, M. A. und Lanham, K. A. (2000) Structural organization of the human gastrointestinal glutathione peroxidase (GPX2) promoter and 3'-nontranscribed region: transcriptional response to exogenous redox agents. *Gene* **248**: 109-16.
- Khan, I., Al-Awadi, F. M., Thomas, N., Haridas, S. und Anim, J. T. (2002) Cyclooxygenase-2 inhibition and experimental colitis: beneficial effects of phosphorothioated antisense oligonucleotide and meloxicam. *Scand J Gastroenterol* **37**: 1428-36.
- Khan, K. N., Masferrer, J. L., Woerner, B. M., Soslow, R. und Koki, A. T. (2001) Enhanced cyclooxygenase-2 expression in sporadic and familial adenomatous polyposis of the human colon. *Scand J Gastroenterol* **36**: 865-9.
- Kilty, I., Logan, A. und Vickers, P. J. (1999) Differential characteristics of human 15-lipoxygenase isozymes and a novel splice variant of 15S-lipoxygenase. *Eur J Biochem* **266**: 83-93.
- Kim, H., Jen, J., Vogelstein, B. und Hamilton, S. R. (1994) Clinical and pathological characteristics of sporadic colorectal carcinomas with DNA replication errors in microsatellite sequences. *Am J Pathol* **145**: 148-56.
- Kimura, H., Hokari, R., Miura, S., Shigematsu, T., Hirokawa, M., Akiba, Y., Kurose, I., Higuchi, H., Fujimori, H., Tsuzuki, Y., Serizawa, H. und Ishii, H. (1998) Increased expression of an inducible isoform of nitric oxide synthase and the formation of peroxynitrite in colonic mucosa of patients with active ulcerative colitis. *Gut* **42**: 180-7.
- Kitamura, T., Kawamori, T., Uchiya, N., Itoh, M., Noda, T., Matsuura, M., Sugimura, T. und Wakabayashi, K. (2002) Inhibitory effects of mofezolac, a cyclooxygenase-1 selective inhibitor, on intestinal carcinogenesis. *Carcinogenesis* **23**: 1463-6.
- Knudson, A. G., Jr. (1971) Mutation and cancer: statistical study of retinoblastoma. *Proc Natl Acad Sci U S A* **68**: 820-3.

- Kobayashi, O., Miwa, H., Watanabe, S., Tsujii, M., Dubois, R. N. und Sato, N. (2001) Cyclooxygenase-2 downregulates inducible nitric oxide synthase in rat intestinal epithelial cells. *Am J Physiol Gastrointest Liver Physiol* **281**: G688-96.
- Komatsu, H., Okayasu, I., Mitomi, H., Imai, H., Nakagawa, Y. und Obata, F. (2001) Immunohistochemical detection of human gastrointestinal glutathione peroxidase in normal tissues and cultured cells with novel mouse monoclonal antibodies. *J Histochem Cytochem* **49**: 759-66.
- Korinek, V., Barker, N., Morin, P. J., van Wichen, D., de Weger, R., Kinzler, K. W., Vogelstein, B. und Clevers, H. (1997) Constitutive transcriptional activation by a beta-catenin-Tcf complex in APC-/- colon carcinoma. *Science* **275**: 1784-7.
- Krämer, K., Hoppe P. und Packer, L., eds. (1997) Nutraceuticals in Health and Disease., Marcel Dekker, Inc, New York.
- Krieg, P., Kinzig, A., Heidt, M., Marks, F. und Furstenberger, G. (1998) cDNA cloning of a 8-lipoxygenase and a novel epidermis-type lipoxygenase from phorbol ester-treated mouse skin. *Biochim Biophys Acta* **1391**: 7-12.
- Kruis, W. (2001) Klinische Bedeutung von Mutaflo® (E. coli Stamm Nissle 1917) in der Therapie der Colitis Ulcerosa und des Morbus Crohn. In *Darmflora in Symbiose und Pathogenität*, Alfred-Nissle-Gesellschaft, e.v. 58089 Hagen.
- Kruschewski, M., Foitzik, T., Perez-Canto, A., Hubotter, A. und Buhr, H. J. (2001) Changes of colonic mucosal microcirculation and histology in two colitis models: an experimental study using intravital microscopy and a new histological scoring system. *Dig Dis Sci* **46**: 2336-43.
- Kubes, P. (2000) Inducible nitric oxide synthase: a little bit of good in all of us. *Gut* **47**: 6-9.
- Kühn, H. (1996) Biosynthesis, metabolism and biological importance of the primary 15-lipoxygenase metabolites 15-hydro(pero)XY-5Z,8Z,11Z,13E-eicosatetraenoic acid and 13-hydro(pero)XY-9Z,11E-octadecadienoic acid. *Prog Lipid Res* **35**: 203-26.
- Kühn, H. und Chan, L. (1997) The role of 15-lipoxygenase in atherogenesis: pro- and antiatherogenic actions. *Curr Opin Lipidol* **8**: 111-7.
- Kühn, H. und Thiele, B. J. (1999) The diversity of the lipoxygenase family. Many sequence data but little information on biological significance. *FEBS Lett* **449**: 7-11.
- Kühn, H. und Borchert, A. (2002) Regulation of enzymatic lipid peroxidation: the interplay of peroxidizing and peroxide reducing enzymes. *Free Radic Biol Med* **33**: 154-72.
- Kyo, K., Parkes, M., Takei, Y., Nishimori, H., Vyas, P., Satsangi, J., Simmons, J., Nagawa, H., Baba, S., Jewell, D., Muto, T., Lathrop, G. M. und Nakamura, Y. (1999) Association of ulcerative colitis with rare VNTR alleles of the human intestinal mucin gene, MUC3. *Hum Mol Genet* **8**: 307-11.
- Landino, L. M., Crews, B. C., Timmons, M. D., Morrow, J. D. und Marnett, L. J. (1996) Peroxynitrite, the coupling product of nitric oxide and superoxide, activates prostaglandin biosynthesis. *Proc Natl Acad Sci U S A* **93**: 15069-74.
- Lane, D. P. (1992) Cancer. p53, guardian of the genome. *Nature* **358**: 15-6.
- Laszlo, F. und Whittle, B. J. (1997) Actions of isoform-selective and non-selective nitric oxide synthase inhibitors on endotoxin-induced vascular leakage in rat colon. *Eur J Pharmacol* **334**: 99-102.
- Leach, F. S., Nicolaides, N. C., Papadopoulos, N., Liu, B., Jen, J., Parsons, R., Peltomaki, P., Sistonen, P., Aaltonen, L. A., Nystrom-Lahti, M. und et al. (1993) Mutations of a mutS homolog in hereditary nonpolyposis colorectal cancer. *Cell* **75**: 1215-25.
- Lehmann, U. (2001) Herstellung und Charakterisierung von resistentem Stärke aus verschiedenen Quellen - Überprüfung gesundheitsfördernder Effekte auf die Darmschleimhaut bei Kontrollpersonen und Patienten mit Colitis ulcerosa. *Biologisch-Pharmazeutische Fakultät der Friedrich-Schiller-Universität Jena, Inauguraldiss.*
- Lehmann, U., Jacobasch, G. und Schmiedl, D. (2002) Characterization of resistant starch type III from banana (*Musa acuminata*). *J Agric Food Chem* **50**: 5236-40.
- Lehmann, U., Rossler, C., Schmiedl, D. und Jacobasch, G. (2003) Production and physicochemical characterization of resistant starch type III derived from pea starch. *Nahrung* **47**: 60-3.

- Leuratti, C., Watson, M. A., Deag, E. J., Welch, A., Singh, R., Gottschalg, E., Marnett, L. J., Atkin, W., Day, N. E., Shuker, D. E. und Bingham, S. A. (2002) Detection of malondialdehyde DNA adducts in human colorectal mucosa: relationship with diet and the presence of adenomas. *Cancer Epidemiol Biomarkers Prev* **11**: 267-73.
- Licinio, J., Prolo, P., McCann, S. M. und Wong, M. L. (1999) Brain iNOS: current understanding and clinical implications. *Mol Med Today* **5**: 225-32.
- Lin, K. M., Shashidharan, M., Thorson, A. G., Ternent, C. A., Blatchford, G. J., Christensen, M. A., Watson, P., Lemon, S. J., Franklin, B., Karr, B., Lynch, J. und Lynch, H. T. (1998) Cumulative incidence of colorectal and extracolonic cancers in MLH1 and MSH2 mutation carriers of hereditary nonpolyposis colorectal cancer. *J Gastrointest Surg* **2**: 67-71.
- Liu, Q., Chan, S. T. und Mahendran, R. (2003) Nitric oxide induces cyclooxygenase expression and inhibits cell growth in colon cancer cell lines. *Carcinogenesis* **24**: 637-42.
- Llor, X., Jacoby, R. F., Teng, B. B., Davidson, N. O., Sitrin, M. D. und Brasitus, T. A. (1991) K-ras mutations in 1,2-dimethylhydrazine-induced colonic tumors: effects of supplemental dietary calcium and vitamin D deficiency. *Cancer Res* **51**: 4305-9.
- Louis, E., Ribbens, C., Godon, A., Franchimont, D., De Groote, D., Hardy, N., Boniver, J., Belaiche, J. und Malaise, M. (2000) Increased production of matrix metalloproteinase-3 and tissue inhibitor of metalloproteinase-1 by inflamed mucosa in inflammatory bowel disease. *Clin Exp Immunol* **120**: 241-6.
- Lundberg, J. O., Hellstrom, P. M., Lundberg, J. M. und Alving, K. (1994) Greatly increased luminal nitric oxide in ulcerative colitis. *Lancet* **344**: 1673-4.
- Lynch, H. T., Smyrk, T., McGinn, T., Lanspa, S., Cavalieri, J., Lynch, J., Slominski-Castor, S., Cayouette, M. C., Priluck, I. und Luce, M. C. (1995) Attenuated familial adenomatous polyposis (AFAP). A phenotypically and genotypically distinctive variant of FAP. *Cancer* **76**: 2427-33.
- Ma, Q. Y., Williamson, K. E. und Rowlands, B. J. (2002) Variability of cell proliferation in the proximal and distal colon of normal rats and rats with dimethylhydrazine induced carcinogenesis. *World J Gastroenterol* **8**: 847-52.
- MacDonald, T. T. und Pender, S. L. (1998) Lamina propria T cells. *Chem Immunol* **71**: 103-17.
- MacDonald, T. T., Monteleone, G. und Pender, S. L. (2000) Recent developments in the immunology of inflammatory bowel disease. *Scand J Immunol* **51**: 2-9.
- Maiorino, M., Aumann, K. D., Brigelius-Flohé, R., Doria, D., van den Heuvel, J., McCarthy, J., Roveri, A., Ursini, F. und Flohé, L. (1995) Probing the presumed catalytic triad of selenium-containing peroxidases by mutational analysis of phospholipid hydroperoxide glutathione peroxidase (PHGPx). *Biol Chem Hoppe Seyler* **376**: 651-60.
- Marinović, M. (2003) Verhalten der PKC-Isoenzymmuster, des NF-kappaB und des beta-Catenins bei der kolorektalen Kanzerogenese und CED - Effekt von RS III auf die Signalübertragung und den Krankheitsverlauf. *Mathematisch-Naturwissenschaftliche Fakultät der Universität Potsdam, Inauguraldiss.*
- Markowitz, S. (2000) TGF-beta receptors and DNA repair genes, coupled targets in a pathway of human colon carcinogenesis. *Biochim Biophys Acta* **1470**: M13-20.
- Marnett, L. J. (1994) DNA adducts of alpha,beta-unsaturated aldehydes and dicarbonyl compounds. *IARC Sci Publ* **125**: 151-63.
- Marnett, L. J., Riggins, J. N. und West, J. D. (2003) Endogenous generation of reactive oxidants and electrophiles and their reactions with DNA and protein. *J Clin Invest* **111**: 583-93.
- Masferrer, J. L., Isakson, P. C. und Seibert, K. (1996) Cyclooxygenase-2 inhibitors: a new class of anti-inflammatory agents that spare the gastrointestinal tract. *Gastroenterol Clin North Am* **25**: 363-72.
- Masferrer, J. L., Leahy, K. M., Koki, A. T., Zweifel, B. S., Settle, S. L., Woerner, B. M., Edwards, D. A., Flickinger, A. G., Moore, R. J. und Seibert, K. (2000) Antiangiogenic and antitumor activities of cyclooxygenase-2 inhibitors. *Cancer Res* **60**: 1306-11.

- Maziere, S., Meflah, K., Tavan, E., Champ, M., Narbonne, J. F. und Cassand, P. (1998) Effect of resistant starch and/or fat-soluble vitamins A and E on the initiation stage of aberrant crypts in rat colon. *Nutr Cancer* **31**: 168-77.
- McCafferty, D. M., Miampamba, M., Sihota, E., Sharkey, K. A. und Kubes, P. (1999) Role of inducible nitric oxide synthase in trinitrobenzene sulphonic acid induced colitis in mice. *Gut* **45**: 864-73.
- McLellan, E. A., Medline, A. und Bird, R. P. (1991) Dose response and proliferative characteristics of aberrant crypt foci: putative preneoplastic lesions in rat colon. *Carcinogenesis* **12**: 2093-8.
- Mei, J. M., Hord, N. G., Winterstein, D. F., Donald, S. P. und Phang, J. M. (2000) Expression of prostaglandin endoperoxide H synthase-2 induced by nitric oxide in conditionally immortalized murine colonic epithelial cells. *Faseb J* **14**: 1188-201.
- Mendeloff, A. I., Calkin, B.M. (1988) The epidemiology of idiopathic inflammatory Bowel disease. In *Inflammatory Bowel Disease*, (eds. Kirsner JB und RG, S.), Lea & Febiger, Philadelphia.
- Miller, C. C. und Ziboh, V. A. (1990) Induction of epidermal hyperproliferation by topical n-3 polyunsaturated fatty acids on guinea pig skin linked to decreased levels of 13-hydroxyoctadecadienoic acid (13-hode). *J Invest Dermatol* **94**: 353-8.
- Monsen, U., Sorstad, J., Hellers, G. und Johansson, C. (1990) Extracolonic diagnoses in ulcerative colitis: an epidemiological study. *Am J Gastroenterol* **85**: 711-6.
- Monteleone, G., Kumberova, A., Croft, N. M., McKenzie, C., Steer, H. W. und MacDonald, T. T. (2001) Blocking Smad7 restores TGF-beta1 signaling in chronic inflammatory bowel disease. *J Clin Invest* **108**: 601-9.
- Monteleone, I., Vavassori, P., Biancone, L., Monteleone, G. und Pallone, F. (2002) Immunoregulation in the gut: success and failures in human disease. *Gut* **50 Suppl 3**: III60-4.
- Mörk, H., al-Taie, O. H., Bahr, K., Zierer, A., Beck, C., Scheurlen, M., Jakob, F. und Kohrle, J. (2000) Inverse mRNA expression of the selenocysteine-containing proteins GI-GPx and SeP in colorectal adenomas compared with adjacent normal mucosa. *Nutr Cancer* **37**: 108-16.
- Morris, G. P., Beck, P. L., Herridge, M. S., Depew, W. T., Szewczuk, M. R. und Wallace, J. L. (1989) Hapten-induced model of chronic inflammation and ulceration in the rat colon. *Gastroenterology* **96**: 795-803.
- Morteau, O. (1999) COX-2: promoting tolerance. *Nat Med* **5**: 867-8.
- Morteau, O., Morham, S. G., Sellon, R., Dieleman, L. A., Langenbach, R., Smithies, O. und Sartor, R. B. (2000) Impaired mucosal defense to acute colonic injury in mice lacking cyclooxygenase-1 or cyclooxygenase-2. *J Clin Invest* **105**: 469-78.
- Müller-Decker, K., Albert, C., Lukanov, T., Winde, G., Marks, F. und Furstenberger, G. (1999) Cellular localization of cyclo-oxygenase isozymes in Crohn's disease and colorectal cancer. *Int J Colorectal Dis* **14**: 212-8.
- Nancey, S., Bienvenu, J., Coffin, B., Andre, F., Descos, L. und Flourie, B. (2002) Butyrate strongly inhibits in vitro stimulated release of cytokines in blood. *Dig Dis Sci* **47**: 921-8.
- Naom, I., Lee, J., Ford, D., Bowman, S. J., Lanchbury, J. S., Haris, I., Hodgson, S. V., Easton, D., Lennard-Jones, J. und Mathew, C. G. (1996) Analysis of the contribution of HLA genes to genetic predisposition in inflammatory bowel disease. *Am J Hum Genet* **59**: 226-33.
- Nathan, C. und Xie, Q. W. (1994) Nitric oxide synthases: roles, tolls, and controls. *Cell* **78**: 915-8.
- Nishiyama, R., Sakaguchi, T., Kinugasa, T., Gu, X., MacDermott, R. P., Podolsky, D. K. und Reinecker, H. C. (2001) Interleukin-2 receptor beta subunit-dependent and -independent regulation of intestinal epithelial tight junctions. *J Biol Chem* **276**: 35571-80.
- Nixon, J. B. und Eling, T. E. (2001) 15-lipoxygenase-1 and 15-lipoxygenase-2 may have different and opposing biological functions. *abrufbar über:* <http://www.oxfordbiomed.com/News&Reviews/invrev4.htm> (Zugriff: 28.09.01).

- Nomura, K., Imai, H., Koumura, T., Arai, M. und Nakagawa, Y. (1999) Mitochondrial phospholipid hydroperoxide glutathione peroxidase suppresses apoptosis mediated by a mitochondrial death pathway. *J Biol Chem* **274**: 29294-302.
- Nozoe, T., Yasuda, M., Honda, M., Inutsuka, S. und Korenaga, D. (2002) Immunohistochemical expression of cytokine induced nitric oxide synthase in colorectal carcinoma. *Oncol Rep* **9**: 521-4.
- O'Donnell, V. B., Taylor, K. B., Parthasarathy, S., Kuhn, H., Koesling, D., Fribe, A., Bloodsworth, A., Darley-Usmar, V. M. und Freeman, B. A. (1999) 15-Lipoxygenase catalytically consumes nitric oxide and impairs activation of guanylate cyclase. *J Biol Chem* **274**: 20083-91.
- Ogura, Y., Bonen, D. K., Inohara, N., Nicolae, D. L., Chen, F. F., Ramos, R., Britton, H., Moran, T., Karaliuskas, R., Duerr, R. H., Achkar, J. P., Brant, S. R., Bayless, T. M., Kirschner, B. S., Hanauer, S. B., Nunez, G. und Cho, J. H. (2001) A frameshift mutation in NOD2 associated with susceptibility to Crohn's disease. *Nature* **411**: 603-6.
- Oshima, M., Dinchuk, J. E., Kargman, S. L., Oshima, H., Hancock, B., Kwong, E., Trzaskos, J. M., Evans, J. F. und Taketo, M. M. (1996) Suppression of intestinal polyposis in Apc delta716 knockout mice by inhibition of cyclooxygenase 2 (COX-2). *Cell* **87**: 803-9.
- Ota, S., Bamba, H., Kato, A., Kawamoto, C., Yoshida, Y. und Fujiwara, K. (2002) Review article: COX-2, prostanoids and colon cancer. *Aliment Pharmacol Ther* **16 Suppl 2**: 102-6.
- Otto, H. F. und Remmeli, W. (1996) Kolon und Rektum. In *Pathologie 2 - Verdauungstrakt*, (ed. Remmeli, W.), pp. 533-676, Springer Verlag, Berlin Heidelberg New York.
- Packham, G., Ashmun, R. A. und Cleveland, J. L. (1996) Cytokines suppress apoptosis independent of increases in reactive oxygen levels. *J Immunol* **156**: 2792-800.
- Palmer, R. M., Ferrige, A. G. und Moncada, S. (1987) Nitric oxide release accounts for the biological activity of endothelium-derived relaxing factor. *Nature* **327**: 524-6.
- Panwala, C. M., Jones, J. C. und Viney, J. L. (1998) A novel model of inflammatory bowel disease: mice deficient for the multiple drug resistance gene, mdr1a, spontaneously develop colitis. *J Immunol* **161**: 5733-44.
- Papadopoulos, N., Nicolaides, N. C., Wei, Y. F., Ruben, S. M., Carter, K. C., Rosen, C. A., Haseltine, W. A., Fleischmann, R. D., Fraser, C. M., Adams, M. D. und et al. (1994) Mutation of a mutL homolog in hereditary colon cancer. *Science* **263**: 1625-9.
- Park, H. S., Goodlad, R. A. und Wright, N. A. (1997) The incidence of aberrant crypt foci and colonic carcinoma in dimethylhydrazine-treated rats varies in a site-specific manner and depends on tumor histology. *Cancer Res* **57**: 4507-10.
- Parks, A. G. und Nicholls, R. J. (1978) Proctocolectomy without ileostomy for ulcerative colitis. *Br Med J* **2**: 85-8.
- Pellizzaro, C., Coradini, D. und Daidone, M. G. (2002) Modulation of angiogenesis-related proteins synthesis by sodium butyrate in colon cancer cell line HT29. *Carcinogenesis* **23**: 735-40.
- Peltomaki, P. (2001) Deficient DNA mismatch repair: a common etiologic factor for colon cancer. *Hum Mol Genet* **10**: 735-40.
- Perner, A., Andresen, L., Normark, M., Fischer-Hansen, B., Sorensen, S., Eugen-Olsen, J. und Rask-Madsen, J. (2001) Expression of nitric oxide synthases and effects of L-arginine and L-NMMA on nitric oxide production and fluid transport in collagenous colitis. *Gut* **49**: 387-94.
- Perner, A., Andresen, L., Normark, M. und Rask-Madsen, J. (2002) Constitutive expression of inducible nitric oxide synthase in the normal human colonic epithelium. *Scand J Gastroenterol* **37**: 944-8.
- Perrin, P., Pierre, F., Patry, Y., Champ, M., Berreur, M., Pradal, G., Bornet, F., Meflah, K. und Menanteau, J. (2001) Only fibres promoting a stable butyrate producing colonic ecosystem decrease the rate of aberrant crypt foci in rats. *Gut* **48**: 53-61.
- Pfeiffer, C. J. und Qiu, B. S. (1995) Effects of chronic nitric oxide synthase inhibition on TNB-induced colitis in rats. *J Pharm Pharmacol* **47**: 827-32.

- Plevy, S. E., Targan, S. R., Yang, H., Fernandez, D., Rotter, J. I. und Toyoda, H. (1996) Tumor necrosis factor microsatellites define a Crohn's disease-associated haplotype on chromosome 6. *Gastroenterology* **110**: 1053-60.
- Polyak, K., Xia, Y., Zweier, J. L., Kinzler, K. W. und Vogelstein, B. (1997) A model for p53-induced apoptosis. *Nature* **389**: 300-5.
- Pomare, E. W., Branch, W. J. und Cummings, J. H. (1985) Carbohydrate fermentation in the human colon and its relation to acetate concentrations in venous blood. *J Clin Invest* **75**: 1448-54.
- Powell, D. W., Mifflin, R. C., Valentich, J. D., Crowe, S. E., Saada, J. I. und West, A. B. (1999) Myofibroblasts. II. Intestinal subepithelial myofibroblasts. *Am J Physiol* **277**: C183-201.
- Rachmilewitz, D., Karmeli, F., Okon, E. und Bursztyn, M. (1995) Experimental colitis is ameliorated by inhibition of nitric oxide synthase activity. *Gut* **37**: 247-55.
- Rao, C. V., Kawamori, T., Hamid, R. und Reddy, B. S. (1999) Chemoprevention of colonic aberrant crypt foci by an inducible nitric oxide synthase-selective inhibitor. *Carcinogenesis* **20**: 641-4.
- Read, N. W., Krejs, G. J., Read, M. G., Santa Ana, C. A., Morawski, S. G. und Fordtran, J. S. (1980) Chronic diarrhea of unknown origin. *Gastroenterology* **78**: 264-71.
- Reiter, C. D., Teng, R. J. und Beckman, J. S. (2000) Superoxide reacts with nitric oxide to nitrate tyrosine at physiological pH via peroxynitrite. *J Biol Chem* **275**: 32460-6.
- Ren, H., Musch, M. W., Kojima, K., Boone, D., Ma, A. und Chang, E. B. (2001) Short-chain fatty acids induce intestinal epithelial heat shock protein 25 expression in rats and IEC 18 cells. *Gastroenterology* **121**: 631-9.
- Reuter, B. K., Asfaha, S., Buret, A., Sharkey, K. A. und Wallace, J. L. (1996) Exacerbation of inflammation-associated colonic injury in rat through inhibition of cyclooxygenase-2. *J Clin Invest* **98**: 2076-85.
- Roberts, P. J., Morgan, K., Miller, R., Hunter, J. O. und Middleton, S. J. (2001a) Neuronal COX-2 expression in human myenteric plexus in active inflammatory bowel disease. *Gut* **48**: 468-72.
- Roberts, P. J., Riley, G. P., Morgan, K., Miller, R., Hunter, J. O. und Middleton, S. J. (2001b) The physiological expression of inducible nitric oxide synthase (iNOS) in the human colon. *J Clin Pathol* **54**: 293-7.
- Roediger, W. E. W. (1980) Role of anaerobic bacteria in the metabolic welfare of the colonic mucosa in man. *Gut* **21**.
- Ropponen, K. M., Kellokoski, J. K., Lipponen, P. K., Eskelinen, M. J., Alanne, L., Alhava, E. M. und Kosma, V. M. (2000) Expression of inducible nitric oxide synthase in colorectal cancer and its association with prognosis. *Scand J Gastroenterol* **35**: 1204-11.
- Ross, C. A., McInnis, M. G., Margolis, R. L. und Li, S. H. (1993) Genes with triplet repeats: candidate mediators of neuropsychiatric disorders. *Trends Neurosci* **16**: 254-60.
- Rössler, C. (2002a) Herstellung und Charakterisierung von resistenter Stärke aus Getreide und ihr Einfluss auf die Gallensäuren bei der TNBS-Rattenkolitis. *Fachbereich Humanmedizin der Freien Universität Berlin, Inauguraldiss.*
- Rössler, C., Jacobasch, G., Schmiedl, D., Gebhardt, E. (2002b) Resistente Stärke Type 3 aus Roggenstärke, ein funktionelles Lebensmittel. *Ernährung/Nutrition* **26**.
- Rotruck, J. T. A., Pope, L., Ganther H.E. et al. (1973) Selenium; biological role as a component of glutathion peroxidase. *Science* **179**: 588-599.
- Sakamoto, C. (1998) Roles of COX-1 and COX-2 in gastrointestinal pathophysiology. *J Gastroenterol* **33**: 618-24.
- Sales, K. J., Katz, A. A., Howard, B., Soeters, R. P., Millar, R. P. und Jabbour, H. N. (2002) Cyclooxygenase-1 is up-regulated in cervical carcinomas: autocrine/paracrine regulation of cyclooxygenase-2, prostaglandin e receptors, and angiogenic factors by cyclooxygenase-1. *Cancer Res* **62**: 424-32.
- Salminen, S., Bouley, C., Boutron-Ruault, M. C., Cummings, J. H., Franck, A., Gibson, G. R., Isolauri, E., Moreau, M. C., Roberfroid, M. und Rowland, I. (1998) Functional food science and gastrointestinal physiology and function. *Br J Nutr* **80 Suppl 1**: S147-71.

- Sanderson, J. D., Moss, M. T., Tizard, M. L. und Hermon-Taylor, J. (1992) Mycobacterium paratuberculosis DNA in Crohn's disease tissue. *Gut* **33**: 890-6.
- Sano, H., Kawahito, Y., Wilder, R. L., Hashiramoto, A., Mukai, S., Asai, K., Kimura, S., Kato, H., Kondo, M. und Hla, T. (1995) Expression of cyclooxygenase-1 and -2 in human colorectal cancer. *Cancer Res* **55**: 3785-9.
- Sansonetti, P. (2002) Host-pathogen interactions: the seduction of molecular cross talk. *Gut* **50 Suppl 3**: III2-8.
- Sato, F., Harpaz, N., Shibata, D., Xu, Y., Yin, J., Mori, Y., Zou, T. T., Wang, S., Desai, K., Leytin, A., Selaru, F. M., Abraham, J. M. und Meltzer, S. J. (2002) Hypermethylation of the p14(ARF) gene in ulcerative colitis-associated colorectal carcinogenesis. *Cancer Res* **62**: 1148-51.
- Sato, M. und Ahnen, D. J. (1992) Regional variability of colonocyte growth and differentiation in the rat. *Anat Rec* **233**: 409-14.
- Satsangi, J., Parkes, M., Louis, E., Hashimoto, L., Kato, N., Welsh, K., Terwilliger, J. D., Lathrop, G. M., Bell, J. I. und Jewell, D. P. (1996a) Two stage genome-wide search in inflammatory bowel disease provides evidence for susceptibility loci on chromosomes 3, 7 and 12. *Nat Genet* **14**: 199-202.
- Satsangi, J., Welsh, K. I., Bunce, M., Julier, C., Farrant, J. M., Bell, J. I. und Jewell, D. P. (1996b) Contribution of genes of the major histocompatibility complex to susceptibility and disease phenotype in inflammatory bowel disease. *Lancet* **347**: 1212-7.
- Saur, D., Neuhuber, W. L., Gengenbach, B., Huber, A., Schusdziarra, V. und Allescher, H. D. (2002) Site-specific gene expression of nNOS variants in distinct functional regions of rat gastrointestinal tract. *Am J Physiol Gastrointest Liver Physiol* **282**: G349-58.
- Scheppach, W., Bartram, P., Richter, A., Richter, F., Liepold, H., Dusel, G., Hofstetter, G., Ruthlein, J. und Kasper, H. (1992a) Effect of short-chain fatty acids on the human colonic mucosa in vitro. *JPEN J Parenter Enteral Nutr* **16**: 43-8.
- Scheppach, W., Sommer, H., Kirchner, T., Paganelli, G. M., Bartram, P., Christl, S., Richter, F., Dusel, G. und Kasper, H. (1992b) Effect of butyrate enemas on the colonic mucosa in distal ulcerative colitis. *Gastroenterology* **103**: 51-6.
- Scheppach, W. (1996) Treatment of distal ulcerative colitis with short-chain fatty acid enemas. A placebo-controlled trial. German-Austrian SCFA Study Group. *Dig Dis Sci* **41**: 2254-9.
- Scheppach W., P., M.E., Richter, F. (1997) Ballaststoffe, kurzkettige Fettsäuren und Kolonkarzinom. *Akt. Ernähr.-Med.* **22**: 321-326.
- Schewe, T. (2002) 15-lipoxygenase-1: a prooxidant enzyme. *Biol Chem* **383**: 365-74.
- Schmiedl, D. (1995) Zum Einfluss der Retrogradation von Stärken und Stärkehydrolyseprodukten auf die Bildung und Struktur resisterter Stärke. *Fachbereich Lebensmittelwissenschaft und Biotechnologie der Technischen Universität Berlin, Inauguraldiss.*
- Schmiedl, D., Jacobasch, G., Bäuerlein, M., Riesmeier, J. und Bengs, H. (1998)  $\alpha$ -Amylase-resistente Polysaccharide, Herstellungsverfahren, Verwendung und Lebensmittel mit diesen Polysacchariden. *Patentanmeldung DE 198 30618.0, angem. 30.06.1998.*
- Schmiedl, D., Jacobasch, G. und König, B. (1999a) Verfahren zur Erhöhung des Gehalts an  $\alpha$ -Amylase-resistenter Stärke (RS-Gehalt) eines Polysaccharids, Polysaccharide, deren Verwendung und Lebensmittel mit diesen Polysacchariden. *Patentanmeldung EP 0012459, angem. 09.12.1999.*
- Schmiedl, D., König, B., Jacobasch, G. und Döring, M. (1999b) Herstellung von RS und deren Charakterisierung aus Kartoffel. *Patentanmeldung DE 199 11001.8-43, angem. 15.06.1999.*
- Schmiedl, D., Bäuerlein, M., Bengs, H. und Jacobasch, G. (2000) Production of heat-stable, butyrogenic resistant starch. *Carbohydrate Polymers* **43**: 183-193.
- Schmutzler, G. (2001) Research Tools for Nitric Oxide. *The Scientist* **15**: 23.
- Schottelius, A. J. und Baldwin, A. S., Jr. (1999) A role for transcription factor NF-kappa B in intestinal inflammation. *Int J Colorectal Dis* **14**: 18-28.
- Schwab, J. M., Schluesener, H. J. und Laufer, S. (2003a) COX-3: just another COX or the solitary elusive target of paracetamol? *Lancet* **361**: 981-2.

- Schwab, M., Schaeffeler, E., Marx, C., Fromm, M. F., Kaskas, B., Metzler, J., Stange, E., Herfarth, H., Schoelmerich, J., Gregor, M., Walker, S., Cascorbi, I., Roots, I., Brinkmann, U., Zanger, U. M. und Eichelbaum, M. (2003b) Association between the C3435T MDR1 gene polymorphism and susceptibility for ulcerative colitis. *Gastroenterology* **124**: 26-33.
- Scott, D. J., Hull, M. A., Cartwright, E. J., Lam, W. K., Tisbury, A., Poulsom, R., Markham, A. F., Bonifer, C. und Coletta, P. L. (2001) Lack of inducible nitric oxide synthase promotes intestinal tumorigenesis in the Apc(Min $^{+/-}$ ) mouse. *Gastroenterology* **121**: 889-99.
- Segain, J. P., Raingeard de la Bletiere, D., Bourreille, A., Leray, V., Gervois, N., Rosales, C., Ferrier, L., Bonnet, C., Blottiere, H. M. und Galmiche, J. P. (2000) Butyrate inhibits inflammatory responses through NFkappaB inhibition: implications for Crohn's disease. *Gut* **47**: 397-403.
- Sellin, J. H. (1999) SCFAs: The Enigma of Weak Electrolyte Transport in the Colon. *News Physiol Sci* **14**: 58-64.
- Shamsuddin, A. K. und Trump, B. F. (1981) Colon epithelium. I. Light microscopic, histochemical, and ultrastructural features of normal colon epithelium of male Fischer 344 rats. *J Natl Cancer Inst* **66**: 375-88.
- Shao, J., Sheng, H., Aramandla, R., Pereira, M. A., Lubet, R. A., Hawk, E., Grogan, L., Kirsch, I. R., Washington, M. K., Beauchamp, R. D. und DuBois, R. N. (1999) Coordinate regulation of cyclooxygenase-2 and TGF-beta1 in replication error-positive colon cancer and azoxymethane-induced rat colonic tumors. *Carcinogenesis* **20**: 185-91.
- Shappell, S. B., Boeglin, W. E., Olson, S. J., Kasper, S. und Brash, A. R. (1999) 15-lipoxygenase-2 (15-LOX-2) is expressed in benign prostatic epithelium and reduced in prostate adenocarcinoma. *Am J Pathol* **155**: 235-45.
- Shappell, S. B., Jack, G. S., Keeney, D. S. und Brash, A. R. (2001) 15-LOX-2: Tissue distribution, enzymatic properties, possible biological function and role of reduced expression in prostate cancer. *abruftbar über: http://www.oxfordbiomed.com/News&Reviews/invrev5.htm* (Zugriff: 28.09.01).
- Shattuck-Brandt, R. L., Varilek, G. W., Radhika, A., Yang, F., Washington, M. K. und DuBois, R. N. (2000) Cyclooxygenase 2 expression is increased in the stroma of colon carcinomas from IL-10(-/-) mice. *Gastroenterology* **118**: 337-45.
- Shivananda, S., Lennard-Jones, J., Logan, R., Fear, N., Price, A., Carpenter, L. und van Blankenstein, M. (1996) Incidence of inflammatory bowel disease across Europe: is there a difference between north and south? Results of the European Collaborative Study on Inflammatory Bowel Disease (EC-IBD). *Gut* **39**: 690-7.
- Shureiqi, I., Wojno, K. J., Poore, J. A., Reddy, R. G., Moussalli, M. J., Spindler, S. A., Greenson, J. K., Normolle, D., Hasan, A. A., Lawrence, T. S. und Brenner, D. E. (1999) Decreased 13-S-hydroxyoctadecadienoic acid levels and 15-lipoxygenase-1 expression in human colon cancers. *Carcinogenesis* **20**: 1985-95.
- Shureiqi, I., Chen, D., Lee, J. J., Yang, P., Newman, R. A., Brenner, D. E., Lotan, R., Fischer, S. M. und Lippman, S. M. (2000) 15-LOX-1: a novel molecular target of nonsteroidal anti-inflammatory drug-induced apoptosis in colorectal cancer cells. *J Natl Cancer Inst* **92**: 1136-42.
- Shureiqi, I. und Lippman, S. M. (2001) Lipoxygenase modulation to reverse carcinogenesis. *Cancer Res* **61**: 6307-12.
- Shureiqi, I., Jiang, W., Fischer, S. M., Xu, X., Chen, D., Lee, J. J., Lotan, R. und Lippman, S. M. (2002) GATA-6 transcriptional regulation of 15-lipoxygenase-1 during NSAID-induced apoptosis in colorectal cancer cells. *Cancer Res* **62**: 1178-83.
- Siegert, A., Rosenberg, C., Schmitt, W. D., Denkert, C. und Hauptmann, S. (2002) Nitric oxide of human colorectal adenocarcinoma cell lines promotes tumour cell invasion. *Br J Cancer* **86**: 1310-5.
- Sies, H., ed. (1985). In *Oxidative stress*, Academic Press Inc., London.

- Singer, II, Kawka, D. W., Scott, S., Weidner, J. R., Mumford, R. A., Riehl, T. E. und Stenson, W. F. (1996) Expression of inducible nitric oxide synthase and nitrotyrosine in colonic epithelium in inflammatory bowel disease. *Gastroenterology* **111**: 871-85.
- Singer, II, Kawka, D. W., Schloemann, S., Tessner, T., Riehl, T. und Stenson, W. F. (1998) Cyclooxygenase 2 is induced in colonic epithelial cells in inflammatory bowel disease. *Gastroenterology* **115**: 297-306.
- Smith, W. L., DeWitt, D. L. und Garavito, R. M. (2000) Cyclooxygenases: structural, cellular, and molecular biology. *Annu Rev Biochem* **69**: 145-82.
- Sonoshita, M., Takaku, K., Oshima, M., Sugihara, K. und Taketo, M. M. (2002) Cyclooxygenase-2 expression in fibroblasts and endothelial cells of intestinal polyps. *Cancer Res* **62**: 6846-9.
- Sowa, Y. und Sakai, T. (2003) [Gene-regulating chemoprevention against cancer--as a model for "molecular-targeting prevention" of cancer]. *Nippon Eiseigaku Zasshi* **58**: 267-74.
- Stopera, S. A., Murphy, L. C. und Bird, R. P. (1992) Evidence for a ras gene mutation in azoxymethane-induced colonic aberrant crypts in Sprague-Dawley rats: earliest recognizable precursor lesions of experimental colon cancer. *Carcinogenesis* **13**: 2081-5.
- Strand, M., Prolla, T. A., Liskay, R. M. und Petes, T. D. (1993) Destabilization of tracts of simple repetitive DNA in yeast by mutations affecting DNA mismatch repair. *Nature* **365**: 274-6.
- Sunter, J. P., Appleton, D. R., Wright, N. A. und Watson, A. J. (1978) Pathological features of the colonic tumours induced in rats by the administration of 1,2-dimethylhydrazine. *Virchows Arch B Cell Pathol* **29**: 211-23.
- Sunter, J. P. (1980) Cell proliferation in the gastrointestinal tract. In *Experimental carcinogenesis and cancer in the rodent gut.*, (ed. Appleton, D. A., Sunter, J.P. and A.J. Watson), pp. 255-277, London: Pitman Medical.
- Swidsinski, A., Ladhoff, A., Pernthaler, A., Swidsinski, S., Loening-Baucke, V., Ortner, M., Weber, J., Hoffmann, U., Schreiber, S., Dietel, M. und Lochs, H. (2002) Mucosal flora in inflammatory bowel disease. *Gastroenterology* **122**: 44-54.
- Tabuchi, Y., Arai, Y., Kondo, T., Takeguchi, N. und Asano, S. (2002) Identification of genes responsive to sodium butyrate in colonic epithelial cells. *Biochem Biophys Res Commun* **293**: 1287-94.
- Takahashi, K., Avissar, N., Whitin, J. und Cohen, H. (1987) Purification and characterization of human plasma glutathione peroxidase: a selenoglycoprotein distinct from the known cellular enzyme. *Arch Biochem Biophys* **256**: 677-86.
- Takahashi, M., Mutoh, M., Kawamori, T., Sugimura, T. und Wakabayashi, K. (2000) Altered expression of beta-catenin, inducible nitric oxide synthase and cyclooxygenase-2 in azoxymethane-induced rat colon carcinogenesis. *Carcinogenesis* **21**: 1319-27.
- Tamir, S. und Tannenbaum, S. R. (1996) The role of nitric oxide (NO.) in the carcinogenic process. *Biochim Biophys Acta* **1288**: F31-6.
- Tamura, G., Yin, J., Wang, S., Fleisher, A. S., Zou, T., Abraham, J. M., Kong, D., Smolinski, K. N., Wilson, K. T., James, S. P., Silverberg, S. G., Nishizuka, S., Terashima, M., Motoyama, T. und Meltzer, S. J. (2000) E-Cadherin gene promoter hypermethylation in primary human gastric carcinomas. *J Natl Cancer Inst* **92**: 569-73.
- Tan, M., Li, S., Swaroop, M., Guan, K., Oberley, L. W. und Sun, Y. (1999) Transcriptional activation of the human glutathione peroxidase promoter by p53. *J Biol Chem* **274**: 12061-6.
- Tang, S., Bhatia, B., Maldonado, C. J., Yang, P., Newman, R. A., Liu, J., Chandra, D., Traag, J., Klein, R. D., Fischer, S. M., Chopra, D., Shen, J., Zhau, H. E., Chung, L. W. und Tang, D. G. (2002) Evidence that arachidonate 15-lipoxygenase 2 is a negative cell cycle regulator in normal prostate epithelial cells. *J Biol Chem* **277**: 16189-201.
- Tarrerias, A. L., Millecamps, M., Alloui, A., Beaughard, C., Kemeny, J. L., Bourdu, S., Bommelaer, G., Eschalier, A., Dapoigny, M. und Ardid, D. (2002) Short-chain fatty acid enemas fail to decrease colonic hypersensitivity and inflammation in TNBS-induced colonic inflammation in rats. *Pain* **100**: 91-7.

- Taylor, B. A. und Dozois, R. R. (1987) The J ileal pouch-anal anastomosis. *World J Surg* **11**: 727-34.
- Tepperman, B. L., Brown, J. F. und Whittle, B. J. (1993) Nitric oxide synthase induction and intestinal epithelial cell viability in rats. *Am J Physiol* **265**: G214-8.
- Tham, D. M., Whitin, J. C., Kim, K. K., Zhu, S. X. und Cohen, H. J. (1998) Expression of extracellular glutathione peroxidase in human and mouse gastrointestinal tract. *Am J Physiol* **275**: G1463-71.
- Thun, M. J., Henley, S. J. und Patrono, C. (2002) Nonsteroidal anti-inflammatory drugs as anticancer agents: mechanistic, pharmacologic, and clinical issues. *J Natl Cancer Inst* **94**: 252-66.
- Topping, D. L., Fukushima, M. und Bird, A. R. (2003) Resistant starch as a prebiotic and symbiotic: state of the art. *Proc Nutr Soc* **62**: 171-6.
- Trayhurn, P. (2003) Nutritional genomics - "Nutrigenomics". *Br J Nutr* **89**: 1-2.
- Tsuji, S., Tsujii, M., Kawano, S. und Hori, M. (2001) Cyclooxygenase-2 upregulation as a perigenetic change in carcinogenesis. *J Exp Clin Cancer Res* **20**: 117-29.
- Tsuji, M. und DuBois, R. N. (1995) Alterations in cellular adhesion and apoptosis in epithelial cells overexpressing prostaglandin endoperoxide synthase 2. *Cell* **83**: 493-501.
- Tucker, E., Buda, A., Janghra, B., Cpoad, J., Moorghen, M., Havler, M., Dettmar, P. und Pignatelli, M. (2003) Abnormalities of the cadherin-catenin complex in chemically-induced colo-rectal carcinogenesis. *Proc Nutr Soc* **62**: 229-36.
- Turini, M. E. und DuBois, R. N. (2002) Cyclooxygenase-2: a therapeutic target. *Annu Rev Med* **53**: 35-57.
- Ursini, F., Maiorino, M. und Gregolin, C. (1985) The selenoenzyme phospholipid hydroperoxide glutathione peroxidase. *Biochim Biophys Acta* **839**: 62-70.
- van Es, J. H., Giles, R. H. und Clevers, H. C. (2001) The many faces of the tumor suppressor gene APC. *Exp Cell Res* **264**: 126-34.
- Vasen, H. F., Mecklin, J. P., Watson, P., Utsunomiya, J., Bertario, L., Lynch, P., Svendsen, L. B., Cristofaro, G., Muller, H., Khan, P. M. und et al. (1993) Surveillance in hereditary nonpolyposis colorectal cancer: an international cooperative study of 165 families. The International Collaborative Group on HNPCC. *Dis Colon Rectum* **36**: 1-4.
- Vasen, H. F., Watson, P., Mecklin, J. P. und Lynch, H. T. (1999) New clinical criteria for hereditary nonpolyposis colorectal cancer (HNPCC, Lynch syndrome) proposed by the International Collaborative group on HNPCC. *Gastroenterology* **116**: 1453-6.
- Velazquez, O. C., Lederer, H. M. und Rombeau, J. L. (1996) Butyrate and the colonocyte. Implications for neoplasia. *Dig Dis Sci* **41**: 727-39.
- Velazquez, O. C., Seto, R. W., Choi, J., Zhou, D., Breen, F., Fisher, J. D., Rombeau, J. L., Bain, A. M. und Fisher, J. (1997) Butyrate inhibits deoxycholate-induced increase in colonic mucosal DNA and protein synthesis in vivo. Deoxycholate inhibits in vivo butyrate-mediated BrDU labeling of the colonic crypt. *Dis Colon Rectum* **40**: 1368-75.
- Venkatraman, A., Ramakrishna, B. S. , Shaji, R.V., Nanda Kumar, N.S., Pulimood, A. und Patra, S. (2003) Amelioration of dextran sulfate colitis by butyrate: role of heat shock protein 70 and NF- $\kappa$ B. *Am J Physiol Gastrointest Liver Physiol* **285**: G177-G184.
- Vivona, A. A., Shpitz, B., Medline, A., Bruce, W. R., Hay, K., Ward, M. A., Stern, H. S. und Gallinger, S. (1993) K-ras mutations in aberrant crypt foci, adenomas and adenocarcinomas during azoxymethane-induced colon carcinogenesis. *Carcinogenesis* **14**: 1777-81.
- Vogelstein, B. und Kinzler, K. W. (1993) The multistep nature of cancer. *Trends Genet* **9**: 138-41.
- Vogiagis, D., Brown, W., Glare, E. M. und O'Brien, P. E. (2001) Rat colorectal tumours treated with a range of non-steroidal anti-inflammatory drugs show altered cyclooxygenase-2 and cyclooxygenase-1 splice variant mRNA expression levels. *Carcinogenesis* **22**: 869-74.
- Wakayama, T. und Iseki, S. (1998) Expression and cellular localization of the mRNA for the 25-kDa heat-shock protein in the mouse. *Cell Biol Int* **22**: 295-304.
- Warner, T. D. und Mitchell, J. A. (2002) Cyclooxygenase-3 (COX-3): filling in the gaps toward a COX continuum? *Proc Natl Acad Sci U S A* **99**: 13371-3.

- Warso, M. A. und Lands, W. E. (1983) Lipid peroxidation in relation to prostacyclin and thromboxane physiology and pathophysiology. *Br Med Bull* **39**: 277-80.
- Watson, P. und Lynch, H. T. (1993) Extracolonic cancer in hereditary nonpolyposis colorectal cancer. *Cancer* **71**: 677-85.
- Weber, J. L. und Wong, C. (1993) Mutation of human short tandem repeats. *Hum Mol Genet* **2**: 1123-8.
- Wilkinson, J. M. und Pollard, I. (1993) Immunohistochemical localisation of the 25 kDa heat shock protein in unstressed rats: possible functional implications. *Anat Rec* **237**: 453-7.
- Williams, E. A., Coxhead, J. M. und Mathers, J. C. (2003) Anti-cancer effects of butyrate: use of micro-array technology to investigate mechanisms. *Proc Nutr Soc* **62**: 107-15.
- Willoughby, D. A., Moore, A. R. und Colville-Nash, P. R. (2000) COX-1, COX-2, and COX-3 and the future treatment of chronic inflammatory disease. *Lancet* **355**: 646-8.
- Wingler, K., Muller, C., Schmehl, K., Florian, S. und Brigelius-Flohe, R. (2000) Gastrointestinal glutathione peroxidase prevents transport of lipid hydroperoxides in CaCo-2 cells. *Gastroenterology* **119**: 420-30.
- Wittekind, C., Tannapfel, A. (1995) Ätiologie, Pathogenese und Pathologie des Rektumkarzinoms. *Onkologe* **1**: 4-9.
- Witthöft, T., Eckmann, L., Kim, J. M. und Kagnoff, M. F. (1998) Enteroinvasive bacteria directly activate expression of iNOS and NO production in human colon epithelial cells. *Am J Physiol* **275**: G564-71.
- Wu, S., Lim, K. C., Huang, J., Saidi, R. F. und Sears, C. L. (1998) *Bacteroides fragilis* enterotoxin cleaves the zonula adherens protein, E-cadherin. *Proc Natl Acad Sci U S A* **95**: 14979-84.
- Young, J., Simms, L. A., Biden, K. G., Wynter, C., Whitehall, V., Karamatic, R., George, J., Goldblatt, J., Walpole, I., Robin, S. A., Borten, M. M., Stitz, R., Searle, J., McKeone, D., Fraser, L., Purdie, D. R., Podger, K., Price, R., Buttenshaw, R., Walsh, M. D., Barker, M., Leggett, B. A. und Jass, J. R. (2001) Features of colorectal cancers with high-level microsatellite instability occurring in familial and sporadic settings: parallel pathways of tumorigenesis. *Am J Pathol* **159**: 2107-16.
- Yue, G., Lai, P. S., Yin, K., Sun, F. F., Nagele, R. G., Liu, X., Linask, K. K., Wang, C., Lin, K. T. und Wong, P. Y. (2001) Colon epithelial cell death in 2,4,6-trinitrobenzenesulfonic acid-induced colitis is associated with increased inducible nitric-oxide synthase expression and peroxynitrite production. *J Pharmacol Exp Ther* **297**: 915-25.
- Zijlstra, F. J., van Dijk, A. P., Wilson, J. H., van Riemsdijk-Overbeeke, I. C., Vincent, J. E. und Ouwendijk, R. J. (1992) 15-HETE is the main eicosanoid formed by human colonic mucosa. *Agents Actions*: C53-9.
- Zouboulis, C. C., Kotter, I., Djawari, D., Kirch, W., Kohl, P.K., Ochsendorf, F.R., Keitel, W., Stadler, R., Wollina, U., Proksch, E., Sohnchen, R., Weber, H., Gollnick, H.P.M., Holzle, E., Fritz, K., Licht, T., Orfanos, C.E. (1997) Epidemiological Features of Adamantiades-Behcet's Disease in Germany and in Europe. *Yonsei Med J* **38**: 411-422.