

# Literaturverzeichnis

- Abe, F., Donelly, D. M. X., Moretti, C., Polonsky, J., 1985. Isoflavonoid constituents from *Dalbergia monetaria*. *Phytochemistry* 24, 1071–1076.
- Agrawal, P. K., Bansal, M. C., 1989. Studies in Organic Chemistry 39. Carbon-13 NMR of Flavonoids. Elsevier Science Publishers B.V., Ch. Isoflavonoids, 183–236.
- Akendengue, B., Ngou-Milama, E., Roblot, F., Laurens, A., Hocquemiller, R., Grellier, P., Frappier, F., 2002. Antiplasmodial activity of *Uvaria klaineana*. *Planta Medica* 68, 167–169.
- Akunyili, D. N., Houghton, P. J., 1993. Meroterpenoids and naphthoquinones from *Kigelia pinnata*. *Phytochemistry* 32, 1015–1018.
- Al Magboul, A. Z. I., Bashir, A. K., Khalid, S. A., Farouk, A., 1997. Antimicrobial activity of vernolepin and vernodalin. *Fitoterapia* 68, 83–84.
- Alemayehu, G., Hailu, A., Abegaz, B. M., 1996. Bianthraquinones from *Senna didymobotrya*. *Phytochemistry* 42, 1423–1425.
- Asaka, Y., Kubota, T., Kulkarni, A. B., 1977. Studies on a bitter principle from *Vernonia anthelmintica*. *Phytochemistry* 16, 1838–1839.
- Asres, K., Mascagni, P., O'Neill, M. J., Phillipson, J. D., 1985. Isoflavonoids from *Bolu-santhus speciosus* (BOLUS) HARMS, Leguminosae. *Zeitschrift für Naturforschung* 40c, 617–620.
- Balick, M. J., Cox, P. A., 1996. Drogen, Kräuter und Kulturen – Pflanzen und die Geschichte des Menschen. Wissenschaftliche Buchgesellschaft Darmstadt.
- Basilico, N., Pagani, E., Monti, D., Olliari, P., Taramelli, D., 1998. A microtitre-based method for measuring the haem polymerisation inhibitory activity (HPIA) of antimalarial drugs. *Journal of Antimicrobial Chemotherapy* 42, 55–60.
- Bastos, J. K., Albuquerque, S., Silva, M. L. A., 1999. Evaluation of the trypanocidal activity of lignans isolated from leaves of *Zanthoxylum naranjillo*. *Planta Medica* 65, 541–544.
- Bastos, J. K., Gottlieb, O. R., Sarti, S. J., Filho, D. S., 1996. Isolation of lignans and sesquiterpenoids from leaves of *Zanthoxylum naranjillo*. *Natural Product Letters* 9, 65–70.

- Benoit, F., Valentin, A., Pelissier, Y., Diafouka, F., Marion, C., Kone-Bamba, D., Kone, M., Mallie, M., Yapo, A., Bastide, J. M., 1996. *In vitro* antimalarial activity of vegetal extracts used in West African traditional medicine. American Journal of Tropical Medicine and Hygiene 54, 67–71.
- Berhanu, E., Fetene, M., Dagne, E., 1986. Anthraquinones as taxonomic markers in Ethiopian *Kniphofia* species. Phytochemistry 25, 847–850.
- Berliner Zeitung, 1998. Treibjagd auf einen cleveren Parasiten. Wissenschaftsrubrik der Berliner Zeitung 269, III.
- Binutu, O. A., Adesogan, K. E., Okogun, J. I., 1996. Antibacterial and antifungal compounds from *Kigelia pinnata*. Planta Medica 62, 352–353.
- Böhlke, M., Guinaudeau, H., Angerhofer, C. K., Wongpanchi, V., Soejarto, D. D., Farnsworth, N. R., Mora, G. A., Poveda, L. J., 1996. Costaricine, a new antiplasmodial bis-benzylisoquinoline alkaloid from *Nectandra salicifolia* trunk bark. Journal of Natural Products 59, 576–580.
- Bohlmann, F., Knoll, K. H., Robinson, H., King, R. M., 1980. Neue Guaianolide aus *Cacosmia rugosa*. Phytochemistry 19, 599–602.
- Bohlmann, F., Zdero, C., 1972. Constituents of *Artemisia afra*. Phytochemistry 11, 2329–2330.
- Bohlmann, F., Zdero, C., 1978. New sesquiterpenes and acetylenes from *Athansia* and *Pentzia* species. Phytochemistry 17, 1595–1599.
- Bohlmann, F., Zdero, C., King, R. M., Robinson, H., 1984. Pseudoguaianolides and other sesquiterpene lactones from *Gaillardia* species. Phytochemistry 23, 1979–1988.
- Bravo, J. A., Sauvain, M., Gimenez, A. T., Muñoz, V. O., Callapa, J., Le Men-Olivier, L., Massiot, G., Lavaud, C., 1999. Bioactive phenolic glycosides from *Amburana cearensis*. Phytochemistry 50, 71–74.
- Brieskorn, C. H., Michel, H., 1968. Flavone aus dem Blatt von *Rosmarinus officinalis* L. Tetrahedron Letters 30, 3447–3448.
- Bringmann, G., Menche, D., Bezabih, M., Abegaz, B. M., Kaminsky, R., 1999. Antiplasmodial activity of knipholone and related phenylanthraquinones. Planta Medica 65, 757–758.
- Bringmann, G., Saeb, W., Aké Assi, L., François, G., Narayanan, A. S. S., Peters, K., Peters, E.-M., 1997. Betulinic acid: isolation from *Triphyophyllum peltatum* and *Anistrocladus heyneanus*, antimalarial activity, and crystal structure of the benzyl ester. Planta Medica 63, 255–257.
- Bringmann, G., Saeb, W., God, R., Schäffer, M., François, G., Peters, K., Peters, E.-V., Proksch, P., Hostettmann, K., Aké Assi, L., 1998. 5'-O-Demethyldioncophylline A, a new antimalarial alkaloid from *Triphyophyllum peltatum*. Phytochemistry 49, 1667–1673.

- Buchbauer, G., Silbernagel, E., 1989. Der südafrikanische Wermut. Deutsche Apotheker Zeitung 129, 2173–2177.
- Caballero, P., Smith, C. M., 1986. Isoflavones from an insect-resistant variety of soybean and the molecular structure of afrormosin. Journal of Natural Products 49, 1126–1129.
- Carvalho, L. H., Rocha, E. M. M., Raslan, D. S., Oliveira, A. B., Krettli, A. U., 1988. *In vitro* activity of natural and synthetic naphthoquinones against erythrocytic stages of *P.falciparum*. Brazil Journal of Medicine and Biological Research 21, 485–487.
- Chan, K. L., O'Neill, M. J., Phillipson, J. D., Warhurst, D. C., 1986. Plants as sources of antimalarial drugs. Part 3. *Eurycoma longifolia*. Planta Medica 52, 105–107.
- Chanphen, R., Thebtaranonth, Y., Wanauppathamkul, S., Yuthavong, Y., 1998. Antimalarial principles from *Artemisia indica*. Journal of Natural Products 61, 1146–1147.
- Chapman & Hall, 1994. Phytochemical Dictionary of the Leguminosae, Volume 1, Plants and their constituents, 1. Auflage. Chapman & Hall, University press, Cambridge, Great Britain, 72.
- Chen, M., Christensen, S. B., Zhai, L., Rasmussen, M. H., Theander, T. G., Frøkjaer, S., Steffansen, B., Davidson, J., Kharazmi, A., 1997. The novel oxygenated chalcone, 2,4-dimethoxy-4'-butoxychalcone, exhibits potent activity against human malaria parasite *Plasmodium falciparum* *in vitro* and rodent parasites *Plasmodium berghei* and *Plasmodium yoelii* *in vivo*. The Journal of Infectious Diseases 176, 1327–1333.
- Chen, M., Theander, T. G., Christensen, S. B., Hviid, L., Zhai, L., Kharazmi, A., 1994. Licochalcone A, a new antimalarial agent, inhibits *in vitro* growth of the human malaria parasite *Plasmodium falciparum* and protects mice from *P. yoelii* infection. Antimicrobial Agents and Chemotherapy 38, 1470–1475.
- Cocker, W., Dahl, T., Dempsey, C., McMurry, T. B. H., 1962. Extractives from woods. Part I. Extractives from *Andira inermis* (WRIGHT) H.B.K. Journal of the Chemical Society Perkin Transactions 1, 4906–4909.
- Cubukcu, B., Bray, D. H., Warhurst, D. C., Mericli, A. H., Ozhatay, N., Sariyar, G., 1990. *In vitro* antimalarial activity of crude extracts and compounds from *Artemisia abrotanum* L. Phytotherapy Research 4, 203–204.
- Da Silva, B. P., Velozo, L. S., Parente, J. P., 2000. Biochanin A trglycoside from *Andira inermis*. Fitoterapia 71, 663–667.
- Dagne, E., Steglich, W., 1984. Knipholone: a unique anthraquinone derivative from *Kniphofia foliosa*. Phytochemistry 23, 1729–1731.
- DAZ-Beilage, 2001. Malariamittel: Artemether/Lumefantrin. Beilage: Neue Arzneimittel. Deutsche Apotheker Zeitung 48, 94–97.
- De Almeida Alves, T. M., Nagem, T. J., de Carvalho, L. H., Krettli, A. U., Zani, C. L., 1997. Antiplasmodial triterpene from *Vernonia brasiliiana*. Planta Medica 63, 554–555.

- Desjardins, R. E., Canfield, C. J., Haynes, J. D., Chulay, J. D., 1979. Quantitative assessment of antimalarial activity *in vitro* by a semiautomated microdilution technique. *Antimicrobial Agents and Chemotherapy* 16, 710–718.
- Dhingra, V., Rao, K. V., Narasu, M. L., 2000. Minireview: Current status of artemisinin and its derivatives as antimalarial drugs. *Life Sciences* 66, 279–300.
- Dluzewski, A. R., Garcia, C. R., 1996. Inhibition of invasion and intraerythrocytic development of *Plasmodium falciparum* by kinase inhibitors. *Experientia* 52, 621–623.
- Duke, J. A., 1975. Ethnobotanical observations on the Cuna Indians. *Economic Botany* 29, 278–293.
- Eggeling, W. J., 1951. The indigenous trees of the Uganda Protectorate, 2. Auflage. Robert MacLehose & Company Limited, The University Press, Glasgow, 42–44.
- Eich, E., Pertz, H., Kaloga, M., Schulz, J., Fesen, M. R., Mazumder, A., Pommier, Y., 1996. (–)-Arctigenin as a lead structure for inhibitors of human immunodeficiency virus type-1 integrase. *Journal of Medicinal Chemistry* 39, 86–95.
- El Sohly, H. N., Joshi, A. S., Nimrod, A. C., 1999. Antigiardial isoflavones from *Machaerium arstulatum*. *Planta Medica* 65, 490.
- Elford, B. C., 1986. L-Glutamin influx in malaria-infected erythrocytes: a target for antimalarials? *Parasitology Today* 2, 309–312.
- Elford, B. C., Roberts, M. F., Phillipson, J. D., Wilson, R. J. M., 1987. Potentiation of the antimalarial activity of qinghaosu by methoxylated flavones. *Transactions of the Royal Society for Tropical Medicine and Hygiene* 81, 434–436.
- Ferreira, M. A., Moir, M., Thomson, R. H., 1975. Naturally occurring quinones. Part XXV. Pterocarpenequinones (6H-Benzofuro [3,2-c] [1] benzopyranquinones) from *Brya ebenus*. *Journal of the Chemical Society Perkin Transactions* 1, 1113–1115.
- Figueiredo, J. N., Räz, B., Séquin, U., 1998. Novel quinone methides from *Salacia kraussii* with *in vitro* antimalarial activity. *Journal of Natural Products* 61, 718–723.
- Fonseca, S. F., Barata, L. E. S., Ruveda, E. A., 1979.  $^{13}\text{C}$  nuclear magnetic resonance spectral and conformational analysis of naturally occurring tetrahydrofuran lignans. *Canadian Journal of Chemistry* 57, 441–443.
- François, G., Bringmann, G., D., P. J., Aké Assi, L., Dochez, C., Rübenacker, M., Schneider, C., Wéry, M., Warhurst, D. C., Kirby, G. C., 1994. Activity of extracts and naphthylisoquinoline alkaloids from *Triphyophyllum peltatum*, *Ancistrocladus abbreviatus* and *A. barteri* against *Plasmodium falciparum* *in vitro*. *Phytochemistry* 35, 1461–1464.
- Frédéric, M., De Pauw, M.-C., Llabrès, G., Tits, M., Hayette, M.-P., Brandt, V., Penelle, J., DeMol, P., Angenot, L., 2000. New antimalarial and cytotoxic sungsucine derivatives from *Strychnos icaja* roots. *Planta Medica* 66, 262–269.
- Frédéric, M., Tits, M., Hayette, M.-P., Brandt, V., Penelle, J., DeMol, P., Llabrès, G., Angenot, L., 1999. 10'-Hydroxyusambarensine, a new antimalarial bisindole alkaloid from the roots of *Strychnos usambarensis*. *Journal of Natural Products* 62, 619–621.

- Fujioka, H., Nishiyama, Y., Furukawa, H., Kumada, N., 1989. *In vitro* and *in vivo* activities of atalaphillinine and related acridone alkaloids against rodent malaria. *Antimicrobial Agents and Chemotherapy* 33, 6–9.
- Fukui, K., Nakayama, M., Tsuzuki, K., 1969. Synthesis of pterocarpans. IV. a new synthesis of 3,8,9-trimethoxypterocarpan. *Bulletin of the Chemical Society of Japan* 42, 2395–2396.
- Ganjian, I., Kubo, I., Fludzinski, P., 1983. Insect antifeedant elemanolide lactones from *Vernonia amygdalina*. *Phytochemistry* 22, 2525–2526.
- Gasquet, M., Bamba, D., Babadjamian, A., Balansard, G., Timon-David, P., Metzger, J., 1985. Action amoebicide et anthelminthique du vernolide et de l'hydroxyvernolide isolés des feuilles de *Vernonia colorata* (WILLD.) DRAKE. *European Journal of Medicinal Chemistry* 20, 111–115.
- Govindachari, T. R., Premila, M. S., 1985. The bezofuran norwedelic acid from *Wedelia calendulacea*. *Phytochemistry* 24, 3068–3069.
- Greger, H., Pacher, T., Vajrodaya, S., Bacher, M., Hofer, O., 2000. Infraspecific variation of sulfur-containing bisamides from *Aglaiia leptantha*. *Journal of Natural Products* 63, 616–620.
- Greger, H., Zdero, C., Bohlmann, F., 1986. Eudesman-12,8 $\beta$ -olides and other terpenes from *Artemisia* species. *Phytochemistry* 25, 891–897.
- Gupta, M., Correa, M. D., Solis, P. N., Jones, A., Galdames, C., Guionneau-Sinclair, F., 1993. Medicinal plant inventory of Kuna Indians: Part 1. *Journal of Ethnopharmacology* 40, 77–109.
- Halabalaki, M., Aligiannis, N., Papoutsi, Z., Mitakou, S., Moutsatsou, P., Sekeris, C., Skaltsounis, A. L., 2000. Three new arylbenzofurans from *Onobrychis ebenoides* and evaluation of their binding affinity for the estrogen receptor. *Journal of Natural Products* 63, 1672–1674.
- Harborne, J. B., 1988. The Flavonoids: Advances in Research Since 1980. Chapman & Hall, London.
- Harborne, J. B., 1994. The Flavonoids: Advances in Research Since 1986. Chapman & Hall, London.
- Harborne, J. B., Williams, C. A., 2000. Review: Advances in flavonoid research since 1992. *Phytochemistry* 55, 481–504.
- Hirschhorn, H. H., 1981. Botanical remedies of South and Central America, and the Caribbean: an archival analysis. Part I. *Journal of Ethnopharmacology* 4, 129–158.
- Hooper, M., Kirby, G. C., Kulkarni, M. M., Kulkarni, S. N., Nagasampagi, B. A., O'Neill, M. J., Phillipson, J. D., Rojatkar, S. R., Warhurst, D. C., 1990. Antimalarial activity of parthenin and its derivatives. *European Journal of Medicinal Chemistry* 25, 717–723.

- Hudson, A. T., Randall, A. W., Fry, M., Ginger, C. D., Hill, B., Latter, V. S., Mc Hardy, N., Williams, R. B., 1985. Novel anti-malarial hydroxynaphthoquinones with potent broad spectrum anti-protozoal activity. *Parasitology* 90, 45–55.
- Hurabielle, M., Eberle, J., Paris, M., 1982. Etude des flavonoïdes d'*Artemisia campestris* sous-espèce *glutinosa*. *Journal of Medicinal Plant Research* 46, 124–128.
- Ingham, J. L., Dewick, P. M., 1984. The structure of desmocarpin, a pterocarpan phytoalexin from *Desmodium gangeticum*. *Zeitschrift für Naturforschung* 39c, 531–534.
- Isaka, M., Punya, J., Lertwerawat, Y., Tanticharoen, M., Thebtaranonth, Y., 1999. Antimalarial activity of macrocyclic trichothecens isolated from the fungus *Myrothecium verrucaria*. *Journal of Natural Products* 62, 329–331.
- Jakupovic, J., Baruah, R. N., Chau Thi, T. V., Bohlmann, F., Msonthi, J. D., Schmedemann, G., 1985. New vernolepin derivatives from *Vernonia glabra* and glaucolides from *Vernonia scorpioides*. *Planta Medica* 5, 378–380.
- Jakupovic, J., Klemeyer, H., Bohlmann, F., Graven, E. H., 1988. Glaucolides and guaianolides from *Artemisia afra*. *Phytochemistry* 27, 1129–1133.
- Jakupovic, J., Tan, R. X., Bohlmann, F., Boldt, P. E., Jia, Z. J., 1991. Sesquiterpene lactones from *Artemisia ludoviciana*. *Phytochemistry* 30, 1573–1577.
- Jakupovic, J., Zdero, C., Boeker, R., Warning, U., Bohlmann, F., Jones, S. B., 1987. Vernocistifolide und andere Sesquiterpenlactone aus *Vernonia* und verwandten Arten. *Liebigs Annalen der Chemie*, 111–123.
- Jenett-Siems, K., 2002. Habilitationsschrift, Fachbereich Biologie, Chemie, Pharmazie, Freie Universität Berlin.
- Jenett-Siems, K., Mockenhaupt, F. P., Bienzle, U., Gupta, M. P., Eich, E., 1999. *In vitro* antiplasmoidal activity of Central American medicinal plants. *Tropical Medicine and International Health* 4, 611–615.
- Jenett-Siems, K., Siems, K., Solis, P. N., Gupta, M. P., Mockenhaupt, F. P., Bienzle, U., Eich, E., 2000. Sipandinolide: a butenolide including a novel type of carbon skeleton from *Siparuna andina*. *Planta Medica* 66, 384–385.
- Johri, R. K., Singh, C., 1997. Medicinal uses of *Vernonia* species. *Journal of Medicinal and Aromatic Plant Sciences* 19, 744–752.
- Joli, L. G., Guerra, S., Septimo, R., Solis, P. N., Correa, M., Gupta, M. P., Levy, S., Sandberg, F., 1987. Ethnobotanical inventory of medicinal plants used by the Guaymi Indians in Western Panama. *Journal of Ethnopharmacology* 20, 145–171.
- Jomaa, H., Wiesner, J., Sanderbrand, S., Altincinek, B., Weidemeyer, C., Hintz, M., Türbachova, I., Eberl, M., Zeidler, J., Lichtenthaler, H. K., Soldati, D., Beck, E., 1999. Inhibitors of the nonmevalonate pathway of isoprenoid biosynthesis as antimalarial drugs. *Science* 285, 1573–1575.

- Joshi, K. C., Singh, P., Taneja, S., Cox, P. J., Howie, R. A., Thomson, R. H., 1982. New terpenoid aldehydes from *Kigelia pinnata*: Crystal structure of pinnatal. *Tetrahedron* 38, 2703–2708.
- Kamchonwongpaisan, S., Nilanonta, C., Tarnchompoo, B., Thebtaranonth, C., Thebtaranonth, Y., Yuthavong, Y., Kongsaeree, P., Clardy, J., 1995. An antimalarial peroxide from *Amomum krexvanh* PIERRE. *Tetrahedron Letters* 36, 1821–1824.
- Kanyinda, B., Diallo, B., Vanhaelen-Fastré, R., Vanhaelen, M., 1989. Benzylisoquinoline alkaloids from *Anisocycla cymosa* roots. *Planta Medica* 55, 394.
- Keawpradub, N., Kirby, G. C., Steele, J. C. P., Houghton, P. J., 1999. Antiplasmodial activity of extracts and alkaloids of three *Alstonia* species from Thailand. *Planta Medica* 65, 690–694.
- Keriko, J. M., Nakajima, S., Baba, N., Isozaki, Y., Ikeda, K., Iwasa, J., Karanjy, P. N., 1995. A plant growth regulator from *Vernonia auriculifera* (Asteraceae). *Zeitschrift für Naturforschung* 50c, 455–458.
- Khan, I. A., Avery, M. A., Burandt, C. L., Goins, D. K., Mikell, J. R., Nash, T. E., Azadegan, A., Walker, L. A., 2000. Antigiardial activity of isoflavones from *Dalbergia frutescens* bark. *Journal of Natural Products* 63, 1414–1416.
- Kirby, G. C., O'Neill, M. J., Phillipson, J. D., Warhurst, D. C., 1989. *In vitro* studies on the mode of action of quassinoids with activity against chloroquine-resistant *Plasmodium falciparum*. *Biochemical Pharmacology* 38, 4367–4374.
- Klayman, D. L., 1985. Quinghaosu (Artemisinin): An antimalarial drug from China. *Science* 228, 1049–1055.
- Kobayashi, A., Yata, S., Kawazu, K., 1988. A  $\beta$ -hydroxychalcone and flavonoids from *Alfalfa* callus stimulated by a fungal naphthoquinone, PO-1. *Agricultural and Biological Chemistry* 52, 3223–3227.
- Kobayashi, M., Noguchi, H., Sankawa, U., 1985. Formation of isoflavones by callus culture of *Glycyrrhiza uralensis* with different production pattern. *Chemical and Pharmaceutical Bulletin* 33, 3811–3816.
- Köhler, I., Jenett-Siems, K., Kraft, C., Siems, K., Abbiw, D., Bienzle, U., Eich, E., 2002. Herbal remedies traditionally used against malaria in ghana: bioassay-guided fractionation of *Microglossa pyrifolia* (Asteraceae). *Zeitschrift für Naturforschung* 57c, 1022–1027.
- Köhler, I., Jenett-Siems, K., Mockenhaupt, F. P., González, J. C., Hernandez, M. A., Ibarra, R. A., Berendsohn, W. G., Bienzle, U., Eich, E., 2001. *In vitro* antiplasmodial activity of 4-phenylcoumarines from *Exostema mexicanum*. *Planta Medica* 67, 89–91.
- König, G. M., Wright, A. D., Sticher, O., Angerhofer, C. K., Pezzuto, J. M., 1994. Biological activities of selected marine natural products. *Planta Medica* 60, 532–537.

- Koumaglo, K., Gbeassor, M., Nikabu, O., de Souza, C., Werner, W., 1992. Effects of three compounds extracted from *Morinda lucida* on *Plasmodium falciparum*. *Planta Medica* 58, 533–534.
- Krungkrai, S. R., Yuthavong, Y., 1987. The antimalarial action of qinghaosu and arte-suante in combination with agents that modulate oxidant stress. *Transactions of the Royal Society for Tropical Medicine and Hygiene* 81, 710–714.
- Kupchan, S. M., Hemingway, R. J., Karim, A., Werner, D., 1969. Tumor inhibitors. XLVII. Vernodalin and vernomygdin, two new cytotoxic sesquiterpene lactones from *Vernonia amygdalina*. *Journal of Organic Chemistry* 34, 3908–3911.
- Lambros, C., Vanderberg, J. P., 1979. Synchronization of *Plasmodium falciparum* erythrocytic stages in culture. *Journal of Parasitology* 65, 418–420.
- Lang, G., Passreiter, C. M., Wright, C. W., Filipowicz, N. H., Addae-Kyereme, J., Medina, B. E., Castillo, J.-J., 2002. Antiplasmoidal activities of sesquiterpene lactones from *Eupatorium semialatum*. *Zeitschrift für Naturforschung* 57c, 282–286.
- Li, R., Kenyon, G. L., Cohen, F. E., Chen, X., Gong, B., Dominguez, J. N., Davidson, E., Kurzban, G., Miller, R. E., Nuzum, E. O., Rosenthal, P. J., Mc Karrow, J. H., 1995. *In vitro* antimalarial activity of chalkones and their derivatives. *Journal of Medicinal Chemistry* 38, 5031–5037.
- Likhithwitayawuid, K., Angerhofer, C. K., Chai, H., Pezzuto, J. M., Cordell, G. A., 1993a. Cytotoxic and antimalarial alkaloids from the tubers of *Stephania pierrei*. *Journal of Natural Products* 56, 1468–1478.
- Likhithwitayawuid, K., Angerhofer, C. K., Cordell, G. A., Pezzuto, J. M., 1993b. Cytotoxic and antimalarial bisbenzylisoquinoline alkaloids from *Stephania erecta*. *Journal of Natural Products* 56, 30–38.
- Likhithwitayawuid, K., Kaewamatawong, R., Ruangrungsi, N., Krungkrai, J., 1998. Antimalarial naphthoquinones from *Nepenthes thorelli*. *Planta Medica* 64, 237–341.
- Lin, L.-Z., Shieh, H.-L., Angerhofer, C. K., Pezzuto, J. M., Cordell, G. A., 1993. Cytotoxic and antimalarial bisbenzylisoquinoline alkaloids from *Cyclea barbata*. *Journal of Natural Products* 56, 22–29.
- Liu, K. C. S., Yang, S. L., Roberts, M. F., Elford, B. C., Phillipson, J. D., 1989. The contribution of flavonoids to the antimalarial activity of *Artemisia annua*. *Planta Medica* 55, 654–655.
- Liu, M., Wilairat, P., Go, M. L., 2001. Antimalarial alkoxylated and hydroxylated chalcones: structure-activity relationship analysis. *Journal of Medicinal Chemistry* 44, 4443–4452.
- Liu, Y., Mabry, T. J., 1981. Flavonoids from *Artemisia frigida*. *Phytochemistry* 20, 1389–1395.
- Lock de Ugaz, O., Costa, J., Sanchez, L., Ubillas Sanchez, R. P., Tempesta, M. S., 1991. Flavonoids from *Andira inermis*. *Fitoterapia* 62, 89–90.

- Loewe, W., Gust, R., Witzel, S., Dietrich, C., 2001. Preparation of 3-hydroxybenzoylindoles for treatment of diseases associated with protein kinase and estrogen activity. Patent DE 19947863.
- MacRae, W. D., Towers, G. H. N., 1985. Non-alkaloidal constituents of *Virola elongata* bark. *Phytochemistry* 24, 561–566.
- Markham, K. R., Mabry, T. J., 1975. The Flavonoids. Chapman & Hall, London, Ch. Mass spectrometry of flavonoids, 78–126.
- Matissek, R., Schnepel, F.-M., Steiner, G., 1989. Matissek, Reinhard: Lebensmittelanalytik: Grundzüge, Methoden, Anwendungen. Springer-Verlag, Berlin/Heidelberg, Ch. 2.2.1 Extraktion nach Ammoniakaufschluß - Methode nach Röse-Gottlieb, 38–40.
- McMurry, T. B. H., Martin, E., 1972. 3-Hydroxy-9-methoxy and 3-methoxy-9-hydroxypterocarpans. *Phytochemistry* 11, 3283–3286.
- Meyer, C. G., 2000. Tropenmedizin – Infektionskrankheiten. ecomed verlagsgesellschaft AG & Co. KG, Ch. Reiseimpfungen und Malaria prophylaxe und Protozoeninfektionen, 25–29 und 183–198.
- Mitscher, L., Park, Y. H., Clark, D., 1980. Antimicrobial agents from higher plants. Antimicrobial isoflavonoids and related substances from *Glycyrrhiza glabra* L. var. *typica*. *Journal of Natural Products* 43, 259–269.
- Mockenhaupt, F. P., 1995. Mefloquine resistance in *Plasmodium falciparum*. *Parasitology Today* 11, 248.
- Mockenhaupt, F. P., Thies, F. L., Meyer, C. G., 1998/99. Malaria im Kindesalter. *Pädiatrische Praxis* 55, 305–318.
- Moideen, S. V. K., Houghton, P. J., Rock, P., Croft, S. L., Aboagye-Nyame, F., 1999. Activity of extracts and naphthoquinones from *Kigelia pinnata* against *Trypanosoma brucei brucei* and *Trypanosoma brucei rhodesiense*. *Planta Medica* 65, 536–540.
- Moretti, C., Sauvain, M., Lavaud, C., Massiot, G., Bravo, J.-A., Muñoz, M., 1998. A novel antiprotozoal aminosteroid from *Saracha punctata*. *Journal of Natural Products* 61, 1390–1393.
- Morton, J. F., 1981. Atlas of Medicinal Plants of Middle America. Charles C. Thomas, Springfield, IL, 275–276.
- Mosmann, T., 1983. Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *Journal of Immunological Methods* 65, 55–63.
- Muñoz, V., Sauvain, M., Mollinedo, P., Callapa, J., Rojas, I., Gimanez, A., Valentin, A., Mallié, M., 1999. Antimalarial activity and cytotoxicity of (-)-roemrefidine isolated from the stem bark of *Sparattanthelium amazonum*. *Planta Medica* 65, 448–449.
- Muangnoicharoen, N., Frahm, A. W., 1981. Arylbenzofurans from *Dalbergia parviflora*. *Phytochemistry* 20, 291–293.

- Müller, M., Heide, L., 2001. Malaria in Afrika: Wie hilfreich sind Arzneipflanzen? Zeitschrift für Phytotherapie 22, 77–81.
- Müller, M., Karhagomba, I. B., Hirt, H. M., Wemakor, E., 2000. The potential of *Artemisia annua* L. as a locally produced remedy for malaria in the tropics: agricultural, chemical and clinical aspects. Journal of Ethnopharmacology 73, 487–493.
- Mutschler, E., Schäfer-Korting, M., 1996. Arzneimittelwirkungen - Lehrbuch der Pharmakologie und Toxikologie. Wissenschaftliche Verlagsgesellschaft mbH Stuttgart, Ch. Prophylaxe und Therapie von Infektionskrankheiten, 717–722.
- Navarrete, N. E., 1996. Fact Sheet: a quick guide to multipurpose trees from around the world. *Andira inermis*: more than a beautiful ornamental tree. <http://www.winrock.org/forestry/FACTPUB/factsh/ainermis.htm>, Zugriff am 27.12.2001.
- Neuwinger, H. D., 2000. African traditional Medicine; A dictionary of plant use and application. Medpharm GmbH Scientific Publisher: Stuttgart.
- Nkunya, M. H. H., Weenen, H., Kinabo, L. S., 1992. Constituents of *Artemisia afra*. Fitoterapia 63, 279–280.
- Nógrádi, M., Kajtár-Peredy, M., 1988. Synthesis of some 3-substituted 2-arylbenzofurans. Acta Chimica Hungaria 125, 497–501.
- Ohigashi, H., Huffman, M. A., Izutsu, D., Koshimizu, K., Kawanaka, M., Sugiyama, H., Kirby, G. C., Warhurst, D. C., Allen, D., Wright, C. W., Phillipson, J. D., Timon-David, P., Delmas, F., Elias, R., Balansard, G., 1994. Toward the chemical ecology of medicinal plant use in chimpanzees: the case of *Vernonia amygdalina*, a plant used by wild chimpanzees possibly for parasite-related diseases. Journal of Chemical Ecology 20, 541–553.
- Oketch-Rabah, A., Christensen, B., Frydenvang, K., Dossaji, S. F., Theander, T. G., Cornett, C., Watkins, W. M., Kharazmi, A., Lemmich, E., 1998. Antiprotozoal properties of 16,17-dihydrobrachycalyxolide from *Vernonia brachycalyx*. Planta Medica 64, 559–562.
- Oketch-Rabah, H. A., Lemmich, E., Dossaji, S. F., Theander, T. G., Olsen, C. E., Cornett, C., Kharazmi, A., Christensen, S. B., 1997. Two new antiprotozoal 5-methylcoumarins from *Vernonia brachycalyx*. Journal of Natural Products 60, 458–461.
- Onogi, B., Kraft, C., Köhler, I., Freund, M., Jenett-Siems, K., Siems, K., Beyer, G., Melzig, M. F., Bienzle, U., Eich, E., 2002. Antiplasmodial activity of naphthoquinones and one anthraquinone from *Stereospermum kunthianum*. Phytochemistry 60, 39–44.
- O'Neill, M. J., Bray, D. H., Boardman, P., Chan, K. L., Phillipson, J. D., 1987. Plants as sources of antimarial drugs. Part 4. Activity of *Brucea javanica* fruits against chloroquine-resistant *Plasmodium falciparum* *in vitro* and against *Plasmodium berghei* *in vivo*. Journal of Natural Products 50, 41–48.
- O'Neill, M. J., Bray, D. H., Boardmann, P., Phillipson, J. D., Warhurst, D. C., 1985. Plants as sources of antimarial drugs. Part 1. *In vitro* test method for the evaluation of crude extracts from plants. Planta Medica 51, 394–398.

- Partridge, S. J., Russell, P. F., Anderson, M. M., Wright, C. W., Phillipson, J. D., Kirby, G. C., Warhurst, D. C., 1990. *In vitro* cytotoxic, antimalarial and antimicrobial activities of protoberberine alkaloids. *Journal of Pharmacy and Pharmacology* 42 (Suppl.), 97P.
- Paulo, A., Gomes, E. T., Steele, J., Warhurst, D. C., Houghton, P. J., 2000. Antiplasmodial activity of *Cryptolepis sanguinolenta* alkaloids from leaves and roots. *Planta Medica* 66, 30–34.
- Petit, S., 2002. Species description: *Stereospermum kunthianum*. <http://www.fao.org/ag/AGP/AGPC/doc/Gbase/DATA/Pf000534.HTM>, Zugriff am 21.08.2002.
- Phillipson, J. D., Wright, C. W., 1991. Can ethnopharmacology contribute to the development of antimalarial agents? *Journal of Ethnopharmacology* 32, 155–165.
- Phillipson, J. D., Wright, C. W., Kirby, G. C., Warhurst, D. C., 1995. Phytochemistry of some plants used in traditional medicine for the treatment of protozoal diseases. In: Hostettmann, K., Marston, A., Maillard, M., Hamburger, M. (Eds.), *Phytochemistry of Plants used in Traditional Medicine*. Clarendorn Press, Oxford, 95–135.
- Pschyrembel, W., 1975. *Klinisches Wörterbuch*. Walter de Gruyter Verlag, Berlin, New York.
- Rabe, T., Mullholland, D., van Staden, J., 2002. Isolation and identification of antibacterial compounds from *Vernonia colorata* leaves. *Journal of Ethnopharmacology* 80, 91–94.
- Rücker, G., Walter, R. D., Manns, D., Mayer, R., 1991. Antimalarial activity of some natural peroxides. *Planta Medica* 57, 295–296.
- Reisch, J., Hussain, R. A., Mester, I., 1984. Flavonoids from *Flindersia australis*. *Phytochemistry* 23, 2114–2115.
- Ridley, R. G., 1999. Planting the seeds of new antimalarial drugs. *Science* 285, 1502–1503.
- Rimpler, H., 1990. Biogene Arzneistoffe / Horst Rimpler. Unter Mitarb. von Eich, E., Förstermann, U., Chlodwig, F., Glasl, H., Illes, P., Wichtl, M. Georg Thieme Verlag, Stuttgart New York.
- Rodriguez, E., Towers, G. H. N., Mitchell, J. C., 1976. Review: Biological activities of sesquiterpene lactones. *Phytochemistry* 15, 1573–1580.
- Rosenthal, P. J., 1998. Proteases of malaria parasites: new targets for chemotherapy. *Emerging Infectious Diseases* 4, 49–56.
- Rücker, D., 1999. Plasmodien-Untermieter ist die Schwachstelle. *Pharmazeutische Zeitung* 36, 2845–2847.
- Rücker, G., 1995. Malariaactive Verbindungen aus Pflanzen, insbesondere Peroxide [1]. *Pharmazie in unserer Zeit* 24, 189–195.
- Sariyar, G., Sari, A., Mat, A., 1994. Quaternary alkaloids of *Papaver spicatum*. *Planta Medica* 60, 293.

- Sarkanen, K. V., Wallis, A. F. A., 1973. PMR analysis and conformation of 2,5-bis-(3,4,5-trimethoxyphenyl)-3,4-dimethyltetrahydrofuran isomers. *Journal of Heterocyclic Chemistry* 10, 1025–1027.
- Sauvain, M., Moretti, C., Bravo, J.-A., Callapa, J., Muñoz, V., Ruiz, E., Richard, B., Le Men-Olivier, L., 1996. Antimalarial activity of alkaloids from *Pogonopus tubulosus*. *Phytotherapy Research* 10, 198–201.
- Schmidt, T. J., Heilmann, J., 2002. Quantitative structure-cytotoxicity relationships of sesquiterpene lactones derived from partial charge (Q)-based fractional accessible surface area descriptors (Q\_frASAs). *Quantitative Structure-Activity Relationships* 21, 276–287.
- Schneider, W., 1989. *Malaria-Neue Aspekte in Diagnostik, Therapie und Prophylaxe*. Editiones <Roche>, Basel/Grenzach-Wyhlen Mayr Miesbach, Druckerei und Verlag GmbH.
- Silfen, J., Yanai, P., Cabantchik, Z. I., 1988. Bioflavonoid effects on *in vitro* cultures of *Plasmodium falciparum*. Inhibition of permeation pathways induced in the host cell membrane by the intraerythrocytic parasite. *Biochemical Pharmacology* 37, 4269–4276.
- Silva, M., Mundaca, J. M., 1971. Flavonoid and triterpene constituents of *Baccharis rhomboidalis*. *Phytochemical Reports* 10, 1942–1943.
- Singh, P., Jakupovic, J., Bohlmann, F., King, R. M., Robinson, H., 1985. Highly oxygenated guianolides from *Bishopanthus soliceps*. *Phytochemistry* 24, 2110–2112.
- Sittie, A. A., Lemmich, E., Olsen, C. E., Hviid, L., Kharazmi, A., Nkrumah, F. K., Christensen, S. B., 1999. Structure-activity studies: *in vitro* antileishmanial and antimalarial activities of anthraquinones from *Morinda lucida*. *Planta Medica* 65, 259–261.
- Steele, J. C. P., Simmonds, M. S. J., Veitch, N. C., Warhurst, D. C., 1999. Evaluation of the anti-plasmodial activity of bisbenzylisoquinoline alkaloids from *Abuta grandifolia*. *Planta Medica* 65, 413–416.
- Stomberg, R., Li, S., Lundquist, K., Wallis, A. F. A., 1997. Crystal structure of 2-(3',4'-dimethoxyphenyl)-3-hydromethyl-7-methoxybenzo[*b*]furan, C<sub>18</sub>H<sub>18</sub>O<sub>5</sub>. *Zeitschrift für Kristallographie - New Crystal structures* 212, 473–474.
- Sumaryono, W., Proksch, P., Wray, V., Witte, L., Hartmann, T., 1991. Qualitative and quantitative analysis of the phenolic constituents from *Orthosiphon aristatus*. *Planta Medica* 57, 176–180.
- Tahara, S., Ingham, J. L., Nakahara, S., Mizutani, J., Harborne, J. B., 1984. Fungitoxic dihydrofuranoisoflavones and related compounds in white lupin, *Lupinus albus*. *Phytochemistry* 23, 1889–1900.
- Talukdar, A. C., Jain, N., De, S., Krishnamurty, H. G., 2000. An isoflavone from *Myristica malabarica*. *Phytochemistry* 53, 155–157.
- Tan, R. X., Zheng, W. F., Tang, H. Q., 1998. Biologically active substances from the genus *Artemisia*. *Planta Medica* 64, 295–302.

- Tanaka, R., Matsunaga, S., 1989. Loliolide and Olean-12-en- $\beta$ ,9 $\alpha$ ,11 $\alpha$ -triol from *Euphorbia supina*. *Phytochemistry* 28, 1699–1702.
- Tchuendem, M. H. K., Mbah, J. A., Tsopmo, A., Ayafor, J. F., Sternér, O., Okunjic, C. C., Iwu, M. M., Schuster, B. M., 1999. Anti-plasmodial sesquiterpenoids from the African *Renealmia cincinnata*. *Phytochemistry* 52, 1095–1099.
- Thebtaranonth, C., Thebtaranonth, Y., Wanauppathamkul, S., Yuthavong, Y., 1995. Antimalarial sesquiterpenes from tubers of *Cyperus rotundus*: structure of 10,12-peroxycalamenene, a sesquiterpen endoperoxid. *Phytochemistry* 40, 125–128.
- Tofern, B., 1999. Dissertation, Fachbereich Pharmazie, Freie Universität Berlin.
- Tofern, B., Jenett-Siems, K., Siems, K., Jakupovic, J., Eich, E., 2000. Bonaspectins and neobonaspectins, first sesquilignans and sesquineolignans from a convolvulaceous species. *Phytochemistry* 53, 119–128.
- Toubiana, R., Gaudemer, A., 1967. Structure du vernalide, nouvel ester sesquiterpenique isolé de *Vernonia colorata*. *Tetrahedron Letters* 14, 1333–1336.
- Trager, W., Jensen, J. B., 1976. Human malaria parasites in continuous culture. *Science* 193, 673–675.
- Tschesche, R., Delhvi, S., Sepúlveda, S., Breitmaier, E., 1979. Eucryphin, a new chromone rhamnoside from the bark of *Eucryphia cordifolia*. *Phytochemistry* 18, 867–869.
- Van Agtmael, M. A., Eggelte, T. A., van Boxtel, C. J., 1999. Artemisinin drugs in the treatment of malaria: from medicinal herb to registered medication. *Trends in Pharmaceutical Sciences* 20, 199–205.
- Van Staden, L. F., Drewes, S. E., 1994. Knipholone from *Bulbine latifolia* and *Bulbine-frutescens*. *Phytochemistry* 35, 685–686.
- Vennerstrom, J. L., Klayman, D. L., 1988. Protoberberine alkaloids as antimalarials. *Journal of Medicinal Chemistry* 31, 1084–1087.
- Watt, J. M., Breyer-Brandwijk, M. G., 1962. The medicinal and poisonous plants of Southern and Eastern Africa, 2. Auflage. E.&S. Livingstone LTD: Edinburgh London.
- Weiss, C. R., Moideen, S. V. K., Croft, S. L., Houghton, P. J., 2000. Activity of extracts and isolated naphthoquinones from *Kigelia pinnata* against *Plasmodium falciparum*. *Journal of Natural Products* 63, 1306–1309.
- WHO, 1997. World malaria situation in 1994. *Weekly Epidemiological Record* 72, 269–292.
- Wichtl, M., 1997. Teedrogen und Phytopharmaka. Ein Handbuch für die Praxis auf wissenschaftlicher Grundlage, 3. Auflage. Wissenschaftliche Verlagsgesellschaft mbH Stuttgart.
- Wiedermann, G., Wernsdorfer, W. H., Kollaritsch, H., 1999. Update zur Malaria prophylaxe. *Österreichische Apotheker Zeitung* 11, 493–499.

- Wollenweber, E., 1974. Flavonoid-Exkretion bei *Betula*-Arten. Biochemie und Physiologie der Pflanzen 166, 425–428.
- Wollenweber, E., Mann, K., 1989. Exudate flavonoids in three essential oil plants from the Ciskei (South Africa). Fitoterapia 60, 249–251.
- Wright, C. W., Allen, D., Phillipson, J. D., Kirby, G. C., Warhurst, D. C., Massiot, G., Le Men-Olivier, L., 1993. *Alstonia* species: are they effective in malaria treatment ? Journal of Ethnopharmacology 40, 41–45.
- Wright, C. W., Bray, D. H., O'Neill, M. J., Warhurst, D. C., Phillipson, J. D., Quettin-Leclercq, J., Angenot, L., 1991. Antiamoebic and antiplasmodial activities of alkaloids isolated from *Strychnos usambarensis*. Planta Medica 57, 337–340.
- Wright, C. W., Marshall, S. J., Russel, P. F., Anderson, M. M., Phillipson, J. D., Kirby, G. C., Warhurst, D. C., Schiff Jr., P. L., 2000. *In vitro* antiplasmodial, antiamoebic, and cytotoxic activities of some monomeric isoquinoline alkaloids. Journal of Natural Products 63, 1638–1640.
- Wright, C. W., Phillipson, J. D., 1990. Natural products and the development of selective antiprotozoal drugs. Phytotherapy Research 4, 127–139.
- Yang, S., Roberts, M. F., O'Neill, M. J., Bucar, F., Phillipson, J. D., 1995. Flavonoids and chromenes from *Artemisia annua*. Phytochemistry 38, 255–257.
- Yang, Y.-Z., Ranz, A., Pan, H.-Z., Zhang, Z.-N., Lin, X.-B., Meshnick, S. R., 1992. Daphnetin: a novel antimalarial agent with *in vitro* and *in vivo* activity. American Journal of Tropical Medicine and Hygiene 46, 15–20.
- Yenjai, C., Sripontan, S., Sripajun, P., Kittakoop, P., Jintasirikul, A., Tanticharoen, M., Thebtaranonth, Y., 2000. Coumarins and carbazoles with antiplasmodial activity from *Clausena harmandiana*. Planta Medica 66, 277–279.
- Yu, H. W., Wright, C. W., Cai, Y., Yang, S. L., Phillipson, J. D., Kirby, G. C., Warhurst, D. C., 1994. Antiprotozoal activities of *Centipeda minima*. Phytotherapy Research 8, 436–438.
- Zdero, C., Bohlmann, F., Wassahusen, D. C., Mungai, M. G., 1991. Glaucolides from old world *Vernonia* species. Phytochemistry 30, 4025–4028.
- Zhang, H. J., Tamez, P. A., Hoang, V. D., Tan, G. T., Hung, N. V., Xuan, L. T., Huong, L. M., Cuong, N. M., Thao, D. T., Soejarto, D. D., Fong, H. H. S., Pezzuto, J. M., 2001. Antimalarial compounds from *Rhaphidophora decursiva*. Journal of Natural Products 64, 772–777.