

7 Literaturverzeichnis

- Addlagatta, A., Quillin, M. L., Omotoso, O., Liu, J. O. and Matthews, B. W. (2005). Identification of an SH3-binding motif in a new class of methionine aminopeptidases from *Mycobacterium tuberculosis* suggests a mode of interaction with the ribosome. *Biochemistry* **44**, 7166-74.
- Agashe, V. R., Guha, S., Chang, H. C., Genevoux, P., Hayer-Hartl, M., Stemp, M., Georgopoulos, C., Hartl, F. U. and Barral, J. M. (2004). Function of trigger factor and DnaK in multidomain protein folding: increase in yield at the expense of folding speed. *Cell* **117**, 199-209.
- Albanese, V., Yam, A. Y., Baughman, J., Parnot, C. and Frydman, J. (2006). Systems analyses reveal two chaperone networks with distinct functions in eukaryotic cells. *Cell* **124**, 75-88.
- Arcus, V. (2002). OB-fold domains: a snapshot of the evolution of sequence, structure and function. *Curr Opin Struct Biol* **12**, 794-801.
- Ban, N., Nissen, P., Hansen, J., Moore, P. B. and Steitz, T. A. (2000). The complete atomic structure of the large ribosomal subunit at 2.4 Å resolution. *Science* **289**, 905-20.
- Beatrix, B., Sakai, H. and Wiedmann, M. (2000). The alpha and beta subunit of the nascent polypeptide-associated complex have distinct functions. *J Biol Chem* **275**, 37838-45.
- Beckmann, R., Spahn, C. M., Eswar, N., Helmers, J., Penczek, P. A., Sali, A., Frank, J. and Blobel, G. (2001). Architecture of the protein-conducting channel associated with the translating 80S ribosome. *Cell* **107**, 361-72.
- Bloss, T. A., Witze, E. S. and Rothman, J. H. (2003). Suppression of CED-3-independent apoptosis by mitochondrial betaNAC in *Caenorhabditis elegans*. *Nature* **424**, 1066-71.
- Bonven, B. and Gullov, K. (1979). Peptide chain elongation rate and ribosomal activity in *Saccharomyces cerevisiae* as a function of the growth rate. *Mol Gen Genet* **170**, 225-30.
- Braakman, I., Hoover-Litty, H., Wagner, K. R. and Helenius, A. (1991). Folding of influenza hemagglutinin in the endoplasmic reticulum. *J Cell Biol* **114**, 401-11.
- Bradford, M. M. (1976). A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* **72**, 248-54.
- Branden, C. and Tooze, J. (1998). *Introduction to protein structure*, 2nd edn. Garland Pub.
- Buchner, J. (1996). Supervising the fold: functional principles of molecular chaperones. *Faseb J* **10**, 10-9.
- Bukau, B. (2005). Ribosomes catch Hsp70s. *Nat Struct Mol Biol* **12**, 472-3.
- Chang, H. C., Kaiser, C. M., Hartl, F. U. and Barral, J. M. (2005). De novo folding of GFP fusion proteins: high efficiency in eukaryotes but not in bacteria. *J Mol Biol* **353**, 397-409.
- Crooke, E. and Wickner, W. (1987). Trigger factor: a soluble protein that folds pro-OmpA into a membrane-assembly-competent form. *Proc Natl Acad Sci U S A* **84**, 5216-20.
- Deng, J. M. and Behringer, R. R. (1995). An insertional mutation in the BTF3 transcription factor gene leads to an early postimplantation lethality in mice. *Transgenic Res* **4**, 264-9.
- Deuerling, E., Patzelt, H., Vorderwulbecke, S., Rauch, T., Kramer, G., Schaffitzel, E., Mogk, A., Schulze-Specking, A., Langen, H. and Bukau, B. (2003). Trigger Factor and DnaK possess overlapping substrate pools and binding specificities. *Mol Microbiol* **47**, 1317-28.

- Deuerling, E., Schulze-Specking, A., Tomoyasu, T., Mogk, A. and Bukau, B. (1999). Trigger factor and DnaK cooperate in folding of newly synthesized proteins. *Nature* **400**, 693-6.
- Deutschbauer, A. M., Jaramillo, D. F., Proctor, M., Kumm, J., Hillenmeyer, M. E., Davis, R. W., Nislow, C. and Giaever, G. (2005). Mechanisms of haploinsufficiency revealed by genome-wide profiling in yeast. *Genetics* **169**, 1915-25.
- Dobson, C. M. and Karplus, M. (1999). The fundamentals of protein folding: bringing together theory and experiment. *Curr Opin Struct Biol* **9**, 92-101.
- Dorman, G. and Prestwich, G. D. (1994). Benzophenone photophores in biochemistry. *Biochemistry* **33**, 5661-73.
- Doudna, J. A. and Batey, R. T. (2004). Structural insights into the signal recognition particle. *Annu Rev Biochem* **73**, 539-57.
- Dunker, A. K., Cortese, M. S., Romero, P., Iakoucheva, L. M. and Uversky, V. N. (2005). Flexible nets. The roles of intrinsic disorder in protein interaction networks. *Febs J* **272**, 5129-48.
- Ellis, R. J. and Hartl, F. U. (1999). Principles of protein folding in the cellular environment. *Curr Opin Struct Biol* **9**, 102-10.
- Ferbitz, L., Maier, T., Patzelt, H., Bukau, B., Deuerling, E. and Ban, N. (2004). Trigger factor in complex with the ribosome forms a molecular cradle for nascent proteins. *Nature* **431**, 590-6.
- Frank, R. (1992). Spot synthesis: an easy technique for the positionally addressable parallel chemical synthesis on a membrane support. *Tetrahedron* **48**, 9217-9232.
- Franke, J., Reimann, B., Hartmann, E., Kohlerl, M. and Wiedmann, B. (2001). Evidence for a nuclear passage of nascent polypeptide-associated complex subunits in yeast. *J Cell Sci* **114**, 2641-8.
- Frydman, J. (2001). Folding of newly translated proteins in vivo: the role of molecular chaperones. *Annu Rev Biochem* **70**, 603-47.
- Frydman, J., Erdjument-Bromage, H., Tempst, P. and Hartl, F. U. (1999). Co-translational domain folding as the structural basis for the rapid de novo folding of firefly luciferase. *Nat Struct Biol* **6**, 697-705.
- Gautschi, M., Just, S., Mun, A., Ross, S., Rucknagel, P., Dubaquié, Y., Ehrenhofer-Murray, A. and Rospert, S. (2003). The yeast N(alpha)-acetyltransferase NatA is quantitatively anchored to the ribosome and interacts with nascent polypeptides. *Mol Cell Biol* **23**, 7403-14.
- Gautschi, M., Lilie, H., Funfschilling, U., Mun, A., Ross, S., Lithgow, T., Rucknagel, P. and Rospert, S. (2001). RAC, a stable ribosome-associated complex in yeast formed by the DnaK-DnaJ homologs Ssz1p and zutin. *Proc Natl Acad Sci U S A* **98**, 3762-7.
- Gautschi, M., Mun, A., Ross, S. and Rospert, S. (2002). A functional chaperone triad on the yeast ribosome. *Proc Natl Acad Sci U S A* **99**, 4209-14.
- Genevaux, P., Keppel, F., Schwager, F., Langendijk-Genevaux, P. S., Hartl, F. U. and Georgopoulos, C. (2004). In vivo analysis of the overlapping functions of DnaK and trigger factor. *EMBO Rep* **5**, 195-200.
- George, R., Beddoe, T., Landl, K. and Lithgow, T. (1998). The yeast nascent polypeptide-associated complex initiates protein targeting to mitochondria in vivo. *Proc Natl Acad Sci U S A* **95**, 2296-301.
- Gilbert, R. J., Fucini, P., Connell, S., Fuller, S. D., Nierhaus, K. H., Robinson, C. V., Dobson, C. M. and Stuart, D. I. (2004). Three-dimensional structures of translating ribosomes by Cryo-EM. *Mol Cell* **14**, 57-66.
- Gilmore, R., Collins, P., Johnson, J., Kellaris, K. and Rapiejko, P. (1991). Transcription of full-length and truncated mRNA transcripts to study protein translocation across the endoplasmic reticulum. *Methods Cell Biol* **34**, 223-39.

- Görlich, D., Kurzchalia, T. V., Wiedmann, M. and Rapoport, T. A. (1991). Probing the molecular environment of translocating polypeptide chains by cross-linking. *Methods Cell Biol* **34**, 241-62.
- Görlich, D., Prehn, S., Hartmann, E., Kalies, K. U. and Rapoport, T. A. (1992). A mammalian homolog of SEC61p and SECYp is associated with ribosomes and nascent polypeptides during translocation. *Cell* **71**, 489-503.
- Gouet, P., Courcelle, E., Stuart, D. I. and Metz, F. (1999). ESPript: analysis of multiple sequence alignments in PostScript. *Bioinformatics* **15**, 305-8.
- Grallath, S., Schwarz, J. P., Bottcher, U. M., Bracher, A., Hartl, F. U. and Siegers, K. (2006). L25 functions as a conserved ribosomal docking site shared by nascent chain-associated complex and signal-recognition particle. *EMBO Rep* **7**, 78-84.
- Gu, S. Q., Peske, F., Wieden, H. J., Rodnina, M. V. and Wintermeyer, W. (2003). The signal recognition particle binds to protein L23 at the peptide exit of the Escherichia coli ribosome. *Rna* **9**, 566-73.
- Halic, M., Becker, T., Pool, M. R., Spahn, C. M., Grassucci, R. A., Frank, J. and Beckmann, R. (2004). Structure of the signal recognition particle interacting with the elongation-arrested ribosome. *Nature* **427**, 808-14.
- Halic, M. and Beckmann, R. (2005). The signal recognition particle and its interactions during protein targeting. *Curr Opin Struct Biol* **15**, 116-25.
- Halic, M., Gartmann, M., Schlenker, O., Mielke, T., Pool, M. R., Sinning, I. and Beckmann, R. (2006). Signal recognition particle receptor exposes the ribosomal translocon binding site. *Science* **312**, 745-7.
- Harms, J., Schluenzen, F., Zarivach, R., Bashan, A., Gat, S., Agmon, I., Bartels, H., Franceschi, F. and Yonath, A. (2001). High resolution structure of the large ribosomal subunit from a mesophilic eubacterium. *Cell* **107**, 679-88.
- Hartl, F. U. and Hayer-Hartl, M. (2002). Molecular chaperones in the cytosol: from nascent chain to folded protein. *Science* **295**, 1852-8.
- Hesterkamp, T., Deuerling, E. and Bukau, B. (1997). The amino-terminal 118 amino acids of Escherichia coli trigger factor constitute a domain that is necessary and sufficient for binding to ribosomes. *J Biol Chem* **272**, 21865-71.
- Hesterkamp, T., Hauser, S., Lutcke, H. and Bukau, B. (1996). Escherichia coli trigger factor is a prolyl isomerase that associates with nascent polypeptide chains. *Proc Natl Acad Sci U S A* **93**, 4437-41.
- Hu, G. Z. and Ronne, H. (1994). Yeast BTF3 protein is encoded by duplicated genes and inhibits the expression of some genes in vivo. *Nucleic Acids Res* **22**, 2740-3.
- Huang, P., Gautschi, M., Walter, W., Rospert, S. and Craig, E. A. (2005). The Hsp70 Ssz1 modulates the function of the ribosome-associated J-protein Zuo1. *Nat Struct Mol Biol* **12**, 497-504.
- Hundley, H., Eisenman, H., Walter, W., Evans, T., Hotokezaka, Y., Wiedmann, M. and Craig, E. (2002). The in vivo function of the ribosome-associated Hsp70, Ssz1, does not require its putative peptide-binding domain. *Proc Natl Acad Sci U S A* **99**, 4203-8.
- Hundley, H. A., Walter, W., Bairstow, S. and Craig, E. A. (2005). Human Mpp11 J protein: ribosome-tethered molecular chaperones are ubiquitous. *Science* **308**, 1032-4.
- Jackson, R. J. and Hunt, T. (1983). Preparation and use of nuclease-treated rabbit reticulocyte lysates for the translation of eukaryotic messenger RNA. *Methods Enzymol* **96**, 50-74.
- Kaeberlein, M., Powers, R. W., 3rd, Steffen, K. K., Westman, E. A., Hu, D., Dang, N., Kerr, E. O., Kirkland, K. T., Fields, S. and Kennedy, B. K. (2005). Regulation of yeast replicative life span by TOR and Sch9 in response to nutrients. *Science* **310**, 1193-6.
- Kalies, K. U., Görlich, D. and Rapoport, T. A. (1994). Binding of ribosomes to the rough endoplasmic reticulum mediated by the Sec61p-complex. *J Cell Biol* **126**, 925-34.

- Kalthoff, C. (2003). A novel strategy for the purification of recombinantly expressed unstructured protein domains. *J Chromatogr B Analyt Technol Biomed Life Sci* **786**, 247-54.
- Kanno, M., Chalut, C. and Egly, J. M. (1992). Genomic structure of the putative BTF3 transcription factor. *Gene* **117**, 219-28.
- Keenan, R. J., Freymann, D. M., Stroud, R. M. and Walter, P. (2001). The signal recognition particle. *Annu Rev Biochem* **70**, 755-75.
- Kelley, W. L. (1999). Molecular chaperones: How J domains turn on Hsp70s. *Curr Biol* **9**, R305-8.
- Kim, S. H., Shim, K. S. and Lubec, G. (2002). Human brain nascent polypeptide-associated complex alpha subunit is decreased in patients with Alzheimer's disease and Down syndrome. *J Investig Med* **50**, 293-301.
- Klein, D. J., Moore, P. B. and Steitz, T. A. (2004). The roles of ribosomal proteins in the structure assembly, and evolution of the large ribosomal subunit. *J Mol Biol* **340**, 141-77.
- Koch, H. G., Moser, M. and Muller, M. (2003). Signal recognition particle-dependent protein targeting, universal to all kingdoms of life. *Rev Physiol Biochem Pharmacol* **146**, 55-94.
- Kolb, V. A., Makeyev, E. V. and Spirin, A. S. (2000). Co-translational folding of an eukaryotic multidomain protein in a prokaryotic translation system. *J Biol Chem* **275**, 16597-601.
- Kramer, G., Patzelt, H., Rauch, T., Kurz, T. A., Vorderwulbecke, S., Bukau, B. and Deuerling, E. (2004a). Trigger factor peptidyl-prolyl cis/trans isomerase activity is not essential for the folding of cytosolic proteins in Escherichia coli. *J Biol Chem* **279**, 14165-70.
- Kramer, G., Rauch, T., Rist, W., Vorderwulbecke, S., Patzelt, H., Schulze-Specking, A., Ban, N., Deuerling, E. and Bukau, B. (2002). L23 protein functions as a chaperone docking site on the ribosome. *Nature* **419**, 171-4.
- Kramer, G., Rutkowska, A., Wegryzn, R. D., Patzelt, H., Kurz, T. A., Merz, F., Rauch, T., Vorderwulbecke, S., Deuerling, E. and Bukau, B. (2004b). Functional dissection of Escherichia coli trigger factor: unraveling the function of individual domains. *J Bacteriol* **186**, 3777-84.
- Kroes, R. A., Jastrow, A., McLone, M. G., Yamamoto, H., Colley, P., Kersey, D. S., Yong, V. W., Mkrdichian, E., Cerullo, L., Leestma, J. and Moskal, J. R. (2000). The identification of novel therapeutic targets for the treatment of malignant brain tumors. *Cancer Lett* **156**, 191-8.
- Laemmli, U. K. (1970). Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* **227**, 680-5.
- Laksanalamai, P., Whitehead, T. A. and Robb, F. T. (2004). Minimal protein-folding systems in hyperthermophilic archaea. *Nat Rev Microbiol* **2**, 315-24.
- Laufen, T., Mayer, M. P., Beisel, C., Klostermeier, D., Mogk, A., Reinstein, J. and Bukau, B. (1999). Mechanism of regulation of hsp70 chaperones by DnaJ cochaperones. *Proc Natl Acad Sci U S A* **96**, 5452-7.
- Lauring, B., Kreibich, G. and Weidmann, M. (1995a). The intrinsic ability of ribosomes to bind to endoplasmic reticulum membranes is regulated by signal recognition particle and nascent-polypeptide-associated complex. *Proc Natl Acad Sci U S A* **92**, 9435-9.
- Lauring, B., Sakai, H., Kreibich, G. and Wiedmann, M. (1995b). Nascent polypeptide-associated complex protein prevents mistargeting of nascent chains to the endoplasmic reticulum. *Proc Natl Acad Sci U S A* **92**, 5411-5.
- Lauring, B., Wang, S., Sakai, H., Davis, T. A., Wiedmann, B., Kreibich, G. and Wiedmann, M. (1995c). Nascent-polypeptide-associated complex: a bridge between ribosome and cytosol. *Cold Spring Harb Symp Quant Biol* **60**, 47-56.
- Lopez, N., Halladay, J., Walter, W. and Craig, E. A. (1999). SSB, encoding a ribosome-associated chaperone, is coordinately regulated with ribosomal protein genes. *J Bacteriol* **181**, 3136-43.

- Macario, A. J. and Conway De Macario, E. (2001). The molecular chaperone system and other anti-stress mechanisms in archaea. *Front Biosci* **6**, D262-83.
- Maier, R., Scholz, C. and Schmid, F. X. (2001). Dynamic association of trigger factor with protein substrates. *J Mol Biol* **314**, 1181-90.
- Makarova, K. S., Aravind, L., Galperin, M. Y., Grishin, N. V., Tatusov, R. L., Wolf, Y. I. and Koonin, E. V. (1999). Comparative genomics of the Archaea (Euryarchaeota): evolution of conserved protein families, the stable core, and the variable shell. *Genome Res* **9**, 608-28.
- Marchler-Bauer, A., Panchenko, A. R., Shoemaker, B. A., Thiessen, P. A., Geer, L. Y. and Bryant, S. H. (2002). CDD: a database of conserved domain alignments with links to domain three-dimensional structure. *Nucleic Acids Res* **30**, 281-3.
- Markesich, D. C., Gajewski, K. M., Nazimiec, M. E. and Beckingham, K. (2000). bicaudal encodes the Drosophila beta NAC homolog, a component of the ribosomal translational machinery*. *Development* **127**, 559-72.
- Möller, I., Beatrix, B., Kreibich, G., Sakai, H., Lauring, B. and Wiedmann, M. (1998a). Unregulated exposure of the ribosomal M-site caused by NAC depletion results in delivery of non-secretory polypeptides to the Sec61 complex. *FEBS Lett* **441**, 1-5.
- Möller, I., Jung, M., Beatrix, B., Levy, R., Kreibich, G., Zimmermann, R., Wiedmann, M. and Lauring, B. (1998b). A general mechanism for regulation of access to the translocon: competition for a membrane attachment site on ribosomes. *Proc Natl Acad Sci U S A* **95**, 13425-30.
- Moncollin, V., Fischer, L., Cavallini, B., Egly, J. M. and Chambon, P. (1992). Class II (B) general transcription factor (TFIIB) that binds to the template-committed preinitiation complex is different from general transcription factor BTF3. *Proc Natl Acad Sci U S A* **89**, 397-401.
- Moncollin, V., Miyamoto, N. G., Zheng, X. M. and Egly, J. M. (1986). Purification of a factor specific for the upstream element of the adenovirus-2 major late promoter. *Embo J* **5**, 2577-84.
- Moreau, A., Yotov, W. V., Glorieux, F. H. and St-Arnaud, R. (1998). Bone-specific expression of the alpha chain of the nascent polypeptide-associated complex, a coactivator potentiating c-Jun-mediated transcription. *Mol Cell Biol* **18**, 1312-21.
- Mueller, T. D., Kamionka, M. and Feigon, J. (2004). Specificity of the interaction between ubiquitin-associated domains and ubiquitin. *J Biol Chem* **279**, 11926-36.
- Neidhardt, F. C. (1987). *Escherichia coli and Salmonella typhimurium : cellular and molecular biology*. American Society for Microbiology.
- Netzer, W. J. and Hartl, F. U. (1997). Recombination of protein domains facilitated by co-translational folding in eukaryotes. *Nature* **388**, 343-9.
- Netzer, W. J. and Hartl, F. U. (1998). Protein folding in the cytosol: chaperonin-dependent and -independent mechanisms. *Trends Biochem Sci* **23**, 68-73.
- Neupert, W. and Lill, R. (1994). Protein synthesis. Cradle at the ribosome. *Nature* **370**, 421-2.
- Newton, C. R. and Graham, A. (1997). *Pcr*, 2nd edn. BIOS Scientific Publishers ; Springer.
- Nicola, A. V., Chen, W. and Helenius, A. (1999). Co-translational folding of an alphavirus capsid protein in the cytosol of living cells. *Nat Cell Biol* **1**, 341-5.
- Osborne, A. R., Rapoport, T. A. and van den Berg, B. (2005). Protein translocation by the Sec61/SecY channel. *Annu Rev Cell Dev Biol* **21**, 529-50.
- Otto, H., Conz, C., Maier, P., Wolffe, T., Suzuki, C. K., Jenö, P., Rucknagel, P., Stahl, J. and Rospert, S. (2005). The chaperones MPP11 and Hsp70L1 form the mammalian ribosome-associated complex. *Proc Natl Acad Sci U S A* **102**, 10064-9.
- Parthun, M. R., Mangus, D. A. and Jaehning, J. A. (1992). The EGD1 product, a yeast homolog of human BTF3, may be involved in GAL4 DNA binding. *Mol Cell Biol* **12**, 5683-9.

- Pfund, C., Lopez-Hoyo, N., Ziegelhoffer, T., Schilke, B. A., Lopez-Buesa, P., Walter, W. A., Wiedmann, M. and Craig, E. A. (1998). The molecular chaperone Ssb from *Saccharomyces cerevisiae* is a component of the ribosome-nascent chain complex. *Embo J* **17**, 3981-9.
- Pool, M. R., Stumm, J., Fulga, T. A., Sinning, I. and Dobberstein, B. (2002). Distinct modes of signal recognition particle interaction with the ribosome. *Science* **297**, 1345-8.
- Potter, M. D., Seiser, R. M. and Nicchitta, C. V. (2001). Ribosome exchange revisited: a mechanism for translation-coupled ribosome detachment from the ER membrane. *Trends Cell Biol* **11**, 112-5.
- Raasi, S. and Pickart, C. M. (2003). Rad23 ubiquitin-associated domains (UBA) inhibit 26 S proteasome-catalyzed proteolysis by sequestering lysine 48-linked polyubiquitin chains. *J Biol Chem* **278**, 8951-9.
- Raasi, S., Varadan, R., Fushman, D. and Pickart, C. M. (2005). Diverse polyubiquitin interaction properties of ubiquitin-associated domains. *Nat Struct Mol Biol* **12**, 708-14.
- Raden, D. and Gilmore, R. (1998). Signal recognition particle-dependent targeting of ribosomes to the rough endoplasmic reticulum in the absence and presence of the nascent polypeptide-associated complex. *Mol Biol Cell* **9**, 117-30.
- Rakwalska, M. and Rospert, S. (2004). The ribosome-bound chaperones RAC and Ssb1/2p are required for accurate translation in *Saccharomyces cerevisiae*. *Mol Cell Biol* **24**, 9186-97.
- Receveur-Brechot, V., Bourhis, J. M., Uversky, V. N., Canard, B. and Longhi, S. (2006). Assessing protein disorder and induced folding. *Proteins* **62**, 24-45.
- Reimann, B., Bradsher, J., Franke, J., Hartmann, E., Wiedmann, M., Prehn, S. and Wiedmann, B. (1999). Initial characterization of the nascent polypeptide-associated complex in yeast. *Yeast* **15**, 397-407.
- Rhee, S. S. and Burke, D. H. (2004). Tris(2-carboxyethyl)phosphine stabilization of RNA: comparison with dithiothreitol for use with nucleic acid and thiophosphoryl chemistry. *Anal Biochem* **325**, 137-43.
- Rost, B. and Sander, C. (1993). Prediction of protein secondary structure at better than 70% accuracy. *J Mol Biol* **232**, 584-99.
- Sambrook, J., Fritsch, E. F. and Maniatis, T. (1989). *Molecular cloning : a laboratory manual*, 2nd edn. Cold Spring Harbor Laboratory.
- Sanger, F., Nicklen, S. and Coulson, A. R. (1977). DNA sequencing with chain-terminating inhibitors. *Proc Natl Acad Sci U S A* **74**, 5463-7.
- Scheuring, U. J., Corbeil, J., Mosier, D. E. and Theofilopoulos, A. N. (1998). Early modification of host cell gene expression induced by HIV-1. *Aids* **12**, 563-70.
- Schlünzen, F., Tocilj, A., Zarivach, R., Harms, J., Gluehmann, M., Janell, D., Bashan, A., Bartels, H., Agmon, I., Franceschi, F. and Yonath, A. (2000). Structure of functionally activated small ribosomal subunit at 3.3 angstroms resolution. *Cell* **102**, 615-23.
- Schneider, B. L., Steiner, B., Seufert, W. and Futcher, A. B. (1996). pMPY-ZAP: a reusable polymerase chain reaction-directed gene disruption cassette for *Saccharomyces cerevisiae*. *Yeast* **12**, 129-34.
- Schubert, U., Anton, L. C., Gibbs, J., Norbury, C. C., Yewdell, J. W. and Bennink, J. R. (2000). Rapid degradation of a large fraction of newly synthesized proteins by proteasomes. *Nature* **404**, 770-4.
- Schuwirth, B. S., Borovinskaya, M. A., Hau, C. W., Zhang, W., Vila-Sanjurjo, A., Holton, J. M. and Cate, J. H. (2005). Structures of the bacterial ribosome at 3.5 Å resolution. *Science* **310**, 827-34.
- Shi, X., Parthun, M. R. and Jaehning, J. A. (1995). The yeast EGD2 gene encodes a homologue of the alpha NAC subunit of the human nascent-polypeptide-associated complex. *Gene* **165**, 199-202.

- Sikorski, R. S. and Hieter, P. (1989). A system of shuttle vectors and yeast host strains designed for efficient manipulation of DNA in *Saccharomyces cerevisiae*. *Genetics* **122**, 19-27.
- Smith, R. L., Baca, O. and Gordon, J. (1976). Co-ordinate synthesis of ribosomes and elongation factors in the liver of immature chicks following a metabolic shift up. *J Mol Biol* **100**, 115-26.
- Spahn, C. M., Beckmann, R., Eswar, N., Penczek, P. A., Sali, A., Blobel, G. and Frank, J. (2001). Structure of the 80S ribosome from *Saccharomyces cerevisiae*--tRNA-ribosome and subunit-subunit interactions. *Cell* **107**, 373-86.
- Spahn, C. M., Gomez-Lorenzo, M. G., Grassucci, R. A., Jorgensen, R., Andersen, G. R., Beckmann, R., Penczek, P. A., Ballesta, J. P. and Frank, J. (2004). Domain movements of elongation factor eEF2 and the eukaryotic 80S ribosome facilitate tRNA translocation. *Embo J* **23**, 1008-19.
- Spreter, T., Pech, M. and Beatrix, B. (2005). The crystal structure of archaeal nascent polypeptide-associated complex (NAC) reveals a unique fold and the presence of a ubiquitin-associated domain. *J Biol Chem* **280**, 15849-54.
- Stoller, G., Rucknagel, K. P., Nierhaus, K. H., Schmid, F. X., Fischer, G. and Rahfeld, J. U. (1995). A ribosome-associated peptidyl-prolyl cis/trans isomerase identified as the trigger factor. *Embo J* **14**, 4939-48.
- Stroh, C. and Schulze-Osthoff, K. (1998). Death by a thousand cuts: an ever increasing list of caspase substrates. *Cell Death Differ* **5**, 997-1000.
- Terzi, L., Pool, M. R., Dobberstein, B. and Strub, K. (2004). Signal recognition particle Alu domain occupies a defined site at the ribosomal subunit interface upon signal sequence recognition. *Biochemistry* **43**, 107-17.
- Teter, S. A., Houry, W. A., Ang, D., Tradler, T., Rockabrand, D., Fischer, G., Blum, P., Georgopoulos, C. and Hartl, F. U. (1999). Polypeptide flux through bacterial Hsp70: DnaK cooperates with trigger factor in chaperoning nascent chains. *Cell* **97**, 755-65.
- Theobald, D. L., Mitton-Fry, R. M. and Wuttke, D. S. (2003). Nucleic acid recognition by OB-fold proteins. *Annu Rev Biophys Biomol Struct* **32**, 115-33.
- Thiede, B., Dimmler, C., Siejak, F. and Rudel, T. (2001). Predominant identification of RNA-binding proteins in Fas-induced apoptosis by proteome analysis. *J Biol Chem* **276**, 26044-50.
- Thomas, B. J. and Rothstein, R. (1989). Elevated recombination rates in transcriptionally active DNA. *Cell* **56**, 619-30.
- Thompson, J. D., Higgins, D. G. and Gibson, T. J. (1994). CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res* **22**, 4673-80.
- Tompa, P. (2002). Intrinsically unstructured proteins. *Trends Biochem Sci* **27**, 527-33.
- Towbin, H., Staehelin, T. and Gordon, J. (1979). Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets: procedure and some applications. *Proc Natl Acad Sci U S A* **76**, 4350-4.
- Turner, G. C. and Varshavsky, A. (2000). Detecting and measuring cotranslational protein degradation in vivo. *Science* **289**, 2117-20.
- Ullers, R. S., Luirink, J., Harms, N., Schwager, F., Georgopoulos, C. and Genevaux, P. (2004). SecB is a bona fide generalized chaperone in *Escherichia coli*. *Proc Natl Acad Sci U S A* **101**, 7583-8.
- Valent, Q. A., Kendall, D. A., High, S., Kusters, R., Oudega, B. and Luirink, J. (1995). Early events in preprotein recognition in *E. coli*: interaction of SRP and trigger factor with nascent polypeptides. *Embo J* **14**, 5494-505.
- Vorderwülbecke, S., Kramer, G., Merz, F., Kurz, T. A., Rauch, T., Zachmann-Brand, B., Bukau, B. and Deuerling, E. (2004). Low temperature or GroEL/ES overproduction permits growth of *Escherichia coli* cells lacking trigger factor and DnaK. *FEBS Lett* **559**, 181-7.

- Wang, S., Sakai, H. and Wiedmann, M. (1995). NAC covers ribosome-associated nascent chains thereby forming a protective environment for regions of nascent chains just emerging from the peptidyl transferase center. *J Cell Biol* **130**, 519-28.
- Ward, J. J., McGuffin, L. J., Bryson, K., Buxton, B. F. and Jones, D. T. (2004). The DISOPRED server for the prediction of protein disorder. *Bioinformatics* **20**, 2138-9.
- Wegrzyn, R. D. and Deuerling, E. (2005). Molecular guardians for newborn proteins: ribosome-associated chaperones and their role in protein folding. *Cell Mol Life Sci* **62**, 2727-38.
- Wegrzyn, R. D., Hofmann, D., Merz, F., Nikolay, R., Rauch, T., Graf, C. and Deuerling, E. (2006). A conserved motif is prerequisite for the interaction of NAC with ribosomal protein L23 and nascent chains. *J Biol Chem* **281**, 2847-57.
- Wenschuh, H., Volkmer-Engert, R., Schmidt, M., Schulz, M., Schneider-Mergener, J. and Reineke, U. (2000). Coherent membrane supports for parallel microsynthesis and screening of bioactive peptides. *Biopolymers* **55**, 188-206.
- Whitehead, T. P., Bullock, D. G. and Carter, T. J. (1979). High-density lipoprotein cholesterol analysis. *Clin Chem* **25**, 2055-7.
- Wickner, W. and Schekman, R. (2005). Protein translocation across biological membranes. *Science* **310**, 1452-6.
- Widmann, M. and Christen, P. (2000). Comparison of folding rates of homologous prokaryotic and eukaryotic proteins. *J Biol Chem* **275**, 18619-22.
- Wiedmann, B., Sakai, H., Davis, T. A. and Wiedmann, M. (1994). A protein complex required for signal-sequence-specific sorting and translocation. *Nature* **370**, 434-40.
- Wilson, D. N. and Nierhaus, K. H. (2003). The ribosome through the looking glass. *Angew Chem Int Ed Engl* **42**, 3464-86.
- Wimberly, B. T., Brodersen, D. E., Clemons, W. M., Jr., Morgan-Warren, R. J., Carter, A. P., Vornrhein, C., Hartsch, T. and Ramakrishnan, V. (2000). Structure of the 30S ribosomal subunit. *Nature* **407**, 327-39.
- Woolhead, C. A., McCormick, P. J. and Johnson, A. E. (2004). Nascent membrane and secretory proteins differ in FRET-detected folding far inside the ribosome and in their exposure to ribosomal proteins. *Cell* **116**, 725-36.
- Yaffe, M. P. and Schatz, G. (1984). Two nuclear mutations that block mitochondrial protein import in yeast. *Proc Natl Acad Sci U S A* **81**, 4819-23.
- Yan, W., Schilke, B., Pfund, C., Walter, W., Kim, S. and Craig, E. A. (1998). Zuo1, a ribosome-associated DnaJ molecular chaperone. *Embo J* **17**, 4809-17.
- Yewdell, J. W. (2001). Not such a dismal science: the economics of protein synthesis, folding, degradation and antigen processing. *Trends Cell Biol* **11**, 294-7.
- Yotov, W. V., Moreau, A. and St-Arnaud, R. (1998). The alpha chain of the nascent polypeptide-associated complex functions as a transcriptional coactivator. *Mol Cell Biol* **18**, 1303-11.
- Yotov, W. V. and St-Arnaud, R. (1996a). Differential splicing-in of a proline-rich exon converts alphaNAC into a muscle-specific transcription factor. *Genes Dev* **10**, 1763-72.
- Yotov, W. V. and St-Arnaud, R. (1996b). Mapping of the human gene for the alpha-NAC/1.9.2 (NACA/1.9.2) transcriptional coactivator to Chromosome 12q23-24.1. *Mamm Genome* **7**, 163-4.
- Young, J. C., Agashe, V. R., Siegers, K. and Hartl, F. U. (2004). Pathways of chaperone-mediated protein folding in the cytosol. *Nat Rev Mol Cell Biol* **5**, 781-91.
- Zarnt, T., Tradler, T., Stoller, G., Scholz, C., Schmid, F. X. and Fischer, G. (1997). Modular structure of the trigger factor required for high activity in protein folding. *J Mol Biol* **271**, 827-37.

- Zheng, X. M., Black, D., Chambon, P. and Egly, J. M. (1990). Sequencing and expression of complementary DNA for the general transcription factor BTF3. *Nature* **344**, 556-9.
- Zheng, X. M., Moncollin, V., Egly, J. M. and Chambon, P. (1987). A general transcription factor forms a stable complex with RNA polymerase B (II). *Cell* **50**, 361-8.