

7 Literaturverzeichnis

1. Alfaro V, Pesquero J, Palacios L. Acid-base disturbance during hemorrhage in rats: significant role of strong inorganic ions. *J Appl Physiol* 1999;86,1617-1625.
2. Andersen JL, Sandgaard NC, Bie P. Volume expansion during acute angiotensin II receptor (AT(1)) blockade and NOS inhibition in conscious dogs. *Am J Physiol* 2002;282,R1140-1148.
3. Arnestad JP, Bengtsson A, Bengtson JP, Tylman M, Redl H, Schlag G. Formation of cytokines by retransfusion of shed whole blood. *Br J Anaesth* 1994;72,422-425.
4. Atcheson R, Lambert DG. Update on opioid receptors. *Br J Anaesth* 1994;73,132-134.
5. Ba ZF, Wang P, Koo DJ, Cioffi WG, Bland KI, Chaudry IH. Alterations in tissue oxygen consumption and extraction after trauma and hemorrhagic shock. *Crit Care Med* 2000;28,2837-2842.
6. Bainbridge FA. The influence of venous filling upon the rate of the heart. *J Physiol* 1915;50,65-84.
7. Baumert JH, Reyle-Hahn M, Hecker K, Tenbrinck R, Kuhlen R, Rossaint R. Increased airway resistance during xenon anaesthesia in pigs is attributed to physical properties of the gas. *Br J Anaesth* 2002;88,540-545.
8. Berthold H, Munter K, Just A, Kirchheim HR, Ehmke H. Stimulation of the renin-angiotensin system by endothelin subtype A receptor blockade in conscious dogs. *Hypertension* 1999;33,1420-1424.
9. Bia MJ, DeFronzo RA. Extrarenal potassium homeostasis. *Am J Physiol* 1981;240,F257-268.
10. Boemke W, Krebs MO, Djalali K, Bünger H, Kaczmarczyk G. Renal nerves are not involved in sodium and water retention during mechanical ventilation in awake dogs. *Anesthesiology* 1998;89,942-953.
11. Boemke W, Hocher B, Schleyer N, Krebs MO, Kaczmarczyk G. Hemodynamic, renal, and endocrine responses to acute ET(A) blockade at different ANGII plasma levels. *Am J Physiol* 2001;280,R1322-1331.
12. Boomsma F, Ruprecht J, Man in 't Veld AJ, de Jong FH, Dzoljic M, Lachmann B. Haemodynamic and neurohumoral effects of xenon anaesthesia: a comparison with nitrous oxide. *Anaesthesia* 1990;45,273-278.
13. Bourque CW, Oliet SHR. Osmoreceptors in central nervous system. *Annu Rev*

- Physiol 1997;59,601-619.
14. Brent New R, Sampson AC, King MK, Hendrick JW, Clair MJ, McElmurray JH, Mandel J, Mukherjee R, Gasparo M, Spinale FG. Effects of combined angiotensin II and endothelin receptor blockade with developing heart failure. *Circulation* 2000;102,1447-1453.
15. Bufalari A, Miller SM, Giannoni C, Short CE. The use of propofol as an induction agent for halothane and isoflurane anesthesia in dogs. *J Am Anim Hosp Assoc* 1998;34,84-91.
16. Burnier M, Brunner HR. Angiotensin II receptor antagonists. *Lancet* 2000;355,637-645.
17. Cahalan MK, Weiskopf RB, Eger EI 2nd, Yasuda N, Ionescu P, Rampil IJ, Lockhart SH, Freire B, Peterson NA. Hemodynamic effects of desflurane/nitrous oxide anesthesia in volunteers. *Anesth Analg* 1991;73,157-164.
18. Calzia E, Stahl W, Handschuh T, Marx T, Fröba G, Bäder S, Georgieff M, Rademacher P. Respiratory mechanics during xenon anesthesia in pigs. *Anesthesiology* 1999;91,1378-1386.
19. Cernacek P, Strmen J, Levy M. Acute renal effects of endothelin-A blockade: interspecies differences. *J Cardivasc Pharmacol* 1998;31,S269-272.
20. Chang HC, Wu JE, Wang SM, Hung CR. Plasma endothelin level changes during hemorrhagic shock. *J Trauma* 1993;35,825-833.
21. Chiu AT, McCall DE, Price WA, Wong PC, Carini DJ, Duncia JV, Wexler RR, Yoo SE, Johnson AL, Timmermans PB. Nonpeptide angiotensin II receptor antagonists. VII. Cellular and biochemical pharmacology of DuP 753, an orally active antihypertensive agent. *J Pharmacol Exp Ther* 1990;252,711-718.
22. Cox SL, Schelb V, Trendelenburg AU, Starke K. Enhancement of noradrenaline release by angiotensin II and bradykinin in mouse atria: evidence for cross-talk between G(q/11) protein- and G(i/o) protein-coupled receptors. *Br J Pharmacol* 2000;129,1095-1102.
23. Cullen SC, Gross EG. The anesthetic properties of xenon in animals and human beings with additional observations on krypton. *Science* 1951;113,580-582.
24. Cullen D: Anesthetic depth and MAC. In: *Anesthesia*. Miller R (ed) New York, Churchill Livingstone,1986,553-80.
25. Culman J, Blume A, Gohlke P, Unger T. The renin-angiotensin system in the brain: possible therapeutic implications for AT(1)-receptor blockers. *J Hum*

- Hypertens 2002;16,S64-70.
26. Darlington DN, Shinsako J, Dallman MF. Responses of ACTH, epinephrine, norepinephrine, and cardiovascular system to hemorrhage. Am J Physiol 1986;251,H612-618.
27. Davenport A, Maguire J. Is endothelin-induced vasoconstriction mediated only by ET_A receptors in humans? Trends Pharmacol Sci 1994;15,9-11.
28. Deegan R, He HB, Wood AJJ, Wood M. Effect of enflurane and isoflurane on norepinephrine kinetics: A new approach to assessment of sympathetic function during anesthesia. Anesth Analg 1993;77,49-54.
29. Donckier JE, Massart PE, Hodeige D, Van Mechelen H, Clozel JP, Laloux O, Ketelslegers JM, Charlier AA, Heyndrickx GR. Additional hypotensive effect of endothelin-1 receptor antagonism in hypertensive dogs under angiotensin-converting enzyme inhibition. Circulation 1995;96,1250-1256.
30. Dupuis J, Stewart DJ, Cernacek P, Gosselin G. Human pulmonary circulation is an important site for both clearance and production of endothelin-1. Circulation 1996;94,1578-1584.
31. Dupuis J, Schwab AJ, Simard A, Cernacek P, Stewart DJ, Goresky CA. Kinetics of endothelin-1 binding in the dog liver microcirculation in vivo. Am J Physiol 1999;277,G905-914.
32. Eger EI 2nd, Brandstater B, Saidman LJ, Regan MJ, Severinghaus JW, Munson ES. Equipotent alveolar concentrations of methoxyflurane, halothane, diethyl ether, fluroxene, cyclopropane, xenon and nitrous oxide. Anesthesiology 1965;26,771-777.
33. Eger EI 2nd, Saidmann LJ, Brandstater B. Minimum alveolar anesthetic concentration: A standard of anesthetic potency. Anesthesiology 1965a;26;756-763.
34. Engeland WC, Dempsher DP, Byrnes GJ, Presnell K, Gann DS. The adrenal medullary response to graded hemorrhage in awake dogs. Endocrinology 1981;109,1539-1544.
35. Evans RG, Ventura S, Dampney RA, Ludbrook J. Neural mechanisms in the cardiovascular responses to acute central hypovolaemia. Clin Exp Pharmacol Physiol 2001;28,479-487.
36. Fejes-Toth G, Brinck-Johnsen T, Naray-Fejes-Toth A. Cardiovascular and hormonal response to hemorrhage in conscious rats. Am J Physiol

- 1988;254,H947-953.
37. Ficai S, Herizi A, Mimran A, Jover B. Endothelin blockade in angiotensin II hypertension: prevention and treatment studies in the rat. *Clin Exp Pharmacol Physiol* 2001;28,1100-1103.
38. Francis RCE, Höhne C, Kaczmarczyk G, Boemke W. Effects of angiotensin II and endothelin-1 blockade on the hemodynamic changes after acute blood loss and retransfusion in conscious dogs. *Acta Physiol Scand* 2004;180,319-328.
39. Franks NP, Dickenson R, de Sousa SL, Hall AC, Lieb WR. How does xenon produce anesthesia? *Nature* 1998;396,324.
40. Galie N, Seeger W, Naeije R, Simonneau G, Rubin LJ. Comparative analysis of clinical trials and evidence-based treatment algorithm in pulmonary hypertension. *J Am Coll Cardiol* 2004;16, S81-88.
41. Gardiner SM, March JE, Kemp PA, Mullins JJ, Bennett T. Hemodynamic effects of losartan and the endothelin antagonists, SB 209670, in conscious, transgenic ((mRen-2)27), hypertensive rats. *Br J Pharmacol* 1995;116,2237-2244.
42. Glass PS, Gan TJ, Howell S. A review of the pharmacokinetics and pharmacodynamics of remifentanil. *Anesth Analg* 1999;89,S7-14.
43. Goa KL, Wagstaff AJ. Losartan potassium: a review of its pharmacology, clinical efficacy and tolerability in the management of hypertension. *Drugs* 1996;51,820-845.
44. Goetz KL, Wang BC, Sundet WD. Comparative effects of cardiac receptors and sinoaortic baroreceptors on elevations of plasma vasopressin and renin activity elicited by haemorrhage. *J Physiol (Paris)* 1984;79,440-445.
45. Goto T, Nakata Y, Morita S. The minimum alveolar concentration of xenon in the elderly is sex-dependent. *Anesthesiology* 2002;97,1129-1132.
46. Gruss M, Bushell TJ, Bright DP, Lieb WR, Mathie A, Franks NP. Two-pore-domain K⁺ channels are a novel target for the anesthetic gases xenon, nitrous oxide, and cyclopropane. *Mol Pharmacol* 2004;65,443-452.
47. Hettrick DA, Pagel PS, Kersten JR, Tessmer JP, Bosnjak ZJ, Georgieff M, Warltier DC. Cardiovascular effects of xenon in isoflurane-anesthetized dogs with dilated cardiomyopathy. *Anesthesiology* 1998;89,1166-1173.
48. Hintze TH, Vatner SF. Cardiac dynamics during hemorrhage. Relative unimportance of adrenergic inotropic responses. *Circ Res* 1982;50,705-713.
49. Hoffman A, Abassi ZA, Brodsky S, Ramadan R, Winaver J. Mechanisms of big

- endothelin-1-induced diuresis and natriuresis: role of ET(B) receptors. Hypertension 2000;35,732-739.
50. Hoke JF, Cunningham F, James MK, Muir KT, Hoffman WE. Comparative pharmacokinetics and pharmacodynamics of remifentanil, its principle metabolite (GR90291) and alfentanil in dogs. J Pharmacol Exp Ther 1997;281,226-232.
51. Hosokawa A, Nagayama T, Yoshida M, Suzuki-Kusaba M, Hisa H, Kimura T, Satoh S. Facilitation and inhibition by endothelin-1 of adrenal catecholamine secretion in anesthetized dogs. Eur J Pharmacol 2000;397,55-61.
52. Hüneke R, Faßl J, Rossaint R, Lückhoff A. Effects of volatile anesthetics on cardiac ion channels. Acta Anaesthesiol Scand 2004;48,547-561.
53. Inoue A, Yanagisawa M, Kimura S, Kasuya Y, Miyauchi T, Goto K, Masaki T. The human endothelin family: three structurally and pharmacologically distinct isopeptides predicted by three separate genes. Proc Natl Acad Sci U S A 1989;86,2863-2867.
54. Ishiguro Y, Goto T, Nakata Y, Terui K, Niimi Y, Morita S. Effect of xenon on autonomic cardiovascular control- comparison with isoflurane and nitrous oxide. J Clin Anesth 2000;12,196-201.
55. Jacobsson J, Heidvall M, Davidson S. The sevoflurane-sparing effect of nitrous oxide:a clinical study. Acta Anaesthesiol Scand 1999;43,411-414.
56. Jezova M, Armando I, Breganzio C, Yu ZX, Qian S, Ferrans VJ, Imboden H, Saavedra JM. Angiotensin II AT(1) and AT(2) receptors contribute to maintain basal adrenomedullary norepinephrine synthesis and tyrosine hydroxylase transcription. Endocrinology 2003;144:2092-101.
57. Jhaveri R, Joshi P, Batenhorst R, Baughman V, Glass PS. Dose comparison of remifentanil and alfentanil for loss of consciousness. Anesthesiology 1997;87,253-259.
58. Jochem J, Zwirska-Korczala K, Gwozdz B, Walichiewicz P, Josko J. Cardiac and regional haemodynamic effects of endothelin-1 in rats subjected to critical haemorrhagic hypotension. J Physiol Pharmacol 2003;54,383-396.
59. Kaczmarczyk G, Christe W, Mohnhaupt R, Reinhardt HW. An attempt to quantitate the contribution of antidiuretic hormone to the diuresis of left atrial distension in conscious dogs. Pflügers Arch 1983;396,101-105.
60. Kaczmarczyk G, Vogel S, Krebs MO, Bünger H, Scholz A. Vasopressin and renin-angiotensin maintain arterial pessure during PEEP in non-expanded, conscious

- dogs. Am J Physiol 1996;271,R1396-1402.
61. Kaschina E, Unger T. Angiotensin AT1/AT2 receptors: regulation, signalling and function. Blood Press 2003;12,70-88.
62. Kataya J, Viinamäki O, Punnonen R, Kaukinen S. Renin-angiotensin-aldosterone system and plasma vasopressin in surgical patients anaesthetized with halothane or isoflurane. Eur J Anaesth 1988;5,121-129.
63. Kawamura R, Stanley TH, English JB, Hill GE, Liu WS, Webster LR. Cardiovascular response to nitrous oxide exposure for two hours in man. Anesth Analg 1980;59,93-99.
64. Kawase M, Komatsu T, Nishiwaki K, Kimura T, Fujiwara Y, Takahashi T, Shimada Y. Heart rate variability during massive hemorrhage and progressive hemorrhagic shock in dogs. Can J Anesth 2000;47,807-814.
65. Kemmotsu O, Hashimoto Y, Shimosato S. Inotropic effects of isoflurane on mechanics of contraction in isolated cat papillary muscles from normal and failing hearts. Anesthesiology 1973;39,470-477.
66. Kienbaum P, Heuter T, Michel MC, Peters J. Racemic ketamine decreases muscle sympathetic activity but maintains the neural response to hypotensive challenges in humans. Anesthesiology 2000;92,94-101.
67. Kim S, Iwao H. Molecular and cellular mechanisms of angiotensin II-mediated cardiovascular and renal diseases. Pharmacol Rev 2000;52,11-34.
68. Kimura T, Suzuki Y, Yoneda H, Suzuki-Kusaba M, Satoh S. Facilitatory role of the renin-angiotensin system in controlling adrenal catecholamine release in hemorrhaged dogs. J Cardiovasc Pharmacol 1992;19,975-981.
69. Kirchheim HR, Ehmke H, Hackenthal E, Lowe W, Persson P. Autoregulation of renal blood flow, glomerular filtration rate and renin release in conscious dogs. Pflügers Arch 1987;410,441-449.
70. Kotrly KJ, Ebert TJ, Vucins E, Igler FO, Barney JA, Kampine JP. Baroreceptor reflex control of heart rate during isoflurane anesthesia in humans. Anesthesiology 1984;60, 173-179.
71. Koyama S, Fujita T, Shibamoto T, Matsuda Y, Hayashi T Jr, Saeki Y, Kawamoto M, Yamaguchi Y. Relative contribution of renal nerve and adrenal gland to renal vascular tone during prolonged canine hemorrhagic hypotension. Circ Shock 1993;39,269-274.
72. Krebs MO, Kröhn T, Boemke W, Mohnhaupt R, Kaczmarczyk G. Renal and

- hemodynamic effects of losartan in conscious dogs during controlled mechanical ventilation. Am J Physiol 1999;276:F425-432.
73. Krum H, Viskoper RJ, Lacourciere Y, Budde M, Charlton V. The effect of an endothelin-receptor antagonist, bosentan, on blood pressure in patients with essential hypertension. Bosentan Hypertension Investigators. N Engl J Med 1998;338:784-790.
74. Krum H, Liew D. Current status of endothelin blockade for the treatment of cardiovascular and pulmonary vascular disease. Curr Opin Investig Drugs 2003;4:298-302.
75. Lang E, Kapila A, Shlugman D, Hoke JF, Sebel PS, Glass PSA. Reduction of isoflurane minimum alveolar concentration by remifentanil. Anesthesiology 1996;85:721-728.
76. Lee WJ, Yang EK, Ahn DK, Park YY, Park JS, Kim HJ. Central ANGII-receptor antagonists impair cardiovascular and vasopressin response to hemorrhage in rats. Am J Physiol 1995;268,R1500-1506.
77. Leskinen H, Rskoaho H, Huttunen P, Leppäluoto J, Voulteneaho O. Hemorrhage effects on plasma ANP, NH₂-terminal pro-ANP, and pressor hormones in anesthetized and conscious rats. Am J Physiol 1994;266,R1933-1943.
78. Lutropp HH, Romner B, Perhag L, Eskilsson J, Fredriksen S, Werner O. Left ventricular performance and cerebral hemodynamics during xenon anaesthesia. A transesophageal cardiography and transcranial doppler sonography study. Anaesthesia 1993;48:1045-1049.
79. Lynch III C. Myocardial excitation-contraction coupling. In: Yaksh TL, Lynch III C, Zapol WM et al. (eds): Anesthesia: Biologic Foundations. Philadelphia, Lippincott-Raven. 1997,1041-1079.
80. Malek AM, Greene, AL, Izumo S. Regulation of endothelin-1 gene by fluid shear stress is transcriptionally mediated and independent of protein kinase C and cAMP. Cell Biol 1993;90,5999-6003.
81. Malhotra V, Diwan S. Anesthesia and the renal and genitourinary system. In: Miller RD, Anesthesia. 5th edition, Churchill, Livingstone, 2000,1934-1959.
82. Malpas SC, Evans RG, Head GA, Lukoshkova EV. Contribution of renal nerves to renal blood flow variability during hemorrhage. Am J Physiol 1998;274,R1283-1294.
83. Marx T, Froeba G, Wagner D, Baeder S, Goertz A, Georgieff M. Effects on

- hemodynamics and catecholamine release of xenon anaesthesia compared with total iv anaesthesia in the pig. *Brit J Anaesth* 1997;78,326-327.
84. Massart PE, Hodeige DG, Van Mechelen H, Charlier AA, Ketelslegers JM, Heyndrickx GR, Donckier JE. Angiotensin II and endothelin-1 receptor antagonists have cumulative hypotensive effects in canine hypertension. *J Hypertens* 1998;16,835-841.
85. Merin RG, Bernard JM, Doursout MF, Cohen M, Celly JE. Comparison of the effects of isoflurane and desflurane on cardiovascular dynamics and regional blood flow in the chronically instrumented dog. *Anaesthesiology* 1991;74,568-574.
86. Michailov ML, Schad H, Dahlheim H, Jacob ICM, Brechtelsbauer H. Renin-angiotensin system responses of acute graded hemorrhage in dogs. *Circ Shock* 1987;21,217-224.
87. Michelsen LG, Salmenpera M, Hug CC Jr, Szlam F, VanderMeer D. Anesthetic potency of remifentanil in dogs. *Anesthesiology* 1996;84,865-872.
88. Morganti A, Marana I, Airoldi F, Alberti C, Nador B, Palatresi S. Plasma endothelin levels: a meaningless number? *J Cardiovasc Pharmacol* 2000;35,S21-23.
89. Morita H, Vatner SF. Effects of hemorrhage on renal nerve activity in conscious dogs. *Circ Res* 1985;57,788-793.
90. Myre K, Raeder J, Rostrup M, Buanes T, Stokland O. Catecholamine release during laparoscopic fundoplication with high and low doses of remifentanil. *Acta Anaesthesiol Scand* 2003;47,267-273.
91. Nagayama T, Kuwakubo F, Matsumoto T, Fukushima Y, Yoshida M, Suzuki-Kusaba M, Hisa H, Matsumura Y, Kimura T, Satoh S. Role of endogenous endothelins in catecholamine secretion in the rat adrenal gland. *Eur J Pharmacol* 2000;406,69-74.
92. Nakamoto H, Suzuki H, Murakami M, Kageyama Y, Naitoh M, Sakamaki Y, Ohishi A, Saruta T. Different effects of low and high doses of endothelin on haemodynamics and hormones in the normotensive conscious dog. *J Hypertens* 1991;9,337-344.
93. Nakamura K, Okada S, Yokotani K. Endothelin ET_A- and ET_B-receptor-mediated inhibition of noradrenaline release from isolated rat stomach. *J Pharmacol Sci* 2003;91,34-40.
94. Nakata Y, Goto T, Morita S. Effects of xenon on hemodynamic responses to skin

- incision in humans. *Anesthesiology* 1999;90,406-410.
95. Nelson LD, Osborn JL. Neurogenic control of renal function in response to graded nonhypotensive hemorrhage in conscious dogs. *Am J Physiol* 1993;264,R661-667.
96. Notarius CF, Erice F, Stewart D, Magder S. Effect of baroreceptor activation and systemic hypotension on plasma endothelin-1 and neuropeptide Y. *Can J Physiol Pharmacol* 1995;73,1136-1143.
97. Öberg B, Thoren P. Increased activity in left ventricular receptors during hemorrhage or occlusion of caval veins in the cat. A possible cause of the vaso-vagal reaction. *Acta Physiol Scand* 1972;85,164-173.
98. Opgenorth TJ, Adler AL, Calzadilla SV, Chiou WJ, Dayton BD, Dixon DB, Gehrke LJ, Hernandez L, Magnuson SR, Marsh KC, Novosad EI, Von Geldern TW, Wessale JL, Winn M, Wu-Wong JR. Pharmacological characterization of A-127722: an orally active and highly potent ETA-selective receptor antagonist. *J Pharmacol Exp Ther* 1996;276,473-481.
99. Pagel PS, Warltier DC. Anesthetics and left ventricular function. In: Warltier DC (ed): *Ventricular function*. Baltimore, Williams and Wilkins. 1995,213-252.
100. Palacios B, Lim SL, Pang CCY. Role of endothelin ET_A- and ET_B- receptors in haemodynamic compensation following haemorrhage in anaesthetized rats. *Br J Pharmacol* 2002;135,876-882.
101. Picker O, Scheeren TWL, Arndt JO. Inhalation anaesthetics increase heart rate by decreasing cardiac vagal activity in dogs. *Br J Anaesth* 2001;87,748-754.
102. Picker O, Schindler AW, Schwarte LA, Preckel B, Schlack W, Scheeren TWL, Thämer V. Xenon increases total body oxygen consumption during isoflurane anaesthesia in dogs. *Br J Anaesth* 2002;88,546-554.
103. Pollock DM, Polakowski JS, Wegner CD, Opgenorth TJ. Beneficial effect of ETA receptor blockade in a rat model of radiocontrast-induced nephropathy. *Ren Fail* 1997;19:753-761.
104. Pollock DM, Derebail VK, Yamamoto T, Pollock JS. Combined effects of AT(1) and ET(A) receptor antagonists, candesartan, and A-127722 in DOCA-salt hypertensive rats. *Gen Pharmacol* 2000;34,337-342.
105. Pollock DM. Endothelin antagonists in the treatment of renal failure. *Curr Opin Investig Drugs* 2001;2:513-520.
106. Preckel B, Ebel D, Mullenheim J, Frassdorf J, Thamer V, Schlack W. The

- direct myocardial effects of xenon in the dog heart in vivo. *Anesth Analg* 2002;94:545-551.
107. Riggleman A, Harvey J, Baylis C. Endothelin mediates some of the renal actions of acutely administered angiotensin II. *Hypertension* 2001;38,105-109.
 108. Rorie DK, Tyce GM, Sill JC. Increased norepinephrine release from dogs pulmonary artery caused by nitrous oxide. *Anesth Analg* 1986;65,560-564.
 109. Rothe CF, Drees JA. Vascular capacitance and fluid shifts in dogs during prolonged hemorrhagic hypotension. *Circ Res* 1976;38,347-356.
 110. Rossaint R, Reyle-Hahn M, Schulte am Esch J, Scholz J, Scherpereel P, Vallet B, Giunta F, Del Turco M, Erdmann W, Tenbrinck R, Hammerle AF, Nagele P. Multicenter randomized comparison of the efficacy and safety of xenon and isoflurane in patients undergoing elective surgery. *Anesthesiology* 2003;98,6-13.
 111. Rossi GP, Albertin G, Neri G, Andreis PG, Hofmann S, Pessina AC, Nussdorfer GG. Endothelin-1 stimulates steroid secretion of human adrenocortical cells ex vivo via both ETA and ETB receptor subtypes. *J Clin Endocrinol Metab* 1997;82:3445-3449.
 112. Ryckwaert F, Colson P, Andre E, Perrigault PF, Guillon G, Barberis C. Haemodynamic effects of an angiotensin-converting enzyme inhibitor and angiotensin receptor antagonist during hypovolaemia in the anaesthetized pig. *Br J Anaesth* 2002;89,599-604.
 113. Sander-Jensen K, Mehlsen J, Stadeager C, Christensen NJ, Fahrenkrug J, Schwartz TW, Warberg J, Bie P. Increase in vagal activity during hypotensive lower-body negative pressure in humans. *Am J Physiol* 1988;255,R149-156.
 114. Schadt JC, Gaddis RR. Endogenous opiate peptides may limit norepinephrine release during hemorrhage. *J Pharmacol Exp Ther* 1985;232,656-660.
 115. Schadt JC, Gaddis RR. Role of adrenal medulla in hemodynamic response to hemorrhage and naloxone. *Am J Physiol* 1988;254,R559-565.
 116. Schadt JC, Gaddis RR. Renin-angiotensin system and opioids during acute hemorrhage in conscious rabbits. *Am J Physiol* 1990;258,R543-551.
 117. Schadt JC, Ludbrook J. Hemodynamic and neurohumoral responses to acute hypovolemia in conscious mammals. *Am J Physiol* 1991;260,H305-318.
 118. Schaffer JP. Modified sequentially rejective multiple test procedures. *J Am Statist Assoc* 1986;81,826-831.
 119. Schalcher C, Cotter G, Reisin L, Bertel O, Kobrin I, Guyene TT, Kiowski W.

- The dual endothelin receptor antagonist tezosentan acutely improves hemodynamic parameters in patients with advanced heart failure. Am Heart J 2001;142:340-9.
120. Scheeren TWL, Schwarte LA, Arndt JO. Metabolic regulation of cardiac output during inhalation anaesthesia in dogs. Acta Anaesthesiol Scand 1999;43,421-430.
 121. Schirmer U. Lachgas. Anästhesist 1998;47,245-255.
 122. Schmidt M, Marx T, Kotzerke J, Lüderwald S, Armbruster S, Topalidis P, Schirmer U, Reinelt H. Cerebral and regional organ perfusion in pigs during xenon anaesthesia. Anaesthesia 2001;56,1154-1159.
 123. Scholz H, Kramer BK, Hamann M, Gotz KH, Kurtz A. Effects of endothelins on renin secretion from rat kidneys. Acta Physiol Scand 1995;155,173-182.
 124. Seagard JL, Elegbe EO, Hopp FA, Bosnjak ZJ, von Colditz JH, Kalbfleisch JH, Kampine JP. Effects of isoflurane on baroreceptorreflex. Anesthesiology 1983;59,511-520.
 125. Shen YT, Knight DR, Thomas JX Jr, Vatner SF. Relative roles of cardiac receptors and arterial baroreceptors during hemorrhage in conscious dogs. Circ Res 1990;66,397-405.
 126. Shiraishi Y, Fujimura S, Handa M, Kimura T, Ota K, Nakada T. Vasopressin and atrial natriuretic peptide release in cardiopulmonary denervated dogs. Am J Physiol 1990;258,R704-710.
 127. Sitbon O, Badesch DB, Channick RN, Frost A, Robbins IM, Simonneau G, Tapson VF, Rubin LJ. Effects of the dual endothelin receptor antagonist bosentan in patients with pulmonary arterial hypertension: a 1-year follow-up study. Chest 2003;124,247-254.
 128. Sousa de SL, Dickinson R, Lieb WR, Franks NP. Contrasting synaptic action of the inhalational general anesthetics isoflurane and xenon. Anesthesiology 2000;92,1055-1066.
 129. Spielman WS, Zatzman ML, Freeman RH, Davis OJ, Johnson JA. Arterial pressure regulation during hemorrhage: homeostatic role of angiotensin II. Proc Soc Exp Biol Med 1975;149,19-22.
 130. Stearns RA, Chakravarty PK, Chen R, Chiu SHL. Biotransformation of losartan to its active carboxylic acid metabolite in human liver microsomes: role of cytochrome P4502C and 3A subfamily members. Drug Metab Dispos 1995;23,207-215.

131. Stowe DF, Rehmert GC, Kwok WM, Weigt HU, Georgieff M, Bosnjak ZJ. Xenon does not alter cardiac function or major cation currents in isolated guinea pig hearts or myocytes. *Anesthesiology* 2000;92,516-520.
132. Stulak JM, Juncos LA, Haas JA, Romero JC. Systemic hemodynamics and renal function in hemorrhaged dogs resuscitated with cross-linked hemoglobin. *Am J Physiol* 2000;278,R28-33.
133. Teerlink JR, Carteaux JP, Sprecher U, Löffler BM, Clozel M, Clozel JP. Role of endogenous endothelin in normal hemodynamic status of anesthetized dogs. *Am J Physiol* 1995;268,H432-440.
134. Thrasher TN, Keil LC. Arterial baroreceptors control blood pressure and vasopressin responses to hemorrhage in conscious dogs. *Am J Physiol* 1998;275:R1843-57.
135. Thrasher TN, Keil LC. Systolic pressure predicts plasma vasopressin response to hemorrhage and vena caval constriction in dogs. *Am J Physiol* 2000;279,R1035-1042.
136. Timmermans PB, Wong PC, Chiu AT, Herblin WF, Benfield P, Carini DJ, Lee RJ, Wexler RR, Saye JA, Smith RD. Angiotensin II receptors and angiotensin II receptor antagonists. *Pharmacol Rev* 1993;45,205-251.
137. Torre-Amione G, Young JB, Colucci WS, Lewis BS, Pratt C, Cotter G, Stangl K, Elkayam U, Teerlink JR, Frey A, Rainisio M, Kobrin I. Hemodynamic and clinical effects of tezosentan, an intravenous dual endothelin receptor antagonist, in patients hospitalized for acute decompensated heart failure. *J Am Coll Cardiol* 2003;42,140-147.
138. Tournadre JP, Chassard D, Muchada R. Overestimation of low cardiac output measured by thermodilution. *Br J Anaesth* 1997;79,514-516.
139. Treeck B, Roald AB, Tenstad O, Aukland K. Effect of exogenous and endogenous angiotensin II on intrarenal distribution of glomerular filtration rate in rats. *J Physiol* 2002;541,1049-1057.
140. Ullman J. Vasopressin and angiotensin II in blood pressure control during isoflurane anesthesia in rats. *Acta Anaesthesiol Scand* 1999;43,860-865.
141. Ullman J. Influence of neurohumoral blockade on heart rate and blood pressure responses to haemorrhage in isoflurane anaesthetized rats. *Acta Physiol Scand* 2000;169,189-194.
142. Ullman J, Eriksson S, Rundgren M. Losartan increases renal blood flow during

- isoflurane anaesthesia in sheep. *Acta Anaesthesiol Scand* 2001;45,1168-1175.
143. Ullman J, Eriksson S, Rundgren M. Effects of losartan, prazosin and a vasopressin V₁ – receptor antagonist on renal and femoral blood flow in conscious sheep. *Acta Physiol Scand* 2001a;171,99-104.
144. Ünlügenc H, Itegin M, Öcal I, Özalevli M, Güler T, Isik G. Remifentanil produces vasorelaxation in isolated rat thoracic aorta strips. *Acta Anaesthesiol Scand* 2003;47,65-69.
145. Van der Linden P, Rausin I, Deltell A, Bekrar Y, Gilbart E, Bakker J, Vincent JL. Detection of tissue hypoxia by arteriovenous gradient for PCO₂ and pH in anesthetized dogs during progressive hemorrhage. *Anesth Analg* 1995;80,269-275.
146. Vatner SF. Effects of hemorrhage on regional blood flow distribution in dogs and primates. *J Clin Invest* 1974;54,225-235.
147. Verhaar MC, Strachan FE, Newby DE, Cruden NL, Koomans HA, Rabelink TJ, Webb DJ. Endothelin-A receptor antagonist-mediated vasodilatation is attenuated by inhibition of nitric oxide synthesis and by endothelin-B receptor blockade. *Circulation* 1998;97,752-756.
148. Verhaar MC, Grahn AY, Van Weerd AW, Honing ML, Morrison PJ, Yang YP, Padley RJ, Rabelink TJ. Pharmacokinetics and pharmacodynamic effects of ABT-627, an oral ETA selective endothelin antagonist, in humans. *Br J Clin Pharmacol* 2000;49,562-573.
149. Victor RG, Thoren P, Morgan DA, Mark AL. Differential control of adrenal and renal sympathetic nerve activity during hemorrhagic hypotension in rats. *Circ Res* 1989;64,686-694.
150. Wall BM, Huch KM, Runyan KR, Williams HH, Gavras H, Cooke CR. Effects of vasopressin V₁-receptor blockade during acute and sustained hypovolemic hypotension. *Am J Physiol* 1997;270,R356-364.
151. Wang BC, Sundet WD, Hakumaki MO, Goetz KL. Vasopressin and renin responses to hemorrhage in conscious, cardiac-denervated dogs. *Am J Physiol* 1983;245,H399-405.
152. Wang BC, Flora-Ginter G, Leadley RJ, Goetz KL. Ventricular receptors stimulate vasopressin release during hemorrhage. *Am J Physiol* 1988;254,R204-211.
153. Webb DJ, Strachan FE. Clinical experience with endothelin antagonists. *Am J*

Hypertens 1998;11,S71-79.

154. Weiskopf RB, Townsley MI, Riordan KK, Chadwick K, Baysinger M, Mahoney E. Comparison of cardiopulmonary responses to graded hemorrhage during enflurane, halothane, isoflurane, and ketamine anesthesia. *Anesth Analg* 1981;60,481-491.
155. Wildsmith JA, Marshall RL. Positive end-expiratory pressure. Immediate haemodynamic effects during artificial ventilation. *Anaesthesia* 1978;33,20-24.
156. Wong PC, Price WA Jr, Chiu AT, Carini DJ, Duncia JV, Johnson AL, Wexler RR, Timmermans PB. Nonpeptide angiotensin II receptor antagonists. Studies with EXP9270 and DuP 753. *Hypertension* 1990;15,823-834.
157. Wong PC, Hart SD, Duncia JV, Timmermans PB. Nonpeptide angiotensin II receptor antagonists. Studies with DuP 753 and EXP3174 in dogs. *Eur J Pharmacol* 1991;202,323-330.
158. Wouters PF, Van de Velde MA, Marcus MA, Deruyter HA, Van Aken H. Hemodynamic changes during induction of anesthesia with eltanalone and propofol in dogs. *Anesth Analg* 1995;81,125-131.
159. Yamaguchi N. Role of ET_A and ET_B receptors in endothelin-1-induced adrenal catecholamin secretion in vivo. *Am J Physiol* 1997;272,R1290-1297.
160. Yanagisawa M, Kurihara H, Kimura S, Tomobe Y. A novel potent vasoconstrictor peptide produced by vascular endothelial cells. *Nature* 1988;332,411-415.
161. Zhang P, Ohara A, Mashimo T, Imanaka H, Uchiyama A, Yoshiya I. Pulmonary resistance in dogs: a comparison of xenon with nitrous oxide. *Can J Anaesth* 1995;42,547-553.
162. Zimmerman RS, Maymind M, Barbee RW. Endothelin blockade lowers total peripheral resistance in hemorrhagic shock recovery. *Hypertension* 1994;23,205-210.