

6 Literaturverzeichnis

- Abbruscato, T.J. und Davis, T.P. (1999). Protein expression of brain endothelial cell E-cadherin after hypoxia/aglycema: influence of astrocyte contact. *Brain Res.* **842**, 277-86.
- Abdul-Khalil, H., Schubert, S., Fischer, T., Böttcher, W., Harke, C., Alexi-Meskishvili, V., Lange, P. und Blasig, I.E. (2000). The effect of continuous treatment of sodium nitroprusside on the serum kinetics of the brain marker protein S-100 β in neonates undergoing corrective cardiac surgery by means of hypothermic cardiopulmonary bypass. *Clin. Chem. Lab. Med.* **38**, 1173-1175.
- Anafi, M., Rosen, M.K., Gish, G.D., Kay, L.E. und Pawson, T. (1996). A potential SH3 domain-binding site in the Crk SH2 domain. *J. Biol. Chem.* **271**, 21365-74.
- Anderson, J.M., Stevenson, B.R., Jesaitis, L.A., Goodenough, D.A. und Mooseker, M.S. (1988). Characterization of ZO-1, a protein component of the tight junction from mouse liver and Madin-Darby canine kidney cells. *J. Cell. Biol.* **106**, 1141-9.
- Anderson, J.M. und van Itallie, C.M. (1995). Tight junctions and the molecular basis for regulation of paracellular permeability. *Am. J. Physiol.* **269**, G467-75.
- Anderson, J.M. (2001). Molecular structure of tight junctions and their role in epithelial transport. *News Physiol. Sci.* **16**, 126-30.
- Ando-Akatsuka, Y. et al. (1996). Interspecies diversity of the occludin sequence: cDNA cloning of human, mouse, dog and rat-kangaroo homologues. *J. Cell Biol.* **133**, 43-47.
- Andreeva, A.Y., Krause, E., Müller, E.-C., Blasig, I.E. und Utepbergenov, D.I. (2001). Protein kinase C regulates the phosphorylation and cellular distribution of occludin. *J. Biol. Chem.* **276**, 38480-86.
- Andreotti, A.H., Bunnell, S.C., Feng, S., Berg, L.J. und Schreiber, S.L. (1997). Regulatory intramolecular association in a tyrosine kinase of the Tec family. *Nature* **385**, 93-97.
- Alberts, B., Bray, D., Lewis, J., Raff, M., Roberts, K., Watson, J.D. (1996). Molekularbiologie der Zelle. VCH Verlagsgesellschaft mbH, Weinheim., 1125.
- Allt, G. und Lawrenson, J.G. (1997). Is the pial microvessel a good model for blood-brain barrier studies. *Brain Res. Rev.* **24**, 67-76.
- Altschuh, D., Dubs, M.C., Weiss, E., Zeder-Lutz, G. und Van Regenmortel, M.H. (1992). Determination of kinetic constants for the interaction between a monoclonal antibody and peptides using surface plasmon resonance. *Biochemistry* **27**, 6298-304.
- Atsumi, S., Kokai, Y und Mori, M. (1999). Occludin modulates organization of perijunctional circumferential actin in rat endothelial cells. *Medical electron microscopy* **32**, 11-19.
- Aurand-Lions, M.A., Duncan, L., Du Pasquier, L. und Imhof, B.A. (2000). Cloning of JAM-2 and JAM-3: an emerging junctional adhesion molecular family? *Curr. Top. Microbiol. Immunol.* **251**, 91-8.
- Balda, M.S., Gonzales-Mariscal, L., Macias-Silva, M.E., Torres-Marquez, J.A., Garcia, S. und Cereijido, C. (1991). Assembly and sealing of tight junctions: possible participation of G-proteins, phospholipase C, protein kinase C and calmodulin. *J. Membr. Biol.* **122**, 193-202.
- Balda, M.S., Gonzales-Mariscal, L., Matter, K., Cereijido, M. und Anderson, J.M. (1993). Assembly of the tight junction: the role of diacylglycerol. *J Cell Biol* **123**, 293-302.
- Balda, M.S. et al. (1996a). Functional dissociation of paracellular permeability and transepithelial resistance and disruption of the apical-basolateral intramembrane diffusion barrier by expression of a mutant tight junction protein. *J. Cell Biol.* **134**, 1031-49.

- Balda, M.S.; Anderson, J.M. und Matter, K. (1996b). The SH3 domain of the tight junction protein ZO-1 binds to serine protein kinase that phosphorylates a region C-terminal of this domain. *FEBS Lett.* **399**, 326-32.
- Balda, M.S. und Matter, K. (1998). Tight junctions. *J. Cell Sci.* **111**, 541-47.
- Balda, M.S. und Matter, K. (2000a). Transmembrane proteins of tight junctions. *Cell Dev. Biol.* **11**, 281-89.
- Balda, M.S. und Matter, K. (2000b). The tight junction protein ZO-1 and an interacting transcription factor regulate ErbB-2 expression. *EMBO J.* **19**, 2024-33.
- Balda, M.S., Flores-Maldonado, C., Cereijido, M. und Matter, K. (2000c). Multiple domains of occludin are involved in the regulation of paracellular permeability. *J. Cell. Biochem.* **78**, 85-96.
- Balgi, G., Leckband, D.E. und Nitsche, J.M. (1995). Transport effects on the kinetics of protein-surface binding. *Biophys. J.* **68**, 2251-60.
- Bamforth, S.D., Kniesel, U., Wolburg, H., Engelhardt, B. und Risau, W. (1999). A dominant mutant of occludin disrupts tight junction structure and function. *J. Cell. Sci.* **112**, 1879-88.
- Barton, E.S., Forrest, J.C., Connolly, J.L., Lin, Y., Schnell, F.J., Nusrat, A., Parkos, C.A. und Dermody, T.S. (2001). Junction adhesion molecule is a receptor for reovirus. *Cell* **104**, 441-51.
- Bauer, H. et al. (1999). Astrocytes and neurons express the tight junction-specific protein occludin in vitro. *Exp. Cell Res.* **250**, 434-38.
- Bauer, H.-C. und Bauer, H. (2000). Neuronal induction of the blood-brain barrier: still an enigma. *Cell. Mol. Neurobiol.* **20**, 13-28.
- Bazzoni, G. et al. (2000a). Homophilic interaction of junctional adhesion molecule. *J. Biol. Chem.* **275**, 30970-30976.
- Bazzoni, G., Martinez-Estrada, O.M., Orsenigo, F., Cordenonsi, M., Citi, S. und Dejana, E. (2000b). Interaction of junctional adhesion molecule with the tight junction components ZO-1, cingulin and occludin. *J. Biol. Chem.* **275**, 20520-20526.
- Beckmann, J.S. und Koppenol, W.H. (1996). Nitric oxide, superoxide, and peroxynitrite: the good, the bad, and the ugly. *Am. J. Physiol.* **271**, C1424-37.
- Benveniste, E.N. (1998). Cytokine actions in the central nervous system. *Cytokine & growth factor reviews* **9**, 259-75.
- Betz, A.L. (1993). Oxygen free radicals and the brain microvasculature. In: *The blood-brain barrier* (W.M. Padridge, Ed.). Raven Press, New York.
- BIAjournal (1995). An introduction to BIATechnology. *BIAjournal* **2**, 8-9
- BIATechnology Handbook (1994). Pharmacia Biosensor AB, Uppsala, Sweden, 3-4, 4-3.
- Blum, M.S. et al. (1997). Cytoskeletal rearrangement mediates human and microvascular endothelial tight junction modulation by cytokines. *Am. J. Physiol.* **273**, H286-94.
- Bradbury, M.W.B. (1985). The blood-brain barrier: transport across the cerebral endothelium. *Circ. Res.* **57**, 213-22.
- Brahms et al. (1980). Determination of protein secondary structure in solution by vacuum ultraviolet circular dichroism. *J. Mol. Biol.* **138**, 138-49.
- Budnik, V. (1996). Synapse maturation and structural plasticity at Drosophila neuromuscular junctions. *Curr. Opin. Neurobiol.* **6**, 958-67.
- Caide, X., Sui, S.F. (1999). Numerical simulations of surface plasmon resonance system for monitoring DNA hybridization and detecting protein-lipid film interactions. *Eur. Biophys. J.* **28**, 151-7.
- Cancilla, P.A., Bready, J. und Berliner, J. (1993). Brain endothelial-astrocyte interactions. In W.M. Padridge(Ed.), *The blood-brain barrier. Cellular and Molecular Biology*. Raven Press New York, 25-46.

- Chen (1972). Determination of the secondary structures of protein by circular dichroism and optical rotatory dispersion. *Biochem.* **11**, 4120-31.
- Chen, Y-H., Merzdorf, C. Paul, D.L. und Goodenough, D.A.(1997). COOH terminus of occludin is required for tight junction barrier function in early *Xenopus* embryos. *JCB* **138**, 891-99.
- Chi, O.Z., Liu, X. und Weiss, H.R. (1999). Effect of cyclic GMP on microvascular permeability of the cerebral cortex. *Microvasc. Res.* **58**, 35-40.
- Choulier, L., Rauffer-Bruyere, N., Ben Khalifa, M., Martin, F., Vernet, T. und Altschuh, D. (1999). Kinetic analysis of the effect on Fab binding of identical substitutions in a peptide and its parents protein. *Biochemistry* **23**, 2530-7.
- Cereijido, M., Gonzales-Mariscal, L. und Contreras, G. (1989). Tight junction: barrier between higher organisms and environment. *News in Physiol. Sci.* **4**, 72.
- Cereijido, M., Shoshani, L. und Contreras, G. (2000). Molecular physiology and pathophysiology of tight junctions. I. Biogenesis of tight junctions and epithelial polarity. *Am. J. Gastrointest. Liver Physiol.* **279**, G477-82.
- Citi, S. Sabanay, H., Jakes, R., Geiger, B. und Kendrick-Jones, J. (1988). Cingulin, a new peripheral component of tight junctions. *Nature* **333**, 272-76.
- Citi, S. (1992). Protein kinase inhibitors prevent junction dissociation induced by low extracellular calcium in MDCK epithelial cells. *J Cell Biol* **117**, 169-78.
- Citi,S. (1993). The molecular organization of tight junctions. *J Cell Biol.* **12**, 485-89.
- Citi, S. (1994). The molecular composition of tight junctions. In Citi, S., ed. *Molecular Mechanisms of Epithelial Cell Junctions: From Development to Disease*. Austin, TX, R.G. Landes, 83-106.
- Citi, S., D'Atri, F. und Parry, D.A. (2000). Human and Xenopus cingulin share a modular organization of Coiled coil rod domain: predictions for intra- and intermolecular assembly. *J Struct Biol* **131**, 135-45.
- Claude, P. und Goodenough, D.A. (1973). Fractures face of zonulae occludentes from „tight“ and „leaky“ epithelia. *J. Cell Biol.* **58**, 490-400.
- Claude, P. (1978). Morphological factors influencing transepithelial permeability. a model for the resistance of the zonula occludens. *J. Membr. Biol.* **10**, 219-32.
- Cohen, A.R., Woods, D.F., Marfatia, S.M., Walther, Z., Chishti, A.H. und Anderson, J.M. (1998). Human CASK/LIN-2 binds syndecan-2 and protein 4.1 and localizes to the basolateral membrane of epithelial cells. *J. Cell. Biol.* **142**, 129-38.
- Collins, J.R. und Rizzolo, L.J. (1998). Protein-binding domains of the tight junction protein, ZO-2, are highly conserved between avian and mammalian species. *Biochem Biophys Res Commun* **252**, 617-22.
- Cordenonsi, M. , Dàtri, F., Hammar, E., Parry, D.A., Kendrick-Jones, J., Shore, D. und Citi, S. (1999). Cingulin contains globular and Coiled coil domains and interact with ZO-1, ZO-2, ZO-3, and myosin. *J. Cell Biol.* **147**, 1569-82.
- Cuzzocrea, S., Mazzon, E., De Sarro, A. und Caputi, A.P. (2000). Role of free radicals and poly(ADP-ribose)synthetase in intestinal tight junction permeability. *Mol. Med.* **6**, 766-78.
- Cuzzocrea, S. und Reiter, R.J. (2001). Pharmacological action of melatonin in shock, inflammation and ischemia/reperfusion injury. *Eur. J. Pharmacol.* **426**, 1-10.
- Dallasta, L.M. et al. (1999). Blood-brain barrier tight junction disruption in human immunodeficiency virus-1 encephalitis. *Am. J. Pathol.* **155**, 1915-27.
- Caide, X. und Sui, S.F.(1999). Numerical simulations of surface plasmon resonance system for monitoring DNA hybridization and detecting protein-lipid film interactions. *Eur. Biophys. J.* **28**, 151-7.

- Deguchi, M., Hata, Y., Takeuchi, M., Ide, N., Hirao, K., Yao, I., Irie, M., Toyoda, A. und Takai, Y. (1998). BEGAIN (brain-enriched guanylate kinase-associated protein), a novel neuronal PSD-95/SAP90-binding protein. *J. Biol. Chem.* **273**, 26269-72.
- Denker, B.M., Saha, S., Khawaja, S. und Nigam, S.K. (1996). Involvement of a heterotrimeric G protein alpha subunit in tight junction biogenesis. *JBC* **271**, 25750-53.
- Denker, B.M. und Nigam, S.K. (1998). Molecular structure and assembly of tight junction. *Am. J. Physiol.* **274**, F1-9.
- Dejana, E., Lampugnani, M.G., Martinez-Estrada, O. und Bazzoni, G. (2000). The molecular organization of endothelial junctions and their functional role in vascular morphogenesis and permeability. *Int J Dev Biol* **44**, 743-8.
- De Vries, H.E., Kuiper, J., De Boer, A.G., Van Berkel, T.J.C. und Breimer, D.D. (1997). The blood-brain barrier in neuroinflammatory disease. *Pharmakol. Reviews* **49**, 143-55.
- Dobrosotskaya, I., Guy, R.K. und James, G.L. (1997). MAGI-1, a membrane-associated guanylate kinase with a unique arrangement of protein-protein interaction domains. *J. Biol. Chem.* **272**, 31589-97.
- Dragsten, P.R., Blumenthal, R. und Handler, J.S. (1981). Membrane assymetryin epithelia: is the tight junction a barrier to diffusion in the plasma membrane? *Nature* **294**, 718-22.
- Draijer, R. et al. (1995). Expression of cGMP-dependent protein kinase 1 and phosphorylation of its substrate, vasodilator-stimulated phosphoprotein, in human endothelial cells of different origin. *Circ. Res.* **77**, 897-905.
- Dunina-Barkovskaya, A. (1998). Tight junctions: facts and models. *Membr. Cell. Biol.* **11**, 555-89.
- Dye, J.F., Leach, L., Clark, P. und Firth, J.A. (2001). Cyclic amp and acidic fibroblast growth factor have opposing effect on tight and adherens junctions in microvasular endothelial cells in vitro. *Microvasc Res* **62**, 94-113.
- Dyson, H.J. und Wright, P.E. (1991). Defining solution conformations of small linear peptides. *Annu. Rev. Biophys. Biophys. Chem.* **20**, 519-38.
- Dyson, H.J. und Wright, P.E. (2002). Coupling of folding andbinding of unstructured proteins. *Curr. Opin. Struct. Biol.* **12**, 54-60.
- Ebnet, K., Schulz C. U., Meyer zu Brickwedde, M.K., Pendle, G.G. und Vestweber, D. (2000). Junctional adhesion molecule interacts with the PDZ domain-containing proteins AF6 and ZO-1. *J. Biol. Chem.* **275**, 27979-88.
- Ehrlich (1902). Über die Beziehung von chemischer Konstitution, Verheilung und pharmakologischer Wirkung. Berlin, Hirschwald.
- Fanning, A.S., Jamson, B.J., Jesaitis, L.A. und Anderson, J.M. (1998). The tight junction protein ZO-1 establishes a link between the transmembrane protein occludin and the actin cytoskeleton. *J. Biol. Chem.* **273**, 29745-29753.
- Fanning, A.S. und Anderson, J.M. (1999).Protein modules as organizer of membrane structure. *Curr. Opin. Cell Biol.* **11**, 432-39.
- Farquhar, M.G. und Palade, G.E. (1963). Junctional complexes in various epithelia. *J.Cell Biol.* **17**, 375-421.
- Federici, C., Camoin, L., Creminon, C., Chaverot, N., Strosberg, A.D. und Couraud, P.O. (1995). Cultured astrocytes release a factor that decreases endothelin-1 secretion by brain microvessel endothelial cells. *J Neurochem.* **64**, 1008-15.
- Fesenko, I., Kurth, T., Sheth, B., Fleming, T.P., Citi, S. und Hausen, P. (2000). Tight junction biogenesis in the early Xenopus embryo. *Mech. Dev.* **96**, 51-65.
- Franke (1992). *Tetrahedron* **48**, 9217.
- Furuse, M. et al. (1993). Occludin: a novel integral membrane protein localizing at tight junctions. *J. Cell Biol.* **123**, 1777-88.

- Furuse, M., Itoh, M., Hirase, T., Nagafuchi, A., Yonemura, S., Tsukita, S. und Tsukita, S. (1994). Direct association of occludin with ZO-1 and its possible involvement in the localisation of occludin at tight junction. *J. Cell Biol.* **127**, 1617-1626.
- Furuse, M., Fujita, K., Hiragi, T., Fujimoto, K. und Tsukita, S. (1998a). Claudin-1 and 2: novel integrale membrane proteins localizing at tight junctions with no sequence similarity to occludin. *J. Cell Biol.* **141**, 1539-50.
- Furuse, M., Sasaki, H., Fujimoto, K. und Tsukita, S. (1998b). A single gene product, claudin-1 or -2, reconstitutes tight junction strands and recruits occludin in fibroblasts. *J. Cell Biol.* **143**, 391-401.
- Furuse, M., Sasaki, H und Tsukita, S. (1999). Manner of interaction of heterogeneous claudin species within and between tight junction strands. *J. Cell Biol.* **147**, 891-903.
- Furuse, M. et al. (2002). Claudin-based tight junctions are crucial for the mammalian epidermal barrier: a lesson from claudin-1-deficient mice. *JCB* **156**, 1099-1111.
- Gerhardt, H., Liebner, S. und Wolburg, H. (2000). Entwicklung und Pathologie der Blut-Hirn-Schranken-Gefäße. *Neuroforum* **4**, 249-56.
- Giepmans, B.N. und Moolenaar, W.H. (1998). The gap junction protein connexin43 interacts with the second PDZ domain of the zona occludens-1 protein. *Curr. Biol.* **8**, 931-4.
- Glaser, R.W. (1993). Antigen-antibody binding and mass transport by convection and diffusion to a surface: a two-dimensional computer model of binding and dissociation kinetics. *Analytical Biochem.* **213**, 152-61.
- Goldmann, E.E. (1909). Die äußere und innere Sekretion des gesunden und kranken Organismus im Lichte der „vitalen Färbung“. *Beitr. Z. Klein. Chir.* **64**, 192-265.
- Goldstein, G.W. (1988). Endothelial cell-astrocyte interactions. A cellular model of the blood-brain barrier. *Ann. N. Y. Acad. Sci.* **529**, 31-9.
- Gonzalez-Mariscal, L. et al. (1999). Molecular characterization of the tight junction protein ZO-1 in MDCK cells. *Exp. Cell Res.* **248**, 97-109.
- Gonzalez-Mariscal, L., Betanzos, A. und Avila-Flores, A. (2000). MAGUK proteins: structure and role in the tight junction. *Cell and Developmental Biol.* **11**, 315-324.
- Grebekämper, K. und Galla, H.-J. (1994). Translational diffusion measurements of a fluorescent phospholipid between MDCK-I cells support the lipid model of the tight junctions. *Chemistry and Physics of lipids* **71**, 133-43.
- Grindstaff, K.K. et al. (1998). Sec6/8 complex is recruited to cell-cell contacts and specific transports vesicle delivery to the basal-lateral membrane in epithelial cells. *Cell* **93**, 731-740.
- Gumbiner, B., Löwenkopf, T. und Apatira, D. (1986). Identifikation of a 160-kDa polypeptide that binds to the tight junction protein ZO-1. *Proc Natl Acad. Sci. USA* **83**, 3460-64.
- Gumbiner, B. (1987). Structure, biochemistry, and assembly of epithelial tight junctions. *Am J Physiol.* **253**, C749-58.
- Gumbiner, B., Lowenkopf, T. und Apatira, D. (1991). Identification of a 160-kDa polypeptide that binds to the tight junction protein ZO-1. *Proc. Natl. Acad. Sci. U.S.A.* **88**, 3460-4.
- Gumbiner, B. (1993). Breaking through the tight junction barrier. *J. Cell Biol.* **123**, 1631-33.
- Gumbiner, B. (1996). Cell adhesion: the molecular basis of tissue architecture and morphogenesis. *Cell* **84**, 345-57.
- Gupta, M.N., Kaul, R., Guoquiang, D., Dissing, U., Mattiasson, B (1996). Affinity precipitation of proteins. *J Mol Recognit* **9**, 356-9.
- Gursoy-Ozdemir, Y., Bolay, H., Saribas, O. und Dalkara, T. (2000). Role of endothelial nitric oxide generation and peroxynitrite formation in reperfusin injury after focal cerebral ischemia. *Stroke* **31**, 1974-81.

- Hamazaki, Y., Itoh, M., Sasaki, H., Furuse, M. und Tsukita, S. (2001). Multi PDZ-containing protein (MUPP1) is concentrated at tight junction through its possible interaction with claudin-1 and junctional adhesion molecule (JAM). *JBC* **277**, 455-61.
- Haseley, S.R., Talaga, P., Kamerling, J.P. und Vliegenthart, J.F. (1999). Characterization of the carbohydrate binding specificity and k-parameters of lectins by using surface plasmon resonance. *Anal Biochem* **274**, 203-10.
- Hashimoto, S. (2000). Principles of BIACORE. In „Real time analysis of biomolecular interactions“, S. 27, ed. Nagata, K. und Handa, H., Springer-Verlag, Tokyo.
- Haskins, J., Gu, L., Wittchen, E.S., Hibbard, J. und Stevenson, B.R. (1998). ZO-3, a novel member of the MAGUK protein family found at the tight junction, interacts with ZO-1 and occludin. *J. Cell Biol.* **141**, 199-208.
- Hata, Y., Nakanishi, H. und Takai, Y. (1998). Synaptic PDZ domain containing proteins. *Neurosci. Res.* **32**, 1-7.
- Hartleib, J. et al. (2000). Protein A is the von Willebrand factor binding protein on *Staphylococcus aureus*. *Blood* **96**, 2149-56.
- Heiskala, M., Peterson, F.A. und Yang, Y. (2001). The roles of Clauudin superfamily proteins in paracellular transport. *Traffic* **2**, 93-8.
- Hirase, T. et al. (1997). Occludin as a possible determinant of tight junction permeability in endothelial cells. *J. Cell. Sci.* **110**, 1603-13.
- Holash, J.A. und Stewart, P.A. (1993a). The relationship of astrocyte-like cells to the vessels that contribute to the blood-ocular barriers. *Brain Res.* **629**, 218-24.
- Holash, J.A., Noden, D.M. und Stewart, P.A. (1993b). Re-evaluating the role of astrocytes in blood-brain barrier induction. *Dev. Dynam.* **197**, 14-25.
- Holtzhauer, M. (1996). Methoden in der Proteinanalytik. Springer-Verlag Berlin, Heidelberg, New York, 235-239.
- Hoover KB, Liao SY, Bryant PJ. (1998). Loss of the tight junction MAGUK ZO-1 in breast cancer: relationship to glandular differentiation and loss of heterozygosity. *Am J Pathol.* **153**, 767-73.
- Hopkins, A.M., Li, D., Mrsny, R.J., Walsh, S.V und Nusrat, A. (2000). Modulation of tight junction function by G protein-coupled events. *Adv Drug Deliv Res* **41**, 329-40.
- Hordijk, P.L. et al (1999). Vascular-endothelial-cadherin modulates endothelial monolayer permeability. *J. Cell Sci.* **112**, 1915-23.
- Hough, C.D., Woods, D.F., Park, S. und Bryant, P.J. (1997). Organizing a functional junctional complex requires specific domains of the Drosophila MAGUK Discs large. *Genes Dev.* **11**, 3242-53.
- Howarth, A.G., Hughes, M.R. und Stevenson, B.R. (1992) Detection of the tight junction-associated protein ZO-1 in astrocytes and other nonepithelial cell types. *Am. J. Physiol.* **262**, C461-9.
- Howarth, A.G. und Stevenson, B.R. (1995) Molecular environment of ZO-1 in epithelial and non-epithelial cells. *Cell. Motil. Cytoskeleton* **31**, 323-32.
- Hsueh, Y.P., Kim, E. und Sheng, M. (1997). Disulfide-linked head-to-head multimerization in the mechanism of ion channel clustering by PSD-95. *Neuron* **18**, 803-14.
- Huber, J.D., Eglington, R.D. und Davis, T.P. (2001). Molecular physiology and pathophysiology of tight junctions in the blood-brain barrier. *Trends Neurosci.* **24**, 719-25.
- Huber, A.H., Stewart, D.B., Laurens, D.V., Nelson, W.J. und Weis, W.I. (2001a). The cadherin cytoplasmic domain is unstructured in the absence of β -catenin-a possible mechanism for regulating cadherin turnover. *JBC* **276**, 12301-9.
- Huber, A.H. und Weis, W.I. (2001b). The structure of β -catenin/E-cadherin complex and the molecular basis of diverse ligand recognition by β -catenin. *Cell* **105**, 391-402.

- Hurst, R.D. und Fritz, I.B. (1996). Nitric oxide-induced perturbations in a cell culture model of the blood-brain barrier. *J. Cell. Physiol.* **167**, 89-94.
- Ide, N. et al. (1999). Localization of membrane-associated guanylate kinase(MAGI)-1/BAI-associated protein (BAP) 1 at tight junctions of epithelial cells. *Oncogene* **18**, 7810-15.
- Itoh, M., Yonemura, S., Nagafuchi, A., Tsukita, S. und Tsukita, S. (1991). A 220-kD undercoat-constitutive protein: its specific localization at cadherin-based cell-cell adhesion sites. *J. Cell. Biol.* **115**, 1449-62.
- Itoh, M. et al. (1993). The 220-kD protein colocalizing with cadherins in nonepithelial cells is identical to ZO-1, a tight junction-associated protein in epithelial cells: cDNA cloning and immunoelectron microscopy. *J. Cell. Biol.* **121**, 491-502.
- Itoh, M., Nagafuchi, A., Moroi, S. und Tsukita, S. (1997). Involvement of ZO-1 in cadherin-based cell adhesion through its direct binding to α -catenin and actin filaments. *J. Cell Biol.* **138**, 181-92.
- Itoh, M. et al. (1999a). Direct binding of three tight junction-associated MAGUKs, ZO-1, ZO-2 and ZO-3, with the COOH termini of claudins. *J. Cell Biol.* **147**, 1351-67.
- Itoh, M., Morita, K. und Tsukita, S. (1999b). Characterization of ZO-2 as a MAGUK family member associated with tight as well as adherence junctions with a binding affinity to occludin and α -catenin. *J. Biol. Chem.* **274**, 5981-5986.
- Izumi, Y. et al. (1998). An atypical PKC directly associates and colocalizes at the epithelial tight junction with ASIP, a mammalian homologue of *Caenorhabditis elegans* polarity protein PAR-3. *J. Cell Biol.* **143**, 95-106.
- Janigro, D., West, G.A., Nguyen, T.S. und Winn, H.R.. (1994). Regulation of blood-brain barrier endothelial cells by nitric oxide. *Circ. Res.* **75**, 528-38.
- Janzer, R.C. und Raff, (1987). Astrocyte induce blood brain barrier proteins in endothelial cells. *Nature* **325**, 253-57.
- Janzer, R.C. (1993). The blood-brain barrier. Cellular basis. *J. Inher. Metab. Dis.* **16**, 639-47.
- Jesaitis, L.A. und Goodenough, D.A. (1994). Molecular characterisation and tissue distribution of ZO-2, a tight junction homologous to ZO-1 and the Drosophila discs-large tumor suppressor protein. *J. Cell. Biol.* **124**, 949-61.
- Jiang, W.G., Bryce, R.P., Horrobin, D.F. und Mansel, R.E. (1998). Regulation of tight junction permeability and occludin expression by polyunsaturated fatty acids. *BBRC* **244**, 414-20.
- Kachar, B. und Reese, T.S. (1982). Evidence for the lipidic nature of tight junction strands. *Nature* **296**, 464-66.
- Karlsson, R., Michaelsson, A. und Mattsson, L. (1991): Kinetic analysis of monoclonal antibody-antigen interactions with a new biosensor based analytical system. *J. Immunol. Methods* **145**, 229-240.
- Katsube, T., Takahisa, M., Ueda, R., Hashimoto, N., Kobayashi, M. und Togashi, S. (1998). Cortactin associates with the cell-cell junction protein ZO-1 in both Drosophila and mouse. *J. Biol. Chem.* **273**, 29672-7.
- Kawabe, H. et al. (2001). PILT: a novel peripheral membrane protein at tight junctions in epithelial cells. *JBC*, **276**, 48350-5.
- Keon, B.H., Schäfer, S., Kuhn, C., Grund, C. und Franke, W. (1996). Symplekin, a novel type of tight junction plaque protein. *J. Cell Biol.* **134**, 1003-18.
- Killian, J.A. und von Heijne, G. (2000). How proteins adapt to a membrane-water interface. *TIBS* **25**, 429-34.
- Kim, E., Niethammer, M., Rothschild, A., Jan, Y.N. und Sheng, M. (1995). Clustering of Shaker-type K⁺ channels by interaction with a family of membrane-associated guanylate kinases. *Nature* **378**, 85-88.

- Kim, E., Naisbitt, S., Hsueh ,Y.P., Rao, A., Rothschild, A., Craig, A.M. und Sheng M. (1997). GKAP, a novel synaptic protein that interacts with the guanylate kinase-like domain of the PSD-95/SAP90 family of channel clustering molecules. *J. Cell. Biol.* **136**, 669-78.
- Kim, A.S., Kakalis, L.T., Abdul-Manan, N., Liu, G.A. und Rosen, M.K. (2000). Autoinhibition and activarion mechanisms of the Wiskott-Aldrich syndrome protein. *Nature* **404**, 151-58.
- Kniesel, U., Risau, W. und Wolburg, H. (1996). Development of blood-brain barrier tight junction in the rat cortex. *Dev. Brain Res.* **96**, 229-40.
- Kniesel, U. und Wolburg, H. (2000). Tight junction of the blood-brain barrier. *Cell Moll. Neurobiol.* **20**, 57-76.
- Kobayashi, J., Inai, T. und Shibata, Y. (2002). Formation of tight junction strands by expression of claudin mutants in their ZO-1 binding site in MDCK cells. *Histochem. Cell. Biol.* **117**, 29-39.
- Kramer, A., Schuster, A., Reinecke, U., Malin, R., Volkmer-Engert, R., Landgraf, C. und Schneider-Mergener, J. (1994). Combinatorial cellulose-bound peptide libraries: screening tool for the identification of peptides that bind ligands with predefined specificity. *Methods (Comp. Methods Enzymol.)* **6**, 388-395.
- Kramer, A. und Schneider-Mergener, J. (1998). Synthesis and screening of peptide libraries on cellulose membrane supports. *Methods Mol. Biol.* **87**, 25-39.
- Kretschmann, E. (1971). Die Bestimmung optischer Konstanten von Metallen durch Anregung von Oberflächenplasmaschwingungen. *Z. Phys.* **B241**, 313-324.
- Kubota, K. et al. (1999). Ca²⁺-independent cell adhesion activity of claudins, integral membrane proteins of tight junctions. *Curr. Biol.* **9**, 1035-38.
- Kuwabara, H., Kokai, Y., Kojima, T., Takakuwa, R., Mori, M. und Sawada, N. (2001). Occludin regulates actin cytoskeleton in endothelial cells. *Cell Struct Funct* **26**, 109-16.
- Lacaz-Vieira, F. (1997). Calcium site specificity. Early Ca²⁺-related tight junction events. *J. Gene Physiol.* **110**, 727-40.
- Lacaz-Vieira, F., Jaeger, M.M., Farshori, P. und Kachar, B. (1999). Small synthetic peptides homologous to segments of the first external loop of occludin impair tight junction resealing. *J Membr Biol.* **168**, 289-97.
- Lacaz-Vieira, F. (2000). Tight junction dynamics: oscillations and the role of protein kinase C. *J Membr Biol* **178**, 151-61.
- Landschulz, W.H., Johnson, P.F. und McKnight, S.L. (1988). The leucine zipper: a hypothetical structure common to a new class of DNA binding proteins. *Science* **240**, 1759-64.
- Larson, S.M. und Davidson, A.R. (2000). The identification of conserved interactions within the SH3 domain by alignment of sequences and structures. *Protein Science* **9**, 2170-80.
- Li, S.-C. und Deber, C.M. (1994). A measure of helical propensity for amino acids in membrane environments. *Struct. Biol.* **1** (6), 368-374.
- Liebner, S., Kniesel, U., Kalbacher, H. und Wolburg, H. (2000). Correlation of tight junction morphology with the expression of tight junction proteins in blood-brain barrier endothelial cells. *Eur. J. Cell Biol.* **79**, 707-17.
- Lippoldt, A., Kniesel, U. et al. (2000). Structural alterations of tight junctions are associated with loss of polarity in stroke-prone spontaneously hypertensive rat blood-brain barrier endothelial cells. *Brain Res.* **885**, 251-61.
- Lottspeich, F. und Zorbas, H. (1998). Bioanalytik. Spektrum. Akademischer Verlag GmbH, Heidelberg. Berlin.
- Lue, R.A.; Marfatia, S.M.; Branton, D. und Chrishti, A.H. (1994). Cloning and characterization of hdlg : the human homologue of the *Drosophila* discs large tumor suppressor binds to protein 4.1. *PNAS USA* **91**, 9818-22.

- Luo, J., Zhou, J., Zou, W. und Shen, P. (1999). Determination of interaction mechanism of sensorgrams by analysis of binding kinetics. *J Protein Chem.* **18**, 709-19.
- Lum, H. und Malik, A.B. (1994). Regulation of vascular endothelial barrier function. *Am. J. Physiol.* **267**, L223-41.
- Lupas, A. (1996). Coiled coils: new structures and new functions. *TIBS* **21**, 375-82.
- Lupas, A. (1997). Predicting Coiled coil regions in proteins. *Curr Opin Struct Biol* **7**, 388-93.
- Ma, T.Y., Tran, D., Hoa, N., Nguyen, D., Merryfield, M. und Tarnawski, A. (2000). Mechanism of extracellular calcium regulation of intestinal epithelial tight junction permeability: role of cytoskeletal involvement. *Microsc. Res. Techn.* **51**, 156-68.
- Madara, J.L. (1998). Regulation of the movement of solutes across tight junctions. *Annu. Rev. Physiol.* **59**, 253, 143-59.
- Malmqvist, M. (1993). Biospecific interaction analysis using biosensor technology. *Nature* **361**, 186-187.
- Marfatia, S.M., Lue, R.A., Branton, D. und Chishti, A.H. (1994). In vitro binding studies suggest a membrane-associated complex between erythroid p55, protein 4.1, and glycophorin C. *J. Biol. Chem.* **269**, 8631-4.
- Marfatia, S.M., Byron, O., Campbell, G., Liu, S.C. und Grishti, A.H. (2000). Human homologue of the Drosophila discs-large tumor suppressor proteins forms an oligomer in solution. Identification of the self-association site. *J. Biol. Chem.* **275**, 13759-70.
- Martinez-Estrada, O.M. Villa, A., Breviario, F., Orsenigo, F., Dejana, E. und Bazzoni-G. (2001). Association of junctional adhesion molecule with calcium/calmodulin dependent serine protein kinase (CASK/LIN-2) in human epithelial caco-2 cells. *JBC* **276**, 9291-6.
- Martinez-Palomo, A. und Erlij, D. (1975). Structure of tight junctions in epithelia with different permeability. *Proc Natl Acad Sci. USA* **72**, 4487-91.
- Martin-Padura, I. et al. (1998). Junctional adhesion molecule, a novel member of the immunoglobulin superfamily that distributes at intercellular junctions and modulates monocyte transmigration. *J. Cell Biol.* **142**, 117-27.
- Marzioni, D., Banita, M., Felici, A., Paradinas, F.J., Newlands, E., De Nictol, M., Muhlhauser, J. und Castellucci, M. (2001). Expression of ZO-1 and occludin in normal human placenta and in hydratidiform moles. *Mol. Hum. Reprod.* **7**, 279-285.
- Masuko, N., Makino, K., Kawahara, H. und Saya, H. (1999). Interaction of NE-dlg/SAP102, a neuronal and endocrine tissue-specific membrane-associated guanylate-kinase protein, with calmodulin and PSD95/SAP90. A possible regulatory role in molecular clustering at synaptic sites. *J. Biol. Chem.* **274**, 5782-90.
- Matter, K und Balda, M.S. (1998). Biogenesis of tight junctions: the C-terminal domain of occludin mediates basolateral targeting. *J. Cell Sci.* **111**, 511-19.
- Mayer, B.J. (2001). SH3 domains: complexity in moderation. *J. Cell Sci.* **114**, 1253-63.
- Mayhan, W.G. (2000). Nitric oxide donor-induced increase in permeability of the blood-brain barrier. *Brain Res.* **866**, 101-108.
- McCarthy, K.M. et al. (1996). Occludin is a functional component of the tight junction. *J. Cell Sci.* **109**, 2287-98.
- McCarthy, K.M., Francis, S.A., McCormack, J.M., Lai, J., Rogers, R.A., Skare, I.B., Lynch, R.D. und Schneeberger, E.E. (2000). Inducible expression of claudin-1-myc but not occludin-VSV-G results in aberrant tight junction strand formation in MDCK cells. *J. Cell Sci.* **113**, 3387-98.
- McGee, A.W. und Bredt, D.S. (1999). Identification of an intramolecular interaction between the SH3 and guanylate kinase domains of PSD-95. *JBC* **274**, 17431-36.
- McGee, A.W., Dakoji, S.R., Olsen, O., Bredt, D.S., Lim, W.A. und Prehoda, K.E. (2001). Structure of the SH3-guanylate kinase module of PSD-95 suggests a mechanism for regulated assembly of MAGUK scaffolding proteins. *Molecular Cell* **8**, 1291-1301.

- Merrill, J.E. und Murphy, S.P. (1997). Inflammatory events at the blood-brain barrier: regulation of adhesion molecules, cytokines, and chemokines by reactive nitrogen and oxygen species. *Brain Behavior. Immunity*, 245-63.
- Mertsch, K., Grune, T., Siems, W.G., Ladhoff, A., Saupe, N., Blasig, I.E. (1995). Hypoxia and reoxygenation of brain endothelial cells in vitro: a comparison of biochemical and morphological response. *Cell. Mol. Biol* **41**, 243-53.
- Mitich, L.L., Van Itallie, C.M. und Anderson, J.M. (2000). Molecular physiology and pathophysiology of the tight junctions. I. Tight junction structure and function: lesson from mutant animals and proteins. *Am. J. Gastrointest. Liver Physiol.* **279**, G250-54.
- Mizuguchi, H., Utoguchi, N. und Mayumi, T. (1997). Preparation of glial extracellular matrix: a novel method to analyze glial-endothelial cell interaction. *Brain Res. Prot.* **1**, 339-43.
- Mollgard, K., Malinowski, D.N. und Saunders, N.R. (1976). Lack of correlation between tight junction morphology and permeability properties in developing choroid plexus. *Nature* **264**, 293-94.
- Morita, K., Furuse, M., Fujimoto, K. und Tsukita, S. (1999). Claudin multigene family encoding four-transmembrane domain protein components of tight junction strands. *Proc. Natl. Acad. Sci. USA* **96**, 511-16.
- Morita, K., Sasaki, H., Furuse, M. und Tsukita, S. (1999). Endothelial claudin: Claudin-5/TMVCF constitutes tight junction strands in endothelial cells. *J. Cell Biol.* **147**, 185-94.
- Moroi, S. et al. (1998). Occludin is concentrated at tight junctions of mouse/rat but not human/guinea pig Sertoli cells in testis. *Am. J. Physiol.* **274**, C1708-17.
- Morton, T.A., Bennett, D.B., Appelbaum, E.R., Cusimano, D.M., Johanson, K.O., Matico, R.E., Young, P.R., Doyle, M. und Chaiken, I.M. (1994). Analysis of the interaction between human interleukin-5 and the soluble domain of its receptor using a surface plasmon resonance biosensor. *J Mol Recognit* **7**, 47-55.
- Morton, T.A., Myszka, D.G. und Chaiken, I.M. (1995). Interpreting complex binding kinetics from optical biosensors: a comparison of analysis by linearization, the integrated rate equation, and numerical integration. *Anal. Biochem.* **227**, 176-85.
- Morton, T.A. und Myszka, D.G. (1998). Kinetic analysis of macromolecular interactions using surface plasmon resonance biosensors. *Methods in Enzymol.* **295**, 268-294.
- Muresan, Z., Paul, D.L. und Goodenough, D.A. (2000). Occludin 1B, a variant of tight junction protein. *Mol. Biol. Cell.* **11**, 627-34.
- Myszka, D.G., Morton, T.A., Doyle, M.L. und Chaiken, I.M. (1997) Kinetic analysis of a protein antigen-antibody interaction limited by mass transport on an optical biosensor. *Biophys. Chem.* **64**, 127-37.
- Myszka, D.G., He, X., Dembo, M., Morton, T.A. und Goldstein, B. (1998). Extending the range of rate constants available from BIACORE: interpreting mass transport-influenced binding data. *Biophys. J.* **75**, 583-94.
- Nag, S., Picard, P. und Stewart D.J. (2001). Expression of nitric oxide synthases and nitrotyrosine during blood-brain barrier breakdown and repair after cold injury. *Lab. Invest.* **81**, 41-49.
- Nakamura, T., Blechman, J., Tada, S., Rozovskaia, T., Itoyama, T., Bullrich, F., Mazo, A., Croce, C.M., Geiger, B. und Canaani, E. (2000) huASH1 protein, a putative transcription factor encoded by a human homologue of the *Drosophila* ash1 gene, localizes to both nuclei and cell-cell tight junctions. *Proc. Natl. Acad. Sci. U S A* **97**, 7284-9.
- Nelson, W.J. (1992). Regulation of cell surface polarity from bacteria to mammals. *Science* **258**, 948-54.

- Nieba, L., Krebber A. und Pluckthun, A. (1996). Competition BIACore for measuring true affinities: large differences from values determined from binding kinetics. *Anal Biochem.* **234**, 155-65.
- Nigam, S.K., Denisenko, N., Rodriguez-Boulan, E. und Citi, S. (1991). The role of phosphorylation in development of tight junction in cultured renal epithelial (MDCK) cells. *BBRC* **181**, 548-53.
- Nigam, S.K., Rodriguez-Boulan, E. und Silver, R.B. (1992). Changes in intracellular calcium during the development of epithelial polarity and junctions. *PNAS USA* **89**, 6162-66.
- Nilsson, M., Fagman, H. und Ericson, L.E. (1996). Ca^{2+} -dependent and Ca^{2+} -independent regulation of the thyroid epithelial junctioncomplex by protein kinases. *Exp. Cell. Res.* **225**, 1-11.
- Nishimura, M. et al. (2001). JEAP: a novel component of tight junction in exocrine cells. *JBC* **277**, 5583-7.
- Nix, S.L., Chishti, A.H., Anderson, J.M., Walther, Z. (2000). hCASK and hDlg associate in epithelia, and their src homology 3 and guanylate kinase domains participate in both intramolecular and intermolecular interactions. *JBC* **275**, 41192-200.
- Notterpek, L., Roux, K.J., Amici, S.A., Yazdanpour, A., Rahner, C. und Fletcher, B.S. (2001) Peripheral myelin protein 22 is a constituent of intercellular junctions in epithelia. *Proc. Natl. Acad. Sci. U S A.* **98**, 14404-9.
- Nusrat, A., Turner, J.R. und Madara J.L. (2000). Molecular physiology and pathophysiology of the tight junctions. IV. Regulation of tight junctions by extracellular stimuli: nutrients, cytokines and immune cells. *Am. J. Physiol. Gastrointest. Liver Physiol.* **279**, G885-57.
- Nusrat, A., Chen, J.A., Foley, C.S., Liang, T.W., Tom, J., Cromwell, M., Quan, C. und Mrsny, R.J. (2000). The Coiled coil domain of occludin can act to organize structural and functional elements of the epithelial tight junction. *J. Biol. Chem.* **275**, 29816-29822.
- O'Brien, M.J., Perez-Luna, V. H., Brueck, S.R. und Lopez, G.P. (2001). A surface plasmon resonance array biosensor based on spectroscopic imaging. *Biosens Bioelectron* **16**, 97-108.
- O'Shannesssey, D.J., Brigham-Burke, M., Soneson, K.K., Hensley, P. und Brooks, I. (1994). Determination of rate and equilibrium binding constants for macromolecular interactions by surface plasmon resonance. *Methods in Enzymology* **240**, 323-349.
- Otto, A. (1968). A new method for exciting non-radioactive surface plasma oscillations. *Phys. Stat. Sol.* **26**, K99-K101.
- Ozawa, M., Baribault, H. und Kemler, R. (1989). The cytoplasmic domain of the cell adhesion molecule uvomorulin associates with three independent proteins structurally related in different species. *EMBO J.* **8**, 1711-7.
- Padridge, W.M. (1999). Blood-brain barrier biology and methodology. *J. Neurovirol.* **5**, 556-69.
- Palmeri, D., van Zante, A., Huang, C.C., Hemmerich, S. und Rosen, S.D. (2000). Vascular endothelial junction-associated molecule, a novel member of the immunoglobulin superfamily, is localizedto intercellular boundaries of endothelial cells. *J. Biol. Chem.* **275**, 19139-45.
- Pekny, M., Stannes, K.A., Eliasson, C., Betsholtz, C. und Janigro, D. (1998). Impaired induction of blood-brain barrier properties in aortic endothelial cells by astrocytes from GFAP-deficient mice. *Glia* **22**, 390-400.
- Pim, D., Thomas, M., Javier, R., Gardiol, D. und Banks, L. (2000). HPV E6 target degradation of the discs large protein: evidence for the involvement of a novel ubiquitin ligase. *Oncogene* **19**, 719-25.

- Plateel, M., Teissier, E. und Cecchelli, R. (1997). Hypoxia dramatically increases the nonspecific transport of blood-borne proteins to the brain. *J. Neurochem.* **68**, 874-77.
- Posern, G., Zheng, J., Knudsen, B.S., Kardinal, C., Muller, K.B., Voss, J., Shishido, T., Cowburn, D., Cheng, G., Wang, B. et al. (1998). Development of highly selective SH3 binding peptides for Crk and CRKL which disrupt Crk-complexes with DOCK180, SoS and C3G. *Oncogene* **16**, 1903-12.
- Raub, T.J., Kuentzel, S.L. und Sawada, G.A. (1992). Permeability of bovine brain microvessel endothelial cells in vitro: barrier tightening by a factor released from astrogloma cells. *Exp. Cell Res.* **199**, 330-40.
- Raub, T.J. (1996). Signal transduction and glial cell modulation of cultured brain microvessel endothelial tight junction. *Am. J. Physiol.* **271**, C495-503.
- Rajasekaran, A.K., Hojo, M., Huima, T. und Rodriguez-Boulan, E. (1996). Catenins and zonula occludens-1 form a complex during early stages in the assembly of tight junctions. *J. Cell. Biol.* **132**, 451-63.
- Reichert, M., Müller, T. und Hunziker, W. (2000). The PDZ domains of zonula occludens-1 induce an epithelial to mesenchymal transition of Madine-Darby canine kidney I cells. *JBC* **275**, 9492-500.
- Reynolds, A.B., Daniel, J., McCrea, P.D., Wheelock, M.J., Wu, J. und Zhang, Z. (1994). Identification of a new catenin: the tyrosine kinase substrate p120cas associates with E-cadherin complexes. *Mol Cell Biol.* **14**, 8333-42.
- Risau, W., Hallmann, R., Albrecht, U. und Henke-Fahle, S. (1986). Brain induce the expression of an early cell surface marker for blood-brain barrier-specific endothelium. *EMBO J.* **5**, 3179-83.
- Risau, W., Engelhardt, B. und Wekerle, H. (1990). Incomplete presentation of protein (auto-) antigens by rat brain microvascular endothelium in vitro. *J. Cell Biol.* **110**, 1757-66.
- Risau, W. (1991). Induction of blood-brain barrier endothelial cell differentiation. *Ann. NY Acad. Sci.* **633**, 405-19.
- Risau, W. (1995). Differentiation of endothelium. *FASEB J.* **9**, 926-33.
- Rodriguez-Boulan, E und Nelson, W.J. (1989). Morphogenesis of the polarized epithelial cell phenotype. *Science* **245**, 718-25.
- Rubin, L.L. (1991a). The blood-brain barrier in and out of cell culture. *Curr. Opin. Neurobiol.* **1**, 360-63.
- Rubin, L.L. et al. (1991b). A cell culture model of the blood-brain barrier. *J. Cell Biol.* **115**, 1725-35.
- Rubin, L.L. (1992). Endothelial cells: adhesion and tight junction. *Curr. Opin. Cell Biol.* **4**, 830-33.
- Rubin, L.L. und Staddon, J.M. (1999). The cell biology of the blood-brain barrier. *Annu. Rev. Neurosci.* **22**, 11-28.
- Rutten, M.J., Hoover, R.L. und Karnovsky, M.J. (1987). Electrical resistance and macromolecular permeability of brain endothelial monolayer cultures. *Brain Res.* **425**, 301-10.
- Ryeom, S.W., Paul, D. und Goodenough, D.A. (2000). Truncation mutants of the tight junction protein ZO-1 disrupt corneal epithelial cell morphology. *Mol. Biol. Cell* **11**, 1687-96.
- Saha, C., Nigam, S.K. und Denker, B.M. (1998). Involvement of Galphai2 in the maintenance and biogenesis of epithelial cell tight junctions. *J. Biol. Chem.* **273**, 21629-33.
- Saha, C., Nigam, S.K. und Denker, B.M. (2001). Expanding role of G-proteins in tight junction regulation: Galphai(s) stimulates TJ assembly. *Biochem Biophys Res Commun* **285**, 250-6.

- Saitou, M. et al. (1997). Mammalian occludin in epithelial cells: ist expression and subcellular distribution. *Eur. J. Cell. Biol.* **73**, 222-31.
- Saitou, M. et al. (1998). Occludin-deficient embryonic stem cells can differentiate into polarized epithelial cells bearing tight junctions. *J. Cell Biol.* **141**, 397-408.
- Saitou, M. et al. (2000). Complex phenotype of mice lacking occludin, a component of tight junction strands. *Mol. Biol. Cell* **11**, 4131-42.
- Sakakibara, A., Furuse, M., Saitou, M., Ando-Akatsuka, Y. und Tsukita, S. (1997). Possible involvement of phosphorylation of occludin in tight junction formation. *J. Cell Biol.* **137**, 1393-1401.
- Sambrook, J., Fritsch E.F. und Maniatis, T. (1989). „Molecular cloning: A laboratory manual.“, Cold Spring Harbour Laboratory Press, USA.
- Saunders, N.R., Dziegielewska, K.M. und Mollgard, K. (1991). The importance of the blood-brain barrier in fetuses and embryos. *Trends Neurosci.* **14**, 14-15.
- Schleinkofer, K., Dingley, A., Tacken, I., Federwisch, M., Muller-Newen, G., Heinrich, P.C., Vusio, P., Jaques, Y. und Grotzinger J. (2001). Identification of the domain in the human interleucin-11 receptor that mediates ligand binding. *J Mol Biol* **306**, 263-74.
- Schloßhauer, B. und Herzog, K.-H. (1990). Neurothelin: an inducible cell surface glycoprotein of blood-brain barrier-specific endothelial cells and distinct neurons. *J. Cell Biol.* **110**, 1261-74.
- Schmidt, A., Utepbergenov, D.I., Krause, G. und Blasig, I.E. (2001). Use of surface plasmon resonance for real-time analysis of the interaction of ZO-1 and occludin. *BBRC* **288**, 1194-1199.
- Schmidt, A., Utepbergenov, D.I., Krause, G. und Blasig, I.E. (in press). Direct demonstration of the association of the blood-brain barrier proteins ZO-1 and occludin using surface plasmon resonance (SPR) spectroscopy-effect of SIN-1. In *Blood-spinal cord and brain barriers in health and disease*. (H.S. Sharma und J. Westmann, Ed.). Academic press, San Diego.
- Schneeberger, E.E. und Lynch, R.D. (1992). Structure, function and regulation of cellular tight junctions. *Am. J. Physiol.* **262**, L647-61.
- Schuck, P. (1996). Kinetics of ligand binding to receptor immobilized in a polymer matrix, as detected with an evanescent wave biosensor. I. a computer simulation of the influence of mass transport. *Biophys. J.* **70**, 1230-49.
- Schulze, C., Smales, C., Rubin, L.L. und Staddon, J.M. (1997). Lysophosphatidic acid increases tight junction permeability in cultured brain endothelial cells. *J. Neurochem.* **68**, 991-1000.
- Sheth, B., Fontaine, J.J., Ponza, E., McCallum, A., Page, A., Citi, S., Louvard, D., Zahraoui, A. und Fleming, T.P. (2000) Differentiation of the epithelial apical junctional complex during mouse preimplantation development: a role for rab13 in the early maturation of the tight junction. *Mech. Dev.* **97**, 93-104.
- Shin, H., Hsueh, Y.P., Yang, F.C., Kim, E. und Sheng, M. (2000). An intramolecular interaction between Src homology 3 domain and guanylate kinase-like domain required for channel clustering by postsynaptic density-95/SAP90. *J. Neurosci.* **20**, 3580-7.
- Simon, D.B., Lu, Y., Choate, K.A., Velazquez, H., Al-Sabban, E., Praga, M., Casari, G., Bettinelli, A., Colussi, G., Rodriguez-Soriano, J., McCredie, D., Milford, D., Sanjad, S. und Lifton, R.P. Paracellin-1, a renal tight junction protein required for paracellular Mg²⁺ resorption. *Science* **285**, 103-6.
- Small, R.K., Watkins, B.A., Munro, P.M. und Liu, D. (1993). Functional properties of retinal Müller cells following transplantation to the anterior eye chamber. *Glia* **7**, 158-69.
- Sporbert, A., Mertsch, K., Smolenski, A., Haseloff, R.F., Schönfelder, G., Paul, M., Ruth, P., Walter, U. und Blasig, I.E. (1999). Phosphorylation of vasodilator-stimulated phospho

- protein: a consequence of nitric oxide- and cGMP-mediated signal transduction in brain capillary endothelial cells and astrocytes. *Mol. Brain Res.* **61**, 258-266.
- Staddon, J.M., Herrenknecht, K., Schulze, C., Smales, C. und Rubin, L.L. (1995a). Signal transduction at the blood-brain barrier. *Biochem. Soc. Trans.* **23**, 475-79.
- Staddon, J.M., Herrenknecht, K., Smales, C. und Rubin, L.L. (1995b). Evidence that tyrosine phosphorylation may increase tight junction permeability. *J. Cell Sci.* **108**, 609-19.
- Staddon, J.M. und Rubin, L.L. (1996). Cell adhesion, cell junction and the blood-brain barrier. *Curr. Opin. Neurobiol.* **6**, 622-27.
- Stahelin, L.A. (1973). Further observations on the fine structure of freeze-cleaved tight junctions. *J. Cell Sci.* **13**, 763-86.
- Stehle, T. und Schulz, G.E. (1990) Three-dimensional structure of the complex of guanylate kinase from yeast with its substrate GMP. *J Mol Biol.* **211**, 249-54.
- Stevenson, B.R., Siliciano, J.D., Mooseker, M.S. und Goodenough, D.A. (1986). Identification of ZO-1: a high molecular weight polypeptide associated with the tight (zonula occludens) in a variety of epithelia. *J. Cell Biol.* **103**, 755-66.
- Stevenson, B.R., Anderson, J.M., Goodenough, D.A. und Mooseker, M.S. (1988). Tight junction structure and ZO-1 content are identical in two strains of Madin-Darby canine kidney cells which differ in transepithelial resistance. *J. Cell Biol.* **107**, 2401-8.
- Stevenson, B.R., Heintzelman, M.B., Anderson, J.M., Citi, S. und Mooseker, M.S. (1989a) ZO-1 and cingulin: tight junction proteins with distinct identities and localizations. *Am. J. Physiol.* **257**, C621-8.
- Stevenson, B.R. und Paul D.L. (1989b) The molecular constituents of intercellular junctions. *Curr Opin Cell Biol.* **1**, 884-91.
- Stevenson, B.R. und Keon, B.H. (1998). The tight junctions: morphology to molecules. *Annu. Rev. Cell. Dev. Biol.* **14**, 89-109.
- Stuart, R.O., Sun, A., Panichas , M., Herbert, B.M. und Nigam, S.K. (1994). Critical role for intracellular calcium in tight junction biogenesis. *J. Cell. Physiol.* **159**, 423-33.
- Stuart, R.O., Sun, A., Bush, K.T. und Nigam, S.K. (1996). Dependence of epithelial intercellular junction biogenesis on thapsigargin-sensitive intracellular calcium stores. *JBC* **271**.
- Takahashi, K., Matsuo, T., Katsume, T., Ueda, R., Yamamoto, D. (1998). Direct binding between two PDZ domain proteins Canoe and ZO-1 and their roles in regulation of the jun N-terminal kinase pathway in Drosophila morphogenesis. *Mech. Dev.* **78**, 97-111.
- Takeichi, M. (1991). Cadherin cell adhesion receptors as a morphogenetic regulator. *Science* **251**, 1451-5.
- Tavares, G.A., Panepucci, E.H. und Brunger, A.T. (2001). Structural characterization of the intramolecular interaction between the SH3 and guanylate kinase domains of PSD-95. *Molecular Cell* **8**, 1313-25.
- Thomas, U., Phannavong, B., Muller, B., Garner, C.C. und Gundelfinger, E.D. (1997). Functional expression of rat synapse-associated proteins SAP97 and SAP102 in Drosophila dlg-1 mutants: effects on tumor suppression and synaptic bouton structure. *Mech.Dev.* **62**, 161-74.
- Thomas, U., Ebitsch, S., Gorczyca, M., Koh, Y.H., Hough, C.D., Woods, D., Gundelfinger, E.D. und Budnik, V. (2000). Synaptic targeting and localization of discs-large is a stepwise process controlled by different domains of the protein. *Curr. Biol.* **10**, 1108-17.
- Toyofuku, T., Yabuki, M., Otsu, K., Kuzuya, T., Hori, M. und Tada M. (1998). Direct association of the gap junction protein connexin-43 with ZO-1 in cardiac myocytes. *J. Biol. Chem.* **273**, 12725-31.

- Tsukamoto, T. und Nigam, S.K. (1997). Tight junction proteins form large complexes and associate with the cytoskeleton in an ATP depletion model for reversible junction assembly. *J. Biol. Chem.* **272**, 16133-9.
- Tsukita, S., Itoh, M., Nagafuchi, A., Yonemura, S. und Tsukita, S. (1993). Submembranous junctional plaque proteins include potential tumor suppressor molecules. *J Cell Biol.* **123**, 1049-53.
- Tsukita, S. und Furuse, M. (1998). Overcoming barriers in the study of tight junction function: from occludin to claudin. *Genes Cells* **3**, 569-73.
- Tsukita, S. und Furuse, M. (1999a). Occludin and claudins in tight junction strands: leading or supporting players? *Trend. Cell Biol.* **9**, 268-73.
- Tsukita, S., Furuse, M. und Itoh, M. (1999b). Structural and signaling molecules come together at tight junction. *Curr. Opin. Cell Biol.* **11**, 628-33.
- Tsukita, S. und Furuse, M. (2000). Pores in the wall: claudins constitute tight junction strands containing aqueous pores. *J. Cell Biol.* **149**, 13-16.
- Tsukita, S., Furuse, M. und Itoh, M. (2001). Multifunctional strands in tight junctions. *Nat Rev Mol Cell Biol* **2**, 285-93.
- Utepbergenov, D.I., Mertsch, K., Sporbert, A., Tenz, K., Haseloff, R.F., Paul, M. und Blasig, I.E. (1998). NO protects blood-brain barrier in vitro from damage induced by hypoxia-reoxygenation. *FEBS Lett.* **424**, 197-201.
- Van Meer, G. und Simon, K. (1986). The function of tight junctions in maintaining differences in lipid composition between the apical and basolateral cell surface domains of MDCK cells. *EMBO J.* **5**, 1455-64.
- Van der Merwe, A.P. und Barclay, N.A. (1996). Analysis of cell-adhesion molecule interactions using surface plasmon resonance. *Curr. Op. Immunol.* **8**, 257-61.
- Van Buul, J.D. et al. (2002). Migration of human hematopoietic progenitor cells across bone marrow endothelium is regulated by vascular endothelial cadherin. *J. Immunol.* **168**, 588-96.
- Van Itallie, C.M. und Anderson, J.M. (1997). Occludin confers the adhesiveness when expressed in fibroblasts. *J. Cell. Sci.* **110**, 1113-21.
- Walsh, S.V., Hopkins, A.M., Chen, J., Narumiya, S., Parkos, C.A. und Nusrat, A. (2001). Rho kinase regulates tight junction function and is necessary for tight junction assembly in polarized epithelia. *Gastroenterology* **121**, 566-79.
- Walsh, S.V., Hopkins, A.M. und Nusrat, A. (2000). Modulation of tight junction structure and function by cytokines. *Adv Drug Deliv Res* **41**, 303-13.
- Wang, W., Dentler, W.L. und Borchardt, R.T. (2001). VEGF increases BMEC monolayer permeability by affecting occludin expression and tight junction permeability. *Am. J. Physiol. Heart Circ. Physiol.* **280**, H434-40.
- Weber, E. et al. (1994). Expression and polarized targeting of a rab3 isoform in epithelial cells. *J. Cell Biol.* **125**, 583-94.
- Willot, E., Balda, M.S., Heintzelmann, M., Jameson, B. und Anderson, J.M. (1992). Localization and differential expression of two isoforms of the tight junction protein ZO-1. *Am J Physiol* **262**, C1119-24.
- Willot, E. et al. (1993). The tight junction protein ZO-1 is homologous to the Drosophila discs-large tumor suppresser protein of septate junctionc. *Proc Natl Acad. Sci. USA* **90**, 7834-38.
- Wittchen, E.S., Haskins, J. und Stevenson, B.R. (1999). Protein interactions at the tight junction. *J. Biol. Chem.* **274**, 35179-35185.
- Wittchen, E.S., Haskins, J. und Stevenson, B.R. (2000). Exogenous expression of the amino-terminal half of the tight junction proteinZO-3 perturbs junctinal complex assembly. *J Cell Biol* **13**, 825-36.

- Wolburg, H. et al. (1994). Modulation of tight junction structure in blood-brain barrier endothelial cells. *J. Cell Sci.* **107**, 1347-57.
- Wolburg, H., Wolburg-Buchholz, K., Liebner, S. und Engelhard, B. (2001). Claudin-1, claudin-2 and claudin-11 are present in tight junctions choroid plexus epithelium of the mouse. *Neurosci Lett.* **307**, 77-80.
- Wong, V. und Gumbiner, B.M. (1997). A synthetic peptide corresponding to the extracellular domain of occludin perturbs the tight junction permeability barrier. *J. Cell Biol.* **136**, 399-409.
- Wong, V. (1997). Phosphorylation of occludin correlates with occludin localization and function at the tight junction. *Am J Physiol.* **273**, C1859-67.
- Woods, D.F. und Bryant, P.J. (1993). ZO-1, DlgA and PSD-95/SAP90: homologous proteins in tight, septate and synaptic cell junctions. *Mech Dev.* **44**, 85-9.
- Woods, D.F., Wu, J.W. und Bryant, P.J. (1997). Localization of proteins to the apico-lateral junctions of Drosophila epithelia. *Dev Genet.* **20**, 111-8.
- Woods, D.F., Hough, C., Peel, D., Callaini, G. und Bryant, P.J. (1996). Dlg protein is required for junction structure, cell polarity, and proliferation control in Drosophila epithelia. *J. Cell Biol.* **134**, 1469-82.
- Wu, H., Reuver, S.M., Kuhlendahl, S., Chung, W.J. und Carner, C.C. (1998). Subcellular targeting and cytoskeletal attachment of SAP97 to the epithelial lateral membrane. *J. Cell Sci.* **111**, 2365-76.
- Wu, H., Reissner, C., Kuhlendahl, S., Coblenz, B., Reuver, S.M., Kindler, S., Gundelfinger, E.D. und Carner, C.C. (2000). Intramolecular interactions regulate SAP97 binding to GKAP. *EMBO J.* **19**, 5740-51.
- Wu, X., Hepner, K., Castelino-Prabhu, S., Do, D., Kaye, M.B., Yuan, X.-Y., Wood, J., Ross, C., Sawyers, C.L. und Whang, Y.E. (2000). Evidence for regulation of the PTEN tumor suppressor by a membrane-localized multi-PDZ domain containing scaffold protein MAGI-2. *Proc. Natl. Acad. Sci. USA* **97**, 4233-38.
- Wu, Y., Dowbenko, D., Spencer, Richard, L., Lee, J., Gu, Q. und Lasky, L.A. (2000). Interaction of the Tumor Suppressor PTEN/MMAC with a PDZ Domain of MAGI3, a Novel Membrane-associated Guanylate Kinase. *J. Biol. Chem.* **275**, 21477-21485.
- Yamagata, K. et al. (1997). Astrocyte-conditioned medium induce blood-brain barrier properties in endothelial cells. *Clin. Exper. Pharmakol. Physiol.* **24**, 710-13.
- Yamamoto, T., Harada, N., Kano, K., Taya, S., Canaani, E., Matsuura, Y., Mizoguchi, A., Ide, C. und Kaibuchi, K. (1997) The Ras target AF-6 interacts with ZO-1 and serves as a peripheral component of tight junctions in epithelial cells. *J. Cell. Biol.* **139**, 785-95.
- Yamamoto, T., Harada, N., Kawano, Y., Taya, S. und Kaibuchi, K. (1999). In vivo interaction of AF-6 with activated Ras and ZO-1. *Biochem. Biophys. Res. Commun.* **259**, 103-7.
- Zahraoui, A., Joberty, G., Arpin, M., Fontaine, J.J., Hellio, R., Tavitian, A.. und Louvard, D. (1994). A small rab GTPase is distributed in cytoplasmic vesicles in non polarized cells but colocalizes with the tight junction marker ZO-1 in polarized epithelial cells. *J. Cell. Biol.* **124**, 101-15.
- Zhong, Y. et al. (1993). monoclonal antibody 7H6 reacts with a novel tight junction-associated protein distinct from ZO-1, cingulin, and ZO-2. *J. Cell Biol.* **120**, 477-83.
- Zoche, M. (1997). Charakterisierung der molekularen Wechselwirkung zwischen Calmodulin und Zieldomänen-eine Oberflächen-Plasmon-Resonanz spektroskopische Studie. Dissertation, FZ Jülich.
- Zöllner, S., Haseloff, R.F., Kirilyuk, I.A., Blasig, I.E. und Rubanyi, G.M. (1997). Nitroxides increase the detectable amount of nitric oxid released from endothelial cells. *J. Biol. Chem.* **272**, 23076-80.