

REFERENCES

- Ahnert-Hilger, G., Nürnberg, B., Exner, T., Schäfer, T., Jahn, R. (1998). The heterotrimeric G protein Go_2 regulates catecholamine uptake by secretory vesicles. *EMBO J.* **17**, 406-413.
- Ahnert Hilger, G., Weller, U. (1993). Comparison of the intracellular effects of clostridial neurotoxins on exocytosis from streptolysin O-permeabilized rat pheochromocytoma (PC 12) and bovine adrenal chromaffin cells. *Neuroscience* **53**, 547-552.
- Ahnert Hilger, G., Bigalke, H. (1995). Molecular aspects of tetanus and botulinum neurotoxin poisoning. *Prog. Neurobiol.* **46**, 83-96.
- Alder, J., Lu, B., Valtorta, F., Greengard, P., Poo, M. (1992a). Calcium-dependent transmitter secretion reconstituted in Xenopus Oocytes: requirement for synaptophysin. *Science* **257**, 657-661.
- Alder, J., Xie, Z.P., Valtorta, F., Greengard, P., Poo, M. (1992b). Antibodies to synaptophysin interfere with transmitter secretion at neuromuscular synapses. *Neuron* **9**, 759-768.
- Alder, J., Kraink, H., Valtorta, F., Greengard, P., Poo, M.M. (1995). Overexpression of synaptophysin enhances neurotransmitter secretion at Xenopus neuromuscular synapse. *J. Neurosci.* **15**, 511-519.
- Archer, B.T., Ozcelik, T., Jahn, R., Francke, U., Südhof, T.C. (1990). Structures and chromosomal localizations of two human genes encoding synaptobrevins 1 and 2. *J. Biol. Chem.* **265**, 17267-17273.
- Arikawa, K., Molday, L.L., Molday, R.S., Williams, D.S. (1992). Localisation of peripherin/rds in the disk membranes of cone and rod photoreceptors: relationship to disk membrane morphogenesis and retinal degeneration. *J. Cell Biol.* **116**, 659-667.
- Banerjee, A., Barry, V.A., DasGupta, B.R., Martin, T.F.J. (1996). N-ethylmaleimide-sensitive factor acts as a prefusion ATP-dependent step in Ca^{2+} -activated exocytosis. *J. Biol. Chem.* **271**, 20223-20226.
- Barnstable, C.J., Hofstein, R., Akagawa, K. (1985). A marker of early amacrine cell development in rat retina. *Dev. Brain Res.* **20**, 286-290.
- Baumert, M., Maycox, P.R., Navone, F., De Camilli, P., Jahn, R. (1989). Synaptobrevin: an integral membrane protein of 18,000 daltons in small synaptic vesicles. *EMBO J.* **8**, 379-384.
- Bean, A.J., Seifert, R., Chen, Y.A., Sachs, R., Scheller, R.H. (1997). Hrs-2 is an ATPase implicated in calcium-regulated secretion. *Nature* **385**, 826-829.
- Becher, A., Drenckhahn, A., Pahner, I., Margittai, M., Jahn, R., Ahnert-Hilger, G. (1999a). The synaptophysin-synaptobrevin complex: a hallmark of synaptic vesicle maturation. *J. Neurosci.* **19**, 1922-1931.

- Becher, A., Drenckhahn, A., Pahner, I., Ahnert-Hilger, G. (1999b). The synaptophysin-synaptobrevin complex is developmentally upregulated in cultivated neurons but is absent in neuroendocrine cells. *Eur. J. Cell Biol.*, in press.
- Bennett, M.K., Calakos, N., Scheller, R.H. (1992). Syntaxin: a synaptic protein implicated in docking of synaptic vesicles at presynaptic active zones. *Science* **257**, 255-259.
- Betz, A., Okamoto, M., Benseler, F., Brose, N. (1997). Direct interaction of the rat unc-13 homologue Munc13-1 with the N terminus of syntaxin. *J. Biol. Chem.* **272**, 2520-2526.
- Bittner, M.A., Holz, R.W. (1992). A temperature-sensitive step in exocytosis. *J. Biol. Chem.* **267**, 16219-16225.
- Bock, E., Helle, K.B. (1977). Localisation of synaptin on synaptic vesicle membranes, synaptosomal plasma membranes and chromaffin granule membranes. *FEBS Lett.* **82**, 175-178.
- Bradford, M.M. (1979). A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* **72**, 248-254.
- Brose, N., Petrenko, A.G., Südhof, T.C., Jahn, R. (1992). Synaptotagmin: a calcium sensor on synaptic vesicle surface. *Science* **256**, 1021-1025.
- Brose, N., Hofmann, K., Hata, Y., Südhof, T.C. (1995). Mammalian homologues of *Caenorhabditis elegans* unc-13 gene define novel family of C2-domain proteins. *J. Biol. Chem.* **270**, 25273-25280.
- Calakos, N., Scheller, R.H. (1994). Vesicle-associated membrane protein and synaptophysin are associated on the synaptic vesicle. *J. Biol. Chem.* **269**, 24534-24537.
- Chapman, E.R., An, S., Barton, N., Jahn, R. (1994). SNAP-25, a t-SNARE which binds to both syntaxin and synaptobrevin via domains that may form coiled coils. *J. Biol. Chem.* **269**, 27427-27432.
- Chapman, E.R., Hanson, P.I., Jahn, R. (1995). Ca^{2+} regulates the interaction between synaptotagmin and syntaxin 1. *J. Biol. Chem.* **270**, 23667-23671.
- Chapman, E.R., Blasi, J., An, S., Brose, N., Johnston, P.A., Südhof, T.C., Jahn, R. (1996). Fatty acylation of synaptotagmin in PC12 cells and synaptosomes. *Biochem. Biophys. Res. Commun.* **225**, 326-332.
- Connell, G.J., Molday, R.S. (1990). Molecular cloning, primary structure and orientation of the vertebrate photoreceptor cell protein peripherin in the rod outer segment disk membrane. *Biochemistry* **29**, 4691-4698.
- Couve, A., Protopopov, V., Gerst, J.E. (1995). Yeast synaptobrevin homologs are modified posttranslationally by the addition of palmitate. *Proc. Natl. Acad. Sci. U.S.A.* **92**, 5987-5991.

6. REFERENCES

- Cowan, D., Linial, M., Scheller, R.H. (1990). *Torpedo* synaptophysin: Evolution of a synaptic vesicle protein. *Brain Res.* **509**, 1-7.
- Daly, C., and Ziff, E.B. (1997). Post-translational regulation of synaptic vesicle protein expression and the developmental control of synaptic vesicle formation. *J. Neurosci.* **17**, 2365-2375.
- Di Antonio, A., Parfitt, K.D., Schwarz, T.L. (1993). Synaptic transmission persists in synaptotagmin mutants of *Drosophila*. *Cell* **73**, 1281-90.
- Dunphy, J.T., Linder, M.E. (1998). Signalling functions of protein palmitoylation. *Biochim. Biophys. Acta* **1436**, 245-261.
- Edelmann, L., Hanson, P.I., Chapman, E.R., Jahn, R. (1995). Synaptobrevin binding to synaptophysin: a potential mechanism for controlling the exocytotic fusion machine. *EMBO J.* **14**, 224-231.
- Elferink, L.A., Trimble, W.S., Scheller, R.H. (1989). Two vesicle-associated membrane protein genes are differentially expressed in rat central nervous system. *J. Biol. Chem.* **264**, 11061-11064.
- Eshkind, L.E., and Leube, R.E. (1995). Mice lacking synaptophysin reproduce and form typical synaptic vesicles. *Cell Tissue Res.* **282**, 423-433.
- Fasshauer, D., Bruns, D., Shen, B., Jahn, R., Brunger, A.T. (1997). A structural change occurs upon binding of syntaxin to SNAP-25. *J. Biol. Chem.* **272**, 4582-4590.
- Fasshauer, D., Eliason, W.K., Brunger, A.T., Jahn, R. (1998). Identification of a minimal core of the synaptic SNARE complex sufficient for reversible assembly and disassembly. *Biochemistry* **37**, 10354-10362.
- Fasshauer, D., Sutton, R.B., Brunger, A.T., Jahn, R. (1998). Conserved structural features of the synaptic fusion complex: SNARE proteins reclassified as Q- and R-SNAREs. *Proc. Natl. Acad. Sci. U.S.A.* **95**, 15781-15786.
- Fujita, Y., Shirataki, H., Sakisaki, T., Asakura, T., Ohya, T., Kotani, H., Yokoyama, S., Nishioka, H., Matsuura, Y., Mizogushi, A., Scheller, R.H., Takai, Y. (1998). Tomosyn: a syntaxin-1-binding protein that forms a novel complex in the neurotransmitter release process. *Neuron* **20**, 905-915.
- Fykse, E.M., Takei, K., Walch-Solimena, C., Geppert, M., Jahn, R., De Camilli, P., Südhof, T.C. (1993). Relative properties and localizations of synaptic vesicle protein isoforms: the case of the synaptophysins. *J. Neurosci.* **13**, 4997-5007.
- Gaardsvoll, H., Obendorf, D., Winkler, H., Bock, E. (1988). Demonstration of immunochemical identity between the synaptic vesicle proteins synaptin and synaptophysin/p38. *FEBS Lett.* **242**, 117-121.

6. REFERENCES

- Galli, T., Garcia, E.P., Mundigl, O., Chilcote, T.J., De Camilli, P. (1995). v- and t-SNAREs in neuronal exocytosis: a need for additional components to define sites of release. *Neuropharmacology* **43**, 1351-1360.
- Galli, T., McPherson, P.S., De Camilli, P. (1996). The V₀ sector of the V-ATPase, synaptobrevin, and synaptophysin are associated on synaptic vesicles in a Triton X-100-resistant, freeze-thawing sensitive, complex. *J.Biol.Chem.* **271**, 2193-2198.
- Glassmeier, G., Herzig, K.H., Hopfner, M., Lemmer, K., Jansen, A., Scherübl, H. (1998). Expression of functional GABA_A receptors in cholecystokinin-secreting gut neuroendocrine murine STC-1 cells. *J. Physiol. Lond.* **510**, 805-814.
- Hata, Y., Slaughter, C.A., Südhof, T.C. (1993). Synaptic vesicle fusion complex contains unc-18 homologue bound to syntaxin. *Nature* **366**, 347-351.
- Hess, D.T., Slater, T.M., Wilson, M.C., Skene, J.H. (1992). The 25 kDa synaptosomal-associated protein SNAP-25 is the major methionine-rich polypeptide in rapid axonal transport and a major substrate for palmitoylation in adult CNS. *J. Neurosci.* **12**, 4634-4641.
- Huttner, W.B., Schiebler, W., Greengard, P., and De Camilli, P. (1983). Synapsin I (Protein I), a nerve terminal-specific phosphoprotein. III. Its association with synaptic vesicles studied in a highly purified synaptic vesicle preparation. *J. Cell Biol.* **96**, 1374-1388.
- Jahn, R., Schiebler, W., Ouimet, C., Greengard, P. (1985). A 38,000 dalton membrane protein (p38) present in synaptic vesicles. *Proc. Natl. Acad. Sci. USA* **82**, 4137-4141.
- Jahn, R., Navone, F., Greengard, P., De Camilli, P. (1987). Biochemical and immunocytochemical characterisation of p38, an integral membrane glycoprotein of small synaptic vesicles. *Ann. N. Y. Acad. Sci.* **493**, 497-499.
- Jahn, R., De Camilli, P. (1991). Membrane proteins of synaptic vesicles: Markers for neurons and neuroendocrine cells; tools for the study of neurosecretion. In: *Markers for neuronal and endocrine cells*, (Gratzl, M., Langley, K. eds.). VCH, Weinheim, pp.65-66.
- Jahn, R., Südhof, T.C. (1994). Synaptic vesicles and exocytosis. *Annu. Rev. Neurosci.* **17**, 219-246.
- Jahn, R., Südhof, T.C. (1999). Membrane fusion and exocytosis. *Annu. Rev. Biochem.* **68**, 863-911.
- Johnston, P.A., Jahn, R., Südhof, T.C. (1989). Transmembrane topography and evolutionary conservation of synaptophysin. *J. Biol. Chem.* **264**, 1268-1273.
- Johnston, P.A., Südhof, T.C. (1990). The multisubunit structure of synaptophysin. Relationship between disulfide bonding and homo-oligomerization. *J. Biol. Chem.* **265**, 8869-8873.
- Kojima, T., Fukuda, M., Aruga, J., Mikoshiba, K. (1996). Calcium-dependent phospholipid binding to the C₂A domain of a ubiquitous form of double C₂ protein (Doc2 beta). *J. Biochem. Tokyo* **120**, 671-676.

- Komada, M., Kitamura, N. (1995). Growth factor-induced tyrosine phosphorylation of Hrs, a novel 115-kilodalton protein with a structurally conserved putative zinc finger domain. *Mol. Cell. Biol.* **15**, 6213-6221.
- Kraszewski, K., Daniell, L., Mundigl, O., De, Camilli, C. (1996). Mobility of synaptic vesicles in nerve endings monitored by recovery from photo-bleaching of synaptic vesicle-associated fluorescence. *J. Neurosci.* **16**, 5905-5913.
- Kretzschmar, S., Volknandt, W., Zimmermann, H. (1996). Colocalization on the same synaptic vesicles of syntaxin and SNAP-25 with synaptic vesicle proteins: a re-evaluation of functional models required? *Neurosci. Res.* **26**, 141-148.
- Laemmli, U.K. (1970). Cleavage of structural proteins during assembly of the head bacteriophage T4. *Nature* **227**, 680-685.
- Laage, R., Langosch, D. (1997). Dimerization of the synaptic vesicle protein synaptobrevin (vesicle-associated membrane protein) II depends on specific residues within the transmembrane segment. *Eur. J. Biochem.* **249**, 540-546.
- Leube, R.E., Kaiser, P., Seiter, A., Zimbelmann, R., Franke, W.W., Rehm, H., Knaus, P., Prior, P., Betz, H., Reinke, H., Beyreuther, K., Wiedenmann, B. (1987). Synaptophysin: molecular organisation and mRNA expression as determined from cloned cDNA. *EMBO J.* **6**, 3261-3268.
- Leube, R.E., Wiedenmann, B., Franke, W.W. (1989). Topogenesis and sorting of synaptophysin: synthesis of a synaptic vesicle protein from a gene transfected into nonneuroendocrine cells. *Cell* **59**, 433-446.
- Leveque, C., Pupier, S., Marqueze, B., Geslin, L., Kataoka, M., Takahashi, M., De Waard, M., Seagar, M. (1998). Interaction of cysteine string proteins with the alpha1A subunit of the P/Q-type calcium channel. *J. Biol. Chem.* **273**, 13488-13492.
- Martin, J.H. (1991). The collective electrical behaviour of cortical neurons: The electroencephalogram and the mechanism of epilepsy. In *Principals of Neuronal Science* 3rd edition. (Kandel, E., Schwartz, J.H., Jessell, T.M. ed.). Elsevier, New York, U.S.A..
- Martin-Moutot, N., Charvin, N., Leveque, C., Sato, K., Nishiki, T., Kozaki, S., Takahashi, M., Seagar, M. (1996). Interaction of SNARE complexes with P/Q-type calcium channels in rat cerebellar synaptosomes. *J. Biol. Chem.* **271**, 6567-6570.
- Martinéz de la Escalera, G., Gallo, F., Choi, A.L., Weiner, R.I. (1992). Dopaminergic regulation of the GT1 gonadotropin-releasing hormone (GnRH) neuronal cell lines: stimulation of GnRH release via D1-receptors positively coupled to adenylate cyclase. *Endocrinology* **131**, 2965-2971.
- McMahon, H.T., Ushkaryov, Y.A., Edelmann, L., Link, E., Binz, T., Niemann, H., Jahn, R., Südhof, T.C. (1993). Cellubrevin is a ubiquitous tetanus-toxin substrate homologous to a putative synaptic vesicle fusion protein. *Nature* **364**, 346-349.

- McMahon, H.T., Bolshakov, V.Y., Janz, R., Hammer, R.E., Siegelbaum, S.A., Südhof, T.C. (1996). Synaptophysin, a major synaptic vesicle protein, is not essential for neurotransmitter release. *Proc. Natl. Acad. Sci. U.S.A.* **93**, 4760-4764.
- Mollinedo, F., Lazo, P.A. (1997). Identification of two isoforms of the vesicle-membrane fusion protein SNAP-23 in human neutrophils and HL-60 cells. *Biochem. Biophys. Res. Commun.* **231**, 808-812.
- Navone, F., DiGioia, G., Matteoli, M., De Camilli, P. (1988). Small synaptic vesicles and large dense-core vesicles of neurons are related to two distinct types of vesicles of neuroendocrine cells. In: *Molecular mechanisms in secretion*. Alfred Benzon Symposium 25 (Thorn, N.A., Treiman, M., Peterson, O.H. eds). Munksgaard, Copenhagen, pp. 433-450.
- Okamoto, M., Südhof, T.C. (1997). Mints, Munc18-interacting proteins in synaptic vesicle exocytosis. *J. Biol. Chem.* **272**, 31459-31464.
- Orita, S., Sasaki, T., Naito, A., Komuro, R., Ohtsuka, T., Maeda, M., Suzuki, H., Igarashi, H., Takai, Y. (1995). Doc2: a novel brain protein having two repeated C₂-like domains. *Biochem. Biophys. Res. Commun.* **206**, 439-448.
- Orita, S., Naito, A., Sakaguchi, G., Maeda, M., Igarashi, H., Sasaki, T., Takai, Y. (1997). Physical and functional interactions of Doc2 and Munc13 in Ca²⁺-dependent exocytotic machinery. *J. Biol. Chem.* **272**, 16081-16084.
- Otto, H., Hanson, P.I., Jahn, R. (1997). Assembly and disassembly of a ternary complex of synaptobrevin, syntaxin, and SNAP-25 in the membrane of synaptic vesicles. *Proc. Natl. Acad. Sci. U.S.A.* **94**, 6197-6201.
- Ovchinnikov, Y.A., Abdulaev, N.G., Zolotarev, A.S., Artamonov, I.D., Bespalov, I.A., Dergachev, A.E., Tsuda, M. (1988). Octopus rhodopsin: amino acid sequence deduced from cDNA. *FEBS Lett.* **232**, 69-72.
- Pang, D.T., Wang, J.K.T., Valtora, F., Benfenati, F., Greengard, P. (1988). Protein tyrosine phosphorylation in synaptic vesicles. *Proc. Natl. Acad. Sci. USA* **85**, 762-766.
- Perin, M.S., Fried, V.A., Mignery, G.A., Jahn, R., Südhof, T.C. (1990). Phospholipid binding by a synaptic vesicle protein homologous to the regulatory region of protein kinase C. *Nature* **345**, 260-263.
- Perin, M.S., Johnston, P.A., Ozcelik, T., Jahn, R., Francke, U., Südhof, T.C. (1991a). Structural and functional conservation of synaptotagmin (p65) in *Drosophila* and humans. *J. Biol. Chem.* **266**, 615-622.
- Perin, M.S., Brose, N., Jahn, R., Südhof, T.C. (1991b). Domain structure of synaptotagmin (p65). *J. Biol. Chem.* **266**, 623-629.
- Prekeris, R., Terrian, D.M. (1997). Brain myosin V is a synaptic vesicle-associated motor protein: evidence for a Ca²⁺-dependent interaction with the synaptobrevin-synaptophysin complex. *J. Cell Biol.* **137**, 1589-1601.

6. REFERENCES

- Ravichandran, V., Chawla, A., Roche, P.A. (1996). Identification of a novel syntaxin- and synaptobrevin/VAMP binding-protein, SNAP-23, expressed in non-neuronal tissue. *J. Biol. Chem.* **271**, 13300-13303.
- Rehm, H., Wiedenmann, B., Betz, H. (1986). Molecular characterisation of synaptophysin, a major calcium-binding protein of synaptic vesicle membrane. *EMBO J.* **5**, 535-541.
- Robinson, L.J., Martin, T.J. (1998). Docking and fusion in neurosecretion. *Current Opinion in Cell Biology* **10**, 483-492.
- Seagar, M., Takahashi, M. (1998). Interactions between presynaptic calcium channels and proteins implicated in synaptic vesicle trafficking and exocytosis. *J. Bioenerg. Biomembr.* **30**, 347-356.
- Shao, X., Li, C., Fernandez, I., Zhang, X., Südhof, T.C., Rizo, J. (1997). Synaptotagmin-syntaxin interaction: the C2 domain as a Ca²⁺-dependent electrostatic switch. *Neuron* **18**, 133-142.
- Sheng, Z.H., Rettig, J., Cook, T., Catterall, W.A. (1996). Calcium-dependent interaction of N-type calcium channels with the synaptic core complex. *Nature* **379**, 451-454.
- Smith, P.K., Krohn, R.I., Hermanson, G.T., Mallia, A.K., Gartner, F.H., Provenzano, M.D., Fujimoto, E.K., Goeke, N.M., Olson, B.J., Klenk, D.C. (1985). Measurement of protein using bicinchoninic acid. *Anal. Biochem.* **150**, 76-85.
- Stenius, K., Janz, R., Südhof, T.C., Jahn, R. (1995). Structure of synaptogyrin (p29) defines novel synaptic vesicle protein. *J. Cell Biol.* **131**, 1801-1809.
- Südhof, T.C., Baumert, M., Perin, M.S., Jahn, R. (1989). A synaptic vesicle membrane protein is conserved from mammals to *Drosophila*. *Neuron* **2**, 1475-1481.
- Südhof, T.C., Jahn, R. (1991). Proteins of synaptic vesicles involved in exocytosis and membrane recycling. *Neuron* **6**, 665-677.
- Sutton, B., Fasshauer, D., Jahn, R., Brünger, A.T. (1998). Crystal structure of a core synaptic fusion complex at 2.4 Å resolution. *Nature* **395**, 347-353.
- Thomas, L., Hartung, K., Langosch, D., Rehm, H., Bamberg, E., Franke, W.W., Betz, H. (1988). Identification of synaptophysin as a hexameric channel protein of the synaptic vesicle membrane. *Science* **242**, 1050-1053.
- Trimble, W.S., Cowan, D.M., Scheller, R.H. (1988). VAMP-1: A synaptic vesicle-associated integral membrane protein. *Proc. Natl. Acad. Sci. USA* **85**, 4538-4542.
- Tu, Y., Wang, J., Ross, E.M. (1997). Inhibition of brain Gz GAP and other RGS proteins by palmitoylation of G protein alpha subunits. *Science* **278**, 1132-1135.
- Veh, R.W., Lichtinghagen, R., Sewing, S., Wunder, F., Grumbach, I.M., Pongs, O. (1995). Immunohistochemical localization of five members of the Kv1 channel subunits: contrasting

6. REFERENCES

- subcellular locations and neuron-specific co-localizations in rat brain. *Eur J. Neurosci.* **7**, 2189-2205.
- Veit, M., Sollner, T.H., Rothman, J.E. (1996). Multiple palmitoylation of synaptotagmin and the t-SNARE SNAP-25. *FEBS Lett.* **385**, 119-123.
- Walch-Solimena, C., Blasi, J., Edelmann, L., Chapman, E.R., von Mollard, G., Jahn, R. (1995). The t-SNAREs syntaxin 1 and SNAP-25 are present on organelles that participate in synaptic vesicle recycling. *J. Cell Biol.* **128**, 637-645.
- Wang, F.S., Wolenski, J.S., Cheney, R.E., Mooseker, M.S., Jay, D.G. (1996). Function of myosin V in filopodia extension of neuronal growth cones. *Science* **273**, 660-663.
- Weinreb, P.H., Zhen, W., Poon, A.W., Conway, K.A., Lansbury, P.T. (1996). NACP, a protein implicated in Alzheimer's disease and learning, is natively unfolded. *Biochemistry* **35**, 13709-13715.
- Wiedenmann, B., Franke, W.W. (1985). Identification and localisation of synaptophysin: An integral membrane glycoprotein of Mr 38,000 characteristic of presynaptic vesicles. *Cell* **41**, 1017-1028.
- Wiedenmann, B., Huttner, W.B. (1989). Synaptophysin and chromogranins/secretogranins-widespread constituents of distinct types of neuroendocrine vesicles and new tools in tumor diagnosis. *Virchows Archiv. B. Cell. Pathol.* **58**, 95-121.
- Yoshida, A., Oho, C., Omori, A., Kuwahara, R., Ito, T., Takahashi, M. (1992). HPC-1 is associated with synaptotagmin and w-conotoxin receptor. *J. Biol. Chem.* **267**, 24925-24928.
- Zhang, J.Z., Davleton, B.A. Südhof, T.C., Anderson, R.G.W. (1994). Synaptotagmin I is a high affinity receptor for clathrin AP-2: implications for membrane recycling. *Cell* **78**, 751-760.