

8. Literaturverzeichnis

Abrams K., Harvell J.D., Shriner D., Wertz P.W., Maibach H., Maibach H.I., Rehdeld S.J.: Effect of organic solvents on in vitro human skin water barrier function. *J Invest Dermatol* 1993; 101: pp. 609-613

Akimoto Y., Akamatsu H., Okano Y., Masaki H., Horio T.: Effects of UV irradiation on the sebaceous gland and sebum secretion in hamsters. *J Dermatol Sci* 2003; 31 (2): pp. 151-159

Alvarez-Roman R., Merino G., Kalia Y.N., Naik A., Guy R.H.: Skin permeability enhancement by low frequency sonophoresis: lipid extraction and transport pathways. *J Pharm Sci* 2003; 92 (6): pp. 1138-1146

Baker K.P.: Hair growth and replacement in the cat. *Br vet J* 1974; 130: pp: 327-335

Barry B.: Penetration enhancers; In: Shroot B., Schaefer H. (eds): *Skin Pharmakokinetics*. © Karger Basel, Freiburg, Paris, London, New Delhi, Bangkok, Singapore, Tokyo, Sidney, 1987; 1: pp. 121-137

Bartek M.J., LaBudde J.A., Maibach H.I.: Skin permeability in vivo: comparison in rat, rabbit, pig and man. *J Invest Dermatol* Oct 1972: pp. 114-123

Bartek M.J, La Budde J.A.: Percutaneous absorption in vitro. In: Maibach H.I. (ed); *Animal models in Dermatology*. Churchill Livingstone, New York, 1975: pp. 103-120

Behl C.R., Wittkowsky A., Barrett M., Pierson C.L., Flynn G.L.: Technique for preparing appendage-free (scar) on hairless mouse. *J Pharm Sci* 1981; 70 (7): pp. 835-837

Behl, C.R., Flynn G.L, Linn E.E., Smith W.M.: Percutaneous absorption of corticosteroids: age, site and skin sectioning influences on rates of permeation of hairless mouse skin by hydrocortisone. *J Pharm Sci* 1984; 73 (9): pp. 1287-1290

Birkby C.S., Wertz P.W., Downing D.T.: The polar lipids from keratinized tissues of some vertebrates. *Comp Biochem Physiol B*. 1982; 73 (2): pp. 239-42

Block K.: Cholesterol: Evolution of structure and function. In Vance D.E., Vance J.E. (eds); *Biochemistry of Lipids, Lipoproteins and Membranes*. Amsterdam, Elsevier, 1991: pp. 363-382

Blume U., Ferracin J., Vershoore M., Czernielewski J.M., Schaefer H.: Physiology of the vellus hair follicle: Hair growth and sebum excretion. Br J Dermatol 1991; 124: pp. 21-28

Bonté F., Saunois A., Pinguet P., Meybeck A.: Existence of a lipid gradient in the upper stratum corneum and its possible significance. Arch Dermatol Res 1997; 289 (2): pp. 78-82

Bouwstra J.A., Dubbelaar F.E.R., Gooris G.S., Ponec M.: The lipid organisation in the skin barrier. Acta Derm Venerol Supp 2000; 208: pp. 23-30

Bouwstra J.A., Gooris G.S., Duppelaar F.E.R., Ponec M.: Phase behavoir of stratum corneum lipid mixtures based on human ceramides. The role of natural and synthetic ceramide 1. J Invest Dermatol 2002; 118: pp. 606-617

Braun-Falko O., Plewig G., Wolff H.H.: In: Dermatologie und Venerologie. Springer Verlag Berlin, Heidelberg, New York, 1996

Breathnach A.S., Goodman T., Stolinksi C., Gross M., Freeze fracture replication of cells in stratum corneum of human epidermis. J Anat 1973; 114: pp. 65-81

Brunsch A.: Vergleichende Untersuchungen am Haarkleid von Wildcaniden und Haushunden. Z Tierzücht 1956; 67: pp. 205-240

Bronaugh R.L., Steward R.F., Congdon E.R.: Methods for in vitro percutaneous absorption studies. II. Animal models for human skin. Toxicology and applied Pharmacology 1982, 62: pp. 481-488

Burck H.C.: In: Histologische Technik. © Georg Thieme Verlag Stuttgart, New York 1982, 5. Aufl.

Claxton J.H.: The hair follicle group in mice. Anat Rec 1966;154: pp. 195-208

Cook H.W.: Fatty acid desaturation and chain elongation in eukaryotics. In Vance D.E., Vance J.E. (eds): Biochemistry of Lipids, Lipoproteins and Membranes. Amsterdam, Elsevier 1991: pp. 141-170

Cooper G., Schiller A.L.: Anatomy of the guinea pig. Cambridge: Harvard University Press, 1975

Corcuff P., Leveque J.L.: Corneocyte changes after acute UV irradiation and chronic solar exposure. Photo Dermatol 1988; 5: 110-115

Dawson H.L.: A study of hair growth in the guinea pig (*Cavia cobaya*) Am J Anat 45; 1930: pp. 462-484

Dick I.P., Scott R.C.: Pig ear skin as an in vitro model for human skin permeability. J Pharm Pharmacol, 1992; 44: pp. 640-645

Downing D.T.: Lipid and protein structures in the permeability barrier of mammalian epidermis. J Lipid Res 1992; Vol. 33: pp. 301-313

Downing D.T.: Photodensitometry in the thin-layer chromatographic analysis of neutral lipids. J Chromatogr 1968; 38: pp. 91-99

Dupuis D., Rougier A., Rouget R., Lotte C., Kalopissis G.: In vivo relationship between horny layer reservoir effect and percutaneous absorption in human and rat. J Invest Dermatol Apr 1984; 82 (4): pp 353-356

Ebling F.J.G., Hale P.A., Randall V.A.: Hormones and hair growth. In: L.A. Goldshmith (ed.) Physiology, Biochemistry and Molecular Biology of the skin. Oxford Press, Oxford 1991: pp. 660-696

Ebling F.: Hormonal control and methods of measuring sebaceous gland activitiy. J Invest Dermatol 1964; 62: pp. 162-171

Elias P.M., Goerke J., Friend D.S.: Mammalian epidermal barrier layer lipids: Composition and influence on structure. J Invest Dermatol 1977; 69: pp. 535-546

Elias P.M., Friend D.S.: The permeability barrier in mammalian epidermis. J Cell Biol 1975; 65: pp. 180-191

Elias P.M., Leventhal M.: Intercellular volume changes and cell surface expansion during cornification. Clin Res 1979; 27: p. 525a

Elias P.M., Williams M.L., Maloney M.E., Bonifas J.A., Brown B.E., Grayson S., Epstein E.H.: Stratum corneum lipids in disorders of cornification: Steroid sulfatase and cholesterol sulphate in normal desquamation and pathogenesis of recessive X-linked Ichthyosis. J Clin Invest 1984; 74: pp. 1414-1421

Elias P.M., Menon G.K., Grayson S., Brown B.E., Rehfeld S.J.: Avian sebokeratocytes and marine mammal lipokeratinocytes: Structural, lipid biochemical and functional considerations. Am J Anat 1987; 180: pp. 161-177

Elias P.M.: Epidermal lipids, barrier function and desquamation. J Invest Dermatol 1983; 80: pp. 44-49

Elias P.M., Cooper E.R., Korc A., Brown B.E.: Percutaneous transport in relation to stratum corneum structure and lipid composition. J Invest Dermatol 1981; 76; pp. 297-301

Elias P.M.: "Lipids and the epidermal permeability barrier", Archs Dermatol Res 1981: pp. 95-117

Essa E.A., Bonner M.C., Barry B.W.: Possible role of shunt route during iontophoretic drug penetration through skin. Perspectives in Percutaneous Penetration 2002; 8 a: p. 54

Fartasch M.: The nature of the epidermal barrier: Structural aspects. Adv Drug Del Rev 1995; 18: pp. 273-282

Fartasch M., Bassukas I.D., Diepgen T.L.: Structural relationship between epidermal lipid lamellae, lamellar bodies and desmosomes in human epidermis: An ultrastructural study. Br J Dermatol 1993; 128: pp. 1-9

Feldmann R.J., Maibach H.I.: Regional variation in percutaneous penetration of 14 C Cortisol in man. J Invest Dermatol 1967; Vol.48 (2): pp. 181-183

Flynn G.L: Physicochemical determinants of skin absorption. In Gerrity T.R., Henry C.J. (eds): Principles of Route-to-Route Extrapolation for Risk Assessment. New York, Elsevier 1990: pp. 93-127

Forslind B.: A domain mosaic model of the skin barrier. Acta Derm Venerol 1994; 74: pp.1-6

Freinkel R.K., Traczyk T.N.: Lipid composition and acid hydrolase content of lamellar granules of fetal rat epidermis. J Invest Dermatol 1985; 85: pp. 295-298

Fritsch P.: Aufbau und Funktionen der Haut. In: Fritsch P. (ed): Dermatologie und Venerologie, Lehrbuch und Atlas, Springer Verlag 1998: pp. 43-52

Futerman A.H.: Ceramide metabolism compartmentalized in the endoplasmatic reticulum and golgi apparatus. Curr Top Membr 1994; 40: pp. 93-110

Galey W.R., Lonsdale H.K., Nacht S.: The in vitro permeability of skin and buccal mucosa to selected drugs and tritiated water. J Invest Dermatol 1976; 67: pp: 713-717

Gibbs H.F.: A study of the postnatal development of the skin and hair of the mouse. Anat Rec 1941; 80: pp. 61-81

Goodridge A.G.: Fatty acid synthesis in eukaryotics. In Vance D.E., Vance J.E. (eds); Biochemistry of Lipids, Lipoproteins and Membranes. Amsterdam, Elsevier 1991: pp. 111-140

Goppel M., Franz G.: Stability control of valerian ground material and extracts – a new HPTLC method for the routine quantification of valerenic acids and lignans. Pharmazie 2004; 59 (6): pp.446-452

Gray G.M., Yardley H.J.: Different population of pig epidermal cells: isolation and lipid composition. J Lipid Res 1975; 16: pp.441-447

Gray G.M., Yardley H.J.: Lipid composition of cells isolated from pig, human and rat epidermis. J Lipid Res 1975; 16: pp. 434-440

Grayson S., Johnson-Winegar A.G., Wintraub B.U., Isserhof R.R., Epstein E.H., Elias P.M.: Lamellar body-enriched fractions from neonatal mice: Preparative techniques and partial characterization. J Invest Dermatol 1985; 85 (4): pp. 285-289

Greene S.A., Downing D.T., Pochi P.E., Strauss J.S.: Anatomical variation in the amount and composition of human surface lipid. J Invest Dermatol 1970; 54: pp. 240-247

Grove G.L., Lavker R.M., Holze E., Kligman A.M.: The use of nonintrusive tests to monitor age-associated changes in human skin. J Soc Cosmet Chem 1970; 54: pp. 240-247

Grubauer G., Elias P.M., Feingold K.R.: Lipid content and lipid type as determinants of the epidermal permeability barrier. J Lipid Res 1989; 30: pp.89-96

Grubauer G., Elias P.M., Feinglod K.R.: Transepidermal water loss: the signal for recovery of barrier structure and function. J Lipid Res; 1989 30 (3): pp. 323-333

Habermehl K.H.: Haut und Hautorgane. In: Nickel R., Schummer A., Seiferle E.: Lehrbuch der Anatomie der Haussäugetiere. Band III Kreislaufsystem, Haut und Hautorgane. 3. Auflage Blackwell Wissenschaftsverlag Berlin, Wien, 1996: pp. 443-507

Hadgraft J., Walters K.A., Guy R.H.: Epidermal lipids and topical drug delivery. Semin Dermatol 1992; 11 (2): pp. 139-144

Hadgraft J.: The epidermal reservoir: A theoretical approach. Internat J Pharmacol 1979; 2: pp. 265-274

Hamanaka S., Hara M., Nishio H., Otsuka F., Suzuki A., Uchida Y.: Human epidermal glycosylceramides are major precursors of stratum corneum ceramides. J Invest Dermatol 2002; 119: pp. 416-423

Harkey M.R.: Anatomy and physiology of hair. Forensic Science International 1993; 63: pp. 9-18

Hawkins G.S., Reifenrath W.G.: Influence of skin source, penetration cell fluid, and partition coefficient on in vitro skin penetration. J Pharm Sci 1986, 75: pp.378-381

Hedberg C.L., Wertz P.W., Downing D.T.: The time course of lipid biosynthesis in pig epidermis. J Invest Dermatol 1988; 91: pp. 169-174

Hieronymy E.: Haut. In: Pathologie der Laboratoriumstiere. I. Cohrs P., Jaffé R., Meessen R. (Hrsg.) Springer Verlag Berlin, 1958: pp. 587-599

Hoffmann R.: Hormonal interaction and hair growth. Ann Dermatol Venerol 2002; 129 (5): pp. 782-792

Holleran W.M., Ginns E.I., Menon G.K. et al: Consequences of beta-glucocerebrosidase deficiency in epidermis. Ultrastructure and permeability alterations in Gaucher disease. J Clin Invest 1994; 93: pp. 1756-1764

Holleran W.M., Uchida Y., Halkier-Sorensen L., Haratake A., Hara M., Epstein J.H., Elias P.M.: Structural and biochemical basis for the UVB-induced alterations in epidermal barrier functions. Photodermat Photoimmunol Photomed 1997; 13 (4): pp. 117-128

Holmes R.L., Williams M., Cuncliffe W.J.: Pilo-Sebaceous duct obstruction and acne. Br J Dermatol 1972; 87: pp. 327-332

Horii I., Nakayama Y., Obata M., Tagami H.: Stratum corneum hydration and amino acid content in xerotic skin. Br J Dermatol Res 1989; 121: pp. 588-592

Hou S.Y.E., Mitra A.K., White S.H., Menon G.K., Ghadially R., Elias P.M.: Membrane structures in normal and essential fatty acid-deficient stratum corneum: Characterization by ruthenium tetroxide staining and X-ray diffraction. J Invest Dermatol 1991; 96: 215-223

Hueber F., Besnard M., Schaefer H., Wepierre J.: Percutaneous absorption of estradiol and progesterone in normal and appendage-free skin of the hairless rat: Lack of importance of nutritial blood flow. Skin Pharmacol 1994; 7: pp. 245-256

Hueber F., Wepierre J., Schaefer H.: Role of transepidermal and transfollicular routes in percutaneous absorption of hydrocortisone and testosterone: In vivo study in the hairless rat. Skin Pharmacol 1992; 5: pp. 99-107

Hussein M.A.F.: The overall pattern of hair follicle arrangement in the rat and mouse. J Anat 1971; 109 (2): pp. 307-316

Illel B.: Formulation of transfollicular drug administration: Some recent advances. Critical Reviews in Therapeutic Drug Carrier Systems 1997; 14 (3): pp. 207-219

Illel B., Schaefer H., Wepierre J., Doucet O.: Follicles play an important role in percutaneous absorption. J Pharm Sci 1991; 80: pp. 424-247

Illel B., Schaefer H.: Transfollicular percutaneous absorption. Skin model for quantitative studies. Acta Derm Venereol 1988; 68 (5): pp. 427-430

Imokawa G., Akasaki S., Hatotori M., Yoshizuka N.: Selective recovery of deranged water-holding properties by stratum corneum lipids. J Invest Dermatol 1986; 87: pp. 758-761

Ingram D.L., Weaver M.E.: A quantitative study of the blood vessels of the pig's skin and the influence of environmental temperature. Anat Rec 1969, 16: pp. 517-524

de Jagger M.W., Gooris G.S., Dolbnya I.P., Ponec M., Bouwstra J.A.: Modelling the stratum corneum lipid organisation with synthetic lipid mixtures: the importance of synthetic ceramide composition. Biochimica et Biophysica Acta 2004; 1684: pp. 132-140

Jung S., Otberg N., Thiede G., Richter H., Sterry W., Panzner S., Lademann J.: Innovative liposomes as a transfollicular drug delivery system: penetration into porcine hair follicles. *J Invest Dermatol* 2006; 19 (3): pp. 159-167

Junqueira L.C., Caneiro J.: Histologie. Springer Verlag Berlin Heidelberg, New York, 1996: pp. 412-417

Kao J., Hall J., Helman G.: In vitro percutaneous absorption in mouse skin: influence of skin appendages. *Toxicol Appl Pharmacol* 1988; 94 (1): pp. 93-103

Kolb G.M.: Vergleichende Histologie, Zytologie und Mikroanatomie der Tiere. Springer Verlag Berlin, Heidelberg 2. Auflage 1991: pp. 176-195

Kristensen S.: A study of skin diseases in dogs and cats. I Histology of the hairy skin of dogs and cats. *Nord Vet-Med* 1975, 27: pp. 593-603

Küster W., Fritsch P., Happel R.: Hereditäre Verhorngungsstörungen. In: Fritsch P. (ed): Dermatologie, Venerologie – Grundlagen. Klinik. Atlas. 2. Auflage, Berlin, Heidelberg, New York, Springer 2004: pp. 550-556

Lademann J., Otberg N., Richter H., Lindemann U., Weigmann H.-J., Schaefer H., Sterry W.: Investigation of follicular penetration of topically applied substances. *Skin Pharmacol Appl Skin Physiol* 2001; 14 [Suppl 1]: pp.17-22

Lademann J., Weigmann H., Rickmeyer C., Barthelmes H., Schaefer H., Mueller G., Sterry W.: Penetration of titanium dioxide microparticles in a sunscreen formulation into the horny layer and the follicular orifice. *Skin Pharmacol Appl Skin Physiol* 1999; 12 (5): pp. 247-256

Lademann J., Richter H., Schaefer U.F., Blume-Peytavi U., Teichmann A., Otberg N., Sterry W.: Hair follicles – A long time reservoir for drug delivery. *Skin Pharmacol Physiol* 2006; 19 (4): pp. 232-236

Lademann J., Otberg N., Richter H., Jacobi U., Schaefer H., Blume-Peytavi U., Sterry W.: Follikuläre Penetration. Ein entscheidender Penetrationsweg von topisch applizierten Substanzen. *Hautarzt* 2003; 54: pp. 321-323

Lampe M.A., Burlingame A.L., Whitney J.A., Williams M.L., Brown B.E., Roitman E., Elias P.M.: Human stratum corneum lipids: characterization and regional variations. *J Lipid Res* 1983; 24: pp. 120-130

Landmann L.: Epidermal permeability barrier: transformation of lamellar-granule disks into intercellular sheets by a membrane-fusion process: A freeze fracture study. *J Invest Dermatol* 1986; 87: pp. 202-209

Landmann L.: Lamellar granules in mammalian, avian and reptilian epidermis. *J Ultrastr Res* 1988; 72: pp. 245-263

Landmann L.: Das Lipidkonzept der Permeabilitätsbarriere. In Landmann L.: Permeabilitätsbarriere der Epidermis. Grosse Scripta 9, Grosse Verlag Berlin, 1985: pp. 157-169

Landmann L.: The epidermal permeability barrier. *Anat Embryol* 1988; 178: pp. 1-13

Lavrijsen A.P.M., Higoumec I.M., Weerheim A., Oestmann E., Tuinenburg E.E., Boddé H.E., Ponec M.: Validation of an in vivo extraction method for stratum corneum ceramides. *Arch Dermatol Res* 1994; 286: pp. 495-503

Law S., Wertz P.W., Swatzendruber D.C., Squier C.A.: Regional variation in content, composition and organization of porcine epithelial barrier lipids revealed by thin layer chromatography and transmission electron microscopy. *Arch Oral Biol* 1995; 40 (20): pp.1085-1091

Lauer A.C., Lieb L.M., Ramachandran G.L.F., Norman D.W.: Transfollicular drug delivery. *Pharm Res* 1995 12 (2): pp. 179-186

Lieb L.M., Flynn G., Weiner N.: Follicular (pilosebaceous unit) deposition and pharmacological behaviour of cimetidine as a function of formulation. *Pharm Res* 1994; 11 (10): pp. 14-19

Liebich H.G.: Funktionelle Histologie der Haussäugetiere. 3. Auflage, Verlag Schattauer, Stuttgart, New York, 1999: pp. 303-315

Lloyd D.H., Garthwaite G.: Epidermal structure and surface topography of canine skin. *Res Vet Sci* 1982; 33 (1): pp. 99-104

Long S.A., Wertz P.W., Strauss J.S., Downing D.T.: Human stratum corneum polar lipids and desquamation. *Arch Dermatol Res* 277, 1984: pp. 284-287

Madison K.C.: Barrier function of the skin: "La Raison d'être" of the epidermis? J. Invest Dermatol, 2003; 121 (2): pp. 231-241

Madison KC., Swartzendruber D.C., Wertz P.W., Downing D.T.: Lamellar granule extrusion and stratum corneum intercellular lamellae in murine keratinocyte cultures. J Invest Dermatol 1988; 90: pp. 110-116

Magnusson B.M., Walters, K.A., Roberts, M.S.: Veterinary drug delivery: Potential for skin penetration enhancement. Advanced Drug Delivery Reviews 2001; 50: pp. 206-227

Maibach H.I., Feldmann R.J., Melby T.H., Serat W.F.: Regional variation in percutaneous penetration in man. Pesticides Arch Environ Health 1971; 23: pp. 208-211

Mangeldorf S., Otberg N., Maibach H.I., Sinkgraven R., Sterry W., Lademann J.: Ethnic variation in vellus hair follicle size and distribution. Skin Pharmacol Physiol 2006; 19 (3): pp. 159-167

Marks R., Dawber R.P.R.: Skin suface biopsy: An improved technique for the examination of the horny layer. Br J Derm 1971; 84: p.117

Marks R., Barton S.P.: The significance of size and shape of corneocytes. In Marks R., Plewig G. (eds); Stratum corneum. Berlin, Springer, 1983, pp. 175-180

Marzulli F.N., Brown D.W.C., Maibach H.I.: Techniques for studying skin penetration. Toxicol Appl Pharmcol Suppl 1969; 3: 79-83

Mc Geesh A.H.: Percutaneous toxicity. Toxicol Appl Pharmacol Suppl 1965; 2: pp.20-26

Melnik B., Hollman J., Erler E., Verhoeven B., Plewig G.: Microanalytical screening of all major stratum corneum lipids by sequential high-performance-thin-layer-chromatography. J Invest Dermatol 1988; 92: pp: 231-234

Menczel E.: Skin Delipidization and percutaneous absorption. In: Bronaugh R.L., Maibach H.I. (eds): Percutaneous absorption. Drugs, Cosmetics, Mechanisms, Methodology. Marcel Dekker, inc. New York, Basel 1999: pp. 133-139

Menon G.K., Feingold K.R., Moser A.H., Brown B.E., Elias P.M.: De novo sterogenesis in the skin. II Regulation by cutaneous barrier requirements. J Lipid Res 1985; 26: pp. 428-427

Meyer W., Schwarz R., Neurand K.: The skin of domestic mammals as a model for the human skin with special reference to the domestic pig. Curr Probl Dermatol Karger Basel 1978; Vol 7: pp.39-52

Meyer W., Zschemisch N.H., Neurand K.: Untersuchungen zur Struktur der Ohrhaut beim Hausschwein, unter besonderer Berücksichtigung einer Verwendung in der humandermatologischen Forschung. 1. Makroskopische Aspekte und Haarkleid. Berl. Münch. Tierärztl. Wschr. 2001; 114: pp. 92-99

Meyer W.: Bemerkungen zur Eignung der Schweinehaut als biologisches Model für die Haut des Menschen. Hautarzt 1996; 47: pp. 178-182

Meyer W.: Die Haut des Schweines. Schlütersche Verlagsanstalt und Druckerei Hannover, 1986,

Meyer W., Hülmann G., Seger H.: SEM-Atlas on the Hair Cuticle Structure of Central European Mammals. Verlag M. & H. Schaper Alfeld – Hannover, 2002

Meyer W., Schwarz R., Neurand K.: Die Haut der Säugetiere. I. Ihre Bedeutung für die dermatologische Forschung – Grundzüge der vergleichenden Morphologie. Tierärztl. Praxis 1978; 6: pp. 153-162

Militzer K.: Haut und Hautanhangsorgane kleiner Laboratoriumssäugetiere. Verlag Paul Parey Berlin und Hamburg, 1982

Mils V., Vincent C., Croute F., Serre G.: The expression of desmosomal and corneodesmosomal antigens shows specific variations during terminal differentiation of epidermis and hair follicle epithelia. J Histochem Cytochem 1992; 40: pp. 1329-1337

Mills O.H., Kligmann A.M.: The follicular biopsy. Dermatologica 1983; 167: pp. 57-63

Montagna W., Hu F.: Advances in biology of skin. Vol. VIII The pigmentary system. Oxford: Pergamon, 1967

Montagna W., Parakkal P.F.: The structure and function of skin. 3. Aufl. New York: Academic press, 1974

Montagna W.: Penetration and local effect of Vitamin A on the skin of the guinea pig. Proc Soc Exp Biol Med 1954; 86: pp. 668-672

Monteiro-Riviere N.A., Bristol D.G., Manning T.O., Rogers R.A., Riviere J.E.: Interspecies and interregional analysis of the comparative histologic thickness and laser Doppler blood flow measurements at five cutaneous sites in nine species. *J Invest Dermatol* 1990; 95: pp. 582-586

Nicolaides N., Fu H.C., Ansari M.N.A.: Diester waxes in surface lipids of animal skin. *Lipids* 1970; 5 (3): pp. 299-307

Nicolaides N., Fu H.C., Rice G.R.: The skin surface lipids of man compared with those of eighteen species. *J Invest Dermatol* 1968; 51: pp. 83-89

Norlén L., Nicander I., Rozell B.L., Ollmar S., Forslind B.: Inter- and intraindividual differences in human stratum corneum lipid content related to physical parameters of skin barrier function in vivo. *J Invest Dermatol* 1999; 112 (1): pp.72 -77

Norlén L.: Skin barrier formation. The membrane folding model. *J Invest Dermatol* 2001a; 117: pp. 823-829

Norlén L.: Skin barrier structure and function: The single gel phase model. *J Invest Dermatol* 2001b; 117: pp. 830-836

Norlén L.: Skin barrier structure and function: The single gel phase model. *J Invest Dermatol* 2001; 117: pp. 830-836

Oberste-Lehn H., Nobis A.: Die Haaranordnung beim Menschen und bei einigen Säugetieren. *Z Anat Entwicklungsgesch* 1963; 123: pp. 589-642

Otberg N., Richter H., Schaefer H., Blume-Peytavi U., Sterry W., Lademann J.: Variations of hair follicle size and distribution in different body sites. *J Invest Dermatol* 2004;122: pp. 14-19

Otberg N., Richter H., Knuttel A., Schaefer H., Sterry W., Lademann J.: Laser spectroscopic methods for the characterization of open and closed follicles. *Laser Phys Lett* 2004; 1 (1): pp. 46-49

Pagnoni A., Kligman A.M., Gammal S. EL, Stoudemayer T.: Determination of density of follicles on various regions of the face by cyanoacrylate biopsy: correlation with sebum output. *Br J Dermatol* 1994; 131: pp. 862-865

Panchagnula R., Stemmer K., Ritschel W.A.: Animal models for transdermal drug delivery. Meth Find Exp Clin Pharmacol 1997; 19 (5): pp. 335-341

Pinkus F.: Über die Hautsinnesorgane neben dem menschlichen Haar (Haarscheiben) und ihre vergleichend-anatomische Bedeutung. Arch Mikroskop Anat 65, 1905: pp.121-179

Plewig G. Kligman M.: Sampling of sebaceous follicles by the cyanoacralat technique. In: Acne: Morphogenesis and treatment. Springer Verlag Berlin, Heidelberg, New York 1975: p.56

Pochi P.E., Strauss J.S., Downing D.T.: Age-related changes in sebaceous gland activity. J Invest Dermatol 1979; 73 (1): 108-111

Ponec M., Weeheim A., Lankhorst P., Wertz P.W.: New acylceramide in native and reconstructed epidermis. J Invest Dermatol 2003; 120: pp. 581-588

Ponec M., Boelsma E., Gibbs S., Mommaas M.: Characterization of reconstructed skin models. Skin Pharmacol Appl Skin Physiol 2002; 15, suppl.I: pp. 4-17

Potts R.O., Francoeur M.L.: The influence of stratum corneum morphology on water permeability. J Invest Dermatol 1991; 96: pp. 495-499

Priestley G.C.: Histological studies of the skin follicle types of the rat with special reference to the structure of the Huxley layer. J Anat 1967; 101: pp. 491-504

Randall V.A., Sundberg J.P., Philpott M.P.: Animal and in vitro models for the study of hair follicles. J Invest Dermatol 2003; 8 (1), pp 39-45

Riviere J.E. , Papich M.G.: Potential and problems of developing transdermal patches for veterinary applications. Advanced Drug Delivery Reviews 2001; 50: 175-203

Rougier A., Dupuis D., Lotte C., Roguet R.: The measurement of the stratum corneum reservoir. A predictive method for in vivo percutaneous absorption studies: Influence of application time. J Invest Dermatol 1985; 84: p. 66

Ryder M.: Skin. In: Studies in Biology. London: Arnhold 1973; 41: pp. 1-56

Salaman R.N.: The inheritance of fur types and hair characters in rabbits. J Genet 1922; 12: pp. 179-209

Saint-Leger D.: Physiology of the pilosebaceous follicle. Rev Prat 1994; 43: pp. 2315-2319

Sarasola P., Jernigan A.D., Walker D.K., Castledine J., Smith D.G., Rowan T.G.: Pharmakokinetics of selamectin following intravenous, oral and topical administration in cats and dogs. J Vet Pharmacol Ther. 2002; 25 (4): pp. 265-272

Sato K., Sugibayashi K., Morimoto Y.: Species differences in percutaneous absorption of Nicorandil. J Pharm Sci 1991; 80 (2): pp. 104-107

Schäfer H., Lademann J.: The role of follicular penetration. Skin Pharmacol Appl Skin Physiol 2001; 14 (Suppl. 1): pp. 23-27

Schaefer H., Watts F., Brod J., Illel B.,: Follicular penetration in prediction of percutaneous penetration. Vol. 2 in Scott R.C., Hadgraft J. (eds) IBC London 1990, p. 103

Schäfer H., Redelmaier T.E. (eds): Skin Barrier. Principles of Percutaneous Absorption. Karger Basel, 1996

Scheuplein R.J.: Mechanism of percutaneous absorption. J Invest Dermatol Vol 1967; 48 (1): pp. 79-88

Scheuplein R.J., Blank I.H. : Permeability of skin. Physiol Rev 1971; 51(4): pp. 702-747

Schwarz R., Le Roux J.M., Neurand K.: Lichtmikroskopische Untersuchungen an der Haut der Katze. Zbl. Vet. Med. C. Anat Histol Embryol 1976; 5: pp. 325-341

Scott R.C., Corrigan M.A., Smith F., Mason H.: The influence of skin structure on permeability: An intersite and interspecies comparison with hydrophilic penetrants. J Invest Dermatol 1990; 96 (6): pp. 921-926

Scott I.R., Harding C.R., Barrett J.G.: Histidine-rich protein of the keratohyalin granules: Source of free amino acids, urocanic acid and pyrrolidone carboxylic acid in the stratum corneum. Biochim Biophys Acta 1982; 719: pp. 110-117

Scott I.R., Harding C.R.: Filaggrin breakdown to water binding compounds during development of the rat stratum corneum is controlled by the water activity of the environment. *Dev Biol* 1986; 115: pp. 84-92

Seelig J., Seelig A.: Lipid conformation in model membranes and biological membranes. *Q Rev Biophys* 1980; 13: pp. 19-61

Shapiro L.J., Weiss R., Webster D., France J.T.: X-linked Ichthyosis due to steroid sulfatase deficiency. *Lancet* I, 1978, pp.70-72

Sidransky E., Sherer D.M., Ginns E.I.: Gaucher disease in the neonate. A distinct Gaucher phenotype is analogous to a mouse model created by targeted disruption of beta-glucocerebrosidase gene. *Pediat Res* 1992; 32: pp. 494-498

Slee J.: Developmental morphology of the skin and hair follicles in normal and in ragged mice. *J Embryol Exp Morphol* 1962; 10: pp. 507-529

Sperling L.C.: Hair anatomy for the clinician. *J Amer Acad Dermatol* 1991; 25: pp. 1-17

Squier C.A., Cox P., Wertz P.W.: Lipid content and water permeability of skin and oral mucosa. *J Invest Dermatol* 96, 1991: pp. 123-126

Stevens A., Lowe J.: In: *Histologie*. © VHC Verlagsgesellschaft mbH. Weinheim, Basel, Cambridge, New York, 1992: pp. 349-362

Steward M.E.: Sebaceous gland lipids. In: Bereiter-Hahn J., Matoltsy A.G., Richards K.S.: *Biology of the Integument. Vertebrates*. Springer Verlag Berlin, Heidelberg, New York, Tokyo 1986; pp. 824-831

Steward M.E., Downing D.T.: A new 6-hydroxy-4-sphingenine-containing ceramide in human skin. *J Lipid Res* 1999; 40: pp. 1434-1439

Straile W.E.: Sensory hair follicles in mammalian skin: The tylotrich follicle. *Am J Anat* 1960; 106: pp. 133-147

Straile W.E.: The morphology of tylotrich follicles in the skin of the rabbit. *Am J Anat* 1961; 109: pp.1-14

Strickland J.H., Calhoun M.L.: The integumentary system of the cat. Am vet Res 1963; 24: pp. 1018-1029

Swartzendruber D.C., Burnett I.H., Wertz P.W., Madison K.C., Squier C.A.: Osmium tetroxide and rhuthenium tetroxide are complementary agents for the preparation of epidermal samples for transmission electron microscopy. J Invest Dermatol 1995; 104: pp. 417-420

Tayar E.L., Tsai N., Testa R.-S., Carrupt P.-A., Hansch C., Leo A.: Percutaneous penetration of drugs: A quantitative structure-permeability relationship study. J Pharm Sci 1991; 80: pp. 744-749

Tenjarla S., Puranajoti P., Kasina R., Mandal T.: Terbutaline transdermal delivery: formulation studies and limitations of in-vitro predictive parameters. J Pharm Pharmacol 1996; 48 (11): pp. 1138-1142

Thody A.J., Shuster S.: Control and functions of sebaceous lipids. Physiol Rev 1989; 69 (2): pp.383-416

Toellner R.: Illustrierte Geschichte der Medizin. Verlag Andreas und Andreas 1992; Band 3: pp. 1513-1540

Toll R., Jacobi U., Richter H., Lademann J., Schaefer H., Blume-Peytavi U.: Penetration profile of microspheres in follicular targeting of terminal hair follicles. J Invest Dermatol 2004; 123 (1); pp. 168-176

Tregear R.T.: Physical function of the skin. Academic, New York 1966

Tur E., Maibach H.I., Guy R.H.: Percutaneous penetration of Methyl Nicotinate at three anatomic sites: Evidence for an appendageal contribution to transport? Skin Pharmacol 1991; 4, pp. 230-234

Warner R.R., Myers M.C., Taylor D.A.: Electron probe analysis of human skin: Determination of the water concentration profile. J Invest Dermatol 1988; 90: pp. 218-224

Weirich E.G., Rufli T., Longauer J.K.: Formationsmuster des Hautepithels bei Mensch und Meerschweinchen. Dermatologica 1976; 152: pp. 263-269

Williams M.L., Elias P.M.: The extracellular matrix of stratum corneum: Role of lipids in normal and pathological function. CRC Crit Rev Ther Drug Carrier Syst 1887; 3: pp. 95-122

Wester R.C., Maibach H.I.: Animal models for transdermal drug delivery. In: Kydonieus A.F., Berner B.: Transdermal Delivery of Drugs. CRC Press Boca Raton, Florida 1987: pp: 61-70

Wester R.C., Maibach H.I.: In vivo animal models for percutaneous absorption. In: Bronaugh R.L., Maibach H.I. (eds) Percutaneous Absorption. Drugs, Cosmetics, Mechanisms, Methodology. Marcel Dekker inc. New York, Basel; 1999: pp. 251-265

Wertz P.W. and Downing D.T.: Integral lipids of human hair. Lipids 1988; 23: pp. 878-881

Wertz P.W. and Downing D.T.: Integral lipids of mammalian hair. Comp Biochem Physiol (B) 1989; 92: pp. 759-761

Wertz P.W.: Lipids of keratinizing tissues. In In: Bereiter-Hahn J., Matoltsy A.G., Richards K.S.: Biology of the Integument. Vertebrates. Springer Verlag Berlin, Heidelberg, New York, Tokyo 1986; pp. 815-823

Wertz P.W., Abraham W., Landmann L., Downing D.T.: Preparation of liposomes from stratum corneum lipids. J Invest Dermatol 1986; 87: pp. 582.584

Wertz P.W. and Downing D.T.: Ceramides of pig epidermis: Structure and determination. J Lipid Res 1983; 24: pp. 759-765

Wertz P.W. and Downing D.T.: Metabolism of linoleic acid in porcine epidermis. J Lipid Res 1990; 31: pp. 839-844

Wertz P.W., Downing D.T., Freinkel R.K., Traczyk T.N.: Sphingolipids of the stratum corneum and lamellar granules of fetal rat epidermis. J Invest Dermatol 1984; 83: pp. 193-195

Wertz P.W., Downing D.T.: Epidermal lipids. In: Goldsmith LA (ed): Physiology, Biochemistry and Molecular Biology of the Skin. 2.ed, Oxford, Oxford University Press 1991: pp.205-235

Wertz P.W., Swartzendruber D.C., Madison K.C., Downing D.T.: Composition and morphology of epidermal cyst lipids. J Invest Dermatol 1987; 89: pp. 419-424

Wertz P.W.: Origin of the species specific lipids in the HPTLC. 2005; persönliche Kommunikation

Wilken K.: Penetration von Pollenallergenen in die Haut. Dissertation Charité Berlin, 2004

Yamamura T., Tezuka T.: The water-holding capacity of the stratum corneum as measured by ¹H-NMR. J Invest Dermatol 1989; 93: p. 342

Yardley H.J.: Epidermal lipids. In: Goldsmith LA (ed): Biochemistry and Physiology of the Skin. New York, Oxford University Press 1983: p.363

Yeagle P.L.: Cholesterol and the cell membrane. Biochim. Biophys. Acta 1985; 822: pp. 267-287

Zellmer S., Lasch J.: Individual variation of human plantar stratum corneum lipids, determined by automated multiple development of high performance thin-layer chromatography plates. J Chromatogr B 1997, 691: p. 321

von Zglinicki T., Lindberg M., Roomans G.M., Forslind B.: Water and ion distribution profiles in human skin. Acta Derm Venerol (Stockh.) 1993; 73: pp. 340-343

Zirra A.-M.: The functional significance of the skin's stratum lucidum. Rev Roum Morphol Embryol Physiol 1976; 26: pp. 9-12

Zouboulis C.C.: Acne and sebaceous gland function. Clin Dermatol 2004, 22 (5): pp. 360-366