

7 Literaturverzeichnis

- Abriel, H., J. Loffing, J. F. Rebhun, J. H. Pratt, L. Schild, J. D. Horisberger, D. Rotin und O. Staub (1999). Defective regulation of the epithelial Na⁺ channel by Nedd4 in Liddle's syndrome. *J Clin Invest* 103(5): 667-73.
- Aidley, D. J. und P. R. Stanfield (1996). Ion channels: molecules in action. Cambridge University Press.
- Al-Awqati, Q. (1986). Proton-translocating ATPases. *Annu Rev Cell Biol* 2: 179-99.
- Alvarez de la Rosa, D., C. M. Canessa, G. K. Fyfe und P. Zhang (2000). Structure and regulation of amiloride-sensitive sodium channels. *Annu Rev Physiol* 62: 573-94.
- Ashcroft, F. M. (2000). Ion Channels and Disease. Academic Press.
- Ausubel, F. M., R. Brent, R. E. Kingston, D. D. Moore, J. D. Seidman, J. A. Smith und K. Struhl (1993). Current protocols in molecular biology. New York: J. Wiley & Sons: 13.0.1-13.13.9.
- Barhanin, J., F. Lesage, E. Guillemaire, M. Fink, M. Lazdunski und G. Romeo (1996). KvLQT1 and IsK (minK) proteins associate to form the I_{Ks} cardiac potassium current. *Nature* 384(6604): 78-80.
- Bauer, C. K., K. Steinmeyer, J. R. Schwarz und T. J. Jentsch (1991). Completely functional double-barreled chloride channel expressed from a single Torpedo cDNA. *Proc Natl Acad Sci U S A* 88(24): 11052-6.
- Biervert, C., B. C. Schroeder, C. Kubisch, S. F. Berkovic, P. Propping, T. J. Jentsch und O. K. Steinlein (1998). A potassium channel mutation in neonatal human epilepsy. *Science* 279(5349): 403-6.
- Birnboim, H. C. und J. Doly (1979). A rapid alkaline extraction procedure for screening recombinant plasmid DNA. *Nucleic Acids Res* 7(6): 1513-23.
- Bixby, K. A., M. H. Nanao, N. V. Shen, A. Kreusch, H. Bellamy, P. J. Pfaffinger und S. Choe (1999). Zn²⁺-binding and molecular determinants of tetramerization in voltage-gated K⁺ channels. *Nat Struct Biol* 6(1): 38-43.
- Borsani, G., E. I. Rugarli, M. Taglialatela, C. Wong und A. Ballabio (1995). Characterization of a human and murine gene (CLCN3) sharing similarities to voltage-gated chloride channels and to a yeast integral membrane protein. *Genomics* 27(1): 131-41.
- Brandt, S. und T. J. Jentsch (1995). CIC-6 and CIC-7 are two novel broadly expressed members of the CLC chloride channel family. *FEBS Lett* 377(1): 15-20.
- Bretag, A. H. (1987). Muscle chloride channels. *Physiol Rev* 67(2): 618-724.
- Brown, D. A. und P. R. Adams (1980). Muscarinic suppression of a novel voltage-sensitive K⁺ current in a vertebrate neurone. *Nature* 283(5748): 673-676.
- Bullock, W. O., J. M. Fernandez und J. M. Short (1987). XL1-blue - a high efficiency plasmid transforming *Escherichia coli* strain with beta-galactosidase selection. *Biotechniques* 5: 376-378.
- Buyse, G., D. Trouet, T. Voets, L. Missiaen, G. Droogmans, B. Nilius und J. Eggermont (1998). Evidence for the intracellular location of chloride

- channel (ClC)-type proteins: co-localization of ClC-6a and ClC-6c with the sarco/endoplasmic-reticulum Ca²⁺ pump SERCA2b. *Biochem J* 330: 1015-21.
- Canessa, C. M., J. D. Horisberger und B. C. Rossier (1993). Epithelial sodium channel related to proteins involved in neurodegeneration. *Nature* 361(6411): 467-70.
- Canessa, C. M., L. Schild, G. Buell, B. Thorens, I. Gautschi, J. D. Horisberger und B. C. Rossier (1994). Amiloride-sensitive epithelial Na⁺ channel is made of three homologous subunits. *Nature* 367(6462): 463-467.
- Charlier, C., N. A. Singh, S. G. Ryan, T. B. Lewis, B. E. Reus, R. J. Leach und M. Leppert (1998). A pore mutation in a novel KQT-like potassium channel gene in an idiopathic epilepsy family. *Nat Genet* 18(1): 53-55.
- Chen, H. I. und M. Sudol (1995). The WW domain of Yes-associated protein binds a proline-rich ligand that differs from the consensus established for Src homology 3-binding modules. *Proc Natl Acad Sci U S A* 92(17): 7819-23.
- Chen, M. S., R. A. Obar, C. C. Schroeder, T. W. Austin, C. A. Poodry, S. C. Wadsworth und R. B. Vallee (1991). Multiple forms of dynamin are encoded by shibire, a *Drosophila* gene involved in endocytosis. *Nature* 351(6327): 583-6.
- Clague, M. J., S. Urbe, F. Aniento und J. Gruenberg (1994). Vacuolar ATPase activity is required for endosomal carrier vesicle formation. *J Biol Chem* 269(1): 21-4.
- Cole, K. S. (1979). Mostly membranes (Kenneth S. Cole). *Annu Rev Physiol* 41: 1-24.
- Cooper, E. C., K. D. Aldape, A. Abosch, N. M. Barbaro, M. S. Berger, W. S. Peacock, Y. N. Jan und L. Y. Jan (2000). Colocalization and coassembly of two human brain M-type potassium channel subunits that are mutated in epilepsy. *Proc Natl Acad Sci U S A* 97(9): 4914-9.
- Covarrubias, M., A. A. Wei und L. Salkoff (1991). Shaker, Shal, Shab, and Shaw express independent K⁺ current systems. *Neuron* 7(5): 763-73.
- Cox, J. P., K. Yamamoto, P. T. Christie, C. Wooding, T. Feest, F. A. Flinter, P. R. Goodyer, E. Leumann, T. Neuhaus, C. Reid, P. F. Williams, O. Wrong und R. V. Thakker (1999). Renal chloride channel, CLCN5, mutations in Dent's disease. *J Bone Miner Res* 14(9): 1536-42.
- Damke, H., T. Baba, D. E. Warnock und S. L. Schmid (1994). Induction of mutant dynamin specifically blocks endocytic coated vesicle formation. *J Cell Biol* 127(4): 915-34.
- Davis, C. G., M. A. Lehrman, D. W. Russell, R. G. Anderson, M. S. Brown und J. L. Goldstein (1986). The J.D. mutation in familial hypercholesterolemia: amino acid substitution in cytoplasmic domain impedes internalization of LDL receptors. *Cell* 45(1): 15-24.
- De Duve, C. (1963). Lysosomes. In: Ciba Foundation Symposium: 411-412.
- Devuyst, O., P. T. Christie, P. J. Courtois, R. Beauwens und R. V. Thakker (1999). Intra-renal and subcellular distribution of the human chloride channel, CLC-5, reveals a pathophysiological basis for Dent's disease. *Hum Mol Genet* 8(2): 247-57.
- Doyle, D. A., J. Morais Cabral, R. A. Pfuetzner, A. Kuo, J. M. Gulbis, S. L. Cohen, B. T. Chait und R. MacKinnon (1998). The structure of the potassium channel: molecular basis of K⁺ conduction and selectivity. *Science* 280(5360): 69-77.
- Duan, D., C. Winter, S. Cowley, J. R. Hume und B. Horowitz (1997). Molecular identification of a volume-regulated

- chloride channel. *Nature* 390(6658): 417-21.
- Duc, C., N. Farman, C. M. Canessa, J. P. Bonvalet und B. C. Rossier (1994). Cell-specific expression of epithelial sodium channel alpha, beta, and gamma subunits in aldosterone-responsive epithelia from the rat: localization by *in situ* hybridization and immunocytochemistry. *J Cell Biol* 127(6 Pt 2): 1907-21.
- Durell, S. R. und H. R. Guy (1992). Atomic scale structure and functional models of voltage-gated potassium channels. *Biophys J* 62(1): 238-47; discussion 247-50.
- Fahlke, C., T. Knittle, C. A. Gurnett, K. P. Campbell und A. L. George, Jr. (1997). Subunit stoichiometry of human muscle chloride channels. *J Gen Physiol* 109(1): 93-104.
- Fahlke, C., H. T. Yu, C. L. Beck, T. H. Rhodes und A. L. George, Jr. (1997). Pore-forming segments in voltage-gated chloride channels. *Nature* 390(6659): 529-32.
- Firsov, D., I. Gautschi, A. M. Merillat, B. C. Rossier und L. Schild (1998). The heterotetrameric architecture of the epithelial sodium channel (ENaC). *Embo J* 17(2): 344-52.
- Firsov, D., L. Schild, I. Gautschi, A. M. Merillat, E. Schneeberger und B. C. Rossier (1996). Cell surface expression of the epithelial Na channel and a mutant causing Liddle syndrome: a quantitative approach. *Proc Natl Acad Sci U S A* 93(26): 15370-5.
- Fisher, S. E., G. C. Black, S. E. Lloyd, E. Hatchwell, O. Wrong, R. V. Thakker und I. W. Craig (1994). Isolation and partial characterization of a chloride channel gene which is expressed in kidney and is a candidate for Dent's disease (an X-linked hereditary nephrolithiasis). *Hum Mol Genet* 3(11): 2053-9.
- Fisher, S. E., I. van Bakel, S. E. Lloyd, S. H. Pearce, R. V. Thakker und I. W. Craig (1995). Cloning and characterization of CLCN5, the human kidney chloride channel gene implicated in Dent disease (an X-linked hereditary nephrolithiasis). *Genomics* 29(3): 598-606.
- Forgac, M. (1992). Structure and properties of the coated vesicle (H⁺)-ATPase. *J Bioenerg Biomembr* 24(4): 341-50.
- Forgac, M. (1996). Regulation of vacuolar acidification. *Soc Gen Physiol Ser* 51: 121-32.
- Friedrich, T., T. Breiderhoff und T. J. Jentsch (1999). Mutational analysis demonstrates that CIC-4 and CIC-5 directly mediate plasma membrane currents. *J Biol Chem* 274(2): 896-902.
- Gekle, M., S. Mildenberger, R. Freudinger und S. Silbernagl (1995). Endosomal alkalinization reduces Jmax and Km of albumin receptor-mediated endocytosis in OK cells. *Am J Physiol* 268(5 Pt 2): F899-906.
- Goldman, D. E. (1943). *J. Gen. Physiol.* 27: 37-60.
- Gründer, S., A. Thiemann, M. Pusch und T. J. Jentsch (1992). Regions involved in the opening of CIC-2 chloride channel by voltage and cell volume. *Nature* 360(6406): 759-62.
- Gulbis, J. M., M. Zhou, S. Mann und R. MacKinnon (2000). Structure of the cytoplasmic beta subunit-T1 assembly of voltage-dependent K⁺ channels. *Science* 289(5476): 123-7.
- Günther, W., A. Lüchow, F. Cluzeaud, A. Vandewalle und T. J. Jentsch (1998). CIC-5, the chloride channel mutated in Dent's disease, colocalizes with the proton pump in endocytotically active kidney cells. *Proc Natl Acad Sci U S A* 95(14): 8075-80.
- Hadley, J. K., M. Noda, A. A. Selyanko, I. C. Wood, F. C. Abogadie und D. A. Brown (2000). Differential tetraethylammonium sensitivity of

- KCNQ1-4 potassium channels. *Br J Pharmacol* 129(3): 413-415.
- Hanahan, D. (1983). Studies on transformation of *Escherichia coli* with plasmids. *J Mol Biol* 166(4): 557-80.
- Hansson, J. H., C. Nelson-Williams, H. Suzuki, L. Schild, R. Shimkets, Y. Lu, C. Canessa, T. Iwasaki, B. Rossier und R. P. Lifton (1995). Hypertension caused by a truncated epithelial sodium channel gamma subunit: genetic heterogeneity of Liddle syndrome. *Nat Genet* 11(1): 76-82.
- Hansson, J. H., L. Schild, Y. Lu, T. A. Wilson, I. Gautschi, R. Shimkets, C. Nelson-Williams, B. C. Rossier und R. P. Lifton (1995). A de novo missense mutation of the beta subunit of the epithelial sodium channel causes hypertension and Liddle syndrome, identifying a proline-rich segment critical for regulation of channel activity. *Proc Natl Acad Sci U S A* 92(25): 11495-9.
- Harvey, K. F. und S. Kumar (1999). Nedd4-like proteins: an emerging family of ubiquitin-protein ligases implicated in diverse cellular functions. *Trends Cell Biol* 9(5): 166-9.
- Heginbotham, L., Z. Lu, T. Abramson und R. MacKinnon (1994). Mutations in the K⁺ channel signature sequence. *Biophys J* 66(4): 1061-7.
- Heginbotham, L. und R. MacKinnon (1992). The aromatic binding site for tetraethylammonium ion on potassium channels. *Neuron* 8(3): 483-491.
- Heinemann, S. H., F. Conti und W. Stuhmer (1992). Recording of gating currents from *Xenopus* oocytes and gating noise analysis. *Methods Enzymol* 207: 353-68.
- Henn, D. K., A. Baumann und U. B. Kaupp (1995). Probing the transmembrane topology of cyclic nucleotide-gated ion channels with a gene fusion approach. *Proc Natl Acad Sci U S A* 92(16): 7425-9.
- Hershko, A. und A. Ciechanover (1998). The ubiquitin system. *Annu Rev Biochem* 67: 425-79.
- Higuchi, R. (1990). Recombinant PCR. In: *PCR Protocols*, M. A. Innis, D. H. Gelfand, J. J. Srinivas und T. J. White, eds. San Diego: Harcourt Brace Jovanovich: 177-183.
- Hille, B. (1992). *Ionic Channels of Excitable Membranes*. Sinauer Associates, Inc., Sunderland Massachusetts.
- Ho, K., C. G. Nichols, W. J. Lederer, J. Lytton, P. M. Vassilev, M. V. Kanazirska und S. C. Hebert (1993). Cloning and expression of an inwardly rectifying ATP-regulated potassium channel. *Nature* 362(6415): 31-8.
- Huibregtse, J. M., M. Scheffner, S. Beaudenon und P. M. Howley (1995). A family of proteins structurally and functionally related to the E6-AP ubiquitin-protein ligase [published erratum appears in *Proc Natl Acad Sci U S A* 1995 May 23;92(11):5249]. *Proc Natl Acad Sci U S A* 92(7): 2563-7.
- Isacoff, E. Y., Y. N. Jan und L. Y. Jan (1990). Evidence for the formation of heteromultimeric potassium channels in *Xenopus* oocytes. *Nature* 345(6275): 530-4.
- Jan, L. Y. und Y. N. Jan (1994). Potassium channels and their evolving gates. *Nature* 371(6493): 119-22.
- Jan, L. Y. und Y. N. Jan (1997). Cloned potassium channels from eukaryotes and prokaryotes. *Annu Rev Neurosci* 20: 91-123.
- Jentsch, T. J. (2000). Neuronal KCNQ potassium channels: Physiology and role in disease. *Nature Reviews Neuroscience* 1(1): 21-30.
- Jentsch, T. J., T. Friedrich, A. Schriever und H. Yamada (1999). The CLC chloride channel family. *Pflugers Arch* 437(6): 783-95.
- Jentsch, T. J. und W. Gunther (1997). Chloride channels: an emerging

- molecular picture. *Bioessays* 19(2): 117-26.
- Jentsch, T. J., W. Gunther, M. Pusch und B. Schwappach (1995). Properties of voltage-gated chloride channels of the CIC gene family. *J Physiol (Lond)* 482: 19S-25S.
- Jentsch, T. J., K. Steinmeyer und G. Schwarz (1990). Primary structure of *Torpedo marmorata* chloride channel isolated by expression cloning in *Xenopus* oocytes. *Nature* 348(6301): 510-4.
- Jones, S. M., K. E. Howell, J. R. Henley, H. Cao und M. A. McNiven (1998). Role of dynamin in the formation of transport vesicles from the trans-Golgi network. *Science* 279(5350): 573-7.
- Kamb, A., L. E. Iverson und M. A. Tanouye (1987). Molecular characterization of Shaker, a *Drosophila* gene that encodes a potassium channel. *Cell* 50(3): 405-13.
- Kawasaki, M., M. Suzuki, S. Uchida, S. Sasaki und F. Marumo (1995). Stable and functional expression of the CIC-3 chloride channel in somatic cell lines. *Neuron* 14(6): 1285-91.
- Kawasaki, M., S. Uchida, T. Monkawa, A. Miyawaki, K. Mikoshiba, F. Marumo und S. Sasaki (1994). Cloning and expression of a protein kinase C-regulated chloride channel abundantly expressed in rat brain neuronal cells. *Neuron* 12(3): 597-604.
- Kellenberger, S., I. Gautschi, B. C. Rossier und L. Schild (1998). Mutations causing Liddle syndrome reduce sodium-dependent downregulation of the epithelial sodium channel in the *Xenopus* oocyte expression system. *J Clin Invest* 101(12): 2741-50.
- Ketchum, K. A., W. J. Joiner, A. J. Sellers, L. K. Kaczmarek und S. A. Goldstein (1995). A new family of outwardly rectifying potassium channel proteins with two pore domains in tandem. *Nature* 376(6542): 690-5.
- Kharkovets, T., J. P. Hardelin, S. Safieddine, M. Schweizer, A. El-Amraoui, C. Petit und T. J. Jentsch (2000). KCNQ4, a K⁺-channel mutated in a form of dominant deafness, is expressed in the inner ear and in the central auditory pathway. *Proc Natl Acad Sci U.S.A.* 97: 4333-4338.
- Kieferle, S., P. Fong, M. Bens, A. Vandewalle und T. J. Jentsch (1994). Two highly homologous members of the CIC chloride channel family in both rat and human kidney. *Proc Natl Acad Sci U.S.A.* 91(15): 6943-7.
- Koch, M. C., K. Steinmeyer, C. Lorenz, K. Ricker, F. Wolf, M. Otto, B. Zoll, F. Lehmann-Horn, K. H. Grzeschik und T. J. Jentsch (1992). The skeletal muscle chloride channel in dominant and recessive human myotonia. *Science* 257(5071): 797-800.
- Kornak, U., D. Kaspar, M. R. Bösl, E. Kaiser, M. Schweizer, A. Schulz, W. Friedrich, G. Delling und T. J. Jentsch (2001). Loss of the CIC-7 chloride channel leads to osteopetrosis in mice and man. *Cell*: in press.
- Kreusch, A., P. J. Pfaffinger, C. F. Stevens und S. Choe (1998). Crystal structure of the tetramerization domain of the Shaker potassium channel. *Nature* 392(6679): 945-8.
- Krieg, P. A. und D. A. Melton (1984). Functional messenger RNAs are produced by SP6 in vitro transcription of cloned cDNAs. *Nucleic Acids Res* 12(18): 7057-70.
- Kubisch, C., B. C. Schroeder, T. Friedrich, B. Lutjohann, A. El-Amraoui, S. Marlin, C. Petit und T. J. Jentsch (1999). KCNQ4, a novel potassium channel expressed in sensory outer hair cells, is mutated in dominant deafness. *Cell* 96(3): 437-46.
- Laemmli, U. K. (1970). Cleavage of structural proteins during the

- assembly of the head of bacteriophage T4. *Nature* 227(259): 680-5.
- Larsson, H. P., O. S. Baker, D. S. Dhillon und E. Y. Isacoff (1996). Transmembrane movement of the shaker K⁺ channel S4. *Neuron* 16(2): 387-97.
- Leppert, M., V. E. Anderson, T. Quattlebaum, D. Stauffer, P. O'Connell, Y. Nakamura, J. M. Lalouel und R. White (1989). Benign familial neonatal convulsions linked to genetic markers on chromosome 20. *Nature* 337(6208): 647-8.
- Lerche, C., C. R. Scherer, G. Seebom, C. Derst, A. D. Wei, A. E. Busch und K. Steinmeyer (2000). Molecular cloning and functional expression of KCNQ5, a potassium channel subunit that may contribute to neuronal M-current diversity. *J. Biol. Chem.* 275: in press.
- Lesage, F., E. Guillemaire, M. Fink, F. Duprat, M. Lazdunski, G. Romey und J. Barhanin (1996). TWIK-1, a ubiquitous human weakly inward rectifying K⁺ channel with a novel structure. *Embo J* 15(5): 1004-11.
- Lewis, T. B., R. J. Leach, K. Ward, P. O'Connell und S. G. Ryan (1993). Genetic heterogeneity in benign familial neonatal convulsions: identification of a new locus on chromosome 8q. *Am J Hum Genet* 53(3): 670-5.
- Li, M., Y. N. Jan und L. Y. Jan (1992). Specification of subunit assembly by the hydrophilic amino-terminal domain of the Shaker potassium channel. *Science* 257(5074): 1225-1230.
- Lloyd, S. E., S. H. Pearce, S. E. Fisher, K. Steinmeyer, B. Schwappach, S. J. Scheinman, B. Harding, A. Bolino, M. Devoto, P. Goodyer, S. P. Rigden, O. Wrong, T. J. Jentsch, I. W. Craig und R. V. Thakker (1996). A common molecular basis for three inherited kidney stone diseases. *Nature* 379(6564): 445-9.
- Lorenz, C., M. Pusch und T. J. Jentsch (1996). Heteromultimeric CLC chloride channels with novel properties. *Proc Natl Acad Sci U S A* 93(23): 13362-6.
- Ludewig, U., M. Pusch und T. J. Jentsch (1996). Two physically distinct pores in the dimeric CLC-0 chloride channel. *Nature* 383(6598): 340-3.
- MacKinnon, R. (1991). Determination of the subunit stoichiometry of a voltage-activated potassium channel. *Nature* 350(6315): 232-5.
- MacKinnon, R., R. W. Aldrich und A. W. Lee (1993). Functional stoichiometry of Shaker potassium channel inactivation. *Science* 262(5134): 757-9.
- MacKinnon, R., S. L. Cohen, A. Kuo, A. Lee und B. T. Chait (1998). Structural conservation in prokaryotic and eukaryotic potassium channels. *Science* 280(5360): 106-9.
- MacKinnon, R. und D. A. Doyle (1997). Prokaryotes offer hope for potassium channel structural studies. *Nat Struct Biol* 4(11): 877-9.
- Marrion, N. V. (1997). Control of M-current. *Annu Rev Physiol* 59: 483-504.
- Matsumura, Y., S. Uchida, Y. Kondo, H. Miyazaki, S. B. Ko, A. Hayama, T. Morimoto, W. Liu, M. Arisawa, S. Sasaki und F. Marumo (1999). Overt nephrogenic diabetes insipidus in mice lacking the CLC-K1 chloride channel. *Nat Genet* 21(1): 95-8.
- Mellman, I. (1996). Endocytosis and molecular sorting. *Annu Rev Cell Dev Biol* 12: 575-625.
- Mellman, I., R. Fuchs und A. Helenius (1986). Acidification of the endocytic and exocytic pathways. *Annu Rev Biochem* 55: 663-700.
- Methfessel, C., V. Witzemann, T. Takahashi, M. Mishina, S. Numa und B. Sakmann (1986). Patch clamp measurements on *Xenopus*

- laevis oocytes: currents through endogenous channels and implanted acetylcholine receptor and sodium channels. *Pflugers Arch* 407(6): 577-88.
- Middleton, R. E., D. J. Pheasant und C. Miller (1996). Homodimeric architecture of a ClC-type chloride ion channel [see comments]. *Nature* 383(6598): 337-40.
- Miller, C. (1992). Ion channel structure and function. *Science* 258(5080): 240-1.
- Minor, D. L., Y. F. Lin, B. C. Mobley, A. Avelar, Y. N. Jan, L. Y. Jan und J. M. Berger (2000). The polar T1 interface is linked to conformational changes that open the voltage-gated potassium channel. *Cell* 102(5): 657-70.
- Mortensen, R. M., M. Zubiaur, E. J. Neer und J. G. Seidman (1991). Embryonic stem cells lacking a functional inhibitory G-protein subunit (alpha i2) produced by gene targeting of both alleles. *Proc Natl Acad Sci U S A* 88(16): 7036-40.
- Mukherjee, S., R. N. Ghosh und F. R. Maxfield (1997). Endocytosis. *Physiol Rev* 77(3): 759-803.
- Mukhopadhyay, A., A. M. Barbieri, K. Funato, R. Roberts und P. D. Stahl (1997). Sequential actions of Rab5 and Rab7 regulate endocytosis in the Xenopus oocyte. *J Cell Biol* 136(6): 1227-37.
- Nakamura, M., H. Watanabe, Y. Kubo, M. Yokoyama, T. Matsumoto, H. Sasai und Y. Nishi (1998). KQT2, a new putative potassium channel family produced by alternative splicing. Isolation, genomic structure, and alternative splicing of the putative potassium channels. *Receptors Channels* 5(5): 255-271.
- Noda, M., S. Shimizu, T. Tanabe, T. Takai, T. Kayano, T. Ikeda, H. Takahashi, H. Nakayama, Y. Kanaoka, N. Minamino und et al. (1984). Primary structure of Electrophorus electricus sodium channel deduced from cDNA sequence. *Nature* 312(5990): 121-7.
- Palmer, S., J. Perry und A. Ashworth (1995). A contravention of Ohno's law in mice. *Nat Genet* 10(4): 472-6.
- Papazian, D. M., T. L. Schwarz, B. L. Tempel, Y. N. Jan und L. Y. Jan (1987). Cloning of genomic and complementary DNA from Shaker, a putative potassium channel gene from *Drosophila*. *Science* 237(4816): 749-53.
- Papazian, D. M., L. C. Timpe, Y. N. Jan und L. Y. Jan (1991). Alteration of voltage-dependence of Shaker potassium channel by mutations in the S4 sequence. *Nature* 349(6307): 305-10.
- Pelham, H. R. (1991). Multiple targets for brefeldin A. *Cell* 67(3): 449-51.
- Pirozzi, G., S. J. McConnell, A. J. Uveges, J. M. Carter, A. B. Sparks, B. K. Kay und D. M. Fowlkes (1997). Identification of novel human WW domain-containing proteins by cloning of ligand targets. *J Biol Chem* 272(23): 14611-6.
- Piwon, N., W. Günther, M. Schwake, M. R. Bosl und T. J. Jentsch (2000). CIC-5 Cl⁻-channel disruption impairs endocytosis in a mouse model for Dent's disease. *Nature* 408(6810): 369-73.
- Pongs, O., N. Kecske, R. Müller, I. Krah-Jentgens, A. Baumann, H. H. Kiltz, I. Canal, S. Llamazares und A. Ferrus (1988). Shaker encodes a family of putative potassium channel proteins in the nervous system of *Drosophila*. *Embo J* 7(4): 1087-96.
- Ponting, C. P. (1997). CBS domains in CIC chloride channels implicated in myotonia and nephrolithiasis (kidney stones). *J Mol Med* 75(3): 160-3.
- Pusch, M., U. Ludewig, A. Rehfeldt und T. J. Jentsch (1995). Gating of the voltage-dependent chloride channel CIC-0 by the permeant anion. *Nature* 373(6514): 527-31.
- Pusch, M., K. Steinmeyer, M. C. Koch und T. J. Jentsch (1995). Mutations in

- dominant human myotonia congenita drastically alter the voltage dependence of the CIC-1 chloride channel. *Neuron* 15(6): 1455-63.
- Renard, S., N. Voilley, F. Bassilana, M. Lazdunski und P. Barbry (1995). Localization and regulation by steroids of the alpha, beta and gamma subunits of the amiloride-sensitive Na⁺ channel in colon, lung and kidney. *Pflugers Arch* 430(3): 299-307.
- Rettig, J., S. H. Heinemann, F. Wunder, C. Lorra, D. N. Parcej, J. O. Dolly und O. Pongs (1994). Inactivation properties of voltage-gated K⁺ channels altered by presence of beta-subunit. *Nature* 369(6478): 289-94.
- Rudy, B. (1988). Diversity and ubiquity of K channels. *Neuroscience* 25(3): 729-49.
- Rugarli, E. I., D. A. Adler, G. Borsani, K. Tsuchiya, B. Franco, X. Hauge, C. Disteché, V. Chapman und A. Ballabio (1995). Different chromosomal localization of the Clcn4 gene in *Mus spreitus* and C57BL/6J mice. *Nat Genet* 10(4): 466-71.
- Ruppertsberg, J. P., K. H. Schroter, B. Sakmann, M. Stocker, S. Sewing und O. Pongs (1990). Heteromultimeric channels formed by rat brain potassium-channel proteins. *Nature* 345(6275): 535-7.
- Sanger, F., S. Nicklen und A. R. Coulson (1977). DNA sequencing with chain-terminating inhibitors. *Proc Natl Acad Sci U S A* 74(12): 5463-7.
- Sanguinetti, M. C., M. E. Curran, A. Zou, J. Shen, P. S. Spector, D. L. Atkinson und M. T. Keating (1996). Coassembly of K(V)LQT1 and minK (IsK) proteins to form cardiac I(Ks) potassium channel. *Nature* 384(6604): 80-3.
- Scheinman, S. J., L. M. Guay-Woodford, R. V. Thakker und D. G. Warnock (1999). Genetic disorders of renal electrolyte transport. *N Engl J Med* 340(15): 1177-87.
- Schild, L., C. M. Canessa, R. A. Shimkets, I. Gautschi, R. P. Lifton und B. C. Rossier (1995). A mutation in the epithelial sodium channel causing Liddle disease increases channel activity in the *Xenopus laevis* oocyte expression system. *Proc Natl Acad Sci U S A* 92(12): 5699-703.
- Schild, L., Y. Lu, I. Gautschi, E. Schneeberger, R. P. Lifton und B. C. Rossier (1996). Identification of a PY motif in the epithelial Na channel subunits as a target sequence for mutations causing channel activation found in Liddle syndrome. *Embo J* 15(10): 2381-7.
- Schmidt-Rose, T. und T. J. Jentsch (1997). Reconstitution of functional voltage-gated chloride channels from complementary fragments of CLC-1. *J Biol Chem* 272(33): 20515-21.
- Schmidt-Rose, T. und T. J. Jentsch (1997). Transmembrane topology of a CLC chloride channel. *Proc Natl Acad Sci U S A* 94(14): 7633-8.
- Schmitt, N., M. Schwarz, A. Peretz, I. Abitbol, B. Attali und O. Pongs (2000). A recessive C-terminal Jervell and Lange-Nielsen mutation of the KCNQ1 channel impairs subunit assembly. *EMBO J* 19(3): 332-340.
- Schroeder, B. C., M. Hechenberger, F. Weinreich, C. Kubisch und T. J. Jentsch (2000). KCNQ5, a novel potassium channel broadly expressed in brain, mediates M-type currents. *J Biol Chem*.
- Schroeder, B. C., C. Kubisch, V. Stein und T. J. Jentsch (1998). Moderate loss of function of cyclic-AMP-modulated KCNQ2/KCNQ3 K⁺ channels causes epilepsy. *Nature* 396(6712): 687-90.
- Schroeder, B. C., S. Waldegger, S. Fehr, M. Bleich, R. Warth, R. Greger und T. J. Jentsch (2000). A constitutively open potassium channel formed by

- KCNQ1 and KCNE3. *Nature* 403(6766): 196-9.
- Schwake, M., M. Pusch, T. Kharkovets und T. J. Jentsch (2000). Surface expression and single channel properties of KCNQ2/KCNQ3, M-type K⁺ channels involved in epilepsy. *J Biol Chem* 275(18): 13343-8.
- Schweizer, A., P. D. Stahl und J. Rohrer (2000). A Di-aromatic motif in the cytosolic tail of the mannose receptor mediates endosomal sorting. *J Biol Chem* 275(38): 29694-700.
- Selyanko, A. A. und D. A. Brown (1993). Effects of membrane potential and muscarine on potassium M-channel kinetics in rat sympathetic neurones. *J Physiol (Lond)* 472: 711-724.
- Selyanko, A. A. und D. A. Brown (1996). Intracellular calcium directly inhibits potassium M channels in excised membrane patches from rat sympathetic neurons. *Neuron* 16(1): 151-162.
- Selyanko, A. A., J. K. Hadley, I. C. Wood, F. C. Abogadie, T. J. Jentsch und D. A. Brwon (2000). Inhibition of KCNQ1-4 potassium channels expressed in mammalian cells via M1 muscarinic receptors. *J. Physiol.* 522.3: 349-355.
- Selyanko, A. A. und J. A. Sim (1998). Ca²⁺-inhibited non-inactivating K⁺ channels in cultured rat hippocampal pyramidal neurones. *J Physiol (Lond)* 510(Pt 1): 71-91.
- Selyanko, A. A., C. E. Stansfeld und D. A. Brown (1992). Closure of potassium M-channels by muscarinic acetylcholine-receptor stimulants requires a diffusible messenger. *Proc R Soc Lond B Biol Sci* 250(1328): 119-125.
- Sewing, S., J. Roeper und O. Pongs (1996). Kv beta 1 subunit binding specific for shaker-related potassium channel alpha subunits. *Neuron* 16(2): 455-63.
- Shapiro, M. S., J. P. Roche, E. J. Kaftan, H. Cruzblanca, K. Macjie und B. Hille (2000). Reconstitution of muscarinic modulation of the KCNQ2/KCNQ3 K⁺ channels that underlie the neuronal M current. *J. Neurosci.* 20: 1710-1721.
- Shen, N. V., X. Chen, M. M. Boyer und P. J. Pfaffinger (1993). Deletion analysis of K⁺ channel assembly. *Neuron* 11(1): 67-76.
- Shimkets, R. A., R. P. Lifton und C. M. Canessa (1997). The activity of the epithelial sodium channel is regulated by clathrin-mediated endocytosis. *J Biol Chem* 272(41): 25537-41.
- Shimkets, R. A., D. G. Warnock, C. M. Bositis, C. Nelson-Williams, J. H. Hansson, M. Schambelan, J. R. Gill, Jr., S. Ulick, R. V. Milora, J. W. Findling und et al. (1994). Liddle's syndrome: heritable human hypertension caused by mutations in the beta subunit of the epithelial sodium channel. *Cell* 79(3): 407-14.
- Simon, D. B., R. S. Bindra, T. A. Mansfield, C. Nelson-Williams, E. Mendonca, R. Stone, S. Schurman, A. Nayir, H. Alpay, A. Bakkaloglu, J. Rodriguez-Soriano, J. M. Morales, S. A. Sanjad, C. M. Taylor, D. Pilz, A. Brem, H. Trachtman, W. Griswold, G. A. Richard, E. John und R. P. Lifton (1997). Mutations in the chloride channel gene, CLCNKB, cause Bartter's syndrome type III. *Nat Genet* 17(2): 171-8.
- Simon, D. B., F. E. Karet, J. M. Hamdan, A. DiPietro, S. A. Sanjad und R. P. Lifton (1996). Bartter's syndrome, hypokalaemic alkalosis with hypercalciuria, is caused by mutations in the Na-K-2Cl cotransporter NKCC2. *Nat Genet* 13(2): 183-8.
- Simon, D. B., F. E. Karet, J. Rodriguez-Soriano, J. H. Hamdan, A. DiPietro, H. Trachtman, S. A. Sanjad und R. P. Lifton (1996). Genetic heterogeneity of Bartter's syndrome revealed by mutations in the K⁺

- channel, ROMK. *Nat Genet* 14(2): 152-6.
- Singh, N. A., C. Charlier, D. Stauffer, B. R. DuPont, R. J. Leach, R. Melis, G. M. Ronen, I. Bjerre, T. Quattlebaum, J. V. Murphy, M. L. McHarg, D. Gagnon, T. O. Rosales, A. Peiffer, V. E. Anderson und M. Leppert (1998). A novel potassium channel gene, KCNQ2, is mutated in an inherited epilepsy of newborns. *Nat Genet* 18(1): 25-29.
- Snyder, P. M., M. P. Price, F. J. McDonald, C. M. Adams, K. A. Volk, B. G. Zeiher, J. B. Stokes und M. J. Welsh (1995). Mechanism by which Liddle's syndrome mutations increase activity of a human epithelial Na⁺ channel. *Cell* 83(6): 969-78.
- Staub, O., S. Dho, P. Henry, J. Correa, T. Ishikawa, J. McGlade und D. Rotin (1996). WW domains of Nedd4 bind to the proline-rich PY motifs in the epithelial Na⁺ channel deleted in Liddle's syndrome. *Embo J* 15(10): 2371-80.
- Staub, O., I. Gautschi, T. Ishikawa, K. Breitschopf, A. Ciechanover, L. Schild und D. Rotin (1997). Regulation of stability and function of the epithelial Na⁺ channel (ENaC) by ubiquitination. *Embo J* 16(21): 6325-36.
- Staub, O. und D. Rotin (1996). WW domains. *Structure* 4(5): 495-9.
- Steinmeyer, K., C. Lorenz, M. Pusch, M. C. Koch und T. J. Jentsch (1994). Multimeric structure of CIC-1 chloride channel revealed by mutations in dominant myotonia congenita (Thomsen). *Embo J* 13(4): 737-43.
- Steinmeyer, K., C. Ortland und T. J. Jentsch (1991). Primary structure and functional expression of a developmentally regulated skeletal muscle chloride channel. *Nature* 354(6351): 301-4.
- Steinmeyer, K., B. Schwappach, M. Bens, A. Vandewalle und T. J. Jentsch (1995). Cloning and functional expression of rat CLC-5, a chloride channel related to kidney disease. *J Biol Chem* 270(52): 31172-7.
- Stenmark, H., R. G. Parton, O. Steele-Mortimer, A. Lutcke, J. Gruenberg und M. Zerial (1994). Inhibition of rab5 GTPase activity stimulates membrane fusion in endocytosis. *Embo J* 13(6): 1287-96.
- Stobrawa, S. M., T. Breiderhoff, T. Takamori, T. Engel, M. Schweizer, A. A. Zdebik, M. R. Bösl, K. Ruether, H. Jahn, A. Draghun, R. Jahn und T. J. Jentsch (2001). Disruption of ClC-3, a chloride channel expressed on synaptic vesicles, leads to a loss of the hippocampus. *Neuron* 29: in press.
- Stühmer, W., H. Terlau und S. H. Heinemann (1992). Xenopus oocytes for two-electrode an patch-clamp recordings. In: *Practical electrophysiological methods*, H. Kettenmann und R. Grantyn, eds., Wiley-Liss, New York.
- Sudol, M., H. I. Chen, C. Bougeret, A. Einbond und P. Bork (1995). Characterization of a novel protein-binding module--the WW domain. *FEBS Lett* 369(1): 67-71.
- Tanabe, T., H. Takeshima, A. Mikami, V. Flockerzi, H. Takahashi, K. Kangawa, M. Kojima, H. Matsuo, T. Hirose und S. Numa (1987). Primary structure of the receptor for calcium channel blockers from skeletal muscle. *Nature* 328(6128): 313-8.
- Thiemann, A., S. Grunder, M. Pusch und T. J. Jentsch (1992). A chloride channel widely expressed in epithelial and non-epithelial cells. *Nature* 356(6364): 57-60.
- Tinel, N., I. Lauritzen, C. Chouabe, M. Lazdunski und M. Borsotto (1998). The KCNQ2 potassium channel: splice variants, functional and developmental expression. Brain localization and comparison with KCNQ3. *FEBS Lett* 438(3): 171-176.

- Trowbridge, I. S., J. F. Collawn und C. R. Hopkins (1993). Signal-dependent membrane protein trafficking in the endocytic pathway. *Annu Rev Cell Biol* 9: 129-61.
- Uchida, S., S. Sasaki, T. Furukawa, M. Hiraoka, T. Imai, Y. Hirata und F. Marumo (1993). Molecular cloning of a chloride channel that is regulated by dehydration and expressed predominantly in kidney medulla. *J Biol Chem* 268(6): 3821-4.
- Uchida, S., S. Sasaki, K. Nitta, K. Uchida, S. Horita, H. Nihei und F. Marumo (1995). Localization and functional characterization of rat kidney-specific chloride channel, ClC-K1. *J Clin Invest* 95(1): 104-13.
- van der Bliek, A. M. und E. M. Meyerowitz (1991). Dynamin-like protein encoded by the Drosophila shibire gene associated with vesicular traffic. *Nature* 351(6325): 411-4.
- van der Bliek, A. M., T. E. Redelmeier, H. Damke, E. J. Tisdale, E. M. Meyerowitz und S. L. Schmid (1993). Mutations in human dynamin block an intermediate stage in coated vesicle formation. *J Cell Biol* 122(3): 553-63.
- van Slegtenhorst, M. A., M. T. Bassi, G. Borsani, M. C. Wapenaar, G. B. Ferrero, L. de Conciliis, E. I. Rugarli, A. Grillo, B. Franco, H. Y. Zoghbi und et al. (1994). A gene from the Xp22.3 region shares homology with voltage-gated chloride channels. *Hum Mol Genet* 3(4): 547-52.
- van Weert, A. W., K. W. Dunn, H. J. Gueze, F. R. Maxfield und W. Stoorvogel (1995). Transport from late endosomes to lysosomes, but not sorting of integral membrane proteins in endosomes, depends on the vacuolar proton pump. *J Cell Biol* 130(4): 821-34.
- VanDongen, A. M., G. C. Frech, J. A. Drewe, R. H. Joho und A. M. Brown (1990). Alteration and restoration of K⁺ channel function by deletions at the N- and C-termini. *Neuron* 5(4): 433-43.
- Waldecker, S. und T. J. Jentsch (2000). From tonus to tonicity: physiology of CLC chloride channels. *J Am Soc Nephrol* 11(7): 1331-9.
- Waldecker, S. und T. J. Jentsch (2000). Functional and structural analysis of ClC-K chloride channels involved in renal disease. *J Biol Chem* 275(32): 24527-33.
- Wang, H. S., Z. Pan, W. Shi, B. S. Brown, R. S. Wymore, I. S. Cohen, J. E. Dixon und D. McKinnon (1998). KCNQ2 and KCNQ3 potassium channel subunits: molecular correlates of the M-channel. *Science* 282(5395): 1890-1893.
- Wang, Q., Q. Chen und J. A. Towbin (1998). Genetics, molecular mechanisms and management of long QT syndrome. *Ann Med* 30(1): 58-65.
- Wang, Q., M. E. Curran, I. Splawski, T. C. Burn, J. M. Millholland, T. J. VanRaay, J. Shen, K. W. Timothy, G. M. Vincent, T. de Jager, P. J. Schwartz, J. A. Toubin, A. J. Moss, D. L. Atkinson, G. M. Landes, T. D. Connors und M. T. Keating (1996). Positional cloning of a novel potassium channel gene: *KVLQT1* mutations cause cardiac arrhythmias. *Nat Genet* 12(1): 17-23.
- Watanabe, H., E. Nagata, A. Kosakai, M. Nakamura, M. Yokoyama, K. Tanaka und H. Sasai (2000). Disruption of the epilepsy KCNQ2 gene results in neural hyperexcitability. *J Neurochem* 75(1): 28-33.
- Wei, A., T. Jegla und L. Salkoff (1996). Eight potassium channel families revealed by the *C. elegans* genome project. *Neuropharmacology* 35(7): 805-29.
- White, M. M. und C. Miller (1979). A voltage-gated anion channel from the electric organ of *Torpedo californica*. *J Biol Chem* 254(20): 10161-6.

Wollnik, B., B. C. Schroeder, C. Kubisch, H. D. Esperer, P. Wieacker und T. J. Jentsch (1997). Pathophysiological mechanisms of dominant and recessive KVLQT1 K^+ channel mutations found in inherited cardiac arrhythmias. *Hum Mol Genet* 6(11): 1943-9.

Yamamoto, K., J. P. Cox, T. Friedrich, P. T. Christie, M. Bald, P. N. Houtman, M. J. Lapsley, L. Patzer, M. Tsimaratos, T. H. W. G. Van, K. Yamaoka, T. J. Jentsch und R. V. Thakker (2000). Characterization of renal chloride channel (CLCN5) mutations in Dent's disease. *J Am Soc Nephrol* 11(8): 1460-8.

Yang, W. P., P. C. Levesque, W. A. Little, M. L. Conder, P. Ramakrishnan, M. G. Neubauer und M. A. Blanar (1998). Functional expression of two KvLQT1-related potassium channels responsible for an inherited idiopathic epilepsy. *J Biol Chem* 273(31): 19419-19423.

Yang, Y. und F. J. Sigworth (1998). Single-channel properties of I_{Ks} potassium channels. *J Gen Physiol* 112(6): 665-678.

Yau, K. W. (1994). Cyclic nucleotide-gated channels: an expanding new family of ion channels. *Proc Natl Acad Sci U S A* 91(9): 3481-3.

Yu, W., J. Xu und M. Li (1996). NAB domain is essential for the subunit assembly of both alpha-alpha and alpha-beta complexes of shaker-like potassium channels. *Neuron* 16(2): 441-53.