

Bibliography

- [1] E. H. Hall, *On a New Action of the Magnet on Electric Currents*, American Journal of Mathematics **2**, 287 (1879).
- [2] K. v. Klitzing, G. Dorda, and M. Pepper, *New Method for High-Accuracy Determination of the Fine-Structure Constant Based on Quantized Hall Resistance*, Phys. Rev. Lett. **45**, 494 (1980).
- [3] N. W. Ashcroft and N. D. Mermin, *Solid State Physics*. Saunders College, Philadelphia, 1976.
- [4] B. Jeckelmann and B. Jeanneret, *The quantum Hall effect as an electrical resistance standard*, Rep. Prog. Phys. **64**, 1389 (2001).
- [5] E. Abrahams, P. W. Anderson, D. C. Licciardello, and T. V. Ramakrishnan, *Scaling theory of Localization: Absence of Quantum Diffusion in Two Dimensions*, Phys. Rev. Lett. **42**, 673 (1979).
- [6] D. Yoshioka, *The Quantum Hall Effect*. Springer Verlag, 2002.
- [7] R. B. Laughlin, *Quantized Hall conductivity in two dimensions*, Phys. Rev. B **23**, 5632 (1981).
- [8] B. I. Halperin, *Quantized Hall conductance, current-carrying edge states, and the existence of extended states in a two-dimensional disordered potential*, Phys. Rev. B **25**, 2185 (1982).
- [9] D. C. Tsui, H. L. Stormer, and A. C. Gossard, *Two-Dimensional Magnetotransport in the Extreme Quantum Limit*, Phys. Rev. Lett. **48**, 1559 (1982).
- [10] R. Willett, J. P. Eisenstein, H. L. Störmer, D. C. Tsui, A. C. Gossard, and J. H. English, *Observation of an even-denominator quantum number in the fractional quantum Hall effect*, Phys. Rev. Lett. **59**, 1776 (1987).
- [11] R. B. Laughlin, *Anomalous Quantum Hall Effect: An Incompressible Quantum Fluid with Fractionally Charged Excitations*, Phys. Rev. Lett. **50**, 1395 (1983).
- [12] F. D. M. Haldane, *Fractional Quantization of the Hall Effect: A Hierarchy of Incompressible Quantum Fluid States*, Phys. Rev. Lett. **51**, 605 (1983).
- [13] N. Byers and C. N. Yang, *Theoretical Considerations Concerning Quantized Magnetic Flux in Superconducting Cylinders*, Phys. Rev. Lett. **7**, 46 (1961).

- [14] D. Arovas, J. R. Schrieffer, and F. Wilczek, *Fractional Statistics and the Quantum Hall Effect*, Phys. Rev. Lett. **53**, 722 (1984).
- [15] O. Heinonen, ed., *Composite fermions. A Unified View of the Quantum Hall Regime*. World Scientific, 1998.
- [16] J. K. Jain, *Composite-fermion approach for the fractional quantum Hall effect*, Phys. Rev. Lett. **63**, 199 (1989).
- [17] J. K. Jain and R. K. Kamilla, “Composite fermions: Particles of the lowest landau level”, in *Composite fermions. A Unified View of the Quantum Hall Regime* (O. Heinonen, ed.), World Scientific, 1998.
- [18] W. Pan, J.-S. Xia, V. Shvarts, D. E. Adams, H. L. Störmer, D. C. Tsui, L. N. Pfeiffer, K. W. Baldwin, and K. W. West, *Exact Quantization of the Even-Denominator Fractional Quantum Hall State at $\nu = 5/2$ Landau Level Filling Factor*, Phys. Rev. Lett. **83**, 3530 (1999).
- [19] F. D. M. Haldane and E. H. Rezayi, *Spin-singlet wave function for the half-integral quantum Hall effect*, Phys. Rev. Lett. **60**, 956 (1988).
- [20] M. Greiter, X.-G. Wen, and F. Wilczek, *Paired Hall state at half filling*, Phys. Rev. Lett. **66**, 3205 (1991).
- [21] G. Moore and N. Read, *Nonabelions in the fractional quantum Hall effect*, Nucl. Phys. B **360**, 362 (1991).
- [22] R. H. Morf, *Transition from Quantum Hall to Compressible States in the Second Landau Level: New Light on the $\nu = 5/2$ Enigma*, Phys. Rev. Lett. **80**, 1505 (1998).
- [23] E. H. Rezayi and F. D. M. Haldane, *Incompressible Paired Hall State, Stripe Order, and the Composite Fermion Liquid Phase in Half-Filled Landau Levels*, Phys. Rev. Lett. **84**, 4685 (2000).
- [24] N. Read and E. Rezayi, *Beyond paired quantum Hall states: Parafermions and incompressible states in the first excited Landau level*, Phys. Rev. B **59**, 8084 (1999).
- [25] I. L. Aleiner and L. I. Glazman, *Two-dimensional electron liquid in a weak magnetic field*, Phys. Rev. B **52**, 11296 (1995).
- [26] A. A. Koulakov, M. M. Fogler, and B. I. Shklovskii, *Charge Density Wave in Two-Dimensional Electron Liquid in Weak Magnetic Field*, Phys. Rev. Lett. **76**, 499 (1996).
- [27] M. M. Fogler, A. A. Koulakov, and B. I. Shklovskii, *Ground state of a two-dimensional electron liquid in a weak magnetic field*, Phys. Rev. B **54**, 1853 (1996).
- [28] M. M. Fogler and A. A. Koulakov, *Laughlin liquid to charge-density-wave transition at high Landau levels*, Phys. Rev. B **55**, 9326 (1997).

- [29] R. Moessner and J. T. Chalker, *Exact results for interacting electrons in high Landau levels*, Phys. Rev. B **54**, 5006 (1996).
- [30] E. H. Rezayi, F. D. M. Haldane, and K. Yang, *Charge-Density-Wave Ordering in Half-Filled High Landau Levels*, Phys. Rev. Lett. **83**, 1219 (1999).
- [31] H. Fukuyama, P. M. Platzman, and P. W. Anderson, *Two-dimensional electron gas in a strong magnetic field*, Phys. Rev. B **19**, 5211 (1979).
- [32] M. P. Lilly, K. B. Cooper, J. P. Eisenstein, L. N. Pfeiffer, and K. W. West, *Evidence for an Anisotropic State of Two-Dimensional Electrons in High Landau Levels*, Phys. Rev. Lett. **82**, 394 (1999).
- [33] W. Pan, R. R. Du, H. L. S. nad D. C. Tsui, L. N. Pfeiffer, K. W. Baldwin, and K. W. West, *Strongly Anisotropic Electronic Transport at Landau Level Filling Factor $\nu = 9/2$ and $\nu = 5/2$ under a Tilted Magnetic Field*, Phys. Rev. Lett. **83**, 820 (1999).
- [34] E. Fradkin and S. A. Kivelson, *Liquid-crystal phases of quantum Hall systems*, Phys. Rev. B **59**, 8065 (1999).
- [35] S. Scheidl and F. von Oppen, *Elastic Theory of quantum Hall Smectics: Effects of disorder*, Europhys. Lett. **55**, 260 (2001).
- [36] R. G. Mani, J. H. Smet, K. von Klitzing, V. Narayanamurti, W. B. Johnson, and V. Umansky, *Zero-resistance states induced by electromagnetic-wave excitation in GaAs/AlGaAs heterostructures*, Nature **420**, 646 (2002).
- [37] M. A. Zudov, R. R. Du, L. N. Pfeiffer, and K. W. West, *Evidence for a New Dissipationless Effect in 2D Electronic Transport*, Phys. Rev. Lett. **90**, 046807 (2003).
- [38] M. A. Zudov, R. R. Du, L. N. Pfeiffer, and K. W. West, *Multiphoton processes in microwave photoresistance of two-dimensional electron systems*, Phys. Rev. B **73**, 041303(R) (2006).
- [39] S. I. Dorozhkin, J. H. Smet, V. Umansky, and K. von Klitzing, *Microwave photoresponse in the two-dimensional electron system caused by intra-Landau-level transitions*, Phys. Rev. B **71**, 201306 (2005). See also preprint: cond-mat/0409228.
- [40] R. G. Mani, *Novel zero-resistance states induced by photoexcitation in the high mobility GaAs/AlGaAs two-dimensional electron system*, Physica E **25**, 189 (2004).
- [41] C. L. Yang, R. R. Du, L. N. Pfeiffer, and K. W. West, *Influence of a parallel magnetic field on the microwave photoconductivity in a high-mobility two-dimensional electron system*, Phys. Rev. B **74**, 045315 (2006).

- [42] M. A. Zudov, R. R. Du, J. A. Simmons, and J. L. Reno, *Shubnikov-deHaas-like oscillations in millimeterwave photoconductivity in a high-mobility two-dimensional electron gas*, Phys. Rev. B **64**, 201311(R) (2001).
- [43] C. L. Yang, M. A. Zudov, T. A. Knuutila, R. R. Du, L. N. Pfeiffer, and K. W. West, *Observation of Microwave-Induced Zero-Conductance State in Corbino Rings of a Two-Dimensional Electron System*, Phys. Rev. Lett. **91**, 096803 (2003).
- [44] S. I. Dorozhkin, *Giant Magnetoresistance Oscillations Caused by Cyclotron Resonance Harmonics*, JETP Lett. **77**, 577 (2003).
- [45] R. L. Willett, K. W. West, and L. N. Pfeiffer, *High mobility 2D electron gas response to RF electric fields at small magnetic fields*, Bull. Am. Phys. Soc. **48**, 459 (2003).
- [46] V. I. Ryzhii, *Photoconductivity characteristics in thin films subjected to crossed electric and magnetic fields*, Fiz. Tverd. Tela **11**, 2577 (1969). [Sov. Phys. Solid State **11**, 2078 (1970)].
- [47] E. Vasiliadou, G. Müller, D. Heitmann, D. Weiss, K. v. Klitzing, H. Nickel, W. Schlapp, and R. Lösch, *Collective response in the microwave photoconductivity of Hall bar structures*, Phys. rev. B **48**, 17145 (1993).
- [48] A. Durst, S. Sachdev, N. Read, and S. M. Girvin, *Radiation-Induced Magnetoresistance Oscillations in a 2D Electron Gas*, Phys. Rev. Lett. **91**, 086803 (2003).
- [49] I. A. Dmitriev, M. G. Vavilov, I. L. Aleiner, A. D. Mirlin, and D. G. Polyakov, *Oscillatory ac conductivity and photoconductivity of a two-dimensional electron gas: Quasiclassical transport beyond the Boltzmann equation*, Phys. Rev. B **70**, 165305 (2004).
- [50] I. A. Dmitriev, M. G. Vavilov, I. L. Aleiner, A. D. Mirlin, and D. G. Polyakov, *Theory of microwave-induced oscillations in the magnetoconductivity of a two-dimensional electron gas*, Phys. Rev. B **71**, 115316 (2005). See also preprint: cond-mat/0310668.
- [51] A. V. Andreev, I. L. Aleiner, and A. J. Millis, *Dynamical Symmetry Breaking as the Origin of the Zero-dc-Resistance State in an ac-Driven System*, Phys. Rev. Lett. **91**, 056803 (2003).
- [52] J. Shi and X. C. Xie, *Radiation-Induced "Zero-Resistance State" and the Photon-Assisted Transport*, Phys. Rev. Lett. **91**, 086801 (2003).
- [53] X. L. Lei and S. Y. Liu, *Radiation-Induced Magnetoresistance Oscillation in a Two-Dimensional Electron Gas in Faraday Geometry*, Phys. Rev. Lett. **91**, 226805 (2003).
- [54] M. G. Vavilov and I. L. Aleiner, *Magnetotransport in a two-dimensional electron gas at large filling factors*, Phys. Rev. B **69**, 035303 (2004).

- [55] V. I. Ryzhii, R. A. Suris, and B. S. Shchamkhalova, *Photoconductivity of a two-dimensional electron gas in a strong magnetic field*, Fiz. Tekh. Poluprovodn. **20**, 2078 (1986). [Sov. Phys. Semicond. **20**, 1299 (1986)].
- [56] I. A. Dmitriev, A. D. Mirlin, and D. G. Polyakov, *Cyclotron-Resonance Harmonics in the ac Response of a 2D Electron Gas with Smooth Disorder*, Phys. Rev. Lett. **91**, 226802 (2003).
- [57] J. Dietel, L. I. Glazman, F. W. J. Hekking, and F. von Oppen, *Microwave Photoconductivity of two-dimensional electron systems with unidirectional periodic modulation*, Phys. Rev. B **71**, 045329 (2005).
- [58] W. Kohn, *Cyclotron Resonance and de Haas-van Alphen Oscillations of an Interacting Electron Gas*, Phys. Rev. **123**, 1242 (1961).
- [59] A. A. Koulakov and M. E. Raikh, *Classical model for the negative dc conductivity of ac-driven two-dimensional electrons near the cyclotron resonance*, Phys. Rev. B **68**, 115324 (2003).
- [60] I. E. Aronov, E. A. Kaner, and A. A. Slutskin, *A parametric resonance in semiconductors*, Solid State Commun. **38**, 245 (1981).
- [61] I. E. Aronov and E. A. Kaner, *Plasmon instability in semiconductors with parametric cyclotron pumping*, JETP Lett. **34**, 325 (1981).
- [62] I. E. Aronov and O. N. Baranetz, *Cyclotron parametric resonance in semiconductors*, Phys. Rep. **197**, 99 (1990).
- [63] C. Joas, M. E. Raikh, and F. von Oppen, *Parametric resonance of a two-dimensional electron gas under bichromatic irradiation*, Phys. Rev. B **70**, 235302 (2004).
- [64] A. J. Lichtenberg and M. A. Lieberman, *Regular and Chaotic Dynamics*. Springer Verlag, 2nd ed., 1992.
- [65] J. J. Thomsen, *Vibrations and Stability – Advanced Theory, Analysis, and Tools*. Springer Verlag, Heidelberg, 2nd ed., 2003.
- [66] C. Joas, J. Dietel, and F. von Oppen, *Microwave photoconductivity of a modulated two-dimensional electron gas due to intra-Landau-level transitions*, Phys. Rev. B **72**, 165323 (2005).
- [67] R. L. Willett, K. W. West, and L. N. Pfeiffer, *Transition in the Correlated 2D Electron System Induced by a Periodic Density Modulation*, Phys. Rev. Lett. **78**, 4478 (1997).
- [68] J. H. Smet, K. von Klitzing, D. Weiss, and W. Wegscheider, *dc Transport of Composite Fermions in Weak Periodic Potentials*, Phys. Rev. Lett. **80**, 4538 (1998).
- [69] F. von Oppen, A. Stern, and B. I. Halperin, *Composite Fermions in Modulated Structures: Transport and Surface Acoustic Waves*, Phys. Rev. Lett. **80**, 4494 (1998).

- [70] A. D. Mirlin, P. Wölfle, Y. Levinson, and O. Enthi-Wohlman, *Velocity Shift of Surface Acoustic Waves due to Interaction with Composite Fermions in a Modulated Structure*, Phys. Rev. Lett. **81**, 1070 (1998).
- [71] D. Weiss, K. von Klitzing, K. Ploog, and G. Weiman, *Magnetoresistance oscillations in a two-dimensional electron gas induced by a submicrometer periodic potential*, Europhys. Lett. **8**, 179 (1989).
- [72] R. R. Gerhardts, D. Weiss, and K. von Klitzing, *Novel magnetoresistance oscillations in a periodically modulated two-dimensional electron gas*, Phys. Rev. Lett. **62**, 1173 (1989).
- [73] C. Zhang and R. R. Gerhardts, *Theory of magnetotransport in two-dimensional electron systems with unidirectional periodic modulation*, Phys. Rev. B **41**, 12850 (1990).
- [74] A. Messiah, *Quantum Mechanics*. Dover Publications, 2001.
- [75] S. Titeica, *Über die Widerstandsänderung von Metallen im Magnetfeld*, Ann. Phys. **22**, 129 (1935).
- [76] M. Abramowitz and I. A. Stegun, eds., *Handbook of mathematical functions*. Dover publications, Inc., New York, 1970.
- [77] S. L. Sondhi, A. Karlhede, S. A. Kivelson, and E. H. Rezayi, *Skyrmions and the crossover from the integer to fractional quantum Hall effect at small Zeeman energies*, Phys. Rev. B **47**, 16419 (1993).
- [78] F. F. Fang and P. J. Stiles, *Effects of a Tilted Magnetic Field on a Two-Dimensional Electron Gas*, Phys. Rev. **174**, 823 (1968).
- [79] M. B. Pogrebinskii, *Mutual drag of carriers in a semiconductor-insulator-semiconductor system*, Fiz. Tekh. Poluprovodn. **11**, 637 (1977).
- [80] P. J. Price, *Hot Electron Effects in Heterolayers*, Physica B&C **117**, 750 (1983).
- [81] J. P. Eisenstein, G. S. Boebinger, L. N. Pfeiffer, K. W. West, and S. He, *New fractional quantum Hall state in double-layer two-dimensional electron systems*, Phys. Rev. Lett. **68**, 1383 (1992).
- [82] X. G. Feng, S. Zelakiewicz, H. Noh, T. J. Ragucci, T. J. Gramila, L. N. Pfeiffer, and K. W. West, *Negative Electron Drag and Holelike Behavior in the Integer Quantum Hall Regime*, Phys. Rev. Lett. **81**, 3219 (1998).
- [83] J. G. S. Lok, S. Kraus, M. Pohlt, W. Dietsche, K. von Klitzing, W. Wegscheider, and M. Bichler, *Spin effects in the magnetodrag between double quantum wells*, Phys. Rev. B **63**, 041305 (2001).
- [84] K. Muraki, J. G. S. Lok, S. Kraus, W. Dietsche, K. von Klitzing, D. Schuh, M. Bichler, and W. Wegscheider, *Coulomb Drag as a Probe of the Nature of Compressible States in a Magnetic Field*, Phys. Rev. Lett. **92**, 246801 (2004).

- [85] F. von Oppen, S. H. Simon, and A. Stern, *Oscillating Sign of Drag in High Landau Levels*, Phys. Rev. Lett. **87**, 106803 (2001).
- [86] I. V. Gornyi, A. D. Mirlin, and F. von Oppen, *Coulomb drag in high Landau levels*, Phys. Rev. B **70**, 245302 (2004).
- [87] R. Bistritzer and A. Stern, *Physical Picture behind the Oscillating Sign of Drag in High Landau Levels*, Phys. Rev. Lett. **96**, 226801 (2006).
- [88] T. J. Gramila, J. P. Eisenstein, A. H. MacDonald, L. N. Pfeiffer, and K. W. West, *Mutual friction between parallel two-dimensional electron systems*, Phys. Rev. Lett. **66**, 1216 (1991).
- [89] K. Flensberg, B. Y. Hu, A.-P. Jauho, and J. M. Kinaret, *Linear-response theory of Coulomb drag in coupled electron systems*, Phys. Rev. B **52**, 14761 (1995).
- [90] T. J. Gramila, J. P. Eisenstein, A. H. MacDonald, L. N. Pfeiffer, and K. W. West, *Evidence for virtual-phonon exchange in semiconductor heterostructures*, Phys. Rev. B **47**, 12957 (1993).
- [91] M. C. Bønsager, K. Flensberg, B. Y.-K. Hu, and A. H. MacDonald, *Fictional drag between quantum wells mediated by phonon exchange*, Phys. Rev. B **57**, 7085 (1998).
- [92] A. V. Khaetskii and Y. V. Nazarov, *Coulomb drag in intermediate magnetic fields*, Phys. Rev. B **59**, 7551 (1999).
- [93] A.-P. Jauho and H. Smith, *Coulomb drag between parallel two-dimensional electron systems*, Phys. Rev. B **47**, 4420 (1993).
- [94] L. Zheng and A. H. MacDonald, *Coulomb drag between disordered two-dimensional electron-gas layers*, Phys. Rev. B **48**, 8203 (1993).
- [95] A. Kamenev and Y. Oreg, *Coulomb drag in normal metals and superconductors: Diagrammatic approach*, Phys. Rev. B **52**, 7516 (1995).
- [96] P. J. Price, *Two-dimensional electron transport in semiconductor layers. I. Phonon scattering*, Ann. Phys. (N.Y.) **133**, 217 (1981).
- [97] S. K. Lyo, *Low-temperature phonon-drag thermoelectric power in heterojunctions*, Phys. Rev. B **38**, 6345 (1988).
- [98] J. D. Zook, *Piezoelectric Scattering in Semiconductors*, Phys. Rev. **136**, A869 (1964).
- [99] H. Noh, S. Zelakiewicz, T. J. Gramila, L. N. Pfeiffer, and K. W. West, *Phonon-mediated drag in double-layer two-dimensional electron systems*, Phys. Rev. B **59**, 13114 (1999).
- [100] T. J. Gramila, J. P. Eisenstein, A. H. MacDonald, L. N. Pfeiffer, and K. W. West, *Measuring electron-electron scattering rates through mutual friction*, Physica B **197**, 442 (1994).

- [101] M. C. Bønsager, K. Flensberg, B. Y.-K. Hu, and A.-P. Jauho, *Magneto-Coulomb Drag: Interplay of Electron-Electron Interactions and Landau Quantization*, Phys. Rev. Lett. **77**, 1366 (1996).
- [102] K. S. Novoselov, A. K. Geim, S. V. Morozov, D. Jiang, M. I. Katsnelson, I. V. Grigorieva, S. V. Dubonos, and A. A. Firsov, *Two-dimensional gas of massless Dirac fermions in graphene*, Nature **438**, 197 (2005).
- [103] R. Kubo, *Statistical-Mechanical Theory of Irreversible Processes. I. General Theory and Simple Applications to Magnetic and Conduction Problems*, J. Phys. Soc. Jpn. **12**, 570 (1957).
- [104] K. A. Benedict and J. T. Chalker, *An exactly solvable model of the disordered two-dimensional electron gas in a strong magnetic field*, J. Phys. C **19**, 3587 (1986).
- [105] T. Ando and Y. Uemura, *Theory of Quantum Transport in a Two-Dimensional Electron System under Magnetic Fields. I. Characteristics of Level Broadening and Transport under Strong Fields*, J. Phys. Soc. Jpn. **36**, 959 (1974).
- [106] G. D. Mahan, “Polarons in heavily doped semiconductors”, in *Polarons in Ionic Crystals and Polar Semiconductors* (J. T. Devreese, ed.), pp. 553–657, North-Holland, Amsterdam, 1972.
- [107] J. Bardeen and W. Shockley, *Deformation Potentials and Mobilities in Non-Polar Crystals*, Phys. Rev. **80**, 72 (1950).
- [108] K. A. Benedict, R. K. Hills, and C. J. Mellor, *Theory of phonon spectroscopy in the fractional quantum Hall regime*, Phys. Rev. B **60**, 10984 (1999).