

5. Literatur

- Adam EJ & Adam SA. (1994) Identification of cytosolic factors required for nuclear location sequence-mediated binding to the nuclear envelope. *J. Cell Biol.* **125**, 547-555
- Allen TP, Cronshaw JM, Bagley S, Kiseleva E & Goldberg MW. (2000) The nuclear pore complex: mediator of the translocation between nucleus and cytoplasm. *J. Cell Sci.* **113**, 1651-1659
- Andrulis ED, Neiman AM, Zappulla DC & Sternglanz R. (1998) Perinuclear localization of chromatin facilitates transcriptional silencing. *Nature* **394**, 592-595
- Azzi A, Boscoboinik D & Hensey C. (1992) The protein kinase C family. *Eur. J. Biochem.* **208**, 547-557
- Baudier J, Delphin C, Grunwald D, Khochbin S & Lawrence JJ. (1992) Characterization of the tumor suppressor protein p53 as a protein kinase C substrate and a S100b-binding protein. *Proc. Natl. Acad. Sci. USA* **89**, 11627-11631
- Beckmann R, Buchner K, Jungblut PR, Eckerskorn C, Weise C, Hilbert R & Hucho F. (1992) Nuclear substrates of protein kinase C. *Eur. J. Biochem.* **210**, 45-51
- Bertrand F, Veissiere D, Hermelin B, Paul A, Capeau J, Picard J, Cherqui G. (1994) Phosphorylation of vimentin is an intermediate step in protein kinase C-mediated glycoconjugate secretion. *Am. J. Physiol.* **266**, C611-621
- Beyer AL, Christensen ME, Walker BW & LeStourgeon WM. (1977) Identification and characterization of the packaging proteins of core 40S hnRNP particles. *Cell* **11**, 127-138
- Bione S, Maestrini E, Rivella S, Mancini M, Regis S, Romeo G & Toniolo D. (1994) Identification of a novel X-linked gene responsible for Emery-Dreifuss muscular dystrophy. *Nat. Genet.* **8**, 323-327
- Birney E, Kumar S & Krainer AR. (1993) Analysis of the RNA-recognition motif and RS and RGG domains: conservation in metazoan pre-mRNA splicing factors. *Nucleic Acid Res.* **21**, 5803-5816
- Buchner K. (1995) Protein kinase C in the transduction of signals toward and within the cell nucleus. *Eur. J. Biochem.* **228**, 211-221
- Buchner K, Adamec E, Beermann ML & Nixon RA. (1999) Isoform specific translocation of protein kinase C following glutamate administration in primary hippocampal neurons. *Brain Res. Mol. Brain Res.* **64**, 222-235
- Buchner K. (2000) The role of protein kinase C in the regulation of cell growth and in signalling to the cell nucleus. *J. Cancer Res. Clin. Oncol.* **126**, 1-11
- Buendia B & Courvalin JC. (1997) Domain-specific disassembly and reassembly of nuclear membranes during mitosis. *Exp. Cell Res.* **230**, 133-144
- Butler AP, Byus CV & Slaga TJ. (1986) Phosphorylation of histones is stimulated by phorbol esters in quiescent Reuber H35 hepatoma cells. *J. Biol. Chem.* **261**, 9421-9425
- Bradford MM. (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* **72**, 248-254

- Caceres JF, Misteli T, Sreaton GR, Spector DL & Krainer AR. (1997) Role of the modular domains of SR proteins in subnuclear localization and alternative splicing specificity. *J. Cell Biol.* **138**, 225-238
- Carmo-Fonseca & David-Ferreira. (1990) Interactions of intermediate filaments with cell structures. *Electron Microscop. Rev.* **3**, 115-141
- Courey AJ & Tjian R. (1988) Analysis of Sp1 in vivo reveals multiple transcriptional domains, including a novel glutamine-rich activation motif. *Cell* **55**, 887-897
- Crimaudo C, Hortsch M, Gausepohl H & Meyer DI. (1987) Human ribophorins I und II: the primary structure and membrane topology of two highly conserved rough endoplasmic reticulum-specific glycoproteins. *EMBO J.* **6**, 75-82
- Decker LV & Parker PJ. (1994) Protein kinase C – a question of specificity. *TIBS* **19**, 73-77
- Dessev G, Iovcheva-Dessev C, Bischoff JR, Beach D & Goldman R. (1991) A complex containing p34cdc2 and cyclin B phosphorylates the nuclear lamin and disassembles nuclei of clam oocytes in vitro. *J. Cell Biol.* **112**, 523-533
- Divecha N, Banfic H & Irvine RF. (1991) The polyphosphoinositide cycle exists in the nuclei of Swiss 3T3 cells under the control of a receptor (for IGF-I) in the plasma membrane, and stimulation of the cycle increases nuclear diacylglycerol and apparently induces translocation of protein kinase C to the nucleus. *EMBO J.* **10**, 3207-3214
- Dreger M, Henning O, Neubauer G, Mann M & Hucho F. (1999) Identification of phosphorylation sites in native lamina-associated polypeptide 2b. *Biochemistry* **38**, 9426-9434
- Dreger M, Bengtsson L, Schöneberg T, Otto H & Hucho F. (2001) Nuclear envelope proteomics: Novel integral membrane proteins of the inner nuclear membrane. *Proc. Natl. Acad. Sci. USA* **98**, 11943-11948
- Dreyfuss G, Matunis MJ, Pinol-Roma S & Burd CG. (1993) HnRNP proteins and the biogenesis of mRNA. *Annu. Rev. Biochem.* **62**, 289-321
- Dundr M & Misteli T. Functional architecture in the cell nucleus. (2001) *Biochem J.* **356**, 297-310
- Dye BT & Patton JG. (2001) An RNA recognition motif (RRM) is required for the localization of PTB-associated splicing factor (PSF) to subnuclear speckles. *Exp. Cell Res.* **263**, 131-144
- Ellenberg J, Siggia ED, Moreira JE, Smith CL, Presley JF, Worman HJ & Lippincott-Schwartz J. (1997) Nuclear membrane dynamics and reassembly in living cells: Targeting of an inner nuclear membrane protein in interphase and mitosis. *J. Cell Biol.* **138**, 1193-1206
- Emig S, Schmalz D, Shakibaei M & Buchner K. (1995) The nuclear pore complex protein p62 is one of several sialic acid-containing proteins of the nuclear envelope. *J. Biol. Chem.* **270**, 13787-13793
- Fey E, Wan KM, Penman S. (1984) Epithelial cytoskeletal framework and nuclear matrix-intermediate filament scaffold: Threedimensional organisation and protein composition. *J. Cell Biol.* **98**, 1973-1984
- Fields AP, Pettit GR & May WS. (1988) Phosphorylation of lamin B at the nuclear membrane by activated protein kinase C. *J. Biol. Chem.* **263**, 8253-8260
- Flores-Rozas H & Hurwitz J. (1993) Characterization of a new RNA helicase from nuclear extracts of HeLa cells which translocates in 5'-3' direction. *J. Biol. Chem.* **268**, 21372-21383
- Foisner R & Gerace L. (1993) Integral membrane proteins of the nuclear envelope interact with lamins and chromosomes, and binding is modulated by mitotic phosphorylation. *Cell* **73**, 1267-1279

- Fontoura BMA, Blobel G & Matunis MJ. (1999) A conserved biogenesis pathway for nucleoporins: proteolytic processing of a 186-kilodalton precursor generates Nup98 and the novel nucleoporin, Nup96. *J. Cell Biol.* **144**, 1097-1112
- French SW, Kawahara H, Katsuma Y, Ohta M & Swierenga SHH. (1989) Interaction of intermediate filaments with nuclear lamina and cell periphery. *Electron Microsc. Rev.* **2**, 17-51
- Fridell RA, Truant R, Thorne L, Benson RE & Cullen BR. (1997) Nuclear import of hnRNP A1 is mediated by a novel cellular cofactor related to karyopherin- β . *J. Cell Sci.* **110**, 1325-1331
- Fu J, Ren M & Kreibich G. (1997) Interactions among subunits of the oligosacaryltransferase complex. *J. Biol. Chem.* **272**, 29687-29692
- Furukawa K, Pante N, Aebi U & Gerace L. (1995) Cloning of a cDNA for lamina-associated polypeptide 2 (LAP2) and identification of regions that specify targeting to the nuclear envelope. *EMBO J.* **14**, 1626-36
- Furukawa K, Fritze CE & Gerace L. (1998) The major nuclear envelope targeting domain of LAP2 coincides with its lamin binding region but is distinct from its chromatin interaction domain. *J. Biol. Chem.* **273**, 4213-4219
- Furukawa K. (1999) LAP2 binding protein 1 (L2BP1/BAF) is a candidate mediator of LAP2-chromatin interaction. *J. Cell Sci.* **112**, 2485-2492
- Garcia-Bermejo ML, Leskow FC, Fujii T, Wang Q, Blumberg PM, Ohba M, Kuroki T, Han KC, Lee J, Marquez VE & Kazanietz MG. (2002) Diacylglycerol (DAG)-lactones, a new class of protein kinase C (PKC) agonists, induce apoptosis in LNCaP prostate cancer cells by selective activation of PKC α . *J. Biol. Chem.* **277**, 645-655
- Georgatos SD, Pырpasopoulou A & Theodoropoulos PA. (1997) Nuclear envelope breakdown in mammalian cells involves stepwise lamina disassembly and microtubule-driven deformation of the nuclear membrane. *J. Cell Sci.* **110**, 2129-2140
- Ghetti A, Pinol-Roma S, Michael WM, Morandi C & Dreyfuss G. (1992) hnRNP I, the polypyrimidine tract-binding protein: distinct nuclear localization and association with hnRNAs. *Nucleic Acids Res.* **20**, 3671-3678
- Giannakakou P, Sackett DL, Ward Y, Webster KR, Blagosklonny MV & Fojo T. (2000) p53 is associated with cellular microtubules and is transported to the nucleus by dynein. *Nat. Cell Biol.* **2**, 709-717
- Giannakakou P, Nakano M, Nicolaou KC, O'Brate A, Yu J, Blagosklonny MV, Greber UF & Fojo T. (2002) Enhanced microtubule-dependent trafficking and p53 nuclear accumulation by suppression of microtubule dynamics. *Proc. Natl. Acad. Sci. USA.* **99**, 10855-10860
- Görlich D, Prehn S, Laskey RA & Hartmann E. (1994) Isolation of a protein that is essential for the first step of nuclear protein import. *Cell* **79**, 767-778
- Görlich D & Kutay U. (1999) Transport between the cell nucleus and the cytoplasm. *Annu. Rev. Cell Dev. Biol.* **15**, 607-660
- Goldberg MW & Allen TD. (1995) Structural and functional organization of the nuclear envelope. *Curr. Opin. Cell Biol.* **7**, 301-309

- Goldberg MW & Allen TD. (1996) The nuclear pore complex and lamina: three-dimensional structures and interactions determined by field emission in-lens scanning electron microscopy. *J. Mol. Biol.* **257**, 848-865
- Goldberg MW, Wiese C, Allen TD & Wilson KL. (1997) Dimples, pores, star-rings, and thin rings on growing nuclear envelopes: evidence for structural intermediates in nuclear pore complex assembly. *J. Cell Sci.* **110**, 409-420
- Goss VL, Hocevar BA, Thompson LJ, Stratton CA, Burns DJ & Fields AP. (1994) Identification of nuclear β_{II} protein kinase C as a mitotic lamin kinase. *J. Biol. Chem.* **269**, 19074-19080
- Gozani O, Patton JG & Reed R. (1994) A novel set of spliceosome-associated proteins and the essential splicing factor PSF bind stably to pre-mRNA prior to catalytic step II of the splicing reaction. *EMBO J.* **13**, 3356-3367
- Gruenbaum Y, Wilson KL, Harel A, Goldberg M & Cohen M. (2000) Nuclear lamins – structural proteins with fundamental functions. *J. Struct. Biol.* **129**, 313-323
- Guan T, Muller S, Klier G, Pante N, Blevitt JM, Haner M, Paschal B, Aebi U & Gerace L. (1995) Structural analysis of the p62 complex, an assembly of O-linked glycoproteins that localizes near the central gated channel of the nuclear pore complex. *Mol. Biol. Cell* **6**, 1591-603
- Haraguchi T, Koujin T, Segura-Totten M, Lee KK, Matsuoka Y, Yoneda Y, Wilson KL & Hiraoka Y. (2001) BAF is required for emerin assembly into the reforming nuclear envelope. *J. Cell Sci.* **114**, 4574-4585
- Hardt B, Aparicio R & Brause E. (2000) The oligosaccharyltransferase complex from pig liver: cDNA cloning, expression and functional characterisation. *Glycoconj. J.* **17**, 767-779
- Hartig R, Shoeman RL, Janetzko A, Tolstonog G & Traub P. (1998) DNA-mediated transport of the intermediate filament protein vimentin into the nucleus of cultured cells. *J. Cell Sci.* **111**, 3573-3584
- Hartinger J, Stenius K, Hogemann D & Jahn R. (1996) 16-BAC/SDS-PAGE: a two-dimensional gel electrophoresis system suitable for the separation of integral membrane proteins. *Anal. Biochem.* **240**, 126-133
- Hedley ML, Amrein H & Maniatis T. (1995) An amino acid sequence motif sufficient for subnuclear localization of an arginine/serine-rich splicing factor. *Proc. Natl. Acad. Sci. USA* **92**, 11524-11528
- Hirano K, Erdödi F, Patton JG & Hartshorne DJ. (1996) Interaction of protein phosphatase type 1 with a splicing factor. *FEBS Lett.* **389**, 191-194
- Hocevar BA, Burns DJ & Fields AP. (1993) Identification of protein kinase C (PKC) phosphorylation sites on human lamin B. *J. Biol. Chem.* **268**, 7545-7552
- Hu T, Guan T & Gerace L. (1996) Molecular and functional characterization of the p62 complex, an assembly of nuclear pore complex glycoproteins. *J. Cell Biol.* **134**, 589-601
- Huang CK, Devanney JF & Kennedy SP. (1988) Vimentin, a cytoskeletal substrate of protein kinase C. *Biochem. Biophys. Res. Commun.* **150**, 1006-1011
- Imamoto N. (2000) Diversity in nucleocytoplasmic transport pathways. *Cell Struct. Funct.* **25**, 207-216

- Inagaki M, Inagaki N, Takahashi T & Takai Y. (1997) Phosphorylation-dependent control of structures of intermediate filaments: A novel approaching site- and phosphorylation state- specific antibodies. *J. Biochem.* **121**, 407-414
- Jaken S. (1996) Protein kinase C isozymes and substrates. *Curr. Opin. Cell Biol.* **8**, 168-173
- Jankowsky E, Gross CH, Shuman S & Pyle AM. (2001) Active disruption of an RNA-protein interaction by a DExH/D RNA Helicase. *Science* **291**, 121-124
- Jenkins H, Hölman T, Lyon C, Lane B, Stick R & Hutchison C. (1993) Nuclei that lack a lamina accumulate karyophilic proteins and assemble a nuclear matrix. *J. Cell Sci.* **106**, 275-285
- Kalderon D, Richardson WD, Markham AF & Smith AE. (1984) Sequence requirements for nuclear location of simian virus 40 large-T antigen. *Nature* **311**, 33-38
- Kenan DJ, Query CC & Keene JD. (1991) RNA recognition: towards identifying determinants of specificity. *Trends Biochem. Sci.* **16**, 214-220
- Kerr SM, Taggart MH, Lee M & Cooke HJ. (1996) Ott, a mouse x-linked multigene family expressed specifically during meiosis. *Hum. Mol. Genet.* **5**, 1139-1148
- Kiledjian M & Dreyfuss G. (1992) Primary structure and binding specificity of the hnRNP U protein: binding RNA through RGG box. *EMBO J.* **11**, 2655-2664
- Klauck TM, Faux MC, Labudda K, Langeberg LK, Jaken S & Scott JD. (1996) Coordination of three signaling enzymes by AKAP79, a mammalian scaffold protein. *Science* **271**, 1589-1592
- Kolch W, Heidecker G, Kochs G, Hummel R, Vahidi H, Mischak H, Finkenzeller G, Marme D & Rapp UR. (1993) Protein kinase C alpha activates RAF-1 by direct phosphorylation. *Nature* **364**, 249-252
- Komeili A & O'Shea EK. (2001) New perspectives on nuclear transport. *Annu. Rev. Genet.* **35**, 341-364
- Krämer A. (1996) The structure and function of proteins involved in mammalian pre-mRNA splicing. *Annu Rev. Biochem.* **65**, 367-409
- Kreibich G, Czako-Graham M, Grebenau R, Mok W, Rodriguez-Boulan E & Sabatini DD. (1978) Characterization of the ribosomal binding site in rat liver rough microsomes: ribophorins I and II, two integral membrane proteins related to ribosome binding. *J. Supramol. Struct.* **8**, 279-302
- Kreibich G, Ojakian G, Rodriguez-Boulan E & Sabatini DD. (1982) Recovery of ribophorins and ribosomes in "inverted rough" vesicles derived from rat liver rough microsomes. *J. Cell Biol.* **93**, 111-21
- Lam MH, Briggs LJ, Hu W, Martin TJ, Gillespie MT & Jans DA. (1999) Importin beta recognizes parathyroid hormone-related protein with high affinity and mediates its nuclear import in the absence of importin alpha. *J. Biol. Chem.* **274**, 7391-7398
- Lam MH, Thomas RJ, Loveland KL, Schilders S, Gu M, Martin TJ, Gillespie MT & Jans DA. (2002) Nuclear transport of parathyroid hormone (PTH)-related protein is dependent on microtubules. *Mol. Endocrinol.* **16**, 390-401
- Laemmli UK. (1970) Cleavage of structural proteins during the assembly of the head of the bacteriophage T4. *Nature* **227**, 680-685

- Leger-Silvestre I, Noaillac-Depeyre J, Faubladiere M, Gas N. (1997) Structural and functional analysis of the nucleolus of the fission yeast *Schizosaccharomyces pombe*. *Eur. J. Cell. Biol.* **72**, 13-23
- Levine AJ. (1997) p53, the cellular gatekeeper for growth and division. *Cell* **88**, 323-331
- Li H & Bingham PM. (1991) Arginine/serine-rich domains of the su(wa) and tra RNA processing regulators target proteins to a subnuclear compartment implicated in splicing. *Cell* **67**, 335-342
- Lowrie DJ, Stickney JT & Ip W. (2000) Properties of the nonhelical end domains of vimentin suggest a role in maintaining intermediate filament network structure. *J. Struct. Biol.* **132**, 83-94
- Mackay K & Mochly-Rosen D. (2001) Localization, anchoring, and functions of protein kinase C isozymes in the heart. *J. Mol. Cell. Cardiol.* **33**, 1301-1307
- Magnelli L & Chiarugi V. (1997) Regulation of p53 by protein kinase C during multi-stage carcinogenesis. *J. Cancer Res. Clin. Oncol.* **123**, 365-369
- Maison C, Pyrpassoulou A, Theodoropoulos PA & Georgatos SD. (1997) The inner nuclear membrane protein LAP1 forms a native complex with B-type lamins and partitions with spindle-associated mitotic vesicles. *EMBO J.* **16**, 4839-4850
- Malatesta M, Fakan S & Fischer U. (1999) The Sm core domain mediates targeting of U1 snRNP to subnuclear compartments involved in transcription and splicing. *Exp. Cell Res.* **249**, 189-198
- Mandil R, Ashkenazi E, Blass M, Kronfeld I, Kazimirsky G, Rosenthal G, Umansky F, Lorenzo PS, Blumberg PM & Brodie C. (2001) Protein kinase C α and protein kinase C δ play opposite roles in the proliferation and apoptosis of glioma cells. *Cancer Res.* **61**, 4612-4619
- Martelli AM, Sang N, Borgatti P, Capitani S, Neri LM. (1999) Multiple biological responses activated by nuclear protein kinase C. *J. Cell. Biochem.* **74**, 499-521
- Martin L, Crimando C & Gerace L. (1995) cDNA cloning and characterization of lamina-associated polypeptide 1C (LAP1C), an integral protein of the inner nuclear membrane. *J. Biol. Chem.* **270**, 8822-8828
- Martins SB, Eide T, Steen RL, Jahnsen T, Skalhegg BS & Collas P. (2000) HA95 is a protein of the chromatin and nuclear matrix regulating nuclear envelope dynamics. *J. Cell Sci.* **113**, 3703-3713
- Matunis MJ, Coutavas E & Blobel G. (1996) A novel ubiquitin-like modification modulates the partitioning of the Ran-GTPase-activating protein RanGAP1 between the cytosol and the nuclear pore complex. *J. Cell Biol.* **135**, 1457-1470
- McFarlane DE. (1983) Use of benzyldimethyl-n-hexadecylammonium chloride ("16-BAC"), a cationic detergent, in an acidic polyacrylamide gel electrophoresis system to detect base labile protein methylation in intact cells. *Anal. Biochem.* **132**, 231-235
- McFarlane DE. (1989) Two dimensional benzyldimethyl-n-hexadecylammonium chloride-sodium dodecyl sulfate preparative polyacrylamide gel electrophoresis: a high capacity high resolution technique for the purification of proteins from complex mixtures. *Anal. Biochem.* **176**, 457-463
- Meek DW. (1998) Multisite phosphorylation and the integration of stress signals at p53. *Cell Signal.* **10**, 159-166
- Melcak I, Cermanova S, Jirsova K, Koberna K, Malinsky J & Raska I. (2000) Nuclear pre-mRNA compartmentalization: trafficking of released transcripts to splicing factor reservoirs. *Mol. Biol. Cell* **11**, 497-510

- Mical TI & Monteiro MJ. (1998) The role of sequences unique to nuclear intermediate filaments in the targeting and assembly of human lamin B: evidence for a lack of interaction of lamin B with its putative receptor. *J. Cell Sci.* **111**, 3471-3485
- Mili S, Shu HJ, Zhao Y & Pinol-Roma S. (2001) Distinct RNP complexes of shuttling hnRNP proteins with pre-mRNA: candidate intermediates in formation and export of mRNA. *Mol. Cell. Biol.* **21**, 7307-7319
- Misteli T & Spector DL. (1998) The cellular organization of gene expression. *Curr. Opin. Cell Biol.* **10**, 323-331
- Mochly-Rosen D, Khaner H & Lopez J. (1991) Identification of intracellular receptor proteins for activated protein kinase C. *Proc. Natl. Acad. Sci. USA* **88**, 3997-4000
- Moosmann P & Rusconi S. (1996) Alpha complementation of LacZ in mammalian cells. *Nucl. Acids Res.* **24**, 1171-1172
- Morris GE & Manilal S. (1999) Heart to heart: from nuclear proteins to Emery-Dreifuss muscular dystrophy. *Hum. Mol. Genet.* **8**, 1847-1851
- Municio MM, Lozano J, Sanchez P, Moscat J, Diaz-Meco MT. (1995) Identification of heterogeneous ribonucleoprotein A1 as a novel substrate for protein kinase C zeta. *J. Biol. Chem.* **270**, 15884-15891
- Nacro K, Sigano DM, Yan S, Nicklaus MC, Pearce LL, Lewin NE, Garfield SH, Blumberg PM & Marquez VE. (2001) An optimized protein kinase C activating diacylglycerol combining high binding affinity (K_i) with reduced lipophilicity (log P). *J. Med. Chem.* **44**, 1892-1904
- Nakielny S, Shaikh S, Burke B & Dreyfuss G. (1999) Nup153 is an M9-containing mobile nucleoporin with a novel Ran-binding domain. *EMBO J.* **18**, 1982-1995
- Neri LM, Billi AM, Manzoli L, Rubbini S, Gilmour RS, Cocco L & Martelli AM. (1994) Selective nuclear translocation of protein kinase C alpha in Swiss 3T3 cells treated with IGF-I, PDGF and EGF. *FEBS Lett.* **347**, 63-68
- Newton AC. (1995) Protein kinase C: Structure, function and regulation. *J. Biol. Chem.* **270**, 28495-28498
- Nikolakaki E, Simos S, Georgatos SD & Giannakouros T. (1996) A nuclear envelope-associated kinase phosphorylates arginine-serine motifs and modulates interactions between the lamin B receptor and other nuclear proteins. *J. Biol. Chem.* **271**, 8365-8372
- Ohtsubo M, Okazaki H & Nishimoto T. (1989) The RCC1 protein, a regulator for the onset of chromosome condensation locates in the nucleus and binds to DNA. *J. Cell Biol.* **109**, 1389-1397
- O'Reilly DR, Miller LK & Luckow VA. (1992) Baculovirus expression vectors: a laboratory manual. W.H. Freeman and company, New York
- Orstavik S, Eide T, Collas P, Han IO, Tasken K, Kieff E, Jahnsen T & Skalhegg BS. (2000) Identification, cloning and characterization of a novel nuclear protein, HA95, homologous to A-kinase anchoring protein 95. *Biol. Cell* **92**, 27-37
- Otto H, Dreger M, Bengtsson L & Hucho F. (2001) Identification of tyrosine-phosphorylated proteins associated with the nuclear envelope. *Eur. J. Biochem.* **268**, 420-428
- Otto H, Buchner K, Beckmann R, Hilbert R & Hucho F. (1992) GTP-binding proteins in bovine brain nuclear membranes. *Neurochem. Int.* **21**, 409-414

- Owen PJ, Johnson GD & Lord JM. (1996) Protein kinase C-delta associates with vimentin intermediate filaments in differentiated HL60 cells. *Exp. Cell Res.* **225**, 366-373
- Paine PL, Moore LC & Horowitz SB. (1975) Nuclear envelope permeability. *Nature* **254**, 109-114
- Patskan G & Baxter CS. (1985) Specific stimulation of histone H2B and H4 phosphorylation in mouse lymphocytes by 12-O-tetra-decanol-13-acetate. *J. Biol. Chem.* **260**, 12899-12903
- Patton JG, Mayer SA, Tempst P & Nadal-Ginard B. (1991) Characterization and molecular cloning of polypyrimidine tract-binding protein: a component of a complex necessary for pre-mRNA splicing. *Genes Dev.* **5**, 1237-1251
- Patton JG, Porro EB, Galceran J, Tempst P & Nadal-Ginard B. (1993) Cloning and characterization of PSF, a novel pre-mRNA splicing factor. *Genes Dev.* **7**, 393-406
- Paulin-Levasseur M, Blake DL, Julien M & Rouleau L. (1996) The MAN antigens are non lamin constituents of the nuclear lamina in vertebrate cells. *Chromosoma* **104**, 367-379
- Perander M, Bjørkøy G, Johansen T. (2001) Nuclear import and export signals enable rapid nucleoplasmic shuttling of the atypical protein kinase C. *J. Biol. Chem.* **276**, 13015-13024
- Perides G, Harter C & Traub P. (1987) Electrostatic and hydrophobic interactions of the intermediate filament protein vimentin and its amino terminus with lipid bilayers. *J. Biol. Chem.* **262**, 13742-13749
- Perlaky L, Valdez BC & Bush H. (1997) Effects of cytotoxic drugs on translocation of nucleolar RNA helicase RH II/Gu. *Exp. Cell Res.* **235**, 413-420
- Peter M, Nakagawa J, Doree M, Labbe JC & Nigg EA. (1990) In vitro disassembly of the nuclear lamina and M phase-specific phosphorylation of lamins by cdc2 kinase. *Cell* **61**, 591-602
- Philbrick WM, Wysolmerski JJ, Galbraith S, Holt E, Orloff JJ, Yang KH, Vasavada RC, Weir EC, Broadus AE & Stewart AF. (1996) Defining the roles of parathyroid hormone-related protein in normal physiology. *Physiol. Rev.* **76**, 127-73
- Pinol-Roma S. & Dreyfuss G. (1992) Shuttling of pre-mRNA binding proteins between nucleus and cytoplasm. *Nature* **355**, 730-732
- Polioudaki H, Kourmouli N, Drosou V, Bakou A, Theodoropoulos PA, Singh PB, Giannakouros T & Georgatos SD. (2001) Histones H3/H4 form a tight complex with the inner nuclear membrane protein LBR and heterochromatin protein 1. *EMBO reports* **2**, 920-925
- Pollard VW, Michael WM, Naklelny S, Siomi MC, Wang F & Dreyfuss G. (1996) A novel receptor-mediated nuclear protein import pathway. *Cell* **86**, 985-994
- Reichelt R, Holzenburg A, Buhle EL, Jarnik M, Engel A & Aebi U. (1990) Correlation between structure and mass distribution of the nuclear pore complex and of distinct pore complex components. *J. Cell Biol.* **110**, 883-894
- Ribbeck K, Kutay U, Paraskeva E & Görlich D. (1999) The translocation of transportin-cargo complexes through nuclear pores is independent of both Ran and energy. *Curr. Biol.* **9**, 47-50
- Robbins J, Dilworth SM, Laskey RA & Dingwall C. (1991) Two interdependent basic domains in nucleoplasmic nuclear targeting sequence: identification of a class of bipartite nuclear targeting sequence. *Cell* **64**, 615-623

- Rolls MM, Stein PA, Taylor SS, Ha E, McKeon F & Rapoport TA. (1999) A visual screen of a GFP-fusion library identifies a new type of nuclear envelope membrane protein. *J. Cell Biol.* **146**, 29-34
- Ron D, Che-Hong Chen, Caldwell J, Jamieson L, Orr E & Mochly-Rosen D. (1994) Cloning of an intracellular receptor for protein kinase C: a homolog of the β subunit of G proteins. *Proc. Natl. Acad. Sci. USA* **91**, 839-843
- Rosenberger U. (1997) Lokalisierung und Verankerung der Proteinkinase C im neuronalen Zellkern. *Dissertation des Fachbereichs Biologie, Chemie Pharmazie FU Berlin*
- Roskams T & Desmet V. (1997) Parathyroid-hormone-related peptides. A new class of multifunctional proteins. *Am. J. Pathol.* **150**, 779-85
- Rout MP & Blobel G. Isolation of the yeast nuclear pore complex. (1993) *J. Cell Biol.* **123**, 771-83
- Rout MP, Aitchison JD, Suprpto A, Hjertaas K, Zhao Y & Chait BT. (2000) The yeast nuclear pore complex: composition, architecture, and transport mechanism. *J. Cell Biol.* **148**, 635-651
- Ruskin B, Zamore PD & Green MR. (1988) A factor, U2AF, is required for U2 snRNP binding and splicing complex assembly. *Cell* **52**, 207-19
- Ryan KJ & Wentz SR. (2000) The nuclear pore complex: a protein machine bridging the nucleus and cytoplasm. *Curr. Opin. Cell. Biol.* **12**, 361-371
- Sanjay A, Fu J & Kreibich G. (1998) DAD1 is required for the function and structural integrity of the oligosaccharyltransferase complex. *J. Biol. Chem.* **273**, 26094-26099
- Schmalz D, Kalkbrenner F, Hucho F & Buchner K. (1996) Transport of protein kinase C alpha into the nucleus requires intact cytoskeleton while the transport of a protein containing a canonical nuclear localization signal does not. *J. Cell Sci.* **109**, 2401-2406
- Schmalz D, Hucho F & Buchner K. (1998) Nuclear import of protein kinase C occurs by a mechanism distinct from the mechanism used by proteins with a classical nuclear localization signal. *J. Cell Sci.* **111**, 1823-1830
- Shevchenko A, Wilm M, Vorm O & Mann M. (1996) Mass spectrometric sequencing of proteins silver-stained polyacrylamide gels. *Anal. Chem.* **68**, 850-858
- Shimizu T, Cao CX, Shao RG & Pommier Y. (1998) Lamin B phosphorylation by protein kinase C α and proteolysis during apoptosis in human leukemia HL60 cells. *J. Biol. Chem.* **273**, 8669-8674
- Simons G & Georgatos SD. (1992) The inner nuclear membrane protein p58 associates in vivo with a p58 kinase and the nuclear lamins. *EMBO J.* **11**, 4027-4036
- Simons G & Georgatos SD. (1994) The lamin B receptor-associated protein p34 shares sequence homology and antigenic determinants with the splicing factor 2-associated protein p32. *FEBS Lett.* **346**, 225-228
- Sleeman JE & Lamond AI. (1999) Nuclear organization of pre-mRNA splicing factors. *Curr. Opin. Cell. Biol.* **11**, 372-377
- Sorg G & Stamminger T. (1999) Mapping of nuclear localisation signals by simultaneous fusion to green fluorescent protein and to β -galactosidase. *BioTechniques* **26**, 858-862

- Sozeri O, Vollmer K, Liyanage M, Frith D, Kour G, Mark GE & Stabel S. (1992) Activation of the c-Raf protein kinase by protein kinase C phosphorylation. *Oncogene* **7**, 2259-2262
- Staknis D & Reed R. (1994) Direct interactions between pre-mRNA and six U2 small nuclear ribonucleoproteins during spliceosome assembly. *Mol. Cell. Biol.* **14**, 2994-3005
- Sukegawa J & Blobel G. (1993) A nuclear pore complex protein that contains zinc finger motifs, binds DNA, and faces the nucleoplasm. *Cell* **72**, 29-38
- Takano M, Takeuchi M, Ito H, Furukawa K, Sugimoto K, Omata S & Horigome T. (2002) The binding of lamin B receptor to chromatin is regulated by phosphorylation in the RS region. *Eur. J. Biochem.* **269**, 943-953
- Tanese N, Pugh BF & Tjian R. (1991) Coactivators for a proline-rich activator purified from the multisubunit human TFIID complex. *Genes Dev.* **5**, 2212-2224
- Traub P & Shoeman RL. (1994a) Intermediate filament proteins: Cytoskeletal elements with gene-regulatory function? *Int. Rev. Cytol.* **154**, 1-103
- Traub P & Shoeman RL. (1994b) Intermediate filament and related proteins: Potential activators of nucleosomes during transcription initiation and elongation? *BioEssays* **16**, 349-355
- Valdez BC, Henning D, Perumal K & Bush H. (1997) RNA unwinding and RNA folding activities of RNA helicase II/Gu: two activities in separate domains of the same protein. *Eur. J. Biochem.* **250**, 800-807
- Valdez BC. (2000) Structural domains involved in the RNA folding activity of RNA helicase II/Gu protein. *Eur. J. Biochem.* **267**, 6395-6402
- Vallentin A, Prévostel C, Fauquier T, Bonnefont X & Joubert D. (2000) Membrane targeting and cytoplasmic sequestration in the spatiotemporal localization of human protein kinase Ca. *J. Biol. Chem.* **275**, 6014-6021
- Vaughn JL, Goodwin RH, Tompkins GJ & McCawley P. (1977) The establishment of two cell lines from the insect *Spodoptera frugiperda* (Lepidoptera; Noctuidae). *In Vitro*, **13**, 213-217
- Wagner S. (1999) Untersuchungen zur subzellulären Verteilung von Proteinkinase C. *Dissertation des Fachbereichs Biologie, Chemie Pharmazie FU Berlin*
- Wagner S, Harteneck C, Hucho F & Buchner K. (2000) Analysis of the subcellular distribution of Protein Kinase Ca using PKC-GFP fusion proteins. *Exp. Cell Res.* **258**, 204-214
- Wang QJ, Bhattacharyya D, Garfield S, Nacro K, Marquez VE & Blumberg PM. (1999) Differential localization of protein kinase C δ by phorbol esters and related compounds using a fusion protein with green fluorescent protein. *J. Biol. Chem.* **274**, 37233-37239
- Wells NJ, Fry AM, Guano F, Norbury C & Hickson ID. (1995) Cell cycle phase-specific phosphorylation of human topoisomerase II α . Evidence of a role for protein kinase C. *J. Biol. Chem.* **270**, 28357-28363
- Williams DS, Park S, Schlamp CL, Newton AC. (1994) Protein kinase C association with the retinal cytoskeleton and phosphorylation of vimentin. *Eye Eye Res.* **58**, 747-59
- Williamson MP. (1994) The structure and function of proline-rich regions in proteins. *Biochem. J.* **297**, 249-260

- Winstel R, Freund S, Krasel C, Hoppe E & Lohse MJ. (1996) Protein kinase cross-talk: membrane targeting of the β -adrenergic receptor kinase by protein kinase C. *Proc. Natl. Acad. Sci. USA* **93**, 2105-2109
- Worman HJ, Yuan J, Blobel G & Georgatos SD. (1988) A lamin B receptor in the nuclear envelope. *Proc. Natl. Acad. Sci. USA* **85**, 8531-8534
- Worman HJ, Evans CD & Blobel G. (1990) The lamin B receptor of the nuclear envelope inner membrane: a polytopic protein with eight potential transmembrane domains. *J. Cell Biol.* **111**, 1535-1542
- Yang L, Guan T & Gerace L. (1997) Integral membrane proteins of the nuclear envelope are dispersed throughout the endoplasmic reticulum during mitosis. *J. Cell Biol.* **137**, 1199-1210
- Yaseen NR & Blobel G. (1999) Two distinct classes of Ran-binding sites on the nucleoporin Nup-358. *Proc. Natl. Acad. Sci. USA* **96**, 5516-5521
- Ye Q & Worman HJ. (1996) Interaction between an integral protein of the nuclear envelope inner membrane and human chromodomain proteins homologous to *Drosophila* HP1. *J. Biol. Chem.* **271**, 14653-14656
- Yokoyama N, Hayashi N, Seki T, Pante N, Ohba T, Nishii K, Kuma K, Hayashida T, Miyata T, Aebi U, et al. (1995) A giant nucleopore protein that binds Ran/TC4. *Nature* **376**, 184-188
- Zolotukhin S, Potter M, Hauswirth WW, Guy J & Muzyczka N. (1996) A "humanized" green fluorescent protein cDNA adapted for high-level expression in mammalian cells. *J. Virol.* **70**, 4646-4654