
7. Literaturverzeichnis

Akiskal HS. Mood disorders: introduction and overview. In Comprehensive Textbook Psychiatry, B.J. Sadock and V.A. Sadock, eds New York:Lippincott, Williams& Wilkins 2000;1284-1298

Auer DP, Putz B, Kraft E, Lipinski B, Schill J, Holsboer F. Reduced glutamate in the anterior cingulate cortex in depression: an in vivo proton magnetic resonance spectroscopy study. Biological Psychiatry 2000;47:305-313

Baghai TC, Frey R, Kasper S, Elektrokonvulsionstherapie, Klinische und wissenschaftliche Aspekte, Kapitel: Repetitive transkranielle Magnetstimulation (rTMS) in der Behandlung psychiatrischer Erkrankungen 2003

Bajbouj M, Heuser I, Antidepressive Stimulationsverfahren, Nervenarzt 2005;76:28-35

Banay-Schwartz M, Lajtha A, Palkovits M. Regional distribution of glutamate and aspartate in adult and old human brain. Brain Res. 1992;594:343-346

Barker AT, Jalinous R, Freeston IL, Non-invasive stimulation of the human motor cortex, Lancet 1985;1106-1107

Beck AT, Ward CH, Mendelson M, et al. An inventory for measuring depression, Archives of General Psychiatry 1961;4:561-71.

Behar KL, den Hollander JA, Stromski ME, et al. High-resolution ¹H nuclear magnetic resonance study of cerebral hypoxia in vivo. Proc Natl Acad Sci USA 1983;80: 4945-4948

Berger M., Psychiatrische Erkrankungen, Klinik und Therapie, Urban und Fischer Verlag 2004

Berger M., Psychiatrische Erkrankungen Klinik und Therapie, 2. überarbeitete Auflage, Urban & Fischer 2003;98-99

Bloch FW, Hansen WW, Packard ME Nuclear Induction. Phys Rev 1946;69: 127

Brakemeier EL, Luborzewski A, Danker-Hopfe H, et al. Positive predictors for antidepressive response to prefrontal repetitive transcranial magnetic stimulation (rTMS)., J Psychiatr Res. 2006;Mar 20

Bunney WE Jr., Davis JM, Norepinephrine in depressive reactions. A review. Arch. Gen. Psychiatry 13 1965;483-494

Burmeister M, Basic concepts in the study of diseases with complex genetics. Biol. Psychiatry 45 1999;522-532

Burt T, Lisanby SH, Sackheim HA, Neuropsychiatric applications of transcranial magnetic stimulation: a meta analysis, Int J Neuropsychopharmacol. 2002;Mar;5(1):73-103

Conca A, Koppi S, Konig P, et al. Transcranial magnetic stimulation: a novel antidepressive strategy?, Neuropsychobiology 1996;34(4):204-7

Cordes J, Mobascher A, Arends M, et al. Ein neues Behandlungsverfahren der Depression: Die repetitive transkranielle Magnetstimulation (rTMS), Dtsch Med Wochenschr 2005;130:889-892

D'Arsonval A. Dispositifs pour la mesure des courants alternatifs de toutes frequences. CR Société Biologique (Paris) 1896;2 May: 450-451

Diehl DJ, Gershon S., The role of dopamine in mood disorders., Compr Psychiatry. 1992;Mar-Apr;33(2):115-20.

Eschweiler GW, Wegerer C, Schlotter W, et al. Left prefrontal activation predicts therapeutic effects of repetitive transcranial magnetic stimulation (rTMS) in major depression. Psychiatry Research 2000;99:161-172

Faraday M. Effects of the production of electricity from magnetism (1831) In: Faraday M, Williams LP, editors. Basic Books, New York, 1965;531-540

Fava M, Kendler KS. Major depressive disorder. *Neron* 2000;335-341

Figiel GS, Epstein C, McDonald WM, et al. The use of rapid-rate transcranial magnetic stimulation (rTMS) in refractory depressed patients., *J Neuropsychiatry Clin Neurosci.* 1998;Winter;10(1):20-5.

Friemel S, Bernert S, Angermeyer MC, et al. The direct costs of depressive disorders in Germany. *Psychiatr Prax.* 2005;Apr;32(3):113-21

Fuster JM, Transmitters in the prefrontal cortex. aus Fuster JM, *The prefrontal Cortex: Anatomy, Physiology and Neuropsychology of the Frontal Lobe*, 3rd Edition. Lippincott-Raven, Philadelphia 1997;45-60

George MS, Wassermann EM, Williams WA, et al. Daily repetitive transcranial magnetic stimulation (rTMS) improves mood in depression. *Neuroreport.* 1995;Oct 2;6(14):1853-6.

George MS, Wassermann EM, Williams WA, et al., Changes in mood and hormone levels after rTMS of the prefrontal cortex, *J Neuropsychiatry Clin Neurosci* 1996; 8:172-180

George MS, Wassermann EM, Kimbrell TA, et al., Mood improvement following daily left prefrontal repetitive magnetic stimulation in patients with depression: a placebo-controlled crossover trial. *Am J Psychiatry* 1997;54:1752-1756

George MS, Speer AM, Molloy Ma, et al. Low-frequency daily left prefrontal rTMS improves mood in bipolar depression: a placebo-controlled case report. *Hum Psychopharmacol* 1998;13:271-275

George MS, Lisanby SH, Sackeim HA, Transcranial magnetic stimulation. Applications in Neuropsychiatry. *Arch Gen Psychiatry* 56 1999;300–311

Gershon AA, Dannon PN, Grunhaus L. Transcranial magnetic stimulation in the treatment of depression. *Am J Psychiatry* 2003;160:835-845

Grachev ID, Swarnkar A, Szeverenyi NM, et al. Aging alters regional multichemical profile of the human brain: in vivo ¹H-MRS study of young versus middle-aged subjects. *J Neurochem.* 2001;Jan;76(2):582-593

Gruber S, Frey R, Mlynárik V, et al. Quantification of Metabolic Differences in the Frontal Brain of Depressive Patients and Controls Obtained by ¹H-MRS at 3 Tesla. *Investigative Radiology* 2003;July; 38 (7):403-408

Grunhaus L, Dannon PN, Schreiber S, et al. Repetitive transcranial magnetic stimulation is as effective as electroconvulsive therapy in the treatment of nondelusional major depressive disorder: an open study., *Biol Psychiatry* 2000;47:314-324

Hamilton M. A rating scale for depression. *Journal of Neurology, Neurosurgery & Psychiatry* 1960;23:56-62.

Holtzheimer III PE, Russo J, Claypoole KH, et al. Shorter duration of depressive episode may predict response to repetitive transcranial magnetic stimulation. *Depression and Anxiety* 2004;19:24-30

Janowsky DS, Overtstreet DH, The role of acetylcholine metabolisms in mood disorders. In: Bloom FE, Kupfer DJ *Psychopharmacology, the fourth generation of progress*, Raven, New York 1978

Kaiser LG, Schuff N, Cashdollar N, et al. Age related glutamate and glutamine concentration changes in normal human brain: ¹H-MR spectroscopy study at 4 T. *Neurobiology of Aging* 2005;26:665-672

Kaufmann GW, Moser E, Sauer R. *Radiologie, 2. erweiterte Auflage*. Urban und Fischer Verlag München 2001;102-118

Klein E, Kreinin I, Chistyakov A, et al. Therapeutic efficacy of right prefrontal slow repetitive transcranial magnetic stimulation in major depression: a double-blind controlled study. *Arch Gen Psychiatry* 1999;Apr 56(4):300-311

Klein J, Membrane breakdown in acute and chronic neurodegeneration: focus on choline-containing phospholipids. *J Neural Transm* 107: 2000;1027-1063

Kolb B., Whishaw J.Q., *Neuropsychologie*, Spektrum Akademischer Verlag, 2. Auflage 1996;257- 261

Lanfermann H, Herminghaus S, Pilatus U, et al. Grundlagen der ¹H-MR-Spektroskopie intrakranieller Tumoren. *Klin Neuroradiol* 12:2000; 1-17

Laubenberger J., Bayer S., Thiel T., et al., *Klinische Anwendungen der Protonen-Magnetresonanzspektroskopie des Gehirns*. *RöFo* 1998;168: 539-549

Lauterbur PC Image formation by induced local interactions. Examples employing NMR. *Nature* 1973;242: 190

Leitner I, Aschauer H, Bailer U, et al. Behandlungsmöglichkeiten der therapieresistenten Depression. *Journal für Neurologie, Neurochirurgie und Psychiatrie* 2004; 5 (1): 28-39

Malhi G.S., Valenzuela M., Wen W., et al. Magnetic resonance spectroscopy and its applications in psychiatry. *The Australian and New Zealand Journal of Psychiatry*, 2002;36: 31-43

Michael N, Gosling M, Reutemann M, et al. Metabolic changes after repetitive transcranial magnetic stimulation (rTMS) of the left prefrontal cortex: a sham-controlled proton magnetic resonance spectroscopy (¹H MRS) study of healthy brain., *Eur J Neurosci*. 2003;Jun;17(11):2462-8.

Montgomery SA, Asberg M. A new depression scale designed to be sensitive to change. *British Journal of Psychiatry* 1979;134:382-9.

Möller HJ, Laux, Kapfhammer, Psychiatrie und Psychotherapie, Hippocrates Verlag, Stuttgart 2001

Möller HJ, Standardisierte psychiatrische Befunddiagnostik. In: Möller H.J., Laux G., Kapfhammer H.P. (Hrsg.) Psychiatrie und Psychotherapie, Hippocrates Verlag, Stuttgart 2001;73-106

Murray CJ, Lopez AD. Alternative Visions of the Future: Projecting Mortality and Disability, 1990-2020. In: Murray CJ, Lopez AD: The global burden of disease: a comprehensive assessment of mortality and disability from diseases, injuries, and risk factors in 1990 and projecting to 2020. Cambridge, Mass: Harvard Univ. Press, 1996; 325- 395.

Murray CJ, Lopez AD. The global burden of disease, 1990-2020. Nature Medicine Nov 1998;4(11) 1241-1243

Nahas Z, Speer AM, Molloy M, et al. Preliminary results concerning the roles of frequency and intensity in the antidepressant effect of daily left prefrontal rTMS (abstract). Biol Psychiatry. 1998;43:94

Padberg F, Zwanzger P, Thoma H, et al. Repetitive transcranial magnetic stimulation (rTMS) in pharmacotherapy-refractory major depression: comparative study of fast, slow and sham rTMS., Psychiatry Res. 1999;Nov 29;88(3):163-71.

Parker G, Hadzi-Pavlovic D, Austin MP, et al. Sub-typing depression, I. Is psychomotor disturbance necessary and sufficient to the definition of melancholia? Psychological Medicine 1995;25: 815-823

Pascual-Leone A, Rubie B, Pallardo F, et al. Rapid-rate transcranial magnetic stimulation of the left dorsolateral prefrontal cortex in drug-resistant depression; Lancet 1996a;348:23-237

Pascual-Leone A, Catala MD, Lateralized effect of rapid-rate transcranial magnetic stimulation of the prefrontal cortex on mood. Neurology. 1996b;Feb;46(2):499-502.

Paus T, Castro-Alamancos MA, Pertrides M., Cortica-cortical connectivity of the human mid-dorsolateral frontal cortex and its modulation by repetitive transcranial magnetic stimulation. *European Journal of Neuroscience* 1997;17 (9): 3178-3184

Paus T, Primate anterior cingulate cortex: where motor control, drive and cognition interface. *Nature Reviews Neuroscience* 2001;14:1405-1411

Paus T, Barrett J. Transcranial magnetic stimulation (TMS) of the human frontal cortex: implications for repetitive TMS treatment of depression. *J Psychiatry Neurosci* 2004;29(4):268-279

Pouwels P, Brockmann K, Kruse B, et al. Regional age dependence of human brain metabolites from infancy to adulthood as detected by quantitative localized proton MR. *Pediatric Research* 1999;46(6): 474-485

Pridmore S., Substitution of rapid transcranial magnetic stimulation treatments for electroconvulsive therapy treatments in a course of electroconvulsive therapy. *Depress Anxiety* 12 2000;118-123

Protonen-Magnetresonanzspektroskopie des Gehirns. *RöFo*, 168: 539-549

Prudic J, Sackeim HA, Devanand DP. Medication resistance and clinical response to electroconvulsive therapy. *Psychiatry Res* 1990;31:287-296

Prudic J, Haskett RF, Mulsant B, et al. Resistance to antidepressant medications and short-term clinical response to ECT. *Am J Psychiatry* 1996;153:985-992

Ross B, Michaelis T. Clinical applications of magnetic resonance spectroscopy. *Magn Reson Q*. 1994;Dec;10(4):191-247

Rossini PM, Barker AT, Berardelli A, et al.: Non-invasive electrical and magnetic stimulation of the brain, spinal cord and roots: basic principles and procedures for routine clinical application. Report of an IFCN committee. *Electroencephalogr Clin Neurophysiol*. 1994;Aug;91(2):79-92.

Rothwell JC, Techniques and mechanisms of action of transcranial stimulation of the human motor cortex. *J Neurosci Methods* 1997;74(2):113-22

Rush AJ, George MA, Sackeim HA, et al. Vagus nerve stimulation (VNS) for treatment-resistant depressions: a multicenter study. *Biol Psychiatry* 2000; 47:276-286

Sackeim HA. Continuation therapy following ECT: directions for the future research. - *Psychopharmacol Bull* 1994; 30:501-521

Sackeim HA, Prudic J, Devanand DP, et al. A prospective, randomized, double-blind comparison of bilateral and right unilateral electroconvulsive therapy at different stimulus intensities, *Arch Gen Psychiatry* 2000;57:425-434

Sahai S, Glutamate in the mammalian CNS, *Eur Arch Psychiatry Clin Neurosci* 1990;240: 121-133

Salze HJ, Stamm K, Schubert M, et al. Behandlungskosten von Patienten mit der Depressionsdiagnose in haus- und fachärztlicher Versorgung in Deutschland. *Psychiat Prax* 2004;31:147-156

Sanacora G et al., Subtype-specific alterations of gamma-aminobutyric acid and glutamate in patients with major depression. *Arch Gen Psychiatry* 2004;Vol 61, 705-713

Sanders JA, *Magnetic Resonance Spectroscopy, Functional Brain* 1995; 419-463,

Schildkaut JJ, The catecholamine hypothesis of affective disorders: a review of supporting evidence. *Am. J. Psychiatry* 1965;122:509-522.

Spectroscopy and its applications in psychiatry. The Australian and New Zealand Springer Berlin Heidelberg New York, 2. Auflage 2003;412- 421

Schubert F, Gallinat J, Seifert F, et al. Glutamat concentrations in human brain using single voxel proton magnetic resonance spectroscopy at 3 Tesla. *NeuroImage* 2004;21:1762-1771

Statistisches Bundesamt, Gesundheitsbericht für Deutschland, 2002

Steward O, Functional Neuroscience. Springer Verlag. 2000

The Avery-George-Holtzheimer database of rTMS depression studies. Available: www.itis.unibe.ch/MSAvery.htm (accessed 2004 June 11)

Thier P., Die funktionelle Architektur des präfrontalen Kortex, In: Neuropsychologie, Karnath H.O., Thier P., Springer-Verlag 2003;498- 503

Trepel M., Neuroanatomie, Urban & Fischer Verlag, 2. überarbeitete Auflage 1999

Wassermann EM, Wedegaertner FR, Ziemann U, et al. Crossed reduction of human motor cortex excitability by 1-Hz transcranial magnetic stimulation. *Neurosci Lett.* 1998;Jul 10;250(3):141-144.

Wassermann EM. Risk and safety of repetitive transcranial magnetic stimulation: report and suggested guidelines from the international workshop on the safety of repetitive transcranial magnetic stimulation, June-7, 1996, *Electroencephalogr. Clin. Neurophysiol.* 1998;108: 1-16

Weber-Fahr W, Braus DF. Magnetresonanztomographie (MRS), In: Funktionelle Bildgebung in Psychiatrie und Psychotherapie, Walter H., Schattauer 2005;114-128

Wittchen HU, Pittrow D. Prevalence, recognition and management of depression in primary care in Germany: the Depression 2000 study. *Human Psychopharmacology Clin Exp* 2002;17:1-11