

V.

LITERATURVERZEICHNIS

- Alani, E., Cao, L., Kleckner, N.** 1987. A method for gene disruption of multiply disrupted yeast strains. *Genetics*. 116: 541-545.
- Anderson, J., Cundiff, M., Schnars, L., Gao, M., Mackenzie, I., Soll, D. R.** 1989. Hypha formation in the white-opaque transition of *Candida albicans*. *Infect. Immun.* 57: 458-467.
- Anderson, J., Mihalik, K., Soll, D. R.** 1990. Ultrastructure and antigenicity of the unique cell and pimple of the *Candida* opaque phenotype. *J. Bacteriol.* 172: 224-235.
- Anthley, P. P. & Hazen, K. C.** 1988. Role of yeast cell growth temperature on *Candida albicans* virulence in mice. *Infect. Immun.* 56: 2884-2890.
- Ash, J., Dominguez, M., Bergeron, J. M., Thomas, D. Y., Bourbonnais, Y.** 1995. The yeast proprotein convertase encoded by *YAP3* is a glycosylphosphatidylinositol-anchored protein that localizes to the plasma membrane. *J. Biol. Chem.* 270: 20847-20854.
- Azaryan, A. V., Wong, M., Friedman, T. C., Cawley, N. X., Estivariz, F. E., Chen, H. C., Loh, Y. P.** 1993. Purification and characterization of a paired basic residue-specific yeast aspartic protease encoded by the *YAP3* gene. Similarity to the mammalian pro-opiomelanocortin-converting enzyme. *J. Biol. Chem.* 268:11968-11975.
- Bader, O., Schaller, M., Klein, S., Kukula, J., Haak, K., Mühlischlegel, F., Korting, H. C., Schäfer, W., Hube, B.** 2001. The *KEX2* gene of *Candida glabrata* is required for cell surface integrity. *Mol. Microbiol.* 41: 1-15.
- Banerjee, A., Ganesan, K., Datta, A.** 1991. Induction of secretory acid proteinase in *Candida albicans*. *J. Gen. Microbiol.* 137:2455-2461.
- Beausejour, A., Grenier, D., Goulet, J.-P., Deslauriers, N.** 1998. Proteolytic activation of the Interleukin-1 β precursor by *Candida albicans*. *Infect. Immun.* 66: 676-681.
- Berka, R. M., Carmona, C. L., Hayenga, K. J., Thompson, S. A., Ward, M.** 1993. Isolation and characterization of the *Aspergillus oryzae* gene encoding aspergillopepsin O. *Gene* 125: 195-198.
- Berka, R. M., Ward, M., Wilson, L. J., Hayenga, K. J., Kodama, K. H., Carlomagno, L. P., Thompson, S. A.** 1990. Molecular cloning and deletion

of the gene encoding aspergillopepsin A from *Aspergillus awamori*. Gene 86: 153-162.

Boeke, J. D., Lacroute, F., Fink, G. R. 1984. A positive selection for mutants lacking orotidine-5'-phosphate decarboxylase activity in yeast: 5-fluoro-orotic acid resistance. Mol. Gen. Genet. 197: 345-346.

Borg-von Zepelin, M., Meyer, I., Thomssen, R., Wurzner, R., Sanglard, D., Telenti, A., Monod, M. 1999. HIV-Protease inhibitors reduce cell adherence of *Candida albicans* strains by inhibition of yeast secreted aspartic proteases. J. Invest. Dermatol. 113: 747-751.

Borg-von Zepelin, M., Beggah, S., Boggian, K., Sanglard, D., Monod, M. 1998. The expression of the secreted aspartyl proteinases Sap4 to Sap6 from *Candida albicans* in murine macrophages. Mol. Microbiol. 28: 543-554.

Bowman, B. H., Taylor, J. W., White, T. J. 1992. Molecular evolution of the fungi: human pathogens. Mol. Biol. Evol. 9: 893-904.

Brown, A. J. P. & Gow, N. A. R. 1999. Regulatory network controlling *Candida albicans* morphogenesis. Trend. Microbiol. 7: 333-338.

Buffo, J., Herman, M. A., Soll, D. R. 1984. A characterization of pH-regulated dimorphism in *Candida albicans*. Mycopathologia 85: 21-30.

Caro, L. H. P., Tettelin, H., Vossen, J. H., Ram, A. F. J., Van den Ende, H., Klis, F. M. 1997. *In silico* identification of glycosyl-phosphatidylinositol-anchored plasma-membrane and cell wall proteins of *Saccharomyces cerevisiae*. Yeast 13: 1477-1489.

Chaffin, W. L., Lopez-Ribot, J. L., Casanova, M., Gozalbo, D., Martinez, J. P. 1998. Cell wall and secreted proteins of *Candida albicans*: identification, function, and expression. Microbiol. Mol. Biol. Rev. 62: 130-180.

Chomczynski, P. & Sacchi, N. 1987. Single step method for RNA isolation by acid guanidinium-thiocyanate-phenol-chloroform extraction. Anal. Biochem. 162:156-159.

Clark, S. J., Templeton, M. D., Sullivan, P. A. 1997. A secreted aspartic proteinase from *Glomerella cingolata*: purification of the enzyme and molecular cloning of the cDNA. Microbiol. 143: 1395-1403.

- Cole, G. T., Halawa, A. A., Anaissie, E. J.** 1996. The role of the gastrointestinal tract in hematogenous candidiasis: from the laboratory to the bedside. *Clin. Infect. Dis.* 22(Suppl. 2): S73-S78.
- Colina, A. R., Aumont, F., Deslauriers, N., Belhumeur, P., De Repentigny, L.** 1996. Evidence for degradation of gastrointestinal mucin by *Candida albicans* secretory aspartyl proteinase. *Infect. Immun.* 64: 4514-4519.
- Crandall, M. & Edwards, J. E. Jr.** 1987. Segregation of proteinase-negative mutants from heterozygous *Candida albicans*. *J. Gen. Microbiol.* 133: 2817-2824.
- Csank, C., Schröppel, K., Leberer, E., Harcus, D., Mohamed, O., Meloche, S., Thomas, D. Y., Whiteway, M.** 1998. Roles of the *Candida albicans* mitogen-activated protein kinase homolog, Cek1p, in hyphal development and systemic candidiasis. *Infect. Immun.* 66: 2713-2721.
- Cutler, J. E.** 1991. Putative virulence factors of *Candida albicans*. *Annu. Rev. Microbiol.* 45: 187-218.
- Davies, D. R.** 1990. The structure and function of the aspartic proteinases. *Ann. Rev. Biophys. Biophys. Chem.* 19: 189-215.
- Davis, D., Wilson, R. B., Mitchell, A. P.** 2000. *RIM101*-dependent and -independent pathways govern pH responses in *Candida albicans*. *Mol. Cell. Biol.* 20: 971-978.
- De Bernardis, F., Arancia, S., Morelli, L., Hube, B., Sanglard, D., Schäfer, W., Cassone, A.** 1999. Evidence that members of the secretory aspartic proteinase gene family, in particular *SAP2*, are virulence factors for *Candida vaginitis*. *J. Infect. Dis.* 179: 201-208.
- De Bernardis, F., Cassone, A., Sturtevant, J., Calderone, R.** 1995. Expression of *Candida albicans* *SAP1* and *SAP2* in experimental vaginitis. *Infect. Immun.* 63: 1887-1892.
- De Bernardis, F., Mühlischlegel, F. A., Cassone, A., Fonzi, W. A.** 1998. The pH of the host niche controls gene expression in and virulence of *Candida albicans*. *Infect. Immun.* 66: 3317-3325.
- De Bernardis, F., Sullivan, P. A., Cassone, A.** 2001. Aspartyl proteinases of *Candida albicans* and their role in pathogenicity. *Med. Microbiol.* 39: 303-

313.

De Repentigny, L., Aumont, F., Bernard, K., Belhumeur, P. 2000. Characterization of binding of *Candida albicans* to small intestinal mucin and its role in adherence to mucosal epithelial cells. *Infect. Immun.* 68: 3172-3179.

De Sampaio, G., Bourdineaud, J.-P., Lauquin, J.-M. 1999. A constitutive role for GPI anchors in *Saccharomyces cerevisiae*: cell wall targeting. *Mol. Microbiol.* 34: 247-256.

De Viragh, P. A., Sanglard, D., Togni, G., Falchetto, R., & Monod, M. 1993. Cloning and sequencing of two *Candida parapsilosis* genes encoding acid proteases. *J. Gen. Microbiol.* 139: 335-342.

Delbrück, S. & Ernst, J. F. 1993. Morphogenesis-independent regulation of actin transcript levels in the pathogenic yeast *Candida albicans*. *Mol. Microbiol.* 10: 859-866.

Egel-Mitani, M., Flygenring, H. P., Hansen, M. T. 1990. A novel aspartyl protease allowing *KEX2*-independent MF alpha propheromone processing in yeast. *Yeast* 6: 127-137.

El Barkani, A., Kurzai, O., Fonzi, W. A., Ramon, A., Porta, A., Frosch, M., Mühschlegel, F. A. 2000. Dominant active alleles of *RIM101 (PRR2)* bypass the pH restriction on filamentation of *Candida albicans*. *Mol. Cell. Biol.* 20: 4635-4647.

Enschete, M. L. & West, B. C. 1980. *Saccharomyces cerevisiae* septicemia. *Arch. Intern. Med.* 140: 1539.

Ernst, J. F. 2000. Transcription factors in *Candida albicans* – environmental control of morphogenesis. *Microbiol.* 146: 1763-1774.

Fallon, K., Bausch, K., Noonan, J., Huguenel, E., Tamburini, P. 1997. Role of aspartic proteases in disseminated *Candida albicans* infection in mice. *Infect. Immun.* 65: 551-556.

Farley, P. C. & Sullivan, P. A. 1998. The *Rhizopus oryzae* secreted aspartic proteinase gene family: an analysis of gene expression. *Microbiol.* 144: 2355-2366.

Felk, A. 1998. Herstellung einer sekretorischen Aspartyl-Protease $\Delta sap8$ -Einzelmutante und einer $\Delta sap1, 3, 8$ -Dreifachmutante. Identifizierung,

Klonierung und Charakterisierung eines sekretorischen Aspartyl-Protease Gens (SAP) 10. Diplomarbeit. Fachbereich Biologie der Universität Hamburg.

Felk, A., Kretschmar, M., Albrecht, A., Schaller, M., Beinhauer, S., Nichterlein, T., Sanglard, D., Korting, H. C., Schäfer, W., Hube, B. 2002. *Candida albicans* hyphal formation and the expression of the Efg1-regulated proteinases Sap4-6 are required for the invasion of parenchymal organs. *Infect. Immun.* 70:3689-3700.

Felk, A., Schäfer, W., Hube, B. 2000. GenBank accession number AF146440.

Feng, Q., Summers, E., Guo, B., Fink, G. 1999. Ras signalling is required for serum-induced hyphal differentiation in *Candida albicans*. *J. Bacteriol.* 181: 6339-6346.

Filler, S. G., Ibe, B. O., Lukett, P. M., Raj, J. U., Edwards, J. E. Jr. 1991. *Candida albicans* stimulates endothelial cell eicosanoid production. *J. Infect. Dis.* 164: 928-935.

Filler, S. G., Pfunder, A. S., Spellberg, B. J., Spellberg, J. P., Edwards, J. E. Jr. 1996. *Candida albicans* stimulates cytokine production and leukocyte adhesion molecule expression by endothelial cells. *Infect. Immun.* 64: 2609-2617.

Fonzi, W. A. & Irwin, M. Y. 1993. Isogenic strain construction and gene mapping in *Candida albicans*. *Genetics* 134: 717-728.

Fox, J. L. 1993. Fungal infection rates are increasing. *ASM News* 10:515-518.

Fridkin, S. K. & Jarvis, W. R. 1996. Epidemiology of nosocomial fungal infections. *Clin. Microbiol. Rev.* 9: 499-511.

Ganesan, K., Banerjee, A., Datta, A. 1991. Molecular cloning of the secretory acid proteinase gene from *Candida albicans* and its use as a species-specific probe. *Infect. Immun.* 59: 2972-2977.

Gerber, L. D., Kodukula, K., Udenfriend, S. 1992. Phosphatidylinositol glycan (PI-G) anchored membrane proteins. Amino acid requirements adjacent to the site of cleavage and PI-G attachment in the COOH-Terminal signal peptide. *J. Biol. Chem.* 267: 12168-12173.

- Ghannoum, M. A., Swairjo, I., Soll, D. R.** 1990. Variation in lipid and sterol contents in *Candida albicans* white and opaque phenotypes. *J. Med. Vet. Mycol.* 28: 103-117.
- Gilfillan, G. D., Sullivan, D. J., Haynes, K., Parkinson, T., Coleman, D. C., Gow, N. A. R.** 1998. *Candida dubliniensis*: phylogeny and putative virulence factors. *Microbiol.* 144: 829-838.
- Gillum, A. M., Tsay, E. Y. H., Kirsch, D. R.** 1984. Isolation of the *Candida albicans* gene for orotidine-5'-phosphate decarboxylase by complementation of *S. cerevisiae ura3* and *E. coli pyrF* mutations. *Mol. Gen. Genet.* 198: 179-182.
- Gimeno, C. J. & Fink, G. R.** 1994. Induction of pseudohyphal growth by overexpression of *PHD1*, a *Saccharomyces cerevisiae* gene related to transcriptional regulators of fungal development. *Mol. Cell. Biol.* 14: 2100-2112.
- Glee, P. M., Cutler, J. E., Benson, E. E., Bargatze, R. F. Hazen, K. C.** 2001. Inhibition of hydrophobic protein-mediated *Candida albicans* attachment to endothelial cells during physiologic shear flow. *Infect. Immun.* 69: 2815-2820.
- Glee, P. M., Sundstrom, P., Hazen, K. C.** 1995. Expression of surface hydrophobic proteins by *Candida albicans in vivo*. *Infect. Immun.* 63: 1373-1379.
- Goldman, R. C., Frost, D. J., Capobianco, J. O., Kadam, S., Rasmussen, R. R., Abad-Zapatero, C.** 1995. Antifungal drug targets: *Candida* secreted aspartyl protease and fungal wall beta-glucan synthesis. *Infect. Agents Dis.* 4: 228-247.
- Gustafson, K. S., Vercellotti, G. M., Bendel, C. M., Hostetter, M. K.** 1991. Molecular mimicry in *Candida albicans*. *J. Clin. Investig.* 87: 1896-1902.
- Hahn, S., Hoar, E. T., Guarente, L.** 1985. Each of three "TATA elements" specifies a subset of the transcription initiation sites at the *CYC-1* promoter of *Saccharomyces cerevisiae*. *Proc. Natl. Acad. Sci. USA* 82: 8562-8566.
- Hawser, S. P. & Douglas, L. J.** 1994. Biofilm formation by *Candida species* on the surface of catheter materials *in vitro*. *Infect. Immun.* 62: 915-921.
- Haynes, K.** 2001. Virulence in *Candida species*. *Trend. Microbiol.* 9: 591-

596.

Hazen, B. W. & Hazen, K. C. 1988. Dynamic expression of cell surface hydrophobicity during initial yeast cell growth and before germ tube formation of *Candida albicans*. *Infect. Immun.* 56: 2521-2525.

Hazen, K. C. 1989. Participation of yeast cell surface hydrophobicity in adherence of *Candida albicans* to human epithelial cells. *Infect. Immun.* 57: 1894-1900.

Hazen, K. C. 1995. New emerging yeast pathogens. *Clin. Microbiol. Rev.* 8: 462-478.

Hazen, K. C., Brawner, D. L., Riesselman, M. H., Cutler, J. E., Jutila, M. A. 1991. Differential adherence of hydrophobic and hydrophilic *Candida albicans* yeast cells to mouse tissues. *Infect. Immun.* 59: 907-912.

Hector, R. F., Domer, J. E., Carrow, E. W. 1982. Immune responses to *Candida albicans* in genetically distinct mice. *Infect. Immun.* 38:1020-1028.

Hector, R. F., Yee, E., Collins, M. S. 1990. Use of DBA/2N mice in models of systemic candidiasis and pulmonary and systemic aspergillosis. *Infect. Immun.* 58:1476-1478.

Hoegl, L., Ollert, M., Korting, H. C. 1996. The role of *Candida albicans* secreted aspartic proteinase in the development of candidoses. *J. Mol. Med.* 74: 135-142.

Hostetter, M. K. 1998. Linkage of adhesion, morphogenesis, and virulence in *Candida albicans*. *J. Lab. Clin. Med.* 312: 258-263.

Hoyer L. L., Clevenger, J., Hecht, J. E., Ehrhart, E. J., Poulet, F. M. 1999. Detection of Als proteins on the cell wall of *Candida albicans* in murine tissues. *Infect. Immun.* 67: 4251-4255.

Hube, B. & Naglik, J. R. 2001. *Candida albicans* proteinases: resolving the mystery of the gene family. *Microbiol.* 147: 1997-2005.

Hube, B. 1996. *Candida albicans* secreted aspartyl proteinases. *Curr. Top. Med. Mycol.* 7: 55-69.

Hube, B. 2000. Extracellular Proteinases of human pathogenic fungi. In Ernst, J. F., Schmidt, A. (eds.): *Dimorphism in human pathogenic and apathogenic yeasts*. *Contrib. Microbiol. Basel, Karger*, vol. 5, pp 126-137.

Hube, B., Monod, M., Schofield, D. A., Brown, A. J. P., Gow, N. A. R.

1994. Expression of seven members of the gene family encoding secretory aspartyl proteinases in *Candida albicans*. Mol. Microbiol. 14: 87-99.
- Hube, B., Sanglard, D., Monod, M., Brown, A. J. P., Gow, N. A. R.** 1997b. Extracellular proteolytic activity of *Candida* species. In Host-fungus interplay. Edited by H. Van den Bossche, D. A. Stevens, F. C. Odds. National foundation for infectious diseases.
- Hube, B., Sanglard, D., Odds, F. C., Hess, D., Monod, M., Schäfer, W., Brown, A. J. P., Gow, N. A. R.** 1997a. Gene disruption of each of the Secretory Aspartyl Proteinase Genes *SAP1*, *SAP2* and *SAP3* in *Candida albicans* attenuates virulence. Infect. Immun. 65: 3529-3538.
- Hube, B., Turver, C. J., Odds, F. C., Eiffert, H., Boulnois, G. J., Kochel, H., Ruchel, R.** 1991. Sequence of the *Candida albicans* gene encoding the secretory aspartate proteinase. J. Med. Vet. Mycol. 29: 129-132.
- Hull, C. M. & Johnson, A. D.** 1999. Identification of mating type-like locus in the asexual pathogenic yeast *Candida albicans*. Science 285: 1271-1275.
- Hull, C. M., Raisner, R. M., Johnson, A. D.** 2000. Evidence for mating of the "asexual" yeast *Candida albicans* in a mammalian host. Science 289: 307-309.
- Ibrahim, A. S., Filler, S. G., Sanglard, D., Edwards, J. E., Jr., Hube, B.** 1998. Secreted aspartyl proteinases and interaction of *Candida albicans* with human endothelial cells. Infect. Immun. 66: 3003-3005.
- Joshi, K. R., Solanki, A., Prakash, P.** 1993. Morphological identification of *Candida* species on glucose agar, rice extract agar and corn meal agar with and without Tween-80. Indian. J. Pathol. Microbiol. 36: 48-52.
- Kaminishi, H., Hamatake, H., Cho, T., Tamaki, T., Suenaga, N., Fujii, T., Hagihara, Y., Maeda, H.** 1994. Activation of blood clotting factors by microbial proteinases. FEMS Microbiol. Lett. 121: 327-332.
- Kanbe, T. & Cutler, J. E.** 1998. Minimum chemical requirements for adhesin activity of the acid-stable part of *Candida albicans* cell wall phosphomannoprotein complex. Infect. Immun. 66: 5812-5818.
- Kennedy, M. J., Rogers, A. L., Hanselman, L. R., Soll, D. R., Yancey, R. J.** 1988. Variation in adhesion and cell surface hydrophobicity in

Candida albicans white and opaque phenotypes. Mycopathologia 102: 149-156.

Koelsch, G., Tang, J., Loy, J. A., Monod, M., Jackson, K., Foundling, S. I., Lin, X. 2000. Enzymic characteristics of secreted aspartic proteases of *Candida albicans*. Biochem. Biophys. Acta. 1480: 117-131.

Köhler, J. R. & Fink, G. R. 1996. *Candida albicans* strains heterozygous and homozygous in mitogen-activated protein kinase signalling components have defects in hyphal development. Proc. Natl. Acad. Sci. USA 93: 13223-13228.

Kolotila, M. P. & Diamond, R. D. 1990. Effects of neutrophils and *in vitro* oxidants on survival and phenotypic switching of *Candida albicans* WO-I. Infect. Immun. 58: 1174-1179.

Komano, H. & Fuller, R. S. 1995. Shared functions *in vivo* of a glycosylphosphatidylinositol-linked aspartyl protease, Mkc7, and the proprotein processing protease Kex2 in yeast. Proc. Natl. Acad. Sci. USA 92: 10752-10756.

Komano, H., Rockwell, N., Wang, G. T., Krafft, G. A., Fuller, R. S. 1999. Purification and characterization of the yeast glycosylphosphatidylinositol-anchored, monobasic-specific aspartyl protease Yapsin2 (Mkc7p). J. Biol. Chem. 274: 24431-24437.

Kozak, M. 1982. Analysis of ribosome binding sites from the s1 message of reovirus. Initiation at the first and second AUG codons. J. Mol. Biol. 156: 807-820.

Kretschmar, M., Felk, A., Staib, P., Schaller, M., Hess, D., Callapina, M., Morschhäuser, J., Schäfer, W., Korting, H. C., Hof, H., Hube, B., Nichterlein, T. 2002. Individual acid aspartic proteinases (Saps) 1-6 of *Candida albicans* are not essential for invasion and colonization of the gastrointestinal tract in mice. Microb. Pathog. 32: 61-70.

Kretschmar, M., Hube, B., Bertsch, T., Sanglard, D., Merker, R., Schröder, M., Hof, H., Nichterlein, T. 1999. Germ tubes and proteinase activity contribute to virulence of *Candida albicans* in murine peritonitis. Infect. Immun. 67: 6637-6642.

Kurtz, M. B. & Marrinan, J. 1989. Isolations of *Hem3* mutants from *Candida*

- albicans* by sequential gene disruption. Mol. Gen. Genet. 217: 47-52
- Kurtz, M. B., Cortelyou, M. W., Kirsch, D. R.** 1986. Integrative transformation of *Candida albicans*, using a cloned *Candida ADE 2* gene. Mol. Cell. Biol. 6: 142-149.
- Kyte, J. & Doolittle, R. F.** 1982. A simple method for displaying the hydrophobic character of a protein. J. Mol. Biol. 157: 105.
- Lane, S., Birse, C., Zhou, S., Matson, R., Liu, H.** 2001. DNA array studies demonstrate convergent regulation of virulence factors by Chp1, Cph2, and Efg1 in *Candida albicans*. J. Biol. Chem. 276: 48988-48996.
- Leberer, E., Harcus, D., Broadbent, I. D., Clark, K. L., Dignard, D., Ziegelbauer, K., Schmidt, A., Gow, N. A. R., Brown, A. J. P., Thomas, D. Y.** 1996. Signal transduction through homologs of the Ste20p and Ste7p protein kinases can trigger hyphal formation in the pathogenic fungus *Candida albicans*. Proc. Natl. Acad. Sci. USA 93: 13217-13222.
- Lee, K. L., Buchley, H. R., Campbell, C. C.** 1975. An amino acid synthetic medium for the development of mycelial and yeast forms of *Candida albicans*. Sabouraudia 13: 148-153.
- Li, R.-K. & Cutler, J. E.** 1993. Chemical definition of an epitope/adhesin molecule on *Candida albicans*. J. Biol. Chem. 268: 18293-18299.
- Liu, H., Köhler, J. R., Fink, G. R.** 1994. Suppression of hyphal formation in *Candida albicans* by mutation of a STE12 homolog. Science 266: 1723-1725.
- Liu, H., Styles, C. A., Fink, G. R.** 1993. Elements of the yeast pheromone response pathway required for filamentous growth of diploids. Science 262: 1741-1744.
- Lo, H.-J., Köhler, J. R., Didomenico, B., Loebenberg, D., Cacciapuoti, A., Fink, G. R.** 1997. Nonfilamentous *C. albicans* mutants are avirulent. Cell 90: 939-949.
- Lott, T. J., Kuykendall, R. J., Reiss, E.** 1993. Nucleotide sequence analysis of the 5.8S rDNA and adjacent ITS2 region of *Candida albicans* and related species. Yeast 9: 1199-1206.
- MacDonald, F. & Odds, F. C.** 1980. Inducible proteinase of *Candida albicans* in diagnostic serology and in the pathogenesis of systemic

candidosis. J. Med. Microbiol. 13: 423-435.

MacDonald, F. & Odds, F. C. 1983. Virulence for mice of a proteinase-secreting strain of *Candida albicans* and proteinase-deficient mutant. J. Gen. Microbiol. 129: 431-438.

MacKay, V. L., Welch, S. K., Insley, M. Y., Manney, T. R., Holly, J., Saari, G. C., Parker, M. L. 1988. The *Saccharomyces cerevisiae* *BAR1* gene encodes an exported protein with homology to pepsin. Proc. Natl. Acad. Sci. U S A 85:55-59.

Madhani, H. D. & Fink, G. R. 1997. Combinatorial control required for the specificity of yeast MAPK signalling. Science 275: 1314-1317.

Magee, B. B. & Magee, P. T. 2000. Induction of mating in *Candida albicans* by construction of *MTLa* and *MTL α* strains. Science 289: 310-312.

Magee, B. B., Hube, B., Wright, R. J., Sullivan, P. J., Magee, P. T. 1993. The genes encoding the secreted aspartyl proteinases of *Candida albicans* constitute a family with at least three members. Infect. Immun. 61: 3240-3243.

Maneu, V., Cervera, A. M., Martinez, J. P., Gozalbo, D. 1996. Molecular cloning and characterization of a *Candida albicans* gene (*EFB1*) coding for the elongation factor EF-1 beta. FEMS Microbiol. Lett. 145: 157-162

Marcantonio, E. E. & Hynes, R. O. 1988. Antibodies to the conserved cytoplasmic domain of the integrin β_1 subunit react with proteins in vertebrates, invertebrates, and fungi. J. Cell. Biol. 106: 1765-1772.

Masuoka, J., Wu, G., Glee, P. M., Hazen, K. C. 1999. Inhibition of *Candida albicans* attachment to extracellular matrix by antibodies which recognize hydrophobic cell wall proteins. FEMS Microbiol. Immunol. 24: 421-429.

Miller, M. E & Nilsson, U. R. 1974. A major role of the fifth component of complement (C5) in the opsonization of yeast particles. Partial dichotomy of function and immunochemical measurement. Clin. Immunol. Immunopathol. 2: 246-255.

Miyasaki, S. H., White, T. C., Agabian, N. 1994. A fourth secreted aspartyl proteinase gene (*SAP4*) and a *CARE2* repetitive element are located upstream of the *SAP1* gene in *Candida albicans*. J. Bacteriol. 176: 1702-1710.

- Monod, M., Hube, B., Hess, D., Sanglard, D.** 1998. Differential regulation of *SAP8* and *SAP9*, which encode two new members of the secreted aspartic proteinase family in *Candida albicans*. *Microbiol.* 144: 2731-2737.
- Monod, M., Togni, G., Hube, B., Sanglard, D.** 1994. Multiplicity of genes encoding secreted aspartic proteinases in *Candida* species. *Mol. Microbiol.* 13: 357-368.
- Montazeri, M. & Hedrick, H. G.** 1984. Factors affecting spore formation in a *Candida albicans* strain. *Appl. Environ. Microbiol.* 47: 1341-1342.
- Morrison, C. J., Hurst, S. F., Bragg, S. L., Kuykendall, R. J., Diaz, H., Pohl, J., Reiss, E.** 1993. Heterogeneity of the purified extracellular aspartyl proteinase from *Candida albicans*: Characterization with monoclonal antibodies and N-terminal amino acid sequence analysis. *Infect. Immun.* 61: 2030-2036.
- Morrow, B., Anderson, J., Wilson, J., Soll, D. R.** 1989. Bidirectional stimulation of the white-opaque transition of *Candida albicans* by ultraviolet irradiation. *J. Gen. Microbiol.* 135:1201-1208.
- Morrow, B., Srikantha, T., Anderson, J., Soll, D. R.** 1993. Coordinate regulation of two opaque-phase-specific genes during white-opaque switching in *Candida albicans*. *Infect. Immun.* 61: 1823-1828.
- Morrow, B., Srikantha, T., Soll, D. R.** 1992. Transcription of the gene for a pepsinogen, *PEP1*, is regulated by white-opaque switching in *Candida albicans*. *Mol. Cell Biol.* 12: 2997-3005.
- Movahedi, S. & Heale, J. B.** 1990. Purification and characterisation of an aspartic proteinase secreted by *Botrytis cinerea* per. ex. pers. in culture and in infected carrots. *Physiol. Mol. Plant. Pathol.* 36: 289-302.
- Mühlschlegel, F. A. & Fonzi, W. A.** 1997. *PHR2* of *Candida albicans* encodes a functional homolog of the pH-regulated gene *PHR1* with an inverted pattern of expression. *Mol. Cell. Biol.* 17: 5960-5967.
- Murad, A. M. A., d'Enfert, C., Gaillardin, C., Tournu, H., Tekaia, F., Talibi, D., Marechal, D., Marchais, V., Cottin, J., Brown, A. J. P.** 2001a. Transcript profiling in *Candida albicans* reveals new cellular functions for the transcriptional repressors CaTup1, CaMig1 and CaNrg1. *Mol. Microbiol.* 42: 981-993.

- Murad, A. M. A., Lee, P. R., Broadbent, I. D., Barelle, C. J., Brown, A. J. P.** 2000. Clp10, an efficient and convenient integrating vector for *C. albicans*. *Yeast* 16: 325-327. GenBank Number AF 181970.
- Murad, A. M. A., Leng, P., Straffon, M., Wishart, J., Macaskill, S., MacCallum, D., Schnell, N., Talibi, D., Marechal, D., Tekai, F., d'Enfert, C., Gaillardin, C., Odds, F. C., Brown, A. J. P.** 2001b. *NRG1* represses yeast-hypha morphogenesis and hypha-specific gene expression in *Candida albicans*. *EMBO J.* 20: 4742-4752.
- Naglik, J. R., Newport, G., White, T. C., Fernandes-Naglik, L. L., Greenspan, J. S., Greenspan, D., Sweet, S. P., Challacombe, S. J., Agabian, N.** 1999. *In vivo* analysis of secreted aspartyl proteinase expression in human oral candidiasis. *Infect. Immun.* 67: 2482-2490.
- Navarro-Garcia, F., Sanchez, M., Nombela, C., Pla, J.** 2001. Virulence genes in the pathogenic yeast *Candida albicans*. *FEMS Microbiol. Rev.* 25: 245-268.
- Negi, M., Tsuboi, R., Matsui, T., Ogawa, H.** 1984. Isolation and characterization of proteinase of *Candida albicans*: substrate specificity. *J. Ingest. Dermatol.* 83: 32-36.
- Newport, G. & Agabian, N.** 1997. Kex2 influences *Candida albicans* proteinase secretion and hyphal formation. *J. Biol. Chem.* 272: 28954-28961.
- Nguyen, M. H., Peacock, J. E., Morris, A. J., Tanner, D. C., Nguyen, M. L., Snyderman, D. R.** 1996. The changing face of candidemia: emergence of non-*Candida albicans* species and antifungal resistance. *Am. J. Med.* 100: 617-623.
- Nicholas, K. B. & Nicholas, H. B. Jr.** 1997. GeneDoc: a tool for editing and annotating multiple sequence alignments. Distribution by autor www.cris.com/~ketchup.shtml.
- Odds, F. C., Gow, N. A., Brown, A. J.** 2001. Fungal virulence studies come of age. *Genome Biol.* 2:1009.
- Odds, F. C.** 1988. *Candida* and Candidosis. London: Bailliere Tindal.
- Odds, F. C.** 1994 (a). *Candida* species and virulence. *ASM News* 60: 313-318.
- Odds, F. C.** 1994 (b). Pathogenesis of *Candida* infections. *J. Am. Acad.*

Dermatol. 31: S2-S5.

Ollert, M. W., Söhnchen, R., Korting, H. C., Ollert, U., Bräutigam, S., Bräutigam, W. 1993. Mechanisms of adherence of *Candida albicans* to cultured human epidermal keratinocytes. Infect. Immun. 61: 4560-4568.

Olsen, V., Guruprasad, K., Cawley, N. X., Chen, H-C., Blundell, T. L., Loh, Y. P. 1998. Cleavage efficiency of the novel aspartic protease Yapsin 1 (Yap3p) enhanced for substrates with arginine residues flanking the P1 site: correlation with electronegative active-site pockets predicted by molecular modeling. Biochemistry 37: 2768-2777.

Oluwatosin, Y. E. & Kane, P. M. 1998. Mutations in the yeast *KEX2* gene cause a Vma-like phenotype: a possible role for the Kex2 endoprotease in vacuolar acidification. Mol. Cell. Biol. 18: 1534-1543.

Prins, T. W., Wagemarkers, C. A. M., van Kan, J. A. L. 1999. GenBank accession number AF121229.

Ray, T. L. & Payne, C. D. 1990. Comparative production and rapid purification of *Candida* acid proteinase from protein-supplemented cultures. Infect. Immun. 58: 508-514.

Ray, T. L. & Payne, C. D. 1991. Degradation of basement membrane matrix proteins laminin, fibronectin, and type IV collagen by *Candida* acid proteinase. J. Invest. Dermatol. 96: 606A.

Ray, T. L., Payne, C. D., Morrow, B. J. 1991. *Candida* acid proteinases: characterization and role in candidiasis. Adv. Exp. Med. Biol. 306: 173-183.

Remold, H., Fasold, H., Staib, F. 1968. Purification and characterisation of a proteolytic enzyme from *Candida albicans*. Biochem Biophys Acta 167: 399-406.

Ritz, K. & Crawford, J. W. 1999. Colony development in nutritionally heterogeneous environments. In The Fungal Colony, Edited by N. A. R. Gow, G. D. Robson, G. M. Gadd. Cambridge: Cambridge University Press, pp. 49-74.

Rockwell, N. C. & Fuller, R. S. 1998. Interplay between S₁ and S₄ subsites in Kex2 protease: Kex2 exhibits dual specificity for the P₄ side chain. Biochemistry 37: 3386-3391.

Rostand, K. S. & Esko, J. D. 1997. Microbial adherence to and invasion

through proteoglycans. *Infect. Immun.* 65: 1-8.

Rüchel, R. & Böning, B. 1983. Detection of *Candida* protease by enzyme immunoassay and interaction of the enzyme with α - 2 -macroglobulin. *J. Immunol. Methods.* 61: 107-116.

Rüchel, R. 1981. Properties of a purified proteinase from the yeast *Candida albicans*. *Biochem. Biophys. Acta.* 659: 99-113.

Rüchel, R. 1983. On the renin-like activity of *Candida* proteinase and activation of blood coagulation *in vitro*. *Zbl. Bakt. I. Abt. Orig. A.* 255: 368-379.

Rüchel, R. 1984. A variety of *Candida* proteinases and their possible targets of proteolytic attack in the host. *Zbl. Bakt. Hyg. A.* 257: 266-274.

Rüchel, R. 1986. Cleavage of immunoglobulins by pathogenic yeast of the genus *Candida*. *Microbiol. Sci.* 3: 316-319.

Rüchel, R., Tegeler, R., Trost, M. 1982. A comparison of secretory proteinase from different strains of *Candida albicans*. *J. Med. Vet. Mycol.* 20: 233-244.

Rüchel, R., Zimmermann, F., Böning-Stutzer, B., Helmchen, U. 1991. Candidiasis visualised by proteinase-directed immunofluorescence. *Virchows Arch. A. Pathol. Anat. Histopathol.* 419:199-202.

Samonis, G., Anaissie, E. J., Bodey, G. P. 1990. Effects of broad-spectrum antimicrobial agents on yeast colonization of the gastrointestinal tracts of mice. *Antimicrob. Agents Chemother.* 34: 2420-2422.

San-Blas, G., Travassos, L. R., Fries, B. C., Goldman, D. L., Casadevall, A., Carmona, A. K., Barros, T. F., Puccia, R., Hostetter, M. K., Shanks, S. G., Copping, V. M. S., Knox, Y., Gow, N. A. R. 2000. Fungal morphogenesis and virulence. *Med. Mycol.* 38 (Suppl. I): 79-86.

Sanglard, D., Hube, B., Monod, M., Odds, F. C., Gow, N. A. R. 1997. A triple deletion in the *SAP4*, *SAP5* and *SAP6* secretory aspartyl proteinase genes of *Candida albicans* causes attenuated virulence. *Infect. Immun.* 65: 3539-3546.

Schaller, M., Korting, H. C., Schäfer, W., Bastert, J., Chen, W. C., Hube, B. 1999. Secreted aspartic proteinase (Sap) activity contributes to

tissue damage in a model of human oral candidosis. *Mol. Microbiol.* 34: 169-180.

Schaller, M., Schäfer, W., Korting, H. C., Hube, B. 1998. Differential expression of secreted aspartyl proteinases in a model of human oral candidosis and patient samples from the oral cavity. *Mol. Microbiol.* 29: 605-615.

Schröppel, K., Sprößler, K., Whiteway, M., Thomas, D. Y., Röllinghoff, M., Csank, C. 2000. Repression of hyphal proteinase expression by the mitogen-activated protein (MAP) kinase phosphatase Cpp1p of *Candida albicans* is dependent of the MAP kinase Cek1p. *Infect. Immun.* 68: 7159-7161.

Schweizer, A., Rupp, S., Taylor, B. N., Röllinghoff, M., Schröppel, K. 2000. The TEA/ATTS transcription factor CaTec1p regulates hyphal development and virulence in *Candida albicans*. *Mol. Microbiol.* 38: 435-445.

Shepherd, M. G., Yin, C. Y., Ram, S. P., Sullivan, P. A. 1980. Germ tube induction in *Candida albicans*. *Can. J. Microbiol.* 26: 21-26.

Silva, T. M. J., Glee, P. M., Hazen, K. C. 1995. Influence of cell surface hydrophobicity on attachment of *Candida albicans* to extracellular matrix proteins. *J. Med. Vet. Mycol.* 33: 117-122.

Slutsky, B., Staebell, M., Anderson, J., Risen, L., Pfaller, M., Soll, D. R. 1987. „White-opaque-transition“: a second high-frequency switching system in *Candida albicans*. *J. Bacteriol.* 169: 189-197.

Smolenski, G., Sullivan, P. A., Cutfield, S. M., & Cutfield, J. F. 1997. Analysis of secreted aspartic proteinases from *Candida albicans*: purification and characterization of individual Sap1, Sap2 and Sap3 isoenzymes. *Microbiol.* 143: 349-356.

Soll, D. R. 1992. High frequency switching in *Candida albicans*. *Clin. Microbiol. Rev.* 5: 183-203.

Soll, D. R. 1997. Gene regulation during high-frequency switching in *Candida albicans*. *Microbiol.* 143: 279-288.

Soll, D. R. 2002. Phenotypic Switching. In *Candida and Candidiasis*. Edited by R. A. Calderone. ASM Press, Washington, D. C., pp. 123-142.

Soll, D. R., Anderson, J. M., Bergen, M. 1991b. The developmental biology

of the white-opaque transition in *Candida albicans*. In *The Molecular Biology of Candida albicans*. Berlin: Springer, pp. 20-45.

Soll, D. R., Galask, R., Schmid, J., Hanna, C., Mac, K., Morrow, B. 1991a. Genetic dissimilarity of commensal strains carried in different anatomical locations of the same healthy women. *J. Clin. Microbiol.* 29: 681-690.

Soll, D. R., Langtimm, C. J., McDowell, J., Hicks, J., Galask, R. 1987. High-frequency switching in *Candida* strains isolated from vaginitis patients. *J. Clin. Microbiol.* 25:1611-1622.

Sonneborn, A., Bockmühl, D., Ernst, J. F. 1999a. Chlamyospore formation in *Candida albicans* requires the Efg1p morphogenetic regulator. *Infect. Immun.* 67: 5514-5517.

Sonneborn, A., Tebarth, B., Ernst, J. F. 1999b. Control of white-opaque phenotypic switching in *Candida albicans* by the Efg1p morphogenetic regulator. *Infect. Immun.* 67: 4655-4660.

Srikantha, T., Chandrasekhar, A., Soll, D. R. 1995. Funktional analysis of the promoter of the phase-specific *WH11* gene of *Candida albicans*. *Mol. Cell. Biol.* 15: 1797-1805.

Srikantha, T., Tsai, L. K., Daniels, K., Soll, D. R. 2000. *EFG1* null mutants of *Candida albicans* switch, but cannot express the complete phenotype of white-phase budding cells. *J. Bacteriol.* 18: 1580-1591.

Staab, J. F. & Sundstrom, P. 1998. Genetic organization and sequence analysis of the hypha-specific cell wall protein gene *HWP1* of *Candida albicans*. *Yeast* 14: 681-686.

Staab, J. F., Bradway, S. D., Fidel, P., Sundstrom, P. 1999. Adhesive and mammalian transglutaminase substrate properties of *Candida albicans* *HWP1*. *Science* 283: 1535-1538.

Staib, F. 1965. Serum-proteins as nitrogen source for yeast-like fungi. *Sabouradia* 4: 187-193.

Staib, P., Kretschmar, M., Nichterlein, T., Hof, H., Morschhäuser, J. 2000. Differential activation of a *Candida albicans* virulence gene family during infection. *Proc. Natl. Acad. Sci. USA* 97: 6102-6107.

Stehr, F. C., Felk, A., Kretschmar, M., Schaller, M., Schäfer, W., Hube, B. 2000. Extrazelluläre hydrolytische Enzyme und ihre Relevanz bei *Candida*

albicans-Infektionen. *Mycoses* 43 (Suppl. 2): 17-21.

Steinmetz, H. T. 1999. Mykosen. Candidainfektionen in der Intensivmedizin. Ausgabe 1. PVV-Verlag.

Stoldt, V. R., Sonneborn, A., Leuker, C., Ernst, J. F. 1997. Efg1 an essential regulator of morphogenesis of the human pathogen *Candida albicans*, is a member of a conserved class of bHLH proteins regulating morphogenetic processes in fungi. *EMBO* 16: 1982-1991.

Sullivan, P. A. & Wright, R. J. 1992. Two genes for secreted aspartate proteinases in *Candida albicans*. Abstr. 8th Int. Symp. Yeasts, G-10, p. 203.

Suzuki, T., Kanbe, T., Kuroiwa, T., Tanaka, K. 1989. High frequency variation of colony morphology and chromosome reorganization in the pathogenic yeast *Candida albicans*. *J. Gen. Microbiol.* 135: 425-434.

Tiraboschi, I. N., Bennett, J. E., Kauffman, C. A., Rex, J. H., Girmenia, C., Sobel, J. D. 2000. Deep *Candida* infections in the neutropenic and non-neutropenic host: an ISHAM symposium. *Med. Mycol.* 38 (Suppl. I): 199-204.

Tsushima, H. & Mine, H. 1995. Cleavage of human big endothelin-1 by *Candida albicans* aspartic proteinase. *FEMS Immun. Med. Microbiol.* 11: 69-72.

Tsushima, H., Mine, H., Kawakami, Y., Hyodoh, F., Ueki, A. 1994. *Candida albicans* aspartic proteinase cleaves and inactivates human epidermal cysteine proteinase inhibitor, cystatin A. *Microbiol.* 140: 167-171.

Tuomanen, E. 1993. Subversion of leukocyte adhesion systems by respiratory pathogens. *ASM News* 59: 292-296.

Tzung, K-W., Williams, R. M., Scherer, S., Federspiel, N., Jones, T., Hansen, N., Bivolarevic, V., Huizar, L., Komp, C., Surzycki, R., Tamse, R., Davis, R. W., Agabian, N. 2001. Genomic evidence for complete sexual cycle in *Candida albicans*. *Proc. Natl. Acad. Sci. USA* 98: 3249-3253.

Udenfried, S. & Kodukula, K. 1995. How glycosylphosphatidylinositol-anchored membrane proteins are made. *Annu. Rev. Biochem.* 64: 563-591.

Von Heijne, G. 1986. A new method for predicting signal sequence cleavage sites. *Nucleic Acids Res.* 14: 4683-4690.

Vossen, J. H., Müller, W. H., Lipke, P. N., Klis, F. M. 1997. Restrictive

glycosylphosphatidylinositol anchor synthesis in *cwh6/gpi3* yeast cells causes aberrant biogenesis of cell wall proteins. *J. Bacteriol.* 179: 2202-2209.

White, T. C. & Agabian, N. 1995. *Candida albicans* secreted aspartyl proteinases: isoenzyme pattern is determined by cell type, and levels are determined by environmental factors. *J. Bacteriol.* 177: 5215-5221.

White, T. C., Miyasaki, S. H., Agabian, N. 1993. Three distinct secreted aspartyl proteinases in *Candida albicans*. *J. Bacteriol.* 175: 6126-6133.

Wright, R. J., Carne, A., Hieber, A. D., Lamont, I. L., Emerson, G. W., Sullivan, P. A. 1992. A second gene for a secreted aspartate proteinase in *Candida albicans*. *J. Bacteriol.* 174: 7848-7853.

Zaugg, C., Borg-von Zeppelin, M., Reichard, U., Sanglard, D., Monod, M. 2001. Secreted aspartic proteinase family of *Candida tropicalis*. *Infect. Immun.* 69: 405-412.